Activin Signaling Regulates Autophagy and Cardiac Aging through mTORC2

Kai Chang^{1§}, Ping Kang^{1§}, Ying Liu¹, Kerui Huang¹, Erika Taylor², Antonia P. Sagona³, Ioannis P. Nezis³, Rolf Bodmer², Karen Ocorr², Hua Bai^{1*}

¹ Department of Genetics, Development, and Cell Biology, Iowa State University, Ames, IA 50011, USA

² Development, Aging, and Regeneration Program, Sanford-Burnham-Prebys Medical Discovery Institute, La Jolla, California 92037, USA

³ School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

***Corresponding Author:**

Hua Bai

Phone: 515-294-9395

Email: hbai@iastate.edu

§ These authors contribute equally to this work.

Running title:

Activin promotes cardiac aging through mTORC2

Abstract

Age-dependent loss of cardiac tissue homeostasis largely impacts heart performance and contributes significantly to cardiovascular diseases later in life. Cellular quality control machinery, such as autophagy/lysosome system, plays a crucial role in maintaining cardiac health and preventing age-induced cardiomyopathy and heart failure. However, how aging alters the autophagy/lysosome system to impact cardiac function remains largely unknown. Here using *Drosophila* heart as a model system, we show that activin signaling, a member of TGF-beta superfamily, negatively regulates cardiac autophagy and cardiac health during aging. We found that cardiac-specific knockdown of *Daw*, an activin-like protein in *Drosophila*, increased cardiac autophagy and prevented age-related cardiac dysfunction, including arrhythmia and bradycardia (slow heart rate). Inhibition of autophagy blocked *Daw* knockdown-mediated cardioprotection. Consistently, cardiac-specific expression of constitutively activated activin type I receptor *Babo* disrupted cardiac function at young ages. Intriguingly, the key autophagy regulator, mechanistic target of rapamycin complex 1 (mTORC1), was not involved in activin-mediated autophagy. Instead, activin signaling genetically interacted with Rictor, the key subunit of mTORC2, to regulate autophagy and cardiac aging. Knockdown of *Daw* increased the mRNA expression of Rictor and the phosphorylation of AKT in fly hearts. Finally, cardiac-specific silencing of *Daw* not only improved cardiac health, but also prolonged lifespan. Thus, our findings highlight an emerging role of activin signaling and mTORC2 in the regulation of autophagy and cardiac aging.

Key words: Activin ligand, Dawdle, Atg8a, TSC1, Rictor, mTOR complex 2, Autophagic flux, Semi-automatic optical heartbeat analysis (SOHA)

Introduction

Aging is associated with an exponential increase of the incidence of cardiovascular diseases (CVD) ^{1, 2}. Age-related changes in cardiovascular structure and output have been linked to increased risk of coronary heart disease, sudden cardiac death and stroke in elderly population ³. During normal aging, the left ventricular wall of human hearts becomes thickened and the diastolic filling rate of left ventricle gradually decreases. On the other hand, the left ventricular systolic performance at rest remains no change with age ³. Several mechanisms underlying these age-associated changes in cardiovascular structure and function have been proposed, for example changes in growth factor signaling, decreased cellular quality control, altered calcium handling, elevated extracellular matrix deposition or fibrosis, increased mitochondria damage, and the production of reactive oxygen species (ROS) ^{2, 4}. Resolving the contributing mechanisms underlying age-dependent decline of cardiovascular function is critical for the development of therapeutic interventions for the treatment of cardiovascular diseases.

Cellular quality control systems, such as macroautophagy (hereinafter referred to as autophagy), are essential to the maintenance of tissue homeostasis during aging ^{5, 6}. Autophagy is a highly conserved process that is responsible for the degradation and recycling of damaged organelles, protein aggregates and other cytoplasmic substances. The autophagic process initiated with the isolated membrane, or phagophore that elongates to form a double-membrane structure, the autophagosome. Then the autophagosome fuses with a lysosome to form an autolysosome to degrade the cargo within the autolysosome ⁷. It is generally accepted that autophagy activity declines with age ⁸⁻¹⁰. Disruption of autophagy pathways often leads to loss of tissue homeostasis and tissue dysfunction. For example, knocking out autophagy gene Atg5 in the mouse heart accelerates cardiac aging, including an increase in left ventricular hypertrophy and decrease in fractional shortening ¹⁰. As a crucial autophagy regulator, mechanistic target of rapamycin (mTOR), especially mTOR complex 1 (mTORC1), plays an important role in the regulation of cardiac tissue homeostasis ². Rapamycin-mediated suppression of mTORC1 induces autophagy and protects mouse cardiomyocytes from oxidative stress ¹¹. In *Drosophila*, mTORC1 is known to promote electrical pacing-induced heart failure ^{12, 13}, and high-fat-diet-induced cardiomyopathy ¹⁴. The mTOR signaling consists of two complexes, mTORC1 and mTORC2. It is not known whether mTORC2 also contributes to the regulation of autophagy ¹⁵. Although inhibition of mTOR is one of the robust longevity interventions ^{16, 17}, it remains unclear whether mTOR pathway plays a

major role in age-dependent decline of autophagy. Notably, a recent study showed that mTORC1 activity does not increase with age in most tissues of C57BL/6J.Nia mice¹⁸. Thus, the mechanistic basis for age-dependent alterations in autophagy and the contribution of mTORC1 remains to be determined.

Autophagy can be activated by various stresses, such as dietary restriction, hypoxia, oxidative stress, and pathogen infection¹⁹. The activation of autophagy is tightly controlled by key cellular signaling pathways, in particular the nutrient signaling pathways like mTORC1 and AMP-activated protein kinase (AMPK)²⁰. These pathways are often involved in control of longevity. We and other groups recently found that activin signaling, a transforming growth factor beta (TGF-beta) subfamily, regulates muscle proteostasis and longevity in *Drosophila*^{21, 22}. As in mammals, TGF-beta family in *Drosophila* has two major branches, bone morphogenetic protein (BMP) and activin signaling pathways²³. In both pathways, signaling starts with ligand binding to a receptor complex composed of type I and type II receptor kinases, followed by phosphorylation of receptor-activated Smad (R-Smad). *Drosophila* activin signaling is known to regulate neuroblast proliferation, axon guidance, and metabolic homeostasis during development (reviewed in²⁴). The regulation of adult physiology by *Drosophila* activin is not well understood. We previously showed that knockdown of one of the three activin ligands, *Daw*, prolongs lifespan and increases autophagosome formation in flies²¹. However, the role of TGF-beta/activin signaling in cardiac aging remains largely unknown.

It has been shown that TGF-beta factors promote cardiac fibrosis during aging²⁵. Mice with heterozygous mutants for activin receptor-like kinases (ALK4) are protected from pressure overload-induced fibrosis and dysfunction^{26, 27}. Inhibition of ALK5 also reduces myocardial infarction-induced systolic dysfunction and left ventricular remodeling in rats²⁸. However, conflicting results are reported in other studies showing that activin A protects hearts from hypoxia/reoxygenation- and ischemia/reperfusion-induced cell death²⁹. In addition, it is unknown whether TGF-beta family proteins, in particular activin, regulate cardiac homeostasis and function through autophagy. Given that the cardiac development and aging are remarkably conserved between *Drosophila* and mammals, we investigated the role of activin signaling in the regulation of cardiac autophagy and cardiac aging using *Drosophila* heart model and cardiac movement detection techniques³⁰. Here we show that activin signaling acts on hearts to regulate autophagy and age-related cardiomyopathy in *Drosophila*. RNAi-mediated knockdown of the activin-like

protein *Daw* preserved cardiac function with age, while expression of activated activin type I receptor *Babo* induced cardiomyopathy early in life. Inhibition of autophagy blocked the cardioprotective effects by *Daw* knockdown. Intriguingly, activin signaling genetically interacted with mTORC2 subunit Rictor (rapamycin-insensitive companion of mTOR), but not mTORC1 to control autophagy and cardiac aging. Thus, our study uncovered a novel interaction between activin and mTORC2 in the regulation of tissue homeostasis, autophagy, and age-related cardiomyopathy.

Results

Heart-specific knockdown of activin-like ligand *Daw* slows cardiac aging

Our previous study demonstrated that the activin-like protein *Daw* regulated tissue homeostasis (especially in flight muscle) and longevity in *Drosophila*²¹. Interestingly, we found that activin signaling increased in aging fly hearts, as indicated by elevated phosphorylation of Smad2 in heart nuclei (Fig. 1a). This finding suggest that activin signaling may play a role in regulating cardiac aging. To investigate this possibility, we knocked down the expression of *Daw* specifically in the heart with a binary GAL4/UAS system where we crossed *UAS-Daw RNAi* lines into a cardiac-specific tissue driver (*Hand-gal4*). We analyzed cardiac performance in young and old flies using a high-speed video imaging system (semi-automatic optical heartbeat analysis, SOHA)³¹. M-Mode traces from the SOHA analysis indicated that knockdown of *Daw* preserved cardiac contractility at advanced ages (Fig. 1b). In wild-type flies, normal aging is associated with increased cardiac arrhythmia, diastolic intervals, heart period (bradycardia, slow heart rate) (Figs. 1c, 1d, 1e), and decreased cardiac output (Fig. 1f). Systolic intervals normally remain unchanged (Fig. 1g). Interestingly, cardiac-specific knockdown of *Daw* attenuated age-dependent increase in arrhythmia, diastolic intervals, and heart period (Figs. 1c, 1d, 1e). *Daw* knockdown also maintained relatively normal cardiac output at advanced ages (Fig. 1f). Similar results were observed from three independent *Daw* RNAi lines (the knockdown efficiency of *Daw* RNAi was verified by qRT-PCR in our previous studies²¹).

The *Hand-gal4* line drives gene expression in both cardiomyocytes and pericardial cells³². To test the tissue-specific role of *Daw* in cardiac aging, we first crossed *Daw* RNAi into a cardiomyocyte-specific driver *TinC-gal4*³³. Similar to the results from *Hand-gal4* (Fig. 1h), cardiomyocyte-specific knockdown of *Daw* prevented age-related increase in cardiac arrhythmia

(Fig. 1i). We also tested whether *Daw* expression in pericardial cells can regulate cardiac aging using a pericardial cell-specific driver *Dot-gal4*³⁴. Interestingly, knockdown of *Daw* in pericardial cells also attenuated age-induced arrhythmia (Fig. 1j), but not the increase in diastolic interval (Fig. 1k). Thus, *Daw* appears to promote cardiac aging, in particular age-induced arrhythmia, through both cardiomyocyte-specific regulation and non-autonomous signaling from pericardial cells.

Cardiomyocyte-specific knockdown of activin receptor *Babo* delays cardiac aging

In *Drosophila*, activin-like ligand *Daw* signals through the type I receptor *Babo*, the fly homolog of mammalian ALK4 (activin receptor type-1B)²³. To investigate whether *Daw* acts through its receptor *Babo* to modulate cardiac aging, we first asked if *Babo* plays a similar role in preventing age-dependent increase in cardiomyopathy. Consistent with the results from *Daw* knockdown, we found that cardiomyocyte-specific knockdown of *Babo* blocked the age-related increase in cardiac arrhythmia, diastolic intervals, and heart period (Figs. 2a, 2b, 2c). Two independent *Babo* RNAi lines were used and both gave similar results. Systolic intervals remained unchanged in both controls and *Babo* RNAi flies (Fig. 2d). Knockdown of *Babo* did not affect age-dependent changes in cardiac output (Data not shown).

In a reciprocal experiment, we used a GeneSwitch heart driver (*Hand-GS-gal4*) to express a constitutive active form of *Babo* (*Babo-Act*, or *Babo.Q302D*) specifically in adult hearts. We found that cardiac expression of activated *Babo* significantly increased cardiac arrhythmia, diastolic intervals, and heart period already at young ages (Figs. 2e, 2f, 2g). Taken together, these results suggest that *Babo* plays an important role in age-related cardiomyopathy, especially cardiac arrhythmia, diastolic function and heart rate.

***Daw* negatively regulates autophagy in fly hearts**

Autophagy is one of the key mechanisms in maintaining tissue hemostasis, and its activity normally declines with age^{6, 9, 10}. Our previous studies showed that activin signaling regulates autophagosome formation and *Atg8a* transcription²¹. It is likely that *Daw* regulates cardiac aging by modulating autophagy. To first verify the role of *Daw* in autophagy-lysosome activity, we used LysoTracker staining to monitor the lysosomal acidification in adult fat body, a well-established autophagy marker in flies³⁵. Consistent with our previous finding, heterozygous

Daw^[11] loss-of-function mutants showed increased LysoTracker staining, suggesting an elevated basal autophagy/lysosome activities (Fig. 3a). Using a lipidation assay, we furthermore showed that mutation of *Daw* increased the levels of lipidated Atg8a, the fly homolog of mammalian LC3/GABARAP family proteins (Fig. 3b). Finally, mosaic analysis revealed that somatic cell clones in larval fat body expressing RNAi against either *Daw* or *Babo* (marked by GFP) showed increased numbers of autophagosome, indicated by mCherry-Atg8a puncta (Fig. 3c). The data suggest that activin signaling negatively regulates autophagy and lysosome activities.

Next, we examined whether *Daw* plays a similar role in modulating autophagy in fly hearts. Using a commercial antibody for *Drosophila* Atg8a protein (Fig. S1 shows the antibody validation), we found that the number of autophagosome in the heart was higher in *Daw*^[11] mutants compared to wild-type (Fig. S2a). Because autophagy is a dynamic process, the increased autophagosome number could result from increased autophagy activities, or blockage of lysosomal degradation³⁶. We next examined the role of *Daw* in the regulation of autophagic flux (the turnover of autophagosome). We exposed semi-intact fly hearts to a V-ATPase inhibitor bafilomycin A1 (Baf A1), which blocks both lysosomal acidification and autophagosome-lysosome fusion³⁷. We found that Baf A1 treatment resulted in a higher increase in the number of Atg8a-positive autophagosome in *Daw*^[11] hearts compared to wild-type hearts (Figs. S2a-S2c). Thus, the results suggest that *Daw* negatively regulate autophagic flux in fly hearts.

Aging tissues show reduced autophagy and accumulated cellular damages. We next examined whether inhibition of activin/*Daw* signaling in fly hearts can rescue age-dependent decline of autophagy. Using a mCherry-Atg8a reporter line, we observed a dramatic accumulation of Atg8a-positive autophagosome in the heart (Fig. 3d, 3e, 3h). The cardiomyocyte-specific driver *TinC-gal4* was used. The accumulation of autophagosome is likely due to the blockage of autophagosome turnover, because increases numbers of Atg8a-positive puncta were only seen in Baf A1-treated young hearts, but not the old hearts (Figs. 3e, 3h, 3j). Consistently, the adaptor protein p62/Ref(2)P, a marker for defective aggrephagy³⁸, was accumulated in aging hearts and Atg8a mutants (Fig. S3). Intriguingly, flies with heart-specific *Daw* knockdown maintained high levels of cardiac autophagosome turnover (or active autophagic flux) at both young and old ages, as indicated by the induction of autophagosome number upon Baf A1 treatment (Figs. 3f, 3h, 3j). Notably, constitutively activated *Babo* repressed the autophagic flux at both young and old ages (Figs. 3g, 3h). Activation of *Babo* also

blocked the accumulation of autophagosome in old fly hearts, which suggests that besides the regulation of autophagosome turnover, activin signaling also play an important role in autophagosome initiation and formation (Figs. 3g, 3h). We were able to verify these phenotypes using another heart-specific driver *Hand-Gal4* (Fig. 3i). Taken together, our findings suggest that inhibition of activin/Daw signaling promotes autophagosome turnover and maintains healthy autophagic activity in old hearts.

Inhibition of autophagy, but not activation of mTORC1, blocks *Daw* knockdown-mediated cardioprotection

Since we observed an induction of autophagy in *Daw* knockdown flies, we next asked whether autophagy activity is required for *Daw*-regulated cardiac aging. First, we fed control and *Daw* knockdown flies with lysosomal inhibitor chloroquine (CQ). As expected, CQ treatment increased cardiac arrhythmia in *Daw* knockdown flies at old age (Fig. 4a). To directly test whether autophagy plays any role in *Daw*-regulated cardiac aging, we generated double knockdown flies by combining *UAS-Daw RNAi* and *UAS-Atg1 RNAi* fly lines. Atg1 is a serine/threonine-protein kinase that is essential for autophagy initiation. Cardiac-specific knockdown of *Atg1* increased both cardiac arrhythmia and diastolic intervals at young ages (Fig. 4b), similar to the fly heart with activated *Babo* (Figs. 2e, 2f). Consistent with CQ treatment, cardiac-specific knockdown of *Atg1* attenuated cardioprotective effects of *Daw* RNAi, since simultaneously knockdown of *Atg1* and *Daw* in the heart led to age-dependent increase in arrhythmia similar to control flies (Fig. 4c). Thus, these results suggest that autophagy and Atg1 are required for *Daw*-regulated cardiac aging.

mTORC1 is a key negative regulator of autophagy³⁵. We then tested whether *Daw* could negatively regulate autophagy and cardiac aging through activation of mTORC1 signaling. To test this possibility, we first generated double knockdown flies by combining *UAS-Daw RNAi* and *UAS-Tsc1 RNAi*. Tsc1 (Tuberous sclerosis protein 1) is a negative regulator of mTORC1 (Fig. S4a). Surprisingly, activation of mTORC1 via *Tsc1* knockdown had no effects on cardiac arrhythmia (Fig. 4d), but reduced diastolic intervals at young age (Fig. 4e), which is very different from the cardiomyopathy resulted from cardiac-specific expression of activated activin receptor *Babo* (Figs. 2e, 2f). Furthermore, knockdown of *Tsc1* did not rescue the cardioprotective effects of *Daw* knockdown during aging (Fig. 4f). Simultaneously knockdown

of *Tsc1* and *Daw* did not show age-dependent increase in arrhythmia. Similarly, loss-of-function mutant *Tsc1*^[12] did not rescue the elevated LysoTracker staining in *Daw*^[11] (Fig. 4g). These results suggest that activation of mTORC1 is not required for Daw-regulated autophagy and cardiac aging.

It is known that inhibition of mTORC1 prolongs lifespan^{16,39} and attenuates stress-induced cardiomyopathies^{12,13,40,41}. However, the effects of mTORC1 induction on cardiac aging are not well understood. Here, we observed unique changes of cardiac function in response to activation of mTORC1 (via *Tsc1* knockdown), including decreased heart period (Tachycardia, fast heart rate), and reduced diastolic intervals (Figs. 4d, 4e). These cardiac functional changes are not the same as age-induced cardiomyopathies, rather they resemble high-fat-diet-induced cardiac complications where mTORC1 is known to play an important role¹⁴. Intriguingly, flies with cardiac-specific knockdown of *Tsc1* showed no age-dependent increases in heart period, diastolic intervals, and arrhythmia (Figs. S4b, S4c, S4d). These unique alterations in cardiac function were also observed in other mTORC1 activation mutants, such as 1) Knockdown of *Reptor* (Repressed by TOR), a recently identified bZIP transcription factor that is negatively regulated by mTORC1⁴² (Figs. S4e, S4f, S4g), and 2) Over-expression of *Rheb* (Ras Homolog Enriched in Brain), a positive regulator of mTORC1 (Figs. S4h, S4i, S4j).

Daw genetically interacts with mTORC2/Rictor to control autophagy and cardiac aging

Although Daw did not interact mTORC1 to regulate autophagy and cardiac aging, cardiac-specific knockdown of *Daw* induced the phosphorylation of AKT, the upstream regulator of mTORC1 (Figs. 5a, 5b). On the other hand, activated receptor Babo reduced cardiac AKT phosphorylation (Figs. 5c, 5d). It is known that AKT can be activated and phosphorylated by two kinases at distinct positions, PDK1 (phosphoinositide-dependent kinase 1) at Thr308 and Rictor (major component of mTORC2) at Ser473 (Ser505 in flies)⁴³. Cardiac-specific over-expression Rictor can also induce AKT phosphorylation (Figs. 5e, 5f). Interestingly, knockdown of *Daw* up-regulated the mRNA expression of *Rictor* in fly hearts (Figs. 5g, 5h). These results suggest that Daw is a negative regulator of mTORC2/Rictor.

Unlike mTORC1, the role of mTORC2 in the regulation of autophagy is unknown. Intriguingly, we found that loss-of-function mutation in *Rictor* repressed the high lysosomal activities in *Daw*^[11] mutants (Figs. 5i, 5j, 5k, 5l). Similarly, mutations in *Sin1*, another core

component of mTORC2, blocked the elevated lysosomal activity in *Daw* mutants. Furthermore, mosaic analysis revealed that over-expression of Rictor induced lysosome activity (Fig. 5m). Interestingly, unlike *Tsc1*^[12] mutants, *Rictor*^[42] mutants did not repress starvation-induced lysosome activity (Fig. S5c). Similarly, activated Babo also did not block starvation-induced lysosome activity (Fig. S5d). These data suggest that activin signaling and mTORC2/Rictor regulate autophagy differently from mTORC1, and Rictor acts downstream of activin signaling in the regulation of autophagy.

Given that Rictor positively regulates autophagy, we predict that Rictor is a cardiac protective factor. Indeed, cardiac-specific over-expression of *Rictor* blocked age-related increases in diastolic intervals and arrhythmia (Figs. 6a, 6b), while knockdown of *Rictor* induced arrhythmia, especially at old ages (Fig. 6c). Interestingly, over-expression of *Rictor* attenuated the elevated diastolic intervals and arrhythmia in fly hearts with constitutively activated *Babo* (Figs. 6d, 6e), suggesting that fly activin signaling regulates cardiac aging through mTORC2/Rictor (Fig. 6f).

Cardiac-specific reduction of activin-like protein *Daw* prolongs lifespan

Tissue-specific hormonal signaling in systemic aging control have been previously reported^{8, 44, 45}. However, whether longevity can be achieved by maintaining healthy hearts remains unclear. Here we tested the effects of cardiac-specific knockdown of activin signaling in longevity control. Knockdown of *Daw* in the heart using two cardiac tissue drivers (*Hand-gal4* and *TinC-gal4*) significantly prolongs lifespan (range from 14% to 32% extension of median lifespan) (Figs. 7a-7e). Compared to *Hand-gal4*, cardiomyocyte-specific driver *TinC-gal4* produced stronger lifespan extension. Two independent lifespan analyses were performed and both showed similar lifespan phenotypes (Fig. 7e). In the first lifespan trial, multiple control lines (dashed lines in Fig. 7a) and two *Daw* RNAi lines (solid lines in Fig. 7a) were included. Because *Daw* is a hormonal factor, it is unclear whether *Daw* regulates lifespan through cell-autonomous or non-autonomous mechanisms. However, our previous study showed that *Daw* regulates longevity in a tissue-specific manner²¹. Flight muscle-specific knockdown of *Daw* prolongs lifespan, while fat body-specific knockdown of *Daw* shortens lifespan. Therefore, it is possible that cardiac-specific *Daw* signaling regulates longevity through maintenance of cardiac health and production of unknown systemic factors from the fly heart.

Discussion

TGF-beta signaling plays vital roles in a wide range of human diseases, including cancer and cardiovascular diseases⁴⁶. Originally identified as a reproductive hormone, activin, a TGF-beta subfamily member, has become an emerging target for the treatment of many human diseases⁴⁷. In the present study, we investigated the role of activin in cardiac aging using a *Drosophila* heart model. Similar to our previous study in *Drosophila* flight muscle²¹, reduction of activin signaling prevented age-related decline of cardiac function. We think that this is partly due to the activation of autophagy. *Daw* mutant flies exhibited high autophagic flux and elevated lysosome activities. Genetic analysis further showed that autophagy gene Atg1 is required for *Daw*-regulated cardiac aging. Importantly, we found that activation of mTORC1 did not rescue the autophagy and cardiac aging phenotypes in *Daw* knockdown flies. Instead, activin/*Daw* signaling interacts with mTORC2 subunit Rictor to regulate autophagy and cardiac function. Thus, our studies uncover an emerging role of activin signaling and mTORC2 in the regulation of autophagy and cardiac tissue homeostasis.

Although previous studies reveal an interesting link between TGF-beta family proteins and animal aging^{21, 48-50}, the role of activin in cardiac aging is not well understood. It's known that activin A serum levels significantly increase with age⁵¹. The mRNA expression of *activin A*, as well as serum activin A levels are positively correlated with heart failure^{52, 53}, and hypertension in the elderly⁵⁴. These findings suggest that activation of activin signaling at old ages might be detrimental to the heart. Studies showed that activin A promotes cardiac fibrosis and myocardial damage after ischemia reperfusion⁵⁵. Consistently, mice with heterozygous mutations of ALK4 are protected from pressure overload-induced fibrosis and dysfunction^{26, 27}. Inhibition of ALK5 also reduced myocardial infarction-induced systolic dysfunction and left ventricular remodeling in rats²⁸. However, other studies suggest that activin A protects hearts from hypoxia/reoxygenation- and ischemia/reperfusion-induced cell death²⁹, even though it can promote cardiac apoptosis at higher concentrations⁵⁶. The findings from the present study support the negative regulatory role of activin signaling in cardiac function, in particular age-related cardiac dysfunction (e.g., arrhythmia). This is consistent with the pro-aging role of TGF-beta and activin^{21, 50}.

Activin signaling has been implicated in the regulation of a wide range of cellular processes, including cell death and proliferation, inflammation, fibrosis, and metabolic homeostasis^{47, 57-59}. The role of activin signaling in the control of autophagy, a key tissue maintenance process, was only identified recently^{21, 60-63}. In *C. elegans*, reduction of activin-like protein Daf-7 suppressed beta-amyloid peptide-induced autophagosome accumulation^{62, 63}. Another recent study found that activin A blocked oxygen-glucose deprivation-induced autophagy through the inhibition of JNK and p38 MAPK pathways in neuronal PC12 cells⁶⁰. Our previous study showed that knockdown of *Drosophila* activin-like protein *Daw* in flight muscle induced autophagosome formation and autophagy gene expression²¹. However, how activin signaling regulates autophagy and how this regulation contributes to tissue homeostasis during aging remain largely unknown. Our findings in the present study suggest that the fly activin-like protein *Daw* promotes cardiac aging through inhibition of autophagy. It seems that activin signaling controls both autophagosome formation and turnover. Importantly, fly hearts with reduced *Daw* expression maintain high levels of autophagosome turnover (autophagic flux) throughout the life, whereas the autophagosome turnover is largely blocked in aged hearts in wild-type flies. The active autophagosome turnover (and degradation capacity) at old ages is likely responsible for the healthy hearts seen in *Daw* knockdown flies.

mTORC1 is a well-known autophagy regulator and it inhibits autophagosome formation by increasing the phosphorylation of the serine/threonine protein kinase Atg1 (ULK1/ULK2 in mammals), the initiator of the autophagy machinery⁷. Intriguingly, activation of mTORC1 did not block the induction of autophagosome formation in *Daw* knockdown flies, suggesting that there might be unknown mTORC1-independent mechanisms involved in *Daw*-regulated autophagy and cardiac aging. mTORC1-independent autophagy regulation has been observed in several recent studies⁶⁴⁻⁶⁶. A genome-wide screen study identified an important role of type III PI3 kinases in autophagy control under normal nutritional conditions⁶⁴. The screen also revealed that growth factor signaling pathways, such as MAPK-ERK and AKT/FOXO3, negatively regulate autophagy by inhibiting the type III PI3 kinase cascade. Interestingly, we noticed that knockdown of *Daw* led to elevated AKT phosphorylation and the expression of mTORC2 subunit *Rictor*, suggesting mTORC2/AKT signaling may be involved in *Daw*-regulated autophagy and cardiac aging. Indeed, we found that mutation of *Rictor* blocked the activation of autophagy in *Daw* knockdown flies. mTORC2 is one of the two mTOR complexes that is

insensitive to short-term rapamycin treatment, and is involved in glucose homeostasis and actin cytoskeleton reorganization^{67, 68}. The direct regulatory role of mTORC2/Rictor in cellular autophagy has not been previously established, although a few studies suggest that Rictor may be involved in autophagy activation under unique circumstances. For example, Rictor is required for resveratrol-induced autophagy in rat myocardium and H9c2 cardiac myoblast cells⁶⁹. In rat kidney NRK-52E cells, silencing Rictor blocked cisplatin induced autophagy⁷⁰. Our findings align with these studies suggesting that Rictor is a positive regulator of autophagy. However, conflicting findings are reported in other studies showing Rictor negatively regulates autophagosome formation in mouse skeletal muscle⁷¹, and LC3-II levels in human senescent endothelial cells⁷². Thus, more studies are needed to better understand the role of mTORC2/Rictor in the regulation of autophagy and its contribution to tissue homeostasis during aging.

The positive role of Rictor in autophagy control identified in the present study is consistent with the cardioprotection effects of Rictor. We found that cardiac-specific over-expression of *Rictor* slows cardiac aging, similar to *Daw* knockdown. Our findings are consistent with two recent studies showing that disruption of mammalian Rictor induced cardiac dysfunction^{73, 74}. Cardiac-specific deletion of Rictor in mice induced dilation and decreased fractional shortening by activating MST1 kinase and hippo pathway⁷⁴. Unlike mTORC1 mutants, Rictor null mutants (in both worms and mice) are short-lived^{75, 76}, suggesting the detrimental role of mTORC2/Rictor in regulating tissue aging. The role of *Drosophila* Rictor in longevity regulation is not yet known, although it has been shown that *Drosophila* mutants of Rictor and Sin1 decreased the tolerance to heat stress⁷⁷ and over-expression of *Rictor* rescued Pink1 knockdown-induced mitochondrial aggregation in flight muscle⁷⁸. The Rictor-Pink1 interaction suggests that Rictor could be a potential positive regulator for mitophagy, a key process for the mitochondria quality control during aging.

During normal aging, the heart undergoes complex phenotypic changes such as progressive myocardial remodeling, reduced myocardial contractile capacity, increased left ventricular wall thickness and chamber size, prolonged diastole as well as increased arrhythmia³. All of these biological changes can gradually alter cardiac functions and confer vulnerability of the heart to various cardiovascular stresses, thus increasing the chance of developing cardiovascular disease dramatically. The present study has uncovered an important role of TGF-

beta/activin signaling pathway in the regulation of autophagy and cardiac aging in *Drosophila*. We also found an intriguing interaction between activin and mTORC2 in the regulation of cardiac autophagy, as well as a new role of Rictor in cardiac aging in *Drosophila*. It remains to be determined how activin regulates Rictor in fly hearts. It is possible that activin signaling regulates Rictor through both transcriptional activation (e.g., through insulin/FOXO signaling⁷⁹) and post-translational modification.

Additionally, cardiac-specific knockdown of activin-like protein *Daw* extended fly lifespan, suggesting that longevity can be achieved by maintaining a healthy heart. Our findings also suggest that *Daw* itself or other unidentified hormonal factors secreted from cardiomyocytes play an important role in systemic aging control. Hence, it remains to be determined the mechanistic basis for activin-mediated control of cardiac aging and longevity, which will eventually help the development of therapeutic interventions targeting activin for the treatment of age-related cardiovascular diseases.

Materials and methods

Fly husbandry and stocks

Flies were maintained at 25 °C, 60% relative humidity and 12-hour light/dark (40% relative humidity was used in the experiments from Figures 1&2, which were performed in Bodmer laboratory). Adults were reared on agar-based diet with 0.8% cornmeal, 10% sugar, and 2.5% yeast (unless otherwise noted). Fly stocks used in the present study are: *UAS-Daw RNAi* (Bloomington *Drosophila* Stock Center (BDSC) #34974, BDSC #50911, and Vienna *Drosophila* Resource Center (VDRC) #105309), *UAS-Babo RNAi* (BDSC #25933, BDSC #40866), *UAS-Babo-Act* (also known as *Babo.Q302D*)⁸⁰, *UAS-Tsc1 RNAi* (BDSC #52931, #54034), *UAS-Reptor RNAi* (BDSC #25983), *UAS-Rheb* (BDSC #9688), *UAS-Atg1 RNAi* (BDSC #26731), *UAS-Rictor RNAi* (BDSC #36699), *Daw*^[11]⁸¹, *Tsc1*^[12]⁸², *Rictor*^[42]⁸³, *Sin1*^[e03756] (BDSC #18188), *UAS-Rictor*⁷⁷, *UAS-HA-Rictor*⁷⁷, *Hand-gal4* (also known as Hand4.2-gal4)³², and *TinC-gal4* (also known as *TinCΔ4-gal4*)³³, *Dot-gal4*³⁴, *Hand-GS-gal4*⁸⁴, *UAS-mCherry-Atg8a* (BDSC #37750), *Atg8a*^{Δ4}⁸⁵.

UAS-Daw RNAi (BDSC #34974), *UAS-Babo RNAi* (BDSC #25933), and *UAS-Myo RNAi* (BDSC #31114) were backcrossed into *yw*^R background for 5~7 generations, and *yw*^R flies were used as control or wild-type (WT) flies in most of the experiments. For other UAS-RNAi lines

that were not backcrossed to yw^R , following genotypes were used as control: $y^l sc^* v^l$; $P[VALIUM20-mCherry]attP2$ (BDSC #35785), $y^l v^l$; $P[CaryP]attP2$ (BDSC #36303), or $y^l v^l$; $P[CaryP]attP40$ (BDSC # 36304). Female flies were used in all experiments. RU486 (mifepristone, Sigma, St. Louis, MO, USA) were used to activate *Hand-GS-Gal4* at a final concentration of 200 μ M mixed in food.

Mosaic analysis

Two FLPout lines were used: 1). *hs-flp; endogenous P-3x mCherry-Atg8a, UAS-GFP/Cyo; Act>CD2>Gal4,UAS-Dcr2/TM6* (kindly provided by Gabor Juhász). 2). *yw, hs-flp, UAS-CD8::GFP; Act>y+>Gal4,UAS-GFP.nls;UAS-Dcr2/SM6::TM6B* (kindly provided by Junyuan Ji, originally generated by Bruce Edgar). To generate clones, freshly laid eggs (within 4-6 hr) were heated in a 37 °C water bath for 45 min. Early L3 larvae (84 hours after egg laying) were used in the experiments.

Fly heartbeat analysis

To measure cardiac function parameters, semi-intact *Drosophila* adult fly hearts were prepared according to previously described protocols³¹. High-speed 3000 frames movies were taken at a rate of 100 frames per second using a Hamamatsu ORCA-Flash4.0 digital CMOS camera (Hamamatsu Photonics, Hamamatsu, Japan) on an Olympus BX51WI microscope with a 10X water immersion lens (Olympus, Waltham, MA, USA) (Hamamatsu EM-CCD 9300 camera was used in the experiments from Figures 1&2, which were performed in Bodmer laboratory). The live images that contain heart tub within abdominal A3 segment were processed using HCI imaging software (Hamamatsu Photonics, Hamamatsu, Japan). M-modes and cardiac parameters were generated using SOHA, a MATLAB-based image application as described previously^{31,86}. The M-mode provides a snapshot of movement of heart wall over time. Four cardiac parameters were analyzed, including heart period, diastolic interval, systolic interval, and arrhythmia index. Diastolic interval (DI) is the duration for each heart relaxation phase (diastole). Heart period (HP) is the pause time between the two consecutive diastole (HP is the reciprocal of heart rate). Systolic interval (SI) was calculated as the HP minus the DI. Arrhythmia index (AI) is the standard deviation of all HP in each fly normalized to the median HP. Cardiac output was

calculated using following equation: $(\pi r(d)^2 - \pi r(s)^2) \times HR$. $r(d)$ is radius of the heart tube at diastolic phase, while $r(s)$ is radius of the heart tube at systolic phase.

Quantitative RT-PCR

RNA extraction and cDNA synthesis were performed using Cells-to-CT kit (Thermo Fisher Scientific, Waltham, MA, USA) from ~15 dissected adult hearts. QRT-PCR was performed with a Quantstudio 3 Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA, USA). mRNA abundance of each gene was normalized to the expression of ribosomal protein L32 (*RpL32* or *rp49*) by the method of comparative C_T . Primer sequences are listed in Supplementary Table S1.

Antibodies and immunostaining

Since commercial antibodies for *Drosophila* Atg8a show non-specific staining in immunostaining of fly tissues (data not shown), we used a GABARAP (E1J4E) antibody (1:300) (Cell Signaling Technology (CST) #13733, Danvers, MA, USA) to stain and detect endogenous *Drosophila* Atg8a. The epitope used to generate GABARAP antibody has identical amino acid sequences to *Drosophila* Atg8a protein (personal communication with Cell Signaling Technology, and see sequence alignment in Fig. S1a). This antibody has been previously used in several *Drosophila* autophagy studies⁸⁷⁻⁸⁹, and verified by Kim et al.⁹⁰. We further verified the specificity of the GABARAP antibody using an Atg8a loss-of-function mutant *Atg8a^{Δ4}* (Figs. S1b-S1d). Other antibodies used for immunostaining are list below: Ref(2)P (1:1000)³⁸, Phospho-Smad2 antibody (1:500) (CST #3108), Phospho-Akt (Ser473) (1:300) (CST # 4060), Alexa Fluor 488-conjugated Phalloidin (or Alexa Fluor 594) for F-actin staining (Thermo Fisher Scientific, Waltham, MA, USA). All fluorescence-conjugated secondary antibodies were from Jackson ImmunoResearch, West Grove, PA, USA).

For immunostaining, adult female flies were collected and dissected in PBS. Hearts were fixed in 4% paraformaldehyde for 15 min at room temperature (RT). After washing in PBS with 0.1% Triton X-100 (PBST), the fixed hearted were blocked in 5% normal goat serum (NGS) diluted in PBST for 1 hour at RT. Hearts were then washed with PBST and incubated overnight at 4 °C with primary antibodies diluted in 5% NGS. After washing with PBST, the samples were incubated for 2 hours at RT with appropriate fluorescence-conjugated secondary antibodies.

Hearts were mounted in ProLong Gold anti-fade reagent (Thermo Fisher Scientific, Waltham, MA, USA) before imaged using an epifluorescence-equipped BX51WI microscope (Olympus, Waltham, MA, USA).

Imaging analysis

For imaging analysis, fluorescence images were first processed using the deconvolution module in Olympus cellSens Dimensions software, and then the number of puncta in a selected perinuclear region ($\sim 707 \mu\text{m}^2$) was measured with the “Measure and Count” module in cellSens software (Olympus, Waltham, MA, USA).

LysoTracker and Magic Red staining

Acidic organelles (including autolysosome) were monitored by staining tissue with 100 nM LysoTracker Red DND-99 (Invitrogen, Grand Island, NY, USA) for 5 minutes at room temperature. Lysosomal cathepsin B activities were monitored using Magic Red Cathepsin-B Assay kit (#938, ImmunoChemistry Technologies, Bloomington, MN, USA) following manufacturer’s manual. Nuclei were stained with either DAPI or Hoechst 33342 (1 $\mu\text{g}/\text{ml}$) (ImmunoChemistry Technologies).

Western blotting for Atg8a lipidation

Antibodies for western blot included: β -actin antibody (1:2000) (CST #4967), GABARAP (E1J4E) antibody (1:2000) (CST #13733), and HRP-conjugated secondary antibodies (Jackson ImmunoResearch, West Grove, PA, USA). Flies were first transferred to centrifuge tube containing 1x laemmli buffer (10 μl buffer for each milligram of fly), and heated at 100 °C for 3 min. The samples were then homogenized and heated again at 100 °C for 3 min. After the centrifuge at 13000x rpm for 5 min, supernatants were collected and loaded onto Mini-PROTEAN precast gels (Bio-Rad, Hercules, CA, USA). Following incubation with primary and secondary antibodies, the blots were visualized with Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific, Waltham, MA, USA).

Bafilomycin A1 and Chloroquine Treatment

For Bafilomycin A1 treatment, semi-intact hearts were incubated with 100 nM of bafilomycin A1 (Alfa Aesar, Haverhill, MA, USA) in artificial hemolymph (buffer receipt in ³¹) for two hours at room temperature prior to the appropriate immunostaining. DMSO was used as a control.

For chloroquine treatments, 100 µl of 20 mM chloroquine diphosphate salt (Sigma-Aldrich, St Louis, MO, USA) was added onto the fly food. Flies were fed with chloroquine for at least 24 hours prior to the cardiac analysis or the western blots.

Demography and survival analysis

Newly enclosed female flies were allowed to mate for two days, then separated from males and assigned to replicate one-liter demography cages at a density of 100-125 flies per cage. Three independent cages were set-up per genotype. Food was changed every two days, at which time dead flies were removed from the cage and counted. Survival analysis was conducted with JMP statistical software (SAS Institute, Cary, NC, USA), and data from replicate cages were combined. Mortality distributions were compared by Log-rank test.

Statistical analysis

GraphPad Prism (GraphPad Software, La Jolla, CA) was used for statistical analysis. To compare the mean value of treatment groups versus that of control, either student t-test or one-way ANOVA was performed using Dunnett's test for multiple comparison. The effects of mutants during aging was analyzed by two-way ANOVA, including Tukey multiple comparisons test. In SOHA analysis, the outliers were identified using Robust regression and Outlier removal (ROUT) method (Q=1%) prior to the data analysis.

Acknowledgements

We thank Bloomington Drosophila Stock Center, Harvard Medical School, and Drosophila Genomics Resource Center for fly stocks and cDNA clones. We thank Michael O'Connor for activin reagents and fly stocks, DuoJia Pan, Stocker Hugo, Jongkyeong Chung, G bor Juh sz, Jun-yuan Ji, Bruce Edgar for fly stocks. Marc Tatar for help with Demography and survival analysis. This work was supported by NIH/NIA R00 AG048016 and AFAR Research Grants to HB.

Author Contributions Statement

Conceived and designed the experiments: KC RB KO HB. Performed the experiments: KC PK YL KH APS IPN HB. Analyzed the data: KC PK YL KH ET APS IPN HB. Wrote the paper: KC RB KO HB. All authors reviewed and approved the manuscript.

Competing Interests

The authors declare that no competing interest exists.

Figure Legends

Figure 1. Heart-specific knockdown of *Daw* slows cardiac aging.

a. Immunostaining of phospho-Smad2 in fly hearts at young (2 weeks) and old ages (6 weeks).

Arrows indicate cardiomyocyte nuclei and positive pSmad2 staining. Scale bar: 20 m.

b. Representative M-mode traces (8 seconds) showing age-dependent movement of heart wall in control and heart-specific *Daw* knockdown flies (*Hand-gal4*>*UAS-Daw*^{RNAi}).

c-g. Age-related changes in arrhythmia index, diastolic intervals, heart period, cardiac output, and systolic intervals in control (*Ctrl*) and cardiac-specific *Daw* knockdown flies (*Daw*^{RNAi}).

Flies were cultured at 40% relative humidity. *Hand-gal4* driver was used to knock down gene expression specifically in cardiac tissues (cardiomyocytes and pericardial cells). Results from three independent *UAS-Daw*^{RNAi} lines are shown (RNAi #1: BDSC #34974, RNAi #2: VDRC #105309, RNAi #3: BDSC #50911). Statistical significance is assessed by two-way ANOVA

followed by Tukey multiple comparisons test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ns = not significant). $N = 7 \sim 20$. The interaction between genotype and age is statistically significant for heart period ($P = 0.0041$) and diastolic interval ($P = 0.0243$).

h-j. Age-dependent changes in cardiac arrhythmia between control and *Daw* knockdown using various tissue drivers, *Hand-gal4* (cardiomyocytes and pericardial cells), *TinC-gal4* (cardiomyocytes), and *Dot-gal4* (pericardial cells). Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 25 \sim 31$.

k. Age-dependent changes in diastolic intervals between control and *Daw* knockdown in pericardial cells (*Dot-gal4*). Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 13 \sim 26$.

Figure 2. Cardiomyocyte-specific knockdown of activin receptor *Babo* delays cardiac aging.

a-d. Age-dependent changes in cardiac arrhythmia, diastolic intervals, heart period, and systolic intervals in control (*Ctrl*) and cardiomyocyte-specific *Babo* knockdown flies (*Babo*^{RNAi}). Flies were cultured at 40% relative humidity. *TinC-gal4* driver was used. Results from two independent *UAS-Babo*^{RNAi} lines are shown (RNAi #1: BDSC #25933, RNAi #2: BDSC #40866). $N = 15 \sim 30$.

e-g. Changes in cardiac arrhythmia, diastolic intervals and heart period in fly hearts expressing constitutively activated *Babo* (*Babo*^{Act}). GeneSwitch heart driver *Hand-GS-gal4* was used to induce adult-onset *Babo* activation. RU: RU486 (Mifepristone). Flies were cultured at 40% relative humidity. $N = 6 \sim 9$. One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant).

Figure 3. *Daw* negatively regulates autophagy in fly hearts.

a. Representative images of LysoTracker staining of young adult fat body between wild-type (WT) and heterozygous *Daw*^[11] mutants. Scale bar: 20 μm .

b. Western blot analysis on Atg8a lipidation between WT and *Daw*^[11] mutants. GABARAP antibodies were used to detect Atg8a. The lower band represents lipidated Atg8a. Flies were fed with chloroquine (CQ, 20 mM) for at least 24 hours prior to western blots.

c. Representative images of mosaic analysis on autophagosome staining in larval fat body. Fat body clones were generated by crossing *Daw*^{RNAi} and *Babo*^{RNAi} lines into FLPout line carrying mCherry-Atg8a reporter (*hs-flp; endogenous P-3x mCherry-Atg8a, UAS-GFP/Cyo*;

Act>CD2>Gal4,UAS-Dcr2). RNAi Clones are GFP-positive cells (dashed lines). Scale bar: 20 μ m.

d. Representative images of fly hearts expressing autophagosome reporter mCherry-Atg8a at both young and old ages. Arrows indicate cardiomyocyte nuclei and surrounding autophagosomes. Abdominal segments A2-A3 are shown. Heart tube is located between two yellow dashed lines. PC: Pericardial cell.

e-g. Representative images of mCherry-Atg8a reporter surrounding cardiomyocyte nuclei in control, *Daw^{RNAi}*, and *Babo^{Act}* flies. Semi-intact hearts were incubated with 100 nM of bafilomycin A1 (Baf A1) for two hours prior to immunostaining. Scale bar: 10 μ m.

h. Quantification of age-dependent changes in mCherry-Atg8a-positive puncta in perinuclear region in control, *Daw^{RNAi}*, and *Babo^{Act}* flies. *TinC-gal4* was used to drive the expression of Atg8a reporter and gene knockdown. One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=5~7.

i. Quantification of age-dependent changes in mCherry-Atg8a-positive puncta in perinuclear region in control and *Daw^{RNAi}*. *Hand-gal4* was used to drive the expression of Atg8a reporter and *Daw* knockdown. One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=5~7.

j. Quantification of Baf A1-induced mCherry-Atg8a-positive puncta between control and *Daw^{RNAi}*.

Figure 4. Inhibition of autophagy, but not activation of mTORC1, blocks *Daw* knockdown-mediated cardioprotection.

a. Cardiac arrhythmia of chloroquine-treated (20 mM, 24 hours) 6-week-old control and *Daw* knockdown flies (*Hand-gal4* used). One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=14~38.

b. Cardiac arrhythmia and diastolic intervals in heart-specific knockdown of *Atg1* at young ages (2-week-old) (*Hand-gal4* used). One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=16~19.

c. Age-dependent changes in cardiac arrhythmia in control, *Daw^{RNAi}*, and *Daw^{RNAi}; Atg1^{RNAi}* flies (*TinC-gal4* used). One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=16~35.

d-e. Cardiac arrhythmia and diastolic intervals of *Tsc1* knockdown flies (*TinC-gal4* used). Two independent *Tsc1* RNAi lines were used (RNAi-1: BDSC #52931, RNAi-2: BDSC #54034). One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N = 16 \sim 18$.

f. Age-dependent changes in cardiac arrhythmia in control, *Daw^{RNAi}*, and *Daw^{RNAi}; Tsc1^{RNAi}* flies (*TinC-gal4* used). One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N = 7 \sim 30$.

g. Representative images of LysoTracker staining in adult fat body of WT, *Daw^[11]*, *Tsc1^[12]* and double mutant *Daw^[11]; Tsc1^[12]*. Scale bar: 10 μm .

Figure 5. Daw genetically interacts with mTORC2/Rictor to control autophagy.

a. Representative images of phospho-AKT staining in cardiomyocytes of control and *Daw* knockdown flies (*Hand-gal4*). Scale bar: 10 μm .

b. Image quantification of Figure 5a. Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 5$.

c. Representative images of phospho-AKT staining in cardiomyocytes of control and *Daw* knockdown flies (*Hand-gal4*). Scale bar: 10 μm .

d. Image quantification of Figure 5c. Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 5$.

e. Representative images of phospho-AKT staining in cardiomyocytes of control and Rictor over-expression (*Hand-gal4*). Two independent Rictor over-expression used (#1: *UAS-Rictor*, #2: *UAS-HA-Rictor⁷⁷*). Scale bar: 10 μm .

f. Image quantification of Figure 5e. One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N = 5$.

g. QRT-PCR analysis of *Rictor* expression in *Daw* knockdown flies (*Hand-gal4*). Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 3$.

h. QRT-PCR analysis of *Daw* expression in *Daw* knockdown flies (*Hand-gal4*). Student t-test (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). $N = 3$.

i. Representative images of LysoTracker staining in adult fat body of WT, *Daw^[11]*, *Rictor^[42]* and double mutant *Rictor^[42]; Daw^[11]*. Scale bar: 20 μm .

j. Image quantification of Figure 5i. One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N = 5$.

k. Representative images of Magic Red staining in adult fat body of WT, *Daw^[11]*, *Rictor^[42]* and double mutant *Rictor^[42]; Daw^[11]*. Scale bar: 20 μm .

l. Image quantification of Figure 5k. One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N=5$.

m. Representative images of mosaic analysis of LysoTracker staining in larval fat body. Fat body clones were generated by crossing *Rictor* overexpression lines into a FLPout line (*yw*, *hs-flp*, *UAS-CD8::GFP*; *Act>y+>Gal4,UAS-GFP.nls;UAS-Dcr2*). Clones with *Rictor* overexpression are GFP-positive cells (dashed lines). Scale bar: 20 μ m.

Figure 6. Heart-specific overexpression of *Rictor* preserved cardiac function during aging.

a-b. Diastolic intervals and arrhythmia in flies with heart-specific overexpression of *Rictor* (*Hand-gal4*). One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N=19\sim 26$.

c. Cardiac arrhythmia in flies with heart-specific knockdown of *Rictor*. One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N=24$.

d-e. Diastolic intervals and arrhythmia in flies with heart-specific expression of *Babo^{Act}*, or both *Babo^{Act}* and *Rictor* (*Hand-gal4*). One-way ANOVA (** $p < 0.001$, * $p < 0.01$, * $p < 0.05$, ns = not significant). $N=11\sim 27$.

f. Proposed model for activin-mTORC2 interaction in the regulation of autophagy and cardiac aging.

Figure 7. Cardiac-specific knockdown of *Daw* prolongs lifespan.

a-b. Survival analysis of cardiomyocyte-specific (*TinC-gal4*) knockdown of *Daw*. Two independent trials and multiple controls (*yw x yw*, *yw x Daw RNAi*, *TinC x mCherry RNAi*) were performed. Log-Rank test, $p < 0.001$.

c-d. Survival analysis of heart-specific (*Hand-gal4*) knockdown of *Daw*. Two independent trials were performed. Log-Rank test, $p < 0.001$.

e. Lifespan table to show sample size and median lifespan of the survival analysis.

Supplemental file

Table S1. Primer list

Figure S1. Validation of GABARAP antibody.

a. Multiple sequence alignment between *Drosophila* Atg8a and human GABARAP using ClustalW (<https://www.genome.jp/tools-bin/clustalw>). The red box indicates the predicted epitope sequences used in producing GABARAP antibody.

b. Representative images of GABARAP immunostaining in fly hearts of *WT* and *Atg8a^{Δ4}* mutants. Heart tube is located between two yellow dashed lines. The panels on the right are the zoomed-in images. Scale bar: 20 μm.

c. Representative images of GABARAP immunostaining in adult fat body of *WT* and *Atg8a^{Δ4}* mutants. The panels on the right are the zoomed-in images. Scale bar: 20 μm.

d. Western blots testing GABARAP antibody using *Atg8a^{Δ4}* mutants.

Figure S2. Daw inhibits autophagy flux in the heart.

a. Representative images of GABARAP immunostaining in fly hearts of *WT* and *Daw^[11]* mutants treated with or without Baf A1. Heart tube is located between two yellow dashed lines. The panels on the right are the zoomed-in images. PC: Pericardial cell. Scale bar: 20 μm.

b. Quantification of Figure S2. One-way ANOVA (*** p<0.001, ** p<0.01, * p<0.05, ns = not significant). N=5.

c. Quantification of Baf A1-induced puncta between *WT* and *Daw^[11]* mutants.

Figure S3. Aging increases the accumulation of Ref(2)P in the heart.

a-b. Representative images of Ref(2)P in young and old hearts. Scale bar: 10 μm.

c. Representative images of Ref(2)P in the heart of *Atg8a^{Δ4}* mutants. Scale bar: 10 μm.

Figure S4. Activation of mTORC1 alters heart period, but not arrhythmia.

a. Schematic diagram for PI3K/AKT/mTOR signaling pathway.

b-d. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and cardiomyocyte-specific *Tsc1* knockdown flies (*TinC-gal4*). One-way ANOVA (*** p<0.001, ** p<0.01, * p<0.05, ns = not significant). N=14~18.

e-g. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and cardiomyocyte-specific *Reptor* knockdown flies (*TinC-gal4*). One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=13~28.

h-j. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and flies overexpressing *Rheb* (*TinC-gal4*). One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=21~31.

Figure S5. Autophagy regulation by mTORC2.

a. Representative images of LysoTracker staining in adult fat body of WT, *Daw*^[11], *Sin1*^[e03756] and double mutant *Sin1*^{[e03756]; *Daw*^[11]. Scale bar: 20 μ m.}

b. Image quantification of Figure S5a. One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=5.

c. Representative images of LysoTracker staining in fed and starved WT, *Tsc1* mutants, and *Rictor* mutants. Scale bar: 20 μ m.

d. Representative images of mosaic analysis on LysoTracker staining in larval fat body of *Babo*^{Act} flies upon starvation. Larval fat body clones were generated using a FLPout line (*yw*, *hs-flp*, *UAS-CD8::GFP*; *Act*>*y+*>*Gal4*, *UAS-GFP.nls*; *UAS-Dcr2*). Clones with *Babo*^{Ac} expression are GFP-positive cells (dashed lines). Scale bar: 20 μ m.

References

1. Dai, D.F., Chen, T., Johnson, S.C., Szeto, H. & Rabinovitch, P.S. Cardiac aging: from molecular mechanisms to significance in human health and disease. *Antioxidants & redox signaling* **16**, 1492-1526 (2012).
2. North, B.J. & Sinclair, D.A. The intersection between aging and cardiovascular disease. *Circulation research* **110**, 1097-1108 (2012).
3. Lakatta, E.G. & Levy, D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part II: the aging heart in health: links to heart disease. *Circulation* **107**, 346-354 (2003).
4. Dai, D.F. *et al.* Overexpression of catalase targeted to mitochondria attenuates murine cardiac aging. *Circulation* **119**, 2789-2797 (2009).
5. Quarles, E.K. *et al.* Quality control systems in cardiac aging. *Ageing research reviews* **23**, 101-115 (2015).
6. Rubinsztein, D.C., Marino, G. & Kroemer, G. Autophagy and aging. *Cell* **146**, 682-695 (2011).
7. Shibutani, S.T. & Yoshimori, T. A current perspective of autophagosome biogenesis. *Cell research* **24**, 58-68 (2014).
8. Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in *Drosophila* muscles regulates organism-wide proteostasis during aging. *Cell* **143**, 813-825 (2010).
9. Chang, J.T., Kumsta, C., Hellman, A.B., Adams, L.M. & Hansen, M. Spatiotemporal regulation of autophagy during *Caenorhabditis elegans* aging. *eLife* **6** (2017).
10. Taneike, M. *et al.* Inhibition of autophagy in the heart induces age-related cardiomyopathy. *Autophagy* **6**, 600-606 (2010).
11. Dutta, D., Xu, J., Kim, J.S., Dunn, W.A., Jr. & Leeuwenburgh, C. Upregulated autophagy protects cardiomyocytes from oxidative stress-induced toxicity. *Autophagy* **9**, 328-344 (2013).
12. Wessells, R. *et al.* d4eBP acts downstream of both dTOR and dFoxo to modulate cardiac functional aging in *Drosophila*. *Aging cell* **8**, 542-552 (2009).
13. Luong, N. *et al.* Activated FOXO-mediated insulin resistance is blocked by reduction of TOR activity. *Cell metabolism* **4**, 133-142 (2006).
14. Birse, R.T. *et al.* High-fat-diet-induced obesity and heart dysfunction are regulated by the TOR pathway in *Drosophila*. *Cell metabolism* **12**, 533-544 (2010).
15. Saxton, R.A. & Sabatini, D.M. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* **168**, 960-976 (2017).
16. Kaeberlein, M. *et al.* Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* **310**, 1193-1196 (2005).
17. Lee, J.H. *et al.* Sestrin as a feedback inhibitor of TOR that prevents age-related pathologies. *Science* **327**, 1223-1228 (2010).
18. Baar, E.L., Carbajal, K.A., Ong, I.M. & Lamming, D.W. Sex- and tissue-specific changes in mTOR signaling with age in C57BL/6J mice. *Aging cell* **15**, 155-166 (2016).
19. Kroemer, G., Marino, G. & Levine, B. Autophagy and the integrated stress response. *Molecular cell* **40**, 280-293 (2010).
20. He, C. & Klionsky, D.J. Regulation mechanisms and signaling pathways of autophagy. *Annual review of genetics* **43**, 67-93 (2009).

21. Bai, H., Kang, P., Hernandez, A.M. & Tatar, M. Activin signaling targeted by insulin/dFOXO regulates aging and muscle proteostasis in *Drosophila*. *PLoS genetics* **9**, e1003941 (2013).
22. Demontis, F., Patel, V.K., Swindell, W.R. & Perrimon, N. Intertissue control of the nucleolus via a myokine-dependent longevity pathway. *Cell reports* **7**, 1481-1494 (2014).
23. Upadhyay, A., Moss-Taylor, L., Kim, M.J., Ghosh, A.C. & O'Connor, M.B. TGF-beta Family Signaling in *Drosophila*. *Cold Spring Harbor perspectives in biology* (2017).
24. Upadhyay, A., Moss-Taylor, L., Kim, M.J., Ghosh, A.C. & O'Connor, M.B. TGF-beta Family Signaling in *Drosophila*. *Cold Spring Harbor perspectives in biology* **9** (2017).
25. Biernacka, A. & Frangogiannis, N.G. Aging and Cardiac Fibrosis. *Aging and disease* **2**, 158-173 (2011).
26. Li, C.Y. *et al.* Partial inhibition of activin receptor-like kinase 4 attenuates pressure overload-induced cardiac fibrosis and improves cardiac function. *Journal of hypertension* **34**, 1766-1777 (2016).
27. Chen, Y.H. *et al.* Haploinsufficiency of activin receptor-like kinase 4 alleviates myocardial infarction-induced cardiac fibrosis and preserves cardiac function. *Journal of molecular and cellular cardiology* **105**, 1-11 (2017).
28. Tan, S.M., Zhang, Y., Connelly, K.A., Gilbert, R.E. & Kelly, D.J. Targeted inhibition of activin receptor-like kinase 5 signaling attenuates cardiac dysfunction following myocardial infarction. *American journal of physiology. Heart and circulatory physiology* **298**, H1415-1425 (2010).
29. Oshima, Y. *et al.* Activin A and follistatin-like 3 determine the susceptibility of heart to ischemic injury. *Circulation* **120**, 1606-1615 (2009).
30. Ocorr, K., Vogler, G. & Bodmer, R. Methods to assess *Drosophila* heart development, function and aging. *Methods* **68**, 265-272 (2014).
31. Ocorr, K., Fink, M., Cammarato, A., Bernstein, S. & Bodmer, R. Semi-automated Optical Heartbeat Analysis of small hearts. *Journal of visualized experiments : JoVE* (2009).
32. Han, Z. & Olson, E.N. Hand is a direct target of Tinman and GATA factors during *Drosophila* cardiogenesis and hematopoiesis. *Development* **132**, 3525-3536 (2005).
33. Lo, P.C. & Frasch, M. A role for the COUP-TF-related gene seven-up in the diversification of cardioblast identities in the dorsal vessel of *Drosophila*. *Mechanisms of development* **104**, 49-60 (2001).
34. Zhang, F., Zhao, Y. & Han, Z. An in vivo functional analysis system for renal gene discovery in *Drosophila* pericardial nephrocytes. *Journal of the American Society of Nephrology : JASN* **24**, 191-197 (2013).
35. Scott, R.C., Schuldiner, O. & Neufeld, T.P. Role and regulation of starvation-induced autophagy in the *Drosophila* fat body. *Developmental cell* **7**, 167-178 (2004).
36. Klionsky, D.J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy* **12**, 1-222 (2016).
37. Mauvezin, C., Nagy, P., Juhasz, G. & Neufeld, T.P. Autophagosome-lysosome fusion is independent of V-ATPase-mediated acidification. *Nature communications* **6**, 7007 (2015).
38. Nezis, I.P. *et al.* Ref(2)P, the *Drosophila melanogaster* homologue of mammalian p62, is required for the formation of protein aggregates in adult brain. *The Journal of cell biology* **180**, 1065-1071 (2008).

39. Johnson, S.C., Rabinovitch, P.S. & Kaeberlein, M. mTOR is a key modulator of ageing and age-related disease. *Nature* **493**, 338-345 (2013).
40. Shioi, T. *et al.* Rapamycin attenuates load-induced cardiac hypertrophy in mice. *Circulation* **107**, 1664-1670 (2003).
41. Ding, Y. *et al.* Haploinsufficiency of target of rapamycin attenuates cardiomyopathies in adult zebrafish. *Circulation research* **109**, 658-669 (2011).
42. Tiebe, M. *et al.* REPTOR and REPTOR-BP Regulate Organismal Metabolism and Transcription Downstream of TORC1. *Developmental cell* **33**, 272-284 (2015).
43. Sarbassov, D.D., Guertin, D.A., Ali, S.M. & Sabatini, D.M. Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science* **307**, 1098-1101 (2005).
44. Hwangbo, D.S., Gershman, B., Tu, M.P., Palmer, M. & Tatar, M. Drosophila dFOXO controls lifespan and regulates insulin signalling in brain and fat body. *Nature* **429**, 562-566 (2004).
45. Zhang, G. *et al.* Hypothalamic programming of systemic ageing involving IKK-beta, NF-kappaB and GnRH. *Nature* **497**, 211-216 (2013).
46. Akhurst, R.J. & Hata, A. Targeting the TGFbeta signalling pathway in disease. *Nature reviews. Drug discovery* **11**, 790-811 (2012).
47. Tsuchida, K. *et al.* Activin signaling as an emerging target for therapeutic interventions. *Cell communication and signaling : CCS* **7**, 15 (2009).
48. Luo, S., Kleemann, G.A., Ashraf, J.M., Shaw, W.M. & Murphy, C.T. TGF-beta and insulin signaling regulate reproductive aging via oocyte and germline quality maintenance. *Cell* **143**, 299-312 (2010).
49. Narasimhan, S.D. *et al.* PDP-1 links the TGF-beta and IIS pathways to regulate longevity, development, and metabolism. *PLoS genetics* **7**, e1001377 (2011).
50. Shaw, W.M., Luo, S., Landis, J., Ashraf, J. & Murphy, C.T. The C. elegans TGF-beta Dauer pathway regulates longevity via insulin signaling. *Current biology : CB* **17**, 1635-1645 (2007).
51. Baccarelli, A. *et al.* Activin A serum levels and aging of the pituitary-gonadal axis: a cross-sectional study in middle-aged and elderly healthy subjects. *Experimental gerontology* **36**, 1403-1412 (2001).
52. Yndestad, A. *et al.* Elevated levels of activin A in heart failure: potential role in myocardial remodeling. *Circulation* **109**, 1379-1385 (2004).
53. Wei, Q. *et al.* The expression and role of activin A and follistatin in heart failure rats after myocardial infarction. *International journal of cardiology* **168**, 2994-2997 (2013).
54. Tsai, Y.L. *et al.* The Association Between Serum Activin A Levels and Hypertension in the Elderly: A Cross-Sectional Analysis From I-Lan Longitudinal Aging Study. *American journal of hypertension* (2017).
55. Chen, Y. *et al.* Regulation and actions of activin A and follistatin in myocardial ischaemia-reperfusion injury. *Cytokine* **69**, 255-262 (2014).
56. Liu, M., Mao, C., Li, J., Han, F. & Yang, P. Effects of the Activin A-Follistatin System on Myocardial Cell Apoptosis through the Endoplasmic Reticulum Stress Pathway in Heart Failure. *International journal of molecular sciences* **18** (2017).
57. Ghosh, A.C. & O'Connor, M.B. Systemic Activin signaling independently regulates sugar homeostasis, cellular metabolism, and pH balance in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 5729-5734 (2014).

58. Chng, W.B., Bou Sleiman, M.S., Schupfer, F. & Lemaitre, B. Transforming growth factor beta/activin signaling functions as a sugar-sensing feedback loop to regulate digestive enzyme expression. *Cell reports* **9**, 336-348 (2014).
59. Werner, S. & Alzheimer, C. Roles of activin in tissue repair, fibrosis, and inflammatory disease. *Cytokine & growth factor reviews* **17**, 157-171 (2006).
60. Xue, L.X. *et al.* Activin A/Smads signaling pathway negatively regulates Oxygen Glucose Deprivation-induced autophagy via suppression of JNK and p38 MAPK pathways in neuronal PC12 cells. *Biochemical and biophysical research communications* **480**, 355-361 (2016).
61. Guo, B. *et al.* Genome-wide screen identifies signaling pathways that regulate autophagy during *Caenorhabditis elegans* development. *EMBO reports* **15**, 705-713 (2014).
62. Florez-McClure, M.L. & Link, C.D. Erratum: "Florez-McClure ML, Hohsfield LA, Fonte G, Bealor MT, Link CD. Decreased insulin-receptor signaling promotes the autophagic degradation of beta-amyloid peptide in *C. elegans*". *Autophagy* 2007; 3:569-80. *Autophagy* **5**, 132-133 (2009).
63. Florez-McClure, M.L., Hohsfield, L.A., Fonte, G., Bealor, M.T. & Link, C.D. Decreased insulin-receptor signaling promotes the autophagic degradation of beta-amyloid peptide in *C. elegans*. *Autophagy* **3**, 569-580 (2007).
64. Lipinski, M.M. *et al.* A genome-wide siRNA screen reveals multiple mTORC1 independent signaling pathways regulating autophagy under normal nutritional conditions. *Developmental cell* **18**, 1041-1052 (2010).
65. Li, Y. *et al.* Protein kinase C controls lysosome biogenesis independently of mTORC1. *Nature cell biology* **18**, 1065-1077 (2016).
66. Henson, S.M. *et al.* p38 signaling inhibits mTORC1-independent autophagy in senescent human CD8(+) T cells. *The Journal of clinical investigation* **124**, 4004-4016 (2014).
67. Oh, W.J. & Jacinto, E. mTOR complex 2 signaling and functions. *Cell cycle* **10**, 2305-2316 (2011).
68. Gaubitz, C., Prouteau, M., Kusmider, B. & Loewith, R. TORC2 Structure and Function. *Trends in biochemical sciences* **41**, 532-545 (2016).
69. Gurusamy, N. *et al.* Cardioprotection by resveratrol: a novel mechanism via autophagy involving the mTORC2 pathway. *Cardiovascular research* **86**, 103-112 (2010).
70. Li, J. *et al.* Rictor/mTORC2 protects against cisplatin-induced tubular cell death and acute kidney injury. *Kidney international* **86**, 86-102 (2014).
71. Mammucari, C. *et al.* FoxO3 controls autophagy in skeletal muscle in vivo. *Cell metabolism* **6**, 458-471 (2007).
72. Xiong, Y. *et al.* ARG2 impairs endothelial autophagy through regulation of MTOR and PRKAA/AMPK signaling in advanced atherosclerosis. *Autophagy* **10**, 2223-2238 (2014).
73. Shende, P. *et al.* Cardiac mTOR complex 2 preserves ventricular function in pressure-overload hypertrophy. *Cardiovascular research* **109**, 103-114 (2016).
74. Sciarretta, S. *et al.* mTORC2 regulates cardiac response to stress by inhibiting MST1. *Cell reports* **11**, 125-136 (2015).
75. Soukas, A.A., Kane, E.A., Carr, C.E., Melo, J.A. & Ruvkun, G. Rictor/TORC2 regulates fat metabolism, feeding, growth, and life span in *Caenorhabditis elegans*. *Genes & development* **23**, 496-511 (2009).
76. Lamming, D.W. *et al.* Depletion of Rictor, an essential protein component of mTORC2, decreases male lifespan. *Aging cell* **13**, 911-917 (2014).

77. Jevtov, I. *et al.* TORC2 mediates the heat stress response in *Drosophila* by promoting the formation of stress granules. *Journal of cell science* **128**, 2497-2508 (2015).
78. Wu, Z. *et al.* Tricornered/NDR kinase signaling mediates PINK1-directed mitochondrial quality control and tissue maintenance. *Genes & development* **27**, 157-162 (2013).
79. Chen, C.C. *et al.* FoxOs inhibit mTORC1 and activate Akt by inducing the expression of Sestrin3 and Rictor. *Developmental cell* **18**, 592-604 (2010).
80. Brummel, T. *et al.* The *Drosophila* activin receptor baboon signals through dSmad2 and controls cell proliferation but not patterning during larval development. *Genes & development* **13**, 98-111 (1999).
81. Serpe, M. & O'Connor, M.B. The metalloprotease tolloid-related and its TGF-beta-like substrate Dawdle regulate *Drosophila* motoneuron axon guidance. *Development* **133**, 4969-4979 (2006).
82. Gao, X. & Pan, D. TSC1 and TSC2 tumor suppressors antagonize insulin signaling in cell growth. *Genes & development* **15**, 1383-1392 (2001).
83. Lee, G. & Chung, J. Discrete functions of rictor and raptor in cell growth regulation in *Drosophila*. *Biochemical and biophysical research communications* **357**, 1154-1159 (2007).
84. Monnier, V. *et al.* dJun and Vri/dNFIL3 are major regulators of cardiac aging in *Drosophila*. *PLoS genetics* **8**, e1003081 (2012).
85. Pircs, K. *et al.* Advantages and limitations of different p62-based assays for estimating autophagic activity in *Drosophila*. *PloS one* **7**, e44214 (2012).
86. Fink, M. *et al.* A new method for detection and quantification of heartbeat parameters in *Drosophila*, zebrafish, and embryonic mouse hearts. *BioTechniques* **46**, 101-113 (2009).
87. Ratliff, E.P., Barekat, A., Lipinski, M.M. & Finley, K.D. Brain trauma and autophagy: What flies and mice can teach us about conserved responses. *Autophagy* **12**, 2256-2257 (2016).
88. Ratliff, E.P. *et al.* Assessing Basal and Acute Autophagic Responses in the Adult *Drosophila* Nervous System: The Impact of Gender, Genetics and Diet on Endogenous Pathway Profiles. *PloS one* **11**, e0164239 (2016).
89. Barekat, A. *et al.* Using *Drosophila* as an integrated model to study mild repetitive traumatic brain injury. *Scientific reports* **6**, 25252 (2016).
90. Kim, M. *et al.* *Drosophila* Gyf/GRB10 interacting GYF protein is an autophagy regulator that controls neuron and muscle homeostasis. *Autophagy* **11**, 1358-1372 (2015).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

e

Genotype	Sample size	Mean lifespan (days)	Median lifespan (days)	Difference RNAi vs. Control	Prob. (Log-Rank)
Trial #1					
<i>TinC>mCherry^{RNAi}</i>	355	39.1	38	/	/
<i>TinC>Daw^{RNAi-1}</i>	389	44.3	46	21%	<0.001
<i>yw/Daw^{RNAi-1}</i>	301	45.2	44	/	/
<i>Hand>Daw^{RNAi-1}</i>	355	52.3	52	18%	<0.001
Trial #2					
<i>TinC>mCherry^{RNAi}</i>	364	45.3	44	/	/
<i>TinC>Daw^{RNAi-1}</i>	340	58	58	32%	<0.001
<i>Hand>mCherry^{RNAi}</i>	322	55.2	56	/	/
<i>Hand>Daw^{RNAi-1}</i>	315	63.3	64	14%	<0.001

Supplementary file

Supplemental Figure S2

Supplementary file

Supplemental Figure S3

Supplementary file

Supplemental Figure S4

Supplementary file

Supplemental Figure S5

Supplementary Table S1.

Primer name	Primer sequence (5'-3')	Symbol	FlyBase ID
RpL32-f	AAGAAGCGCACCAAGCACTTCATC	RpL32	FBgn0002626
RpL32-r	TCTGTTGTCGATACCCTTGGGCTT	RpL32	FBgn0002626
Daw-f	TGAGCCACCTCATCCAAATCACCT	Daw	FBgn0031461
Daw-r	TCGATCACGATGAATGGCCGGTAA	Daw	FBgn0031461
Rictor-f	CCCAACTGGCTAGCGTTTAT	Rictor	FBgn0031006
Rictor-r	GGCCTTTGCTTTCGTTTGAG	Rictor	FBgn0031006