

1 **Effects of radiofrequency electromagnetic radiation (RF-EMF) on** 2 **honey bee queen development and mating success**

3 Richard ODEMER^{1*}, Franziska ODEMER¹

4
5 ¹University of Hohenheim, Apicultural State Institute, 70593 Stuttgart, Germany

6
7 *Corresponding Author: richard.odemer@uni-hohenheim.de

8 phone: +49 1575 1 302 602

9 **ABSTRACT**

10 Mobile phones can be found almost everywhere across the globe, upholding a direct
11 point-to-point connection between the device and the broadcast tower. The emission of
12 radiofrequency electromagnetic radiation (RF-EMF) puts the surrounding environment
13 inevitably into contact with this pollutant. We have therefore exposed honey bee queen
14 larvae to the radiation of a common mobile phone device (GSM) during all stages of
15 their pre-adult development including pupation. After 14 days of exposure, hatching of
16 adult queens was assessed and mating success after further 11 days, respectively.
17 Moreover, full colonies were established of five of the untreated and four of the treated
18 queens to contrast population dynamics. We found that mobile phone radiation had
19 significantly reduced the hatching ratio but not the mating success. If treated queens
20 were successfully mated, colony development was not adversely affected. We provide
21 evidence that RF-EMF only acts detrimental within the sensitivity of pupal
22 development, once succeeded this point, no further impairment has manifested in
23 adulthood. Our results are discussed against the background of long-lasting
24 consequences for colony performance and the possible implication on periodic colony
25 losses.

26
27 Keywords: radiofrequency electromagnetic radiation (RF-EMF); honey bee queen
28 development; mating success; *Apis mellifera*; mobile phone devices; colony collapse
29 disorder (CCD)

30


31

32 HIGHLIGHTS

- 33 • Chronic RF-EMF exposure significantly reduced hatching of honey bee queens
- 34 • Mortalities occurred during pupation, not at the larval stages
- 35 • Mating success was not adversely affected by the irradiation
- 36 • After the exposure, surviving queens were able to establish intact colonies

37

38 GRAPHICAL ABSTRACT



39

40 Illustrations after Gullan & Cranston 2014

41

42

43

44

45 **1 INTRODUCTION**

46 The modern world turns around technological achievements and it is simply not
47 possible to imagine our everyday life without them. With an estimated 6.9 billion
48 subscriptions globally, mobile phone devices such as smart phones have established
49 their position in our society (WHO, 2014). In many countries, cell phones are important
50 tools not only for communication but also for bank transfers, newscast, social media and
51 numerous other conveniences with an increasing tendency. Provided that this market
52 will be further growing in the future, concerns are rising about the emission of
53 radiofrequency electromagnetic fields (RF-EMF) from these devices and their
54 broadcasting network, i.e. antennas and base stations, perceived as environmental
55 pollution (Balmori 2015).

56 Radiofrequency waves are electromagnetic fields, and unlike ionizing radiation such as
57 X-rays or gamma rays, they can neither break chemical bonds nor cause ionization in
58 the living tissue (Genuis & Lipp 2012). They are usually ranging from 30 kHz-300 GHz
59 with cell phones operating mainly between 800 MHz and 3 GHz, pulsed at low
60 frequencies (Hardell 2017). As a consequence, they are often strictly forbidden in
61 medical facilities and on airplanes, as the radiofrequency signals may interfere with
62 certain electro-medical devices and navigation systems.

63 In the last decade field and laboratory studies have furthermore demonstrated that RF-
64 EMF exposure is of ecological relevance. The radiation may have an impact on
65 surrounding flora as well as vertebrate and invertebrate organisms (Cucurachi et al.
66 2013). Effects have manifested in different ways and some of them are a cause of
67 concern. A large scale monitoring study (> 10 years) revealed that in trees, a closer
68 range to phone masts resulted in significant damages in the side facing the mast in
69 contrast to the opposite side (Waldmann-Selsam et al. 2016) whereas Roux et al. (2006,
70 2008) found exposed tomato plants to show similar consequences when wounded,
71 trimmed or burnt. In chicken eggs, Batellier et al. 2008 found an increased mortality
72 when exposed to cell phone radiation over the entire incubation period. Very similar to
73 previous study results from Bastide et al. (2001) and Grigoryev (2003), this

74 developmental stage seems to be particularly vulnerable for non-thermal radiation. A
75 proportional relationship between the intensity of the electromagnetic field and the
76 negative effects, however, could not be established (Batellier et al. 2008).

77 In fruit flies, reproduction and reproductive organs were also significantly affected by
78 mobile phone radiation (Panagopoulos et al. 2004, Panagopoulos 2012) unlike to the
79 findings of Weisbrot et al. (2003) where a beneficial effect on the reproductive success
80 was reported. In their study, the number of offspring increased by up to 50 % compared
81 to control, demonstrating controversial outcomes. Studies in insects have shown that
82 reproduction cycles and change of generations are quick, making this test system
83 suitable for the detection of possible consequences of RF-EMF exposure. Important
84 biological endpoints such as fertility, reproduction, behavior and development are rather
85 easy to implement, especially in a laboratory setting.

86 Besides the fruit fly as model organism, special ecological relevance is outlined by
87 pollinators, in particular by the honey bee *Apis mellifera*. They provide critical
88 pollination services valued at over \$200 billion worldwide (Lautenbach et al. 2012),
89 representing 9.5 % of the total human food production (Gallai et al. 2009). However,
90 bees have suffered periodic losses within the last century, and in the US a phenomenon
91 called colony collapse disorder (CCD) made headlines in the first decade of the new
92 millennium (vanEngelsdorp et al. 2009). Several causative factors have been outlined in
93 the past, among others, pathogens, malnutrition, management, and pesticides have been
94 narrowly focused as main culprits (Steinhauer et al. 2018).

95 Many other factors were also considered to have an impact on honey bee health,
96 however with a rather insignificant regard. A few to name are air pollution (Girling et
97 al., 2013; McFrederick et al., 2008), nanomaterials (Milivojevic et al., 2015), solar
98 radiation (Ferrari, 2014), robbing insects (Core et al., 2012) and global warming (Le
99 Conte & Navajas, 2008). Worthy of mention, in 2007 a story in an UK newspaper
100 brought to the fore that CCD can be linked to RF-EMF with drastic consequences for
101 bee behavior and homing success (Kimmel et al. 2007; Carreck, 2014). Subsequent
102 studies seem to provide supporting evidence of impaired behavior (Favre, 2011) and
103 affected homing ability (Ferrari, 2014), bearing a potential risk to other bee species such

104 as bumblebees (*Bombus terrestris*), when interacting with floral electric fields and
105 electric field sensing as important sensory modality (Clarke et al., 2013).

106 However, there are far too few scientific publications to draw a clear conclusion in
107 regard if and to which extent mobile phone radiation represents a real threat to honey
108 bees. A current review actually goes as far as stating that all examined studies were
109 characterized by substantial shortcomings which were sometimes even admitted by their
110 authors upfront (Verschaeve, 2014).

111 For a honey bee colony, health and productivity is directly linked to its queen. She
112 represents the growth potential expressed as productivity, being the only egg layer in
113 the collective and therefore responsible for a positive turnover of workers to increase in
114 size at the beginning of each bee season (Moore et al. 2015). In an US survey of winter
115 colony losses, the fourth most important factor identified was due to queen failure
116 (vanEngelsdorp et al., 2008). Given the importance of this individual, our experiments
117 therefore strictly focused on ontogenetic development and further mating success of
118 young queens. We have created a worst case scenario, where mobile phone radiation
119 was adopted by natural means of human exposure. To our knowledge this is the first
120 study that analyzes the effect of a chronic application of mobile phone radiation on
121 honey bee queens. We wanted to prove (i) if under field conditions and good apicultural
122 practice the radiation has any effect at all and to what extent, in addition (ii) we wanted
123 to follow queens which developed under chronic RF-EMF exposure to assess potential
124 risks for the bee colony.

125

126

127

128

129 **2 MATERIALS & METHODS**

130 **2.1 Field sites and weather conditions**

131 The field sites were located near the Apicultural State Institute in Stuttgart-Hohenheim,
132 Southern Germany (48°42'31.8"N 9°12'38.2"E). At the time present, natural food
133 sources consisted mainly of nectar from diverse local flora such as *Taraxacum*
134 *officinale*, *Rubus section*, *Tilia spp.* and others. The average temperature during the
135 experiment ranged from 15.2 to 20.1 °C with a precipitation of 90 to 45 L/m². Overall,
136 good weather conditions prevailed for both, mating and foraging (DWD 2018).


137 **2.2 Experimental setup**

138 This study was performed from May until August in 2018 with healthy queenright
139 colonies from the stock of our apiary. Two replications were employed simultaneously,
140 consisting of two collector colonies: Rep1 (Control1 + EMF1) and Rep2= (Control2 +
141 EMF2). For both approaches, one brood frame with almost fully covered areas of sealed
142 brood and attached bees from eight random colonies were taken out on D-9 and placed
143 in a new ten-frame box, respectively. This box was supplied with two frames of food, as
144 well as a second box on top with ten food frames to ensure sustenance and sufficient
145 room for the hatching bees. Nine days after this procedure (D0), the hive was inspected,
146 and where appropriate, supersedure cells were removed to prevent the introduction of a
147 young queen. Further, 18 frames then were split homogeneously but random into two
148 boxes with nine frames each, complemented with a grafting frame in the center. L1
149 larvae from a selected colony were grafted and introduced, respectively. Again, grafting
150 of the larvae was randomized by using both sides of the brood comb (A and B). Per
151 replication, 26 larvae (13 A, 13 B) were assigned to each treatment, i.e. control and
152 EMF.

153 The two boxes then were placed at a different location in approximately 3 km distance
154 to prevent worker bees to return to their original position. Subsequently, at different
155 intervals, assessments were performed to check the no. of accepted larvae after grafting
156 (D1), to protect the capped cells before hatching (D10), to check the hatching rate (D13)
157 and the mating success (D24). After the young queens have hatched, they were

158 transferred to mating units consisting of one of the former brood frames with
159 approximately 1,000 bees attached and one food comb.

160 Successful mating was confirmed on D24 by the presence of eggs, young larvae and
161 capped brood and queens from each treatment (five from the control, four from the
162 treatment) were re-accommodated in new 10-frame boxes to develop into full colonies.
163 After approximately twelve weeks (D88), a colony assessment was performed to record
164 the number of bees and brood. See Fig. 1 for a detailed timeline.




165

166 [Fig. 1 Timeline of the experiment. At D-9, eight brood frames with attached bees were taken out from
167 respective donor colonies and placed in one collector colony. At D0, supersedure cells were removed and
168 the collector colony was split in two sub-colonies. In addition, a grafting frame with L1 larvae was
169 inserted. RF-EMF exposure lasted until D13, when queens were about to hatch. Young queens were
170 subsequently inserted into mating units where mating success was checked at D24. Successfully mated
171 queens with one frame of approximately 1,000 bees were relocated into new boxes where they were able
172 to establish a new colony. Finally, at D88 the condition of these colonies was assessed.]

173 2.3 EMF treatment

174 Queen larvae/pupae were treated with a mobile phone (AEG M1220, GSM quad band:
175 800/900/1800/1900 MHz, China) attached to the grafting frame holding 26 queen cups
176 (Nicot, NICOTPLAST SAS, Maisod, France), this device was turned off in the control
177 group for sham exposure. To ensure power supply, the phone was equipped with a
178 power bank (PLOCHY 24,000 mAh Solar, China), the battery status was frequently
179 checked. After the larvae were grafted into the cups by using an appropriate tool, 15
180 telephone calls with a two minute duration were applied daily for a total of two weeks
181 (non-modulated emission) at random. The radiation was measured three times in three
182 different distances to the mobile phone with a fixed instrument illustrated in Fig. 2 (Pyle
183 PMD74, Calibration: 2450 MHz, measurement range: 0-15 mW/cm², China) to verify
184 an adequate EMF output.



185 [Fig. 2 Grafting frame placed in the EMF treatment colony containing 26 queen cups. The mobile phone
186 device was attached in the center of the frame, its radiation intensity is indicated with the differently
187 colored sections in the illustration (dark > light)]
188

189 2.4 Colony assessment

190 The amount of bees and brood cells (open and sealed) were estimated with the Liebefeld
191 Method (Imdorf et al., 1987), which is a feasible tool to provide accurate and reliable
192 evaluation of colony strength (measuring error +/- 10 %). Care was taken that all

193 colonies were evaluated by the same person to minimize variation and colony
194 assessment was conducted in the morning before bee flight.

195 **2.5 Statistical analysis**


196 We evaluated the mortality data with a Kaplan-Meier-Survival analysis. Survivorship
197 between control and treatment was compared pairwise and tested for significance with a
198 Log-Rank Tests (Cox-Mantel). Individuals collected at the end of the experiment were
199 considered censored, as were those observed but not collected on the final day.
200 Furthermore, larvae that disappeared during the experiment were considered dead on the
201 last day they were seen. Both treatment groups and the two replicates (Rep1= Control1
202 + EMF1; Rep2= Control2 + EMF2) were additionally compared with a Cox
203 proportional hazards model to determine the hazard ratio (HR). Possible inter-colony
204 effects were evaluated as covariate to justify pooling data of the same treatments. The
205 estimated number of bees and brood cells were checked with a Shapiro-Wilk test for
206 normal distribution. If data was normal, a one-way ANOVA was performed on the two
207 experimental groups, respectively. For all tests RStudio (R Core Team, 2018) and
208 significance level of $\alpha=0.05$ was used.

209

210 **3 RESULTS**

211 **3.1 Honey bee queen survival**

212 The Kaplan-Meier-Survival analysis of both groups showed a significant difference
213 indicating a higher mortality of the EMF treated bees when compared to the control
214 group ($p=0.0054$) (Fig. 3). In addition, a Cox proportional hazards model was applied to
215 determine the hazard ratio (HR) displayed as forest plot (Fig. 4). With a HR of 2.3 the
216 EMF treated queens had a significantly increased risk of dying when compared to the
217 control ($p=0.003$). Moreover, the two replicates (Rep1 and Rep2) were compared as
218 covariate to display possible inter-colony effects. However, with a HR of 1.7 queens in
219 Rep2 did not have a higher risk of dying when compared to Rep1 ($p=0.062$), therefore
220 data of both replicates were pooled.




221

222 [Fig. 3 Both groups were compared with a Kaplan-Meier-Survival analysis. A post-hoc Log-Rank test
 223 (Cox-Mantel) revealed a significant higher mortality in the EMF treatment when compared to the control
 224 (Log-Rank $p=0.0054$), where a significant decrease of individuals occurred during the pupation phase of
 225 the experiment (see also Fig. 5)]

226

227




228

229 [Fig. 4 Both treatment groups and the two replicates (Rep1 and Rep2) were additionally compared with a
230 Cox proportional hazards model to determine the hazard ratio (HR) displayed as forest plot. With a HR of
231 2.3 the EMF treated queens had a significantly increased risk of dying when compared to the control
232 ($p=0.003$). And with a HR of 1.7 queens in Rep2 did not have a higher risk of dying when compared to
233 Rep1 ($p=0.062$)]

234 3.2 Hatching and mating success

235 The acceptance rate of grafted larvae on D1 was 76.9 % and identical in both
236 treatments. As shown in Fig. 3, a significant decrease of individuals in the EMF
237 treatment occurred during the pupation phase of the experiment. At D10, queen cells
238 were protected with a cage to prevent hatching queens from killing each other. We
239 observed a ratio of 73.1 (control) to 65.4 % (treatment) at this stage compared to the
240 initially grafted cells. The hatch of adult queens at D13 revealed a significant decrease
241 of formerly treated queens during pupation with a proportion of 69.2 % in the control to
242 38.5 % in the treatment, levelling out with a similar decrease of both groups at the
243 assessment of mating success at D24 (control 57.7 %, treatment 28.8 %) (for statistical
244 evaluations see also Fig. 3).




245

246 [Fig. 5 Total number of L1 larvae grafted and followed through their ontogenetic development from pupa
247 to adult. In the EMF treatment a significant decrease of individuals came into effect within the pupation
248 phase of the experiment (see also Fig. 3). Illustrations after Gullan & Cranston 2014]

249


250 3.3 Colony assessment

251 The population of bees and brood cells was estimated at D88. The results are shown in
252 Fig. 6A for the number of bees and in Fig. 6B for the number of brood cells. We
253 compared the two treatment groups with a one-way ANOVA but could not see
254 significant differences for the number of bees ($p=0.688$) or the amount of brood cells
255 ($p=0.768$).



256

257 [Fig. 6A Number of bees estimated at D88 in the colonies of the control (n=5) and of the EMF treatment
258 (n=4). Same letters indicate no statistically significantly differences ($p=0.688$, ANOVA).]



259

260 [Fig. 6B Number of brood cells estimated at D88 in the colonies of the control (n=5) and of the EMF
261 treatment (n=4). Same letters indicate no statistically significant differences (p=0.768, ANOVA).]

262

263

264

265 **4 DISCUSSION**

266 The emission of radiofrequency electromagnetic fields (RF-EMF) and their negative
267 effects towards honey bee health has been controversially discussed in the past (Carreck
268 2014; Verschaeve 2014; Panagopoulos et al. 2016). Here, we could demonstrate for the
269 first time that RF-EMF exposure has significantly affected ontogenetic queen
270 development under field conditions. We observed an increased mortality during
271 pupation resulting in a reduced hatching rate of the later queens. This is in line with a
272 reduced reproductive capacity found in fruit flies (*Drosophila melanogaster*)
273 (Panagopoulos et al. 2004, Margaritis et al. 2014), where a linear decrease of fecundity
274 was reported with the frequency of exposure (Panagopoulos and Margaritis 2010). This
275 decrease was further associated with the distance to the mobile phone device showing
276 the strongest effects at < 10 cm (Panagopoulos et al. 2010). In our setup, the most
277 distant queen cups were approximately 21 cm away from the radiation source and we
278 therefore assume that for all larvae a worst case scenario came into effect. In addition,
279 the impairment of fruit flies seemingly depended on field intensity (Panagopoulos et al.
280 2007) not only reducing the offspring but also the ovarian size of the exposed subjects
281 (Panagopoulos 2012).

282 At present, only a few studies have investigated the influence of irradiation on insect
283 development. As an example, larvae and pupae of the dried fruit beetle (*Carpophilus*
284 *hemipterus*) and the yellow fever mosquito (*Aedes aegypti*) were exposed to Gamma
285 radiation (ionizing radiation). The radiotherapy did not cause acute death in larvae but
286 decreased pupation significantly, no effects however could be observed when either
287 young or old pupae were exposed (Johnson 1987, Akter & Khan 2014). It seems likely
288 that RF-EMF had a similar effect in our study, as larval mortality was not elevated.
289 However, this should be further underpinned by exposing larvae and pupae separately.
290 Moreover, Vilić et al. (2017) found honey bee worker larvae significantly affected when
291 exposed to modulated but not to non-modulated RF-EMF radiation, resulting in DNA
292 damage and further corroborating our hypothesis as we only have used non-modulated
293 fields.

294 In addition, we could show that mating success remained unaffected suggesting that
295 navigation and the possible disruption of magnetoreception came not into effect or was
296 at least not long-lasting (Vácha et al. 2009). Interestingly, we provide evidence that
297 developing honey bee queens once they have survived RF-EMF exposure seem to retain
298 the ability to establish an intact colony. This is indicated by similarly strong numbers of
299 bees and the amount of brood in both our treatment groups with the absence of any
300 signs of impairment (e.g. patchy brood pattern). As a further critical step of colony
301 survival however, overwintering should also be assessed to elucidate possible long term
302 effects from the irradiation (Smart et al. 2016).

303 The social entity as a whole is able to buffer environmental stressor of various kinds as
304 an expression of social resilience (Straub et al. 2015). Worker bees are nursing eggs and
305 feeding larvae of different casts in their social state, potentially contributing to this
306 mechanism. Here we focused on the development of individual queens from larvae to
307 adult, however, the outcome of our study could also be influenced by the condition of
308 the collector colonies that we have created but not further assessed. Eggs, larvae and
309 pupae are very sensitive stages of development and intensive care is taken to supply
310 their substantial needs in terms of nutrition and environmental conditions, i.e.
311 maintaining a constant temperature and humidity (Wang et al. 2015, Eouzan et al.
312 2018). RF-EMF radiation is known to affect bees behavior in different ways (Favre
313 2011, Ferrari 2014), which makes it plausible that brood care could also be adversely
314 affected. This important factor should be further investigated and included in future
315 experiments.

316 With an increasing number of mobile phone devices and as a consequence of good
317 accessibility a higher density of phone masts, not only urban but also rural areas in
318 particular are more and more exposed to irradiation (Balmori 2009). A measurement of
319 RF-EMF intensities across different European cities revealed maximum radiation values
320 ranging from 0.84 to 0.59 V/m corresponding to 92.33 and 187.16 nW/cm² (Urbinello et
321 al. 2014a), respectively, with a maximum value of 127 nW/cm² in public transport
322 (Sagar et al. 2016). In contrast, the power flux density measured in our study seemed to
323 be way beyond these values, demonstrating that the intermittent stress on the test
324 subject(s) can be many fold higher than average levels measured in the surroundings,

325 emitted from generators or found in agglomerations. Our findings confirm that there is a
326 high variability in mobile phone emission (Frei et al. 2009), representing an important
327 feature in terms of bioactivity towards living organism's defense against environmental
328 stressors (Panagopoulos et al. 2015). The authors therefore suggest not using simulated
329 but real mobile phone emissions in an experimental setup, which we have considered. In
330 addition, we have tried to apply a human exposure scenario in terms of average number
331 of calls and average call duration performed with mobile phone devices. The mobile call
332 duration reported by the German Federal Network Agency (2011) was 2.5 min per day,
333 in Shum et al. (2011) ranging from 2.1 min (self-reports) to 2.8 min (billing records)
334 and < 2 min in Friebel & Seabright (2011). Further, the average number of calls per day
335 ranging from 4.1 (Shum et al. 2011) to 5 per day in adults (Lenhart 2010). In contrast,
336 an average of 33.1 min was reported for total mobile phone call duration from
337 undergraduate college students per day in the US (Roberts et al. 2014). We therefore
338 decided to employ 2 min per call and 15 calls per day resulting in 30 min exposure per
339 day in our experiment, representing a realistic human exposure.

340 Different exposure scenarios were applied in honey bees and a broad range of effects
341 are reported (Cucurachi et al. 2013). Some studies even claimed with RF-EMF to have
342 found the major cause for CCD (Carreck 2014). However, many of these studies had
343 substantial deficits such as a very low sample size (Sharma & Kumar 2010),
344 intransparent methods (Sahib 2011, Kumar et al. 2011, Dalio 2015) or were even
345 preliminary and did not undergo peer-review (Kimmel et al. 2007). Therefore, findings
346 of this quality were generally not considered reliable in their contribution to colony
347 losses and are far from conclusive (Carreck 2014). To achieve a broader understanding
348 how RF-EMF potentially influences the honey bee superorganism, it is mandatory to
349 emphasize the conditions under which the study was conducted, particularly the level
350 and duration of exposure, in the presence of the relevant environmental situation
351 (Verschaeve 2014).

352 As a trend of the last decades, beekeeping became famous with the life style of
353 townsmen all across the globe (Lorenz & Stark 2015, Kohsaka et al. 2017, Stange et al.
354 2017). Therefore, density of bee colonies held in urban areas has dramatically increased
355 and may favor the spread of diseases or pathogens (Youngsteadt et al. 2015). However,

356 following this trend also bears the risk of a higher exposure to RF-EMF emission, which
357 seems to be continuously increasing in major cities (Urbinello et al. 2014b), potentially
358 affecting bee health in a future scenario. It might also be worthy to look into parasite-
359 host-interactions of the honey bee, *Varroa destructor* in particular, where a disturbance
360 through RF-EMF in host-finding could actually be a benefit (Frey et al. 2013).
361 Surprisingly, not many studies are available that are investigating the influence of such
362 irradiation on bees and other important pollinators. It has even been suggested to create
363 pollinator reservoirs beneath power corridors for an optimal land use and as a benefit for
364 many insects (Russel et al. 2018). Yet, it remains unclear to what extent
365 electromagnetic fields can possibly influence these microenvironments.

366 **Conclusion**

367 Even though detrimental effects on ontogenetic queen development were revealed by
368 the outcome of our study, caution is needed in interpreting these results. We have
369 created by far a worst case scenario to which honey bee colonies would not be exposed
370 under realistic conditions. Duration and level were similar to average human exposure
371 by the use of a mobile phone, but not to those present at an apiary, neither in rural nor in
372 urban areas. And yet, queens that survived the treatment were able to establish full
373 functional colonies, demonstrating an immense recovering potential. Therefore we do
374 not assume any acute negative effects on bee health in the mid-term. However, we do
375 not rule out an influence through lower doses of permanent irradiation, in particular on a
376 chronic sublethal level. Hence, we urgently suggest further research should be carried
377 out in the long-term to ascertain what impacts are to be expected.

378

379

380

381 **ACKNOWLEDGEMENTS**

382 Many thanks to Simon Hummel for providing adequate breeding lines to conduct
383 research with extremely calm honey bees.

384 This research did not receive any specific grant from funding agencies in the public,
385 commercial, or not-for-profit sectors.

386

387 Declarations of interest: none.

388

389 Ethical approval: This article does not contain any studies with human participants or
390 animals performed by any of the authors.

391

392

393

394

395 **REFERENCES**

396

397 Akter H, Khan SA (2014) Sensitivity of Immature Stages of Dengue Causing Mosquito,
398 *Aedes aegypti* (L.) to Gamma Radiation. *J Entomol* 11:56–67. doi:
399 10.3923/je.2014.56.67

400 Balmori A (2009) Electromagnetic pollution from phone masts. Effects on wildlife.
401 *Pathophysiology* 16:191–199. doi: 10.1016/j.pathophys.2009.01.007

402 Balmori A (2015) Anthropogenic radiofrequency electromagnetic fields as an emerging
403 threat to wildlife orientation. *Sci Total Environ* 518–519:58–60. doi:
404 10.1016/j.scitotenv.2015.02.077

405 Bastide M, Youbicier-Simo BJ, Lebecq JC, Giaimis J. (2001) Toxicological study of
406 electromagnetic radiation emitted by television and video display screens and cellular
407 telephones on chickens and mice. *Indoor Built Environ*;10:291–8.

408 Batellier F, Couty I, Picard D, Brillard JP (2008) Effects of exposing chicken eggs to a
409 cell phone in ‘call’ position over the entire incubation period. *Theriogenology*;69: 737–
410 45.

411 Carreck N (2014) Electromagnetic Radiation and Bees, Again... *Bee World* 91:4, 101-
412 102. doi: : 10.1080/0005772X.2014.11417624

413 Clarke D, Whitney H, Sutton G, Robert D (2013) Detection and Learning of Floral
414 Electric Fields by Bumblebees. *Science* (80-) 340:66–69. doi: 10.1126/science.1230883

415 Core A, Runckel C, Ivers J, Quock C, Siapno T, DeNault S, Brown B, DeRisi J, Smith
416 CD, Hafernik J (2012) A New Threat to Honey Bees, the Parasitic Phorid Fly
417 *Apocephalus borealis*. *PLoS One* 7:e29639. doi: 10.1371/journal.pone.0029639

418 Cucurachi S, Tamis WLM, Vijver MG, Peijnenburg WJGM, Bolte JFB, de Snoo GR
419 (2013) A review of the ecological effects of radiofrequency electromagnetic fields (RF-
420 EMF). *Environ Int* 51:116–140. doi: 10.1016/j.envint.2012.10.009

- 421 Dalio, J. S. (2015) Effect of electromagnetic (cell phone) radiations on *Apis mellifera*,
422 Journal of Res in Agriculture and Animal science 2(12): 06-10.
- 423 Deutscher Wetterdienst DWD (2018) Deutschlandwetter im Sommer 2018
424 Pressemitteilung. www.dwd.de, Accessed 26 September 2018
- 425 Eouzan I, Garnery L, Pinto MA, Delalande D, Neves CJ (2018) Hygroregulation , a key
426 ability for eusocial insects : Native Western European bees as a case study. bioRxiv.
427 doi: 10.1101/351874
- 428 Favre D (2011) Mobile phone-induced honeybee worker piping. *Apidologie* 42:270–
429 279. doi: 10.1007/s13592-011-0016-x
- 430 Federal Network Agency (2011) Press Release 28 July 2011: Average duration of 2.5
431 minutes for mobile calls, Bonn, Germany, [http:// www.bundesnetzagentur.de](http://www.bundesnetzagentur.de), Accessed
432 24 August 2018
- 433 Ferrari TE (2014) Magnets, magnetic field fluctuations and geomagnetic disturbances
434 impair the homing ability of honey bees (*Apis mellifera*). *J Apic Res* 53:452–465. doi:
435 10.3896/IBRA.1.53.4.15
- 436 Frei P, Mohler E, Neubauer G, Theis G, Bürgi A, Fröhlich J, Braun-Fahrlander C, Bolte
437 J, Egger M, Rösli M (2009) Temporal and spatial variability of personal exposure to
438 radio frequency electromagnetic fields. *Environ Res* 109:779–785. doi:
439 10.1016/j.envres.2009.04.015
- 440 Frey E, Odemer R, Blum T, Rosenkranz P (2013) Activation and interruption of the
441 reproduction of *Varroa destructor* is triggered by host signals (*Apis mellifera*). *J*
442 *Invertebr Pathol* 113:56–62. doi: 10.1016/j.jip.2013.01.007
- 443 Friebel G, Seabright P (2011) Do women have longer conversations? Telephone
444 evidence of gendered communication strategies. *J Econ Psychol* 32:348–356. doi:
445 10.1016/j.joep.2010.12.008
- 446 Gallai N, Salles JM, Settele J, Vaissière BE (2009) Economic valuation of the
447 vulnerability of world agriculture confronted with pollinator decline. *Ecol Econ* 68:810–
448 821. doi: 10.1016/j.ecolecon.2008.06.014

- 449 Genuis SJ, Lipp CT (2012) Electromagnetic hypersensitivity: Fact or fiction? *Sci Total*
450 *Environ* 414:103–112. doi: 10.1016/j.scitotenv.2011.11.008
- 451 Girling RD, Lusebrink I, Farthing E, Newman TA, Poppy GM (2013) Diesel exhaust
452 rapidly degrades floral odours used by honeybees. *Sci Rep* 3:2779. doi:
453 10.1038/srep02779
- 454 Grigoryev Y. (2003) Biological effects of mobile phone electromagnetic field on chick
455 embryo (risk assessment using the mortality rate). *Radiats Biol Radioecol*;43:541–3
- 456 Gullan P.J., Cranston P.S. (2014) *The Insects: An Outline of Entomology*, Fifth Edition,
457 John Wiley & Sons Blackwell, Oxford ISBN: 1-4051-1113-5
- 458 Hardell L (2017) World Health Organization, radiofrequency radiation and health - a
459 hard nut to crack (Review). *Int J Oncol* 51:405–413. doi: 10.3892/ijo.2017.4046
- 460 Imdorf A, Buehlmann G, Gerig L, Kilchenmann V (1987) Überprüfung der
461 Schätzmethode zur Ermittlung der Brutfläche und der Anzahl Arbeiterinnen in
462 freifliegenden Bienenvölkern. *Apidologie* 18:137–146. doi:
463 <http://dx.doi.org/10.1051/apido:19870204>
- 464 Johnson JA (1987) Sensitivity of Larvae, Pupae, and Adults of the Driedfruit Beetle
465 (Coleoptera : Nitidulidae) to Gamma Radiation. *J Econ Entomol* 80:1302–1305.
- 466 Kimmel S, Kuhn J, Harst W, Stever H (2007) *Electromagnetic Radiation : Influences on*
467 *Honeybees (Apis mellifera)*, unpublished 1–6.
- 468 Kohsaka R, Park MS, Uchiyama Y (2017) Beekeeping and honey production in Japan
469 and South Korea: past and present. *J Ethn Foods* 4:72–79. doi:
470 10.1016/j.jef.2017.05.002
- 471 Kumar NR, Sangwan S, Badotra P. (2011) Exposure to cell phone radiations produces
472 biochemical changes in worker honey bees. *Toxicol Int*;18(1):70–2.
- 473 Lautenbach S, Seppelt R, Liebscher J, Dormann CF (2012) Spatial and temporal trends
474 of global pollination benefit. *PLoS One*. doi: 10.1371/journal.pone.0035954

- 475 Le Conte Y, Navajas M (2008) Climate change: impact on honey bee populations and
476 diseases. *Rev Sci Tech* 27:485–97, 499–510.
- 477 Lenhart A (2010) Cell phones and American adults: They make just as many calls, but
478 text less often than teens. *Pew Res Cent* 1–42.
- 479 Lorenz S, Stark K (2015) Saving the honeybees in Berlin? A case study of the urban
480 beekeeping boom, *Environmental Sociology*, 1:2, 116-126, DOI:
481 10.1080/23251042.2015.1008383
- 482 Margaritis LH, Manta AK, Kokkaliaris KD, Schiza D, Alimisis K, Barkas G, Georgiou
483 E, Giannakopoulou O, Kollia I, Kontogianni G, Kourouzidou A, Myari A, Roumelioti
484 F, Skouroliakou A, Sykioti V, Varda G, Xenos K, Ziomas K (2014) Drosophila
485 oogenesis as a bio-marker responding to EMF sources. *Electromagn Biol Med* 33:165–
486 89. doi: 10.3109/15368378.2013.800102
- 487 McFrederick QS, Kathilankal JC, Fuentes JD (2008) Air pollution modifies floral scent
488 trails. *Atmos Environ* 42:2336–2348. doi: 10.1016/j.atmosenv.2007.12.033
- 489 Milivojević T, Glavan G, Božič J, Sepčić K, Mesarič T, Drobne D (2015) Neurotoxic
490 potential of ingested ZnO nanomaterials on bees. *Chemosphere* 120:547–554. doi:
491 10.1016/j.chemosphere.2014.07.054
- 492 Moore P, Wilson ME, Skinner JA (2015) Honey Bee Queens: Evaluating the Most
493 Important Colony Member, Department of Entomology and Plant Pathology, The
494 University of Tennessee, Knoxville TN, www.extension.org, Accessed 26 September
495 2018
- 496 Panagopoulos DJ (2012) Effect of Microwave Exposure on the Ovarian Development of
497 *Drosophila melanogaster*. *Cell Biochem Biophys* 63:121–132. doi: 10.1007/s12013-
498 012-9347-0
- 499 Panagopoulos DJ, Cammaerts M-C, Favre D, Balmori A (2016) Comments on
500 environmental impact of radiofrequency fields from mobile phone base stations. *Crit*
501 *Rev Environ Sci Technol* 46:885–903. doi: 10.1080/10643389.2016.1182107

- 502 Panagopoulos DJ, Chavdoula ED, Karabarbounis A, Margaritis LH (2007) Comparison
503 of bioactivity between GSM 900 MHz and DCS 1800 MHz mobile telephony radiation.
504 *Electromagn Biol Med* 26:33–44. doi: 10.1080/15368370701205644
- 505 Panagopoulos DJ, Chavdoula ED, Margaritis LH (2010) Bioeffects of mobile telephony
506 radiation in relation to its intensity or distance from the antenna. *Int J Radiat Biol*
507 86:345–357. doi: 10.3109/09553000903567961
- 508 Panagopoulos DJ, Johansson O, Carlo GL (2015) Real versus simulated mobile phone
509 exposures in experimental studies. *Biomed Res Int*. doi: 10.1155/2015/607053
- 510 Panagopoulos DJ, Karabarbounis A, Margaritis LH (2004) Effect of GSM 900-MHz
511 mobile phone radiation on the reproductive capacity of *Drosophila melanogaster*.
512 *Electromagn Biol Med* 23:29–43. doi: 10.1081/JBC-120039350
- 513 Panagopoulos DJ, Margaritis LH (2010) The effect of exposure duration on the
514 biological activity of mobile telephony radiation. *Mutat Res - Genet Toxicol Environ*
515 *Mutagen* 699:17–22. doi: 10.1016/j.mrgentox.2010.04.010
- 516 R Core Team (2018) R: A language and environment for statistical computing. R
517 Foundation for Statistical Computing, Vienna, Austria.
- 518 Roberts J, Yaya L, Manolis C (2014) The invisible addiction: Cell-phone activities and
519 addiction among male and female college students. *J Behav Addict* 3:254–265. doi:
520 10.1556/JBA.3.2014.015
- 521 Roux, D., Vian, A., Girard, S., Bonnet, P., Paladian, F., Davies, E., and Ledoigt, G.
522 (2006). Electromagnetic fields (900 MHz) evoke consistent molecular responses in
523 tomato plants. *Physiologia Plantarum*, 128, 283–288.
- 524 Roux, D., Vian, A., Girard, S., Bonnet, P., Paladian, F., Davies, E., and Ledoigt, G.
525 (2008). High frequency (900 MHz) low amplitude (5 V m⁻¹) electromagnetic field: A
526 genuine environmental stimulus that affects transcription, translation, calcium and
527 energy charge in tomato. *Planta*, 227, 883–891.
- 528 Russell KN, Russell GJ, Kaplan KL, Mian S, Kornbluth S (2018) Increasing the
529 conservation value of powerline corridors for wild bees through vegetation

- 530 management: an experimental approach. *Biodivers Conserv* 27:2541–2565. doi:
531 10.1007/s10531-018-1552-8
- 532 Sagar S, Struchen B, Finta V, Eeftens M, Rössli M (2016) Use of portable exposimeters
533 to monitor radiofrequency electromagnetic field exposure in the everyday environment.
534 *Environ Res* 150:289–298. doi: 10.1016/j.envres.2016.06.020
- 535 Sahib, S. (2011). Electromagnetic radiation (EMR) clashes with honey bees.
536 *International Journal of Environmental Sciences* 1, 897–900.
- 537 Sharma MC, Kumar N (2010) Changes in honeybee behaviour and biology under the
538 influence of cellphone radiations. *Vet Parasitol* 154:98–102. doi:
539 10.1016/j.vetpar.2008.03.001
- 540 Shum M, Kelsh MA, Sheppard AR, Zhao K (2011) An evaluation of self-reported
541 mobile phone use compared to billing records among a group of engineers and
542 scientists. *Bioelectromagnetics* 32:37–48. doi: 10.1002/bem.20613
- 543 Smart M, Pettis J, Rice N, Browning Z, Spivak M (2016) Linking Measures of Colony
544 and Individual Honey Bee Health to Survival among Apiaries Exposed to Varying
545 Agricultural Land Use. *PLoS One* 11:e0152685. doi: 10.1371/journal.pone.0152685
- 546 Stange E, Zulian G, Rusch G, Barton D, Nowell M (2017) Ecosystem services mapping
547 for municipal policy: ESTIMAP and zoning for urban beekeeping. *One Ecosyst*
548 2:e14014. doi: 10.3897/oneeco.2.e14014
- 549 Steinhauer N, Kulhanek K, Antúnez K, Human H, Chantawannakul P, Chauzat M-P,
550 VanEngelsdorp D (2018) Drivers of colony losses. *Curr Opin Insect Sci* 26:142–148.
551 doi: 10.1016/j.cois.2018.02.004
- 552 Straub L, Williams GR, Pettis J, Fries I, Neumann P (2015) Superorganism resilience:
553 eusociality and susceptibility of ecosystem service providing insects to stressors. *Curr*
554 *Opin Insect Sci* 12:109–112. doi: 10.1016/j.cois.2015.10.010
- 555 Urbinello D, Joseph W, Huss A, Verloock L, Beekhuizen J, Vermeulen R, Martens L,
556 Rössli M (2014) Radio-frequency electromagnetic field (RF-EMF) exposure levels in

- 557 different European outdoor urban environments in comparison with regulatory limits.
558 *Environ Int* 68:49–54. doi: 10.1016/j.envint.2014.03.007
- 559 Urbinello D, Joseph W, Verloock L, Martens L, Rösli M (2014) Temporal trends of
560 radio-frequency electromagnetic field (RF-EMF) exposure in everyday environments
561 across European cities. *Environ Res* 134:134–142. doi: 10.1016/j.envres.2014.07.003
- 562 Vacha M, Puzova T, Kviclova M (2009) Radio frequency magnetic fields disrupt
563 magnetoreception in American cockroach. *J Exp Biol* 212:3473–3477. doi:
564 10.1242/jeb.028670
- 565 van Engelsdorp D, Hayes J, Underwood RM, Pettis J (2008) A survey of honey bee
566 colony losses in the U.S., Fall 2007 to Spring 2008. *PLoS One* 3:8–13. doi:
567 10.1371/journal.pone.0004071
- 568 Vanengelsdorp D, Evans JD, Saegerman C, Mullin C, Haubruge E, Nguyen BK, Frazier
569 M, Frazier J, Cox-Foster D, Chen Y, Underwood R, Tarpay DR, Pettis JS (2009) Colony
570 collapse disorder: a descriptive study. *PLoS One* 4:e6481. doi:
571 10.1371/journal.pone.0006481
- 572 Verschaeve L (2014) Environmental Impact of Radiofrequency Fields from Mobile
573 Phone Base Stations. *Crit Rev Environ Sci Technol* 44:1313–1369. doi:
574 10.1080/10643389.2013.781935
- 575 Vilić M, Tlak Gajger I, Tucak P, Štambuk A, Šrut M, Klobučar G, Malarić K, Žura Žaja
576 I, Pavelić A, Manger M, Tkalec M (2017) Effects of short-term exposure to mobile
577 phone radiofrequency (900 MHz) on the oxidative response and genotoxicity in honey
578 bee larvae. *J Apic Res* 56:430–438. doi: 10.1080/00218839.2017.1329798
- 579 Waldmann-Selsam C, Balmori-de la Puente A, Breunig H, Balmori A (2016)
580 Radiofrequency radiation injures trees around mobile phone base stations. *Sci Total*
581 *Environ* 572:554–569. doi: 10.1016/j.scitotenv.2016.08.045
- 582 Wang Y, Ma L-T, Xu B-H (2015) Diversity in life history of queen and worker honey
583 bees, *Apis mellifera* L. *J Asia Pac Entomol* 18:145–149. doi:
584 10.1016/j.aspen.2014.11.005

- 585 Weisbrot D, Lin H, Ye L, Blank M, Goodman R (2003) Effects of mobile phone
586 radiation on reproduction and development in *Drosophila melanogaster*. *J Cell*
587 *Biochem*;89: 48–55
- 588 World Health Organization WHO (2014) Electromagnetic fields and public health:
589 mobile phones, www.who.int, Accessed 16 August 2018
- 590 Youngsteadt E, Appler RH, López-Urbe MM, Tarpy DR, Frank SD (2015)
591 Urbanization increases pathogen pressure on feral and managed honey bees. *PLoS One*.
592 doi: 10.1371/journal.pone.0142031