

1 **Quantifying point-mutations in shotgun metagenomic data**

2 Shruthi Magesh^{1,2}, Viktor Jonsson³, Johan Bengtsson-Palme^{1,4,5,*}

3

4 *Affiliations*

5 ¹ Wisconsin Institute of Discovery, University of Wisconsin-Madison, 330 North Orchard Street,
6 Madison WI 53715, USA

7 ² Department of Biotechnology, School of Bioengineering, SRM Institute of Science and
8 Technology, Kattankulathur 603203, Tamil Nadu, India

9 ³ Chalmers Computational Systems Biology Infrastructure, Chalmers University of Technology,
10 SE-412 96, Gothenburg, Sweden

11 ⁴ Department of Infectious Diseases, Institute of Biomedicine, The Sahlgrenska Academy,
12 University of Gothenburg, Guldhedsgatan 10, SE-413 46, Gothenburg, Sweden

13 ⁵ Centre for Antibiotic Resistance research (CARE) at University of Gothenburg, Gothenburg,
14 Sweden

15 * Corresponding author: Johan Bengtsson-Palme, Department of Infectious Diseases,
16 Guldhedsgatan 10, SE-413 46, Gothenburg, Sweden. Phone: +46 31 342 46 26. Fax: +46 31 84
17 61 13. E-mail address: johan.bengtsson-palme@microbiology.se

18

19

20 **Abstract**

21 Metagenomics has emerged as a central technique for studying the structure and function of
22 microbial communities. Often the functional analysis is restricted to classification into broad
23 functional categories. However, important phenotypic differences, such as resistance to
24 antibiotics, are often the result of just one or a few point mutations in otherwise identical
25 sequences. Bioinformatic methods for metagenomic analysis have generally been poor at
26 accounting for this fact, resulting in a somewhat limited picture of important aspects of microbial
27 communities. Here, we address this problem by providing a software tool called Mumame, which
28 can distinguish between wildtype and mutated sequences in shotgun metagenomic data and
29 quantify their relative abundances. We demonstrate the utility of the tool by quantifying antibiotic
30 resistance mutations in several publicly available metagenomic data sets. We also identified that
31 sequencing depth is a key factor to detect rare mutations. Therefore, much larger numbers of
32 sequences may be required for reliable detection of mutations than for most other applications of
33 shotgun metagenomics. Mumame is freely available from
34 <http://microbiology.se/software/mumame>

35

36 **Keywords**

37 Antibiotic resistance, Bioinformatic tools, Metagenomics, Mutation frequencies, Mutation
38 mapping, Statistical methods

39

40 **Introduction**

41 The revolution in sequencing capacity has created an unprecedented ability to glimpse into the
42 functionality of microbial communities, using large-scale shotgun metagenomic techniques
43 (Quince et al. 2017). This has yielded important insights into broad functional patterns of
44 microbial consortia (Yooseph et al. 2007; Human Microbiome Project Consortium 2012;
45 Sunagawa et al. 2015). However, while overall pathway abundances inferred from metagenomic
46 data can tell us much about the general functions of communities and how they change with e.g.
47 environmental gradients (Bengtsson-Palme 2018; Bahram et al. 2018), there are many important
48 functional differences that are hidden in the subtleties of these communities (Österlund et al.
49 2017). For example, many antibiotic resistance phenotypes are the results of single point
50 mutations rather than acquisition of novel pathways or genes (Johnning et al. 2013). This
51 complicates the studies of selection pressures in environmental communities, as analysis of such
52 mutations is generally limited to a narrow range of species (Johnning et al. 2015b; Johnning et al.
53 2015a; Kraupner et al. 2018).

54 Because of the immense increase in available sequence data, it would be desirable to study these
55 mutations from shotgun metagenomic libraries, much as other traits have been studied at a large
56 scale (Pal et al. 2016). However, attempts to quantify point mutations in metagenomic sequencing
57 data often go wrong because the methods do not sufficiently well distinguish between mutated
58 and wildtype variants of the same gene. For example, a sequenced read may map to a region
59 identical in the mutated and wildtype variant of a gene, causing problems for quantifying their
60 relative proportions (Bengtsson-Palme et al. 2017). In addition, because the sought-after
61 mutations generally are rare in most types of sample, and metagenomic studies are often under-

62 sampled in terms of replicates (Jonsson et al. 2016a), commonly applied statistics methods may
63 not be sufficiently sensitive to reliably detect differences between samples (Jonsson et al. 2016b).
64 In this study, we attempt to provide a partial remedy to these problems through the introduction
65 of a software tool – Mumame – that can quantify and distinguish between wildtype and mutated
66 gene variants in metagenomic data, and through suggesting a statistical framework for handling
67 the output data of the software. We further demonstrate the ability of the method to detect
68 relevant differences between environmental sample types, estimate the sequencing depths
69 required for the method to perform reliably through simulations, and exemplify the utility of the
70 software on detecting resistance mutations in publicly available metagenomes. The Mumame
71 software package is open-source and freely available from
72 <http://microbiology.se/software/mumame>

73 **Methods**

74 *Software implementation*

75 Mumame is implemented in Perl and consists of two commands: mumame, which performs
76 mapping to database of mutations, and mumame_build which builds the database for the former
77 command. The mumame_build command takes a FASTA sequence file and a list of mutations
78 (CSV format) as input. For each entry in the mutation list, it finds the corresponding sequence(s)
79 in the FASTA file, either by sequence identifier or by CARD ARO accessions (Jia et al. 2016). It
80 then excerpts a number of residues upstream and downstream of the mutation position (by
81 default 20 residues for proteins and 55 for nucleotide sequences) and creates one wildtype
82 version and one mutated version of the sequence excerpt with unique sequence IDs. For cases

83 where multiple mutations can occur close to each other on the same sequence, the software
84 attempts to create all possible combinations of mutations (if memory permits – in some situations
85 this is not possible because the number of combinations increase exponentially). The software
86 tool also generates a mapping file between sequence IDs in the database and mutation
87 information from the list.

88 The main `mumame` command takes any number of input files containing DNA sequence reads
89 in FASTA or FASTQ format and maps those against the Mumame database using Usearch
90 (Edgar 2010). For this mapping, the software runs Usearch in `search_global` mode with target
91 coverage set to 0.55 (by default; any value ≥ 0.51 should be feasible for target coverage). The
92 output is then mapped to the wildtype or mutation information in the Mumame database, and
93 data is collected for each input file and combined into one single output table.

94 The output table generated by Mumame can then be analyzed using the R script (R Core Team
95 2016) supplied with the Mumame package. The script reads the read counts for all mutation
96 positions detected, both for wildtype and mutated sequences, and assesses if there are
97 significantly different proportions of mutations between different sample groups directly through
98 a generalized linear model. Alternatively, an overdispersed Poisson generalized linear model
99 accounting for the discrete nature of the data and the differences in sequencing depth can be
100 used (Jonsson et al. 2016b; Bengtsson-Palme et al. 2017). The Poisson model is preferable when
101 the number of counts for a targeted gene is low in all sample groups.

102 *Quantification of mutations in metagenomes*

103 To quantify the abundances of fluoroquinolone resistance mutations in the *gyrA* and *parC* genes
104 (Johnning et al. 2015b), we downloaded the CARD database on 2018-05-24 (Jia et al. 2016). We
5

105 extracted all mutation information regarding the *gyrA* and *parC* genes from the “snps.txt” file and
106 created a new file with that information. We then created a new Mumame database, with the
107 following command: “mumame_build -i card-data/protein_fasta_protein_variant_model.fasta -m
108 gyrA_parC_snps.txt -o gyrA_parC”. That database was used to map all the reads from the
109 samples generated by Kraupner et al. (2018) to the database using Mumame in the Usearch mode
110 (Edgar 2010) and the following options “-d gyrA_parC -c 0.95”. We did this both for the
111 shotgun metagenomics data as well as for the amplicon sequences derived specifically from
112 Enterobacteriaceae *gyrA* and *parC* genes. Prior to this sequence mapping raw reads were quality
113 filtered using Trim Galore! (Babraham Bioinformatics 2012) with the settings “-e 0.1 -q 28 -O 1”.
114 We then used the R script (R Core Team 2016) provided with the Mumame software to compare
115 the matches to mutated and wildtype sequences in the database. The same database and method
116 combination was used to quantify fluoroquinolone resistance mutations in sequence data from an
117 Indian lake exposed to ciprofloxacin pollution (Bengtsson-Palme et al. 2014), as well as in an
118 Indian river upstream and downstream of a wastewater treatment plant processing
119 pharmaceutical waste (Kristiansson et al. 2011; Pal et al. 2016). These samples were preprocessed
120 in the same way as in the Indian lake study (Bengtsson-Palme et al. 2014).

121 To quantify resistance mutations to tetracycline in the sequence data generated by (Lundström et
122 al. 2016), we created a Mumame database for tetracycline resistance mutations in the 16S rRNA
123 gene. We extracted the mutational information related to tetracycline from the CARD “snps.txt”
124 file and then built the database using the following command: “mumame_build -i card-
125 data/nucleotide_fasta_rRNA_gene_variant_model.fasta -m Tet_snps.txt -o Tet -n”. We then
126 mapped all reads from the Lundström et al. (2016) data to the Mumame database using the

127 options “-d Tet -c 0.95 -n”. Reads were quality filtered and statistical differences were assessed as
128 above.

129 *Software evaluation*


130 To assess the limitations of the method in terms of sequencing depth, the samples from the
131 highest and lowest ciprofloxacin concentrations generated by Kraupner et al. (2018; 10 µg/L and
132 0 µg/L, respectively) were downsampled to 1, 5, 10, 20, 30, 40 and 50 million reads. Thereafter,
133 the reads from the downsampled libraries were mapped to the fluoroquinolone resistance
134 mutation database using Mumame as above. Statistical differences were assessed at all simulated
135 sequencing depths and average effect sizes calculated for the significantly altered genes.

136 **Results**

137 *Mumame can quantify point mutation frequencies in metagenomic data*

138 As a proof-of-concept that our method to identify point mutations in metagenomic sequence
139 data is functional, we used Mumame to quantify the mutations in amplicon data from the the
140 *gyrA* and *parC* genes. These genes are targets of fluoroquinolone antibiotics, and often acquire
141 resistance mutations attaining high levels of resistance. We quantified such mutations in an
142 amplicon data set specifically targeting these two genes in *Escherichia coli*. This data set derives
143 from an exposure study with increasing ciprofloxacin concentrations, and enrichments of
144 mutations in the classical fluoroquinolone resistance determining positions S83 and D87 (*gyrA*)
145 and S80 and E84 (*parC*) have previously been verified using other bioinformatic methods
146 (Kraupner et al. 2018). This data set therefore serves as an ideal positive control for our novel
147 method. We found that Mumame were able to identify the difference between the highest

148 concentration (10 µg/L) and the lower ones reported in the original study (Figure 1). However,
149 Mumame only reported an average frequency of mutations of around 11-12% for *gyrA* mutations
150 (Figure 1A), while the original paper finds frequencies of 60-85% (S83) and 30-40% (D87). The
151 A67 position was not quantified in the original paper. The reason for the discrepancies is
152 unknown, but it is likely caused by a taxonomic filtration step that selects for *E. coli* reads used in
153 the Kraupner et al. study, while Mumame does not perform prior filtering. The decision to
154 exclude filtering was made in order to mimic a situation with true metagenomic data where
155 several target species may co-exist. For *parC*, Mumame only quantified the S80 position (Figure
156 1C), because the E84 mutations were not included in the version of the CARD database used for
157 this study. For position S80, Mumame identified around 35% mutated sequences at the highest
158 concentration of ciprofloxacin, while the original study reported around 50%.



159

160 **Figure 1.** Total mutation frequencies quantified using Mumame for three known mutations conferring
 161 resistance to fluoroquinolone in the *E. coli gyrA* gene based on amplicon sequencing (A) and shotgun
 162 metagenomic data (B) from the same samples. Corresponding data for the S80 mutation in *parC* is shown
 163 in (C) for amplicon data and (D) for shotgun data.

164


165 We next evaluated the performance of Mumame on the real shotgun data that was also generated
 166 from the same samples as the amplicon libraries. Ideally, this analysis should generate virtually the
 167 same result as the amplicon analysis. Indeed, we found similar results for the A67 and S83 *gyrA*
 168 mutations (Figure 1B). For the D87 mutation, the frequencies were much lower than for the
 169 other two mutations, albeit still significantly larger than at the lower concentrations ($p < 0.01$).

9

170 For the *parC* gene, the shotgun metagenomic analysis was too noisy to generate a statistically
171 significant result, which was highly surprising to us (Figure 1D). Taken together, these results
172 indicate the high noise levels present for individual gene variants even in deeply sequenced
173 shotgun metagenomes from controlled exposure studies.

174 *The limits to quantification*

175 Noting the much more instable levels of mutations in the shotgun metagenomes, we next
176 investigated the effects of sequencing depth on the ability of our method to detect significantly
177 altered mutation frequencies. For this analysis, we used downsampled data from the shotgun
178 metagenomic library of the ciprofloxacin exposure study (Figure 2). As expected, we found that
179 the number of significantly altered mutation frequencies detected increased with larger
180 sequencing depth (Figure 2A). In addition, the average effect size of the significant mutations
181 became gradually lower with larger sequence depth, also in accordance with expectations (Figure
182 2B). Importantly, the average effect size of detectable mutation frequency differences seems to
183 decrease linearly with sequencing depth. This means that we can calculate an expected detection
184 limit for the method given the characteristics of the data and experimental setup. At 10 million
185 reads, we expect that the proportion of reads with mutation must be 30-40% higher in the
186 exposed sample in order for it to be detected as significant. The required effect decreases to, on
187 average, 10% higher at 50 million reads (Figure 2B). These numbers are of course also dependent
188 on other factors, such as the number of replicates per treatment, but nevertheless they can be
189 used as ballpark numbers to aid the design of metagenomic studies or to interpret non-significant
190 results derived from Mumame analyses.




191

192 **Figure 2.** Relationship between the number of investigated reads and number of mutations with
193 significantly altered frequencies (A) and the average effect size for those mutations (B); as assessed using
194 Mumame on shotgun metagenomic data from a ciprofloxacin exposure experiment.

195

196 *Tetracycline-exposed Escherichia coli populations do not harbor higher abundances of resistance mutations*

197 After performing the validation and limitation testing of the method, we next used Mumame to
198 quantify resistance mutations in a similar controlled aquarium setup under exposure to the
199 antibiotic tetracycline (Lundström et al. 2016). In this study, no amplicon sequencing of the target
200 gene for tetracycline – the 23S rRNA – was performed, and thus there was no *a priori* true result
201 that we could compare to. While Mumame was able to successfully detect tetracycline resistance
202 mutations in the data, we somewhat surprisingly found no enrichment of tetracycline resistance
203 mutations in this data (Figure 3). Notably, this result was obtained despite a very high sequencing
204 depth (on average 181,595,072 paired-end sequences per library). Obtaining a negative result at
205 this sequencing depth suggests that there actually is no enrichment of known *E. coli* resistance
206 mutations in the samples.




207

208 **Figure 3.** Frequencies of *E. coli* tetracycline resistance mutations at exposure to different concentrations
209 of tetracycline, based on shotgun metagenomic data.

210

211 *Fluoroquinolone resistance mutations in ciprofloxacin-polluted sediments*


212 As a final investigation of the performance of the method, we also let Mumame quantify the
213 fluoroquinolone resistance mutations in river and lake sediments polluted by antibiotic
214 manufacturing waste, primarily ciprofloxacin (Kristiansson et al. 2011; Bengtsson-Palme et al.
215 2014; Pal et al. 2016). These libraries are fairly old and were not as deeply sequenced as the other
216 data sets we investigated. While the experimental setup of these studies in terms of number of
217 samples does not allow for proper statistical testing, we did find an enrichment of the
218 fluoroquinolone resistance mutation frequencies downstream of the pollution source, at least for
219 the *E. coli gyrA* and *parC* genes (Figure 4). We also detected a few such mutations in other species,
220 but the counts of those were low and the results largely non-informative due to the small number
221 of detections per mutation (Figure 5).



222

223 **Figure 4.** Relative frequency of *gyrA* and *parC* sequences with resistance mutations in samples taken
224 downstream, at or upstream of the pharmaceutical production wastewater treatment plant, as well as in a
225 lake polluted by dumping of pharmaceutical production waste. The numbers at the top of the bars shows
226 the total number of sequences (wildtype or mutated) identified in each sample.

227



228

229 **Figure 5.** Relative frequency of sequences with resistance mutations in samples taken downstream, at or
 230 upstream of the pharmaceutical production wastewater treatment plant, as well as in a lake polluted by
 231 dumping of pharmaceutical production waste, for *Acinetobacter baumannii* (A), *Salmonella* species (B) and
 232 *Haemophilus parainfluenzae*, *Mycobacterium tuberculosis* and *Neisseria gonorrhoeae* (C).

233

234 Discussion

235 Metagenomics often becomes restricted to investigate gross compositional changes to the
 236 taxonomy and function of microbial communities. Unfortunately, this obscures important
 237 variation between individual sequence variants that may have large outcomes on phenotypes

238 (Österlund et al. 2017; Bengtsson-Palme 2018). One example of such point mutations inducing
239 strong phenotypic changes is resistance mutations in the target genes of antibiotics (Kraupner et
240 al. 2018). However, including mutated sequence variants in the antibiotic resistance gene
241 databases is complicated, and can lead to gross misinterpretations of the data (see for example
242 (Ma et al. 2014). Still, understanding relevant variation between sequences and linking that to
243 phenotypes is somewhat of a holy Grail of metagenomics. This study has made clear that we are
244 not yet at that point in terms of bioinformatic methods and the sequencing depth required to
245 draw firm conclusions. That said, we show in this work that identifying significant and relevant
246 differences in resistance mutation frequencies between sample groups from shotgun
247 metagenomic data is possible, given a sufficiently large sequence depth. However, the
248 quantitative estimates still seem to be highly variable, even at very large sequencing depths.

249 The results of the Mumame evaluation also provides a few other important clues on potential
250 pitfalls with inferring mutation frequencies from shotgun metagenomic data. An important such
251 aspect is the disparity between mutation frequencies described by amplicon sequencing and
252 shotgun data. Particularly, the ability to relatively consistently identify the A67 and S83 mutations
253 in *parC*, while the D87 mutation is seemingly less frequent in the shotgun data is somewhat
254 troubling if the goal is to identify the actual abundances of such mutations. At the same time, the
255 statistical significance of those differences could still be identified. For the A67 and S83
256 mutations, only 5 million reads were required for a significant effect to be detected, while for the
257 D87 mutations a sequencing depth of 50 million reads was required. This is not necessarily a
258 shortcoming of the Mumame software, but may just as well be due to the much noisier nature of

259 counts from metagenomic sequence data compared to the large number of reads corresponding
260 to the same genes deriving from amplicon data (Jonsson et al. 2016a).

261 Another important potential problem highlighted by our evaluation is the need to produce very
262 large sequence data sets to be able to identify and quantify mutations (and wildtype) sequences
263 with any certainty. As a rule of thumb, the targeted regions represent less than 0.004% of the
264 bacterial genome, and each bacterial strain may correspond to only a fraction of a percent of the
265 reads in the shotgun sequence data (depending on its abundance). This means that to identify a
266 single read from a resistance region in the data, one would – on average – need to sequence more
267 than five million reads. To get a reasonably confident measure of reads stemming from wildtype
268 versus strains with mutations, approximately 10 reads from each group would be needed per
269 sample (or, say, 20 reads in total). That would, as a rough estimate, correspond to a hundred
270 million reads per sample. This is, unfortunately, way more sequences than what is typically
271 generated per sample by shotgun metagenomic sequencing projects. In this study, only the
272 samples from the tetracycline exposure study corresponded to such a high sequencing depth.
273 Naturally, these numbers would depend on the proportions of the targeted microorganisms as
274 well as their genome sizes, but ultimately this still presents the largest limitation to mutation
275 studies based on metagenomic sequence data. Potentially, this problem could be partially
276 alleviated by analyzing sufficiently large cohorts and perform the statistical analysis for general
277 trends, but even large cohorts would be insufficient for mutations rare enough to pass below the
278 detection limit.

279 In terms of interpreting the results from the exposure experiments, it is interesting to note the
280 overall clear increase of fluoroquinolone resistance mutations at the highest ciprofloxacin

281 concentration, which nearly perfectly correspond to increases in mobile *qnr* fluoroquinolone
282 genes in the same samples (Kraupner et al. 2018). This is contrasted by the trend seen in the
283 tetracycline exposure experiments, where tetracycline resistance genes – specifically efflux pumps
284 – were enriched at higher tetracycline concentrations (Lundström et al. 2016), while tetracycline
285 resistance mutation abundances were not significantly altered. This non-significant result was
286 obtained despite the exceptionally high sequencing depth of those samples.

287 While we did not have data from a proper experimental setup to address differences between
288 sediments exposed to different degrees of fluoroquinolone pollution, the quantification of
289 resistance mutations seems to provide an important piece of information to explain the results of
290 previous studies of resistance gene abundances in these river samples (Kristiansson et al. 2011).
291 In the original paper, the abundance of mobile fluoroquinolone resistance genes (*qnr* genes) were
292 shown to be enriched in the low-level polluted upstream samples, compared to the highly
293 polluted downstream samples. Importantly, the *qnr* genes only provide resistance to relatively low
294 levels of fluoroquinolones (Hooper and Jacoby 2015), and the authors of hypothesize that
295 chromosomal mutations of the target genes are probably necessary to survive the selection
296 pressure from antibiotics downstream of the pollution source. In this work, we show that this
297 assumption is likely correct. Only a limited number of reads were mapping to these resistance
298 regions and the number of samples unfortunately prevents us from properly assessing a statistical
299 difference between the upstream and downstream samples. Still, the proportion of resistance
300 mutations seems to be systematically higher in the samples downstream of the pollution source,
301 at least for *E. coli*. This indicates that the method we present here can provide important

302 additional information to metagenomic studies of resistance patterns in different environment
303 types, given that a sufficient sequencing depth is achieved.

304 We have here shown the utility of the Mumame tool for finding resistance mutations in shotgun
305 metagenomic data. In this paper, we have used the CARD database (Jia et al. 2016) as the
306 information source for resistance mutation, but the tool is flexible to use any source of such data.
307 It is also not in any means restricted to the mutations investigated in this paper but is
308 fundamentally agnostic to the input data. It can also be used in open screening for mutations in
309 any gene present in the database in parallel, and can handle different mutations in both RNA and
310 protein coding genes. The tool is flexible and fast and can therefore be implemented as a part
311 nearly any screening pipeline for antibiotic resistance data in metagenomic data sets.

312 **Conclusion**

313 This paper presents a software tool called Mumame to analyze shotgun metagenomic data for
314 point mutations, such as those conferring antibiotic resistance to bacteria. Mumame distinguish
315 between wildtype and mutated gene variants in metagenomic data and quantify them, given a
316 sufficient sequencing depth. We also provide a statistical framework for handling the generated
317 count data and account for factors such as differences in sequencing depth. Importantly, our
318 study also reveals the importance of a high sequencing depth – preferably more than 50 million
319 sequenced reads per sample – in order to get reasonably accurate estimates of mutation
320 frequencies, particularly for rare genes or species. The Mumame software package is freely
321 available from <http://microbiology.se/software/mumame>. We expect Mumame to be a useful

322 addition to metagenomic studies of e.g. antibiotic resistance, and to increase the detail by which
323 metagenomes can be screened for phenotypically important differences.

324

325 **Acknowledgements**

326 This work was funded by the Swedish Research Council for Environment, Agricultural Sciences
327 and Spatial Planning (FORMAS; grant 2016-00768).

328 **Author contributions**

329 JBP conceived of the study. SM and JBP collected and analyzed the data. JBP designed and wrote
330 the software package. VJ provided statistical guidance for the R implementation. JBP wrote the
331 draft manuscript. All authors interpreted the data and contributed to the writing of the paper.

332 **Conflict of interest**

333 The authors have no conflicts of interest to declare.

334 **References**

- 335 Babraham Bioinformatics (2012) Trim Galore!
336 https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
- 337 Bahram M, Hildebrand F, Forslund SK, et al (2018) Structure and function of the global topsoil
338 microbiome. *Nature* 320:1039. doi: 10.1038/s41586-018-0386-6

- 339 Bengtsson-Palme J (2018) **Strategies for Taxonomic and Functional Annotation of**
340 **Metagenomes**. In: Nagarajan M (ed) *Metagenomics: Perspectives, Methods, and*
341 *Applications*. Academic Press, Elsevier, Oxford, UK,
- 342 Bengtsson-Palme J, Boulund F, Fick J, et al (2014) Shotgun metagenomics reveals a wide array of
343 antibiotic resistance genes and mobile elements in a polluted lake in India. *Front Microbiol*
344 5:648. doi: 10.3389/fmicb.2014.00648
- 345 Bengtsson-Palme J, Larsson DGJ, Kristiansson E (2017) Using metagenomics to investigate
346 human and environmental resistomes. *Journal of Antimicrobial Chemotherapy* 72:2690–
347 2703. doi: 10.1093/jac/dkx199
- 348 Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
349 26:2460–2461. doi: 10.1093/bioinformatics/btq461
- 350 Hooper DC, Jacoby GA (2015) Mechanisms of drug resistance: quinolone resistance. *Ann N Y*
351 *Acad Sci* 1354:12–31. doi: 10.1111/nyas.12830
- 352 Human Microbiome Project Consortium (2012) Structure, function and diversity of the healthy
353 human microbiome. *Nature* 486:207–214. doi: 10.1038/nature11234
- 354 Jia B, Raphenya AR, Alcock B, et al (2016) CARD 2017: expansion and model-centric curation of
355 the comprehensive antibiotic resistance database. *Nucleic Acids Res* gkw1004. doi:
356 10.1093/nar/gkw1004
- 357 Johnning A, Kristiansson E, Angelin M, et al (2015a) Quinolone resistance mutations in the
358 faecal microbiota of Swedish travellers to India. *BMC Microbiol* 15:235. doi:
359 10.1186/s12866-015-0574-6
- 360 Johnning A, Kristiansson E, Fick J, et al (2015b) Resistance Mutations in *gyrA* and *parC* are
361 Common in *Escherichia* Communities of both Fluoroquinolone-Polluted and
362 Uncontaminated Aquatic Environments. *Front Microbiol* 6:1355. doi:
363 10.3389/fmicb.2015.01355
- 364 Johnning A, Moore ERB, Svensson-Stadler L, et al (2013) Acquired genetic mechanisms of a
365 multiresistant bacterium isolated from a treatment plant receiving wastewater from antibiotic
366 production. *Appl Environ Microbiol* 79:7256–7263. doi: 10.1128/AEM.02141-13
- 367 Jonsson V, Österlund T, Nerman O, Kristiansson E (2016a) Variability in Metagenomic Count
368 Data and Its Influence on the Identification of Differentially Abundant Genes. *Journal of*
369 *Computational Biology* cmb.2016.0180. doi: 10.1089/cmb.2016.0180
- 370 Jonsson V, Österlund T, Nerman O, Kristiansson E (2016b) Statistical evaluation of methods for
371 identification of differentially abundant genes in comparative metagenomics. *BMC*
372 *Genomics* 17:78. doi: 10.1186/s12864-016-2386-y

- 373 Kraupner N, Ebmeyer S, Bengtsson-Palme J, et al (2018) Selective concentration for
374 ciprofloxacin resistance in *Escherichia coli* grown in complex aquatic bacterial biofilms.
375 *Environ Int* 116:255–268. doi: 10.1016/j.envint.2018.04.029
- 376 Kristiansson E, Fick J, Janzon A, et al (2011) Pyrosequencing of antibiotic-contaminated river
377 sediments reveals high levels of resistance and gene transfer elements. *6*:e17038. doi:
378 10.1371/journal.pone.0017038
- 379 Lundström SV, Östman M, Bengtsson-Palme J, et al (2016) Minimal selective concentrations of
380 tetracycline in complex aquatic bacterial biofilms. *Sci Total Environ* 553:587–595. doi:
381 10.1016/j.scitotenv.2016.02.103
- 382 Ma L, Li B, Zhang T (2014) Abundant rifampin resistance genes and significant correlations of
383 antibiotic resistance genes and plasmids in various environments revealed by metagenomic
384 analysis. *Appl Microbiol Biotechnol* 98:5195–5204. doi: 10.1007/s00253-014-5511-3
- 385 Österlund T, Jonsson V, Kristiansson E (2017) HirBin: high-resolution identification of
386 differentially abundant functions in metagenomes. *BMC Genomics* 18:316. doi:
387 10.1186/s12864-017-3686-6
- 388 Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DGJ (2016) The structure and diversity of
389 human, animal and environmental resistomes. *Microbiome* 4:54. doi: 10.1186/s40168-016-
390 0199-5
- 391 Quince C, Walker AW, Simpson JT, et al (2017) Shotgun metagenomics, from sampling to
392 analysis. *Nat Biotechnol* 35:833–844. doi: 10.1038/nbt.3935
- 393 R Core Team (2016) R: A Language and Environment for Statistical Computing. R Foundation
394 for Statistical Computing, Vienna, Austria
- 395 Sunagawa S, Coelho LP, Chaffron S, et al (2015) Ocean plankton. Structure and function of the
396 global ocean microbiome. *Science* 348:1261359. doi: 10.1126/science.1261359
- 397 Yooseph S, Sutton G, Rusch DB, et al (2007) The Sorcerer II Global Ocean Sampling
398 expedition: expanding the universe of protein families. *5*:e16. doi:
399 10.1371/journal.pbio.0050016

400

401

402