1 ***Drosophila melanogaster* infected with *Wolbachia* strain**

2 **wMelCS prefer cooler temperatures**

3

4 Pieter A. Arnold^{*,†,a}, Samantha C. Levin[‡], Aleksej L. Stevanovic, Karyn N. Johnson^b

5

6 School of Biological Sciences, The University of Queensland, Brisbane QLD 4072,

7 Australia

8

9 [†] Present address: Research School of Biology, The Australian National University,

10 Acton ACT

11 2601, Australia

12 [‡] Present address: School of Biomedical Sciences, The University of Queensland,

13 Brisbane QLD 4072, Australia

14 ^a ORCID: 0000-0002-6158-7752

15 ^b ORCID: 0000-0001-8647-8985

16

17 ^{*} Corresponding author:

18 Pieter A. Arnold

19 Email: pieter.arnold@anu.edu.au

20 Phone: +61 2 6125 2543

21 Address: Research School of Biology, 46 Sullivans Creek Road, The

22 Australian National University, Acton ACT 2601, Australia

Abstract

1. Temperature plays a fundamental role in the dynamics of host-pathogen interactions.
2. *Wolbachia* is an endosymbiotic bacteria that infects about 40% of arthropod species, which can affect host behaviour and reproduction. Yet, the effect of *Wolbachia* on host thermoregulatory behaviour is largely unknown, despite its use in disease vector control programs in thermally variable environments.
3. Here, we used a thermal gradient to test whether *Drosophila melanogaster* infected with *Wolbachia* strain wMelCS exhibit different temperature preferences (T_p) to uninfected flies.
4. We found that *Wolbachia*-infected flies preferred a cooler mean temperature ($T_p = 25.06 \pm 0.25^\circ\text{C}$) than uninfected flies ($T_p = 25.78 \pm 0.24^\circ\text{C}$).
5. This finding suggests that *Wolbachia*-infected hosts might seek out cooler microclimates to reduce exposure to and lessen the consequences of high temperatures. This finding has generated hypotheses that will be fruitful in areas of research for exploring the mechanisms by which the change in T_p occurs in this complex and significant host-pathogen-environment interaction.

Keywords: host behaviour; host-pathogen interaction; temperature preference; thermal gradient; vector control; *Wolbachia pipientis*

Running title: *Wolbachia* alters temperature preference

45 Introduction

46 *Wolbachia pipientis* is an endosymbiont bacteria that infects an estimated 40% of
 47 terrestrial arthropod species (Zug & Hammerstein, 2012). The association between
 48 *Wolbachia* and its hosts has been the subject of a wide array of studies, including
 49 the alteration of host behaviours and reproduction (Weeks et al., 2002), cytoplasmic
 50 incompatibility for disease vector control (Clancy & Hoffmann, 1998), and
 51 environmental factors mediating host-pathogen interactions (Murdock et al., 2012).
 52 Temperature is a key environmental modulator of host-pathogen interactions, which
 53 constrains the rate of biological reactions and sets limits to performance and survival
 54 (Thomas & Blanford, 2003).

55 For the insect host, there is little physiological capacity to differentiate their
 56 body temperature (T_b) from the ambient temperature of their surrounding
 57 environment (Angilletta, 2009). Physiological rates and performance are strongly
 58 affected by T_b in ectotherms, so organisms should aim to maintain their T_b across a
 59 range of temperatures that correspond to adequate performance (Huey &
 60 Kingsolver, 1989). One strategy that ectotherms can employ to avoid exposure to
 61 unsuitable temperatures is to modify their behaviour to seek more suitable
 62 microclimates, such as in shade, to find their preferred temperature (T_p) (Sunday et
 63 al., 2014). Temperature preference is the perception and neural integration of
 64 thermal information, resulting in this crucial thermoregulatory behaviour (Abram et
 65 al., 2017). *Drosophila melanogaster* exhibits strong circadian and neurally controlled
 66 temperature preference behaviour, which centres around 24–27°C (Arnold et al.,
 67 2015; Kaneko et al., 2012).

68 The thermal biology of *Wolbachia* and host associations is complex and
 69 variable. High temperatures are unfavourable for some strains, where *Wolbachia*

density is much higher at lower temperatures (e.g., 13–19°C (Moghadam et al., 2017)) and is reduced at higher temperatures (e.g., 26°C (Clancy & Hoffmann, 1998)). *Wolbachia* can also be mostly or completely eliminated by exposure to cyclic heat stress or temperatures above 30°C, reducing vertical transmission of the symbiont (Corbin et al., 2016; Ross et al., 2017). However, higher temperatures do not always reduce *Wolbachia* titre and responses to temperature vary greatly among study systems (e.g., Mouton et al., 2006; Murdock et al., 2014). The *Wolbachia* strain wMelCS that we use here infects natural *D. melanogaster* populations, and at high titres, confers some virus protection but also reduces host lifespan (Chrostek et al., 2013; Hedges et al., 2008).

As temperatures ideally suited to *D. melanogaster* are generally higher than those suited to *Wolbachia*, we predict that manipulating their host's behaviour to seek cooler temperatures would be beneficial. Thus, our objective here is to use an established behavioural assay to test the capacity for *Wolbachia* infection to alter host T_p .

Materials and Methods

Drosophila melanogaster from the Oregon RC line were infected with the wMelCS line of *Wolbachia pipientis* (hereafter +Wol) (Hedges et al., 2008). All flies were reared in 25°C incubators on a standard cornmeal diet and 12 h light/dark cycle. The paired *Wolbachia*-free fly line (hereafter control) was generated from the same wMelCS-infected fly line by treating flies with 0.03% tetracycline. We confirmed that the *Wolbachia* strain was wMelCS by PCR using two primer sets as described in Riegler et al. (2012). These flies were reared on a standard cornmeal diet for at least five generations before use to recover after tetracycline treatment, and microbiota

95 was reconstituted and standardised following standard procedures. Here,
96 temperature preference assays were conducted on adult male flies aged 4–7 days,
97 but both male and female flies of various ages have been shown to exhibit very
98 similar temperature preferences even when infected by *Wolbachia* (Truitt et al.,
99 2018).

100 Temperature preference assays used an identical thermal gradient apparatus
101 to that previously described in (Arnold et al., 2015). The apparatus achieved a stable
102 linear gradient of 0.2°C per cm across a temperature range of 17.5–33.5°C (Arnold
103 et al., 2015). Temperatures were measured throughout the experiment by five K-type
104 thermocouples suspended in the gradient airspace, held by bungs that were fitted
105 into an acrylic cover, recorded by a Squirrel 2040 temperature meter.

106 For each trial, five flies were gently tipped into the centre of the apparatus
107 (25°C) and allowed to freely move about the apparatus for 30 minutes. At the end of
108 the trial period, flies were anaesthetised by CO₂ that was introduced into both ends
109 of the gradient at a low-flow rate to prevent changes to the position of flies. Distance
110 along the gradient was then used to determine the preferred temperature at the
111 position of rest for each fly in the gradient, T_p . A total of 58 flies for each treatment
112 were assessed for T_p , across 24 replicate trials.

113 As circadian rhythm affects T_p in *Drosophila*, we always conducted
114 temperature preference assays between 09:30 and 13:30, a time period across
115 which T_p is stable (Kaneko et al., 2012). Assays were also conducted in darkness by
116 covering the apparatus in black material to prevent phototactic behaviour affecting
117 positioning.

118 Exploratory data analyses where flies in each trial were treated as non-
119 independent in a mixed-effects model (trial as a random variable) found the variance

component of trial to be essentially zero. We therefore assumed that T_p of individual flies were independent and not affected by social interactions during the trial itself (see supplementary material of Arnold et al., 2015). Data were not normally distributed, therefore we conducted a non-parametric Mann-Whitney U test to compare the preferred temperatures of control and +*Wol* flies. We then calculated Cohen's d with 95% confidence intervals (CIs) as a measure of effect size, given that p is not always robust. All analyses were conducted in the R environment for statistical computing (v3.4.1).

Results and discussion

We found that flies infected with *Wolbachia* strain *wMelCS* preferred cooler temperatures compared to those without any *Wolbachia* (Fig. 1). The difference between populations was significant even at $\alpha = 0.01$ (Mann-Whitney U test; $W = 2179$, $p = 0.006$), which was supported by an effect size and 95% CIs that did not overlap with zero (Cohen's $d = 0.394$ [0.189 – 0.563]). Both populations exhibited large variance in T_p , ranging between 18.5 and 29°C (Fig. 1A). There is some overlap in T_p distributions between the populations (Fig. 1B). In absolute terms, +*Wol* flies preferred a cooler mean ($\pm$ SE) temperature of $25.06 \pm 0.25^\circ\text{C}$ compared to the control flies, which preferred $25.78 \pm 0.24^\circ\text{C}$.

Our finding is one of the first empirical accounts of *Wolbachia*-infected flies exhibiting a preference for cooler temperatures. Recently, Truitt et al. (2018) demonstrated that three strains of *Wolbachia* (*wMel*, *wMelCS*, and *wMelPop*) all significantly reduce the T_p of *D. melanogaster* (strain w^{1118}). The authors found that w^{1118} flies infected with *wMelCS* had a 4°C lower mean T_p compared to uninfected flies, although the T_p of uninfected w^{1118} flies was 24.4°C, which is 1.4°C lower than

the Oregon RC flies from the present study. We have identified a similar T_p shift of *Wolbachia*-infected flies in the same direction (although lower magnitude) in a different host strain, using a different temperature preference apparatus. Our results not only support the findings of Truitt et al. (2018), but also suggest that this biologically-interesting phenomenon could be reasonably common.

One explanation for our findings is that *Wolbachia* could potentially manipulate an important aspect of host thermoregulatory behaviour. Pathogens can improve their transmission probability and reproductive capacity by inducing host behavioural changes (Lefèvre & Thomas, 2008), and *Wolbachia* can affect host mate choice and activity levels (van Houte et al., 2013). Changes in thermoregulation behaviour has been well studied from the perspective of the infected host, particularly behavioural fever, where the host elevates its T_b by behavioural means to rid itself of the pathogen (Kluger, 1979). However, it is less clear whether pathogens manipulate host T_p , especially when the pathogen is not parasitic (i.e., commensalistic or mutualistic) and for decreases to T_p .

Variance of temperatures in nature may lead to populations with mixed or incomplete *Wolbachia* infection (van Opijnen & Breeuwer, 1999), but it is possible that *Wolbachia* could manipulate host T_p to maximise its own fitness without negatively affecting the host. Cyclic heat stress fluctuating between 26°C and 37°C at 12 h intervals significantly reduced *Wolbachia* density and cytoplasmic incompatibility of wMel and wMelPop-CLA, but not wAlbB in *Aedes aegypti* (Ross et al., 2017). The wMelCS strain used in the present study likely shares a recent field origin with wMel (Riegler et al., 2005), which is a widely used dengue-suppressing *Wolbachia* strain that itself also alters host T_p to prefer cooler temperatures (Truitt et al., 2018). Temperature fluctuations like the cyclic heat stress experiment by Ross et

al. (2017) might well be experienced naturally in tropical regions. This would likely result in incomplete infection and reduced transmission success, which could explain the erratic temporal and spatial dynamics of *Wolbachia* spread in controlled infected-vector release programs (e.g., Schmidt et al., 2017). Our finding suggests that *Wolbachia*-infected hosts prefer cooler temperatures and might be likely to seek out cooler microclimates, which would reduce exposure to and lessen the fitness consequences of high temperatures.

An alternative hypothesis is that flies infected with *Wolbachia* exhibit a behavioural chill to restrict pathogen replication. Infection with *Pseudomonas aeruginosa* lowered the T_p of *D. melanogaster*, but this behavioural chill response was inefficient and did not reach the critical temperature that would increase survival and limit bacteria growth rate (Fedorka et al., 2016). The change T_p that we observed in flies infected with wMelCS could also be a host behavioural response. However, as discussed earlier, *Wolbachia* tends to perform worse at higher temperatures than are optimal for *D. melanogaster*, and therefore we suggest that manipulation of the host T_p by the pathogen is more likely.

The absolute decrease in T_p of less than 1°C that we observed in flies infected with *Wolbachia* wMelCS provides little buffer to the predicted 2–4°C increase by 2100 due to climate change. However, *Wolbachia* are maternally inherited and exposure to high temperatures can reduce vertical transmission in only a few generations (Corbin et al., 2016). If the infected host prefers cooler temperatures, then this behaviour would confer a selective advantage for *Wolbachia*. Arthropod hosts of *Wolbachia* have rapid generation times relative to the forecast rate of temperature increase, therefore it is conceivable that a minor change in T_p could be

enhanced by selection across generations to allow continued transmission and mitigate fitness consequences.

This study paves the way for discovering the mechanisms by which *Wolbachia* - infection alters host T_p . Whether the observed phenomenon is due to *Wolbachia* directly manipulating host behaviour, a host defense response, or a by-product of infection will need to be determined. The efficacy of introductions of populations of *Wolbachia*-infected vectors may hinge upon a better understanding of complex host-pathogen-environment interactions. Testing for *Wolbachia*-induced changes in thermal preference across multiple host and pathogen strains will elucidate whether unexpected ecological and evolutionary responses might occur in planned vector releases in a changing climate.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Competing interests

The authors have no competing interests to declare.

Acknowledgements

We thank Prof. Craig R. White for providing laboratory space and equipment for this study, and constructive comments from three anonymous reviewers that improved this manuscript.

218 **Contribution of authors**

219 PAA established the methodology, lead the analyses, interpreted results, and wrote
 220 the manuscript; SL and ALS collected the data and contributed to the manuscript
 221 draft; KNJ supervised the project, interpreted results, and contributed substantially to
 222 the manuscript draft.

References

- Abram, P.K., Boivin, G., Moiroux, J., & Brodeur, J. (2017) Behavioural effects of temperature on ectothermic animals: unifying thermal physiology and behavioural plasticity. *Biological Reviews*, **92**, 1859-1876.
- Angilletta, M.J. (2009) Thermal adaptation: a theoretical and empirical synthesis Oxford University Press, New York, U.S.A.
- Arnold, P.A., White, C.R., & Johnson, K.N. (2015) *Drosophila melanogaster* does not exhibit a behavioural fever response when infected with *Drosophila C* virus. *Journal of General Virology*, **96**, 3667-3671.
- Chrostek, E., Marialva, M.S.P., Esteves, S.S., Weinert, L.A., Martinez, J., Jiggins, F.M., & Teixeira, L. (2013) *Wolbachia* variants induce differential protection to viruses in *Drosophila melanogaster*: a phenotypic and phylogenomic analysis. *PLoS Genetics*, **9**, e1003896.
- Clancy, D.J. & Hoffmann, A.A. (1998) Environmental effects on cytoplasmic incompatibility and bacterial load in *Wolbachia*-infected *Drosophila simulans*. *Entomologia Experimentalis et Applicata*, **86**, 13-24.
- Corbin, C., Heyworth, E.R., Ferrari, J., & Hurst, G.D.D. (2016) Heritable symbionts in a world of varying temperature. *Heredity*, **118**, 10-20.
- Fedorka, K.M., Kutch, I.C., Collins, L., & Musto, E. (2016) Cold temperature preference in bacterially infected *Drosophila melanogaster* improves survival but is remarkably suboptimal. *Journal of Insect Physiology*, **93-94**, 36-41.

243 Hedges, L.M., Brownlie, J.C., O'Neill, S.L., & Johnson, K.N. (2008) *Wolbachia* and virus
244 protection in insects. *Science*, **322**, 702.

245 Huey, R.B. & Kingsolver, J.G. (1989) Evolution of thermal sensitivity of ectotherm
246 performance. *Trends in Ecology & Evolution*, **4**, 131-135.

247 Kaneko, H., Head, L.M., Ling, J., Tang, X., Liu, Y., Hardin, P.E., Emery, P., & Hamada, F.N.
248 (2012) Circadian rhythm of temperature preference and its neural control in *Drosophila*.
249 *Current Biology*, **22**, 1851-1857.

250 Kluger, M.J. (1979) Fever in ectotherms: evolutionary implications. *American Zoologist*, **19**,
251 295-304.

252 Lefèvre, T. & Thomas, F. (2008) Behind the scene, something else is pulling the strings:
253 emphasizing parasitic manipulation in vector-borne diseases. *Infection, Genetics and*
254 *Evolution*, **8**, 504-519.

255 Moghadam, N.N., Thorshauge, P.M., Kristensen, T.N., de Jonge, N., Bahrndorff, S., Kjeldal,
256 H., & Nielsen, J.L. (2017) Strong responses of *Drosophila melanogaster* microbiota to
257 developmental temperature. *Fly*, **12**, 1-12.

258 Mouton, L., Henri, H., Bouletreau, M., & Vavre, F. (2006) Effect of temperature on
259 *Wolbachia* density and impact on cytoplasmic incompatibility. *Parasitology*, **132**, 49-56.

260 Murdock, C.C., Blanford, S., Hughes, G.L., Rasgon, J.L., & Thomas, M.B. (2014) Temperature
261 alters *Plasmodium* blocking by *Wolbachia*. *Scientific Reports*, **4**, 3932.

262 Murdock, C.C., Paaijmans, K.P., Cox-Foster, D., Read, A.F., & Thomas, M.B. (2012) Rethinking
263 vector immunology: the role of environmental temperature in shaping resistance. *Nature*
264 *Reviews Microbiology*, **10**, 869-876.

265 Riegler, M., Iturbe-Ormaetxe, I., Woolfit, M., Miller, W.J., & O'Neill, S.L. (2012) Tandem
266 repeat markers as novel diagnostic tools for high resolution fingerprinting of *Wolbachia*.
267 *BMC Microbiology*, **12**, S12.

268 Riegler, M., Sidhu, M., Miller, W.J., & O'Neill, S.L. (2005) Evidence for a global *Wolbachia*
269 replacement in *Drosophila melanogaster*. *Current Biology*, **15**, 1428-1433.

270 Ross, P.A., Wiwatanaratnabutr, I., Axford, J.K., White, V.L., Endersby-Harshman, N.M., &
271 Hoffmann, A.A. (2017) *Wolbachia* infections in *Aedes aegypti* differ markedly in their
272 response to cyclical heat stress. *PLoS Pathogens*, **13**, e1006006.

273 Schmidt, T.L., Barton, N.H., Rašić, G., Turley, A.P., Montgomery, B.L., Iturbe-Ormaetxe, I.,
274 Cook, P.E., Ryan, P.A., Ritchie, S.A., Hoffmann, A.A., O'Neill, S.L., & Turelli, M. (2017) Local
275 introduction and heterogeneous spatial spread of dengue-suppressing *Wolbachia* through
276 an urban population of *Aedes aegypti*. *PLoS Biology*, **15**, e2001894.

277 Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T., & Huey, R.B.
278 (2014) Thermal-safety margins and the necessity of thermoregulatory behavior across
279 latitude and elevation. *Proceedings of the National Academy of Sciences*, **111**, 5610-5615.

280 Thomas, M.B. & Blanford, S. (2003) Thermal biology in insect-parasite interactions. *Trends in*
281 *Ecology & Evolution*, **18**, 344-350.

282 Truitt, A.M., Kapun, M., Kaur, R., & Miller, W.J. (2018) *Wolbachia* modifies thermal
283 preference in *Drosophila melanogaster*. *Environmental Microbiology*, **10.1111/1462-**
284 **2920.14347**.

285 van Houte, S., Ros, V.I.D., & van Oers, M.M. (2013) Walking with insects: molecular
286 mechanisms behind parasitic manipulation of host behaviour. *Molecular Ecology*, **22**, 3458-
287 3475.

288 van Opijnen, T. & Breeuwer, J.A.J. (1999) High temperatures eliminate *Wolbachia*, a
289 cytoplasmic incompatibility inducing endosymbiont, from the two-spotted spider mite.
290 *Experimental and Applied Acarology*, **23**, 871-881.

291 Weeks, A.R., Reynolds, K.T., & Hoffmann, A.A. (2002) *Wolbachia* dynamics and host effects:
292 what has (and has not) been demonstrated? *Trends in Ecology & Evolution*, **17**, 257-262.

293 Zug, R. & Hammerstein, P. (2012) Still a host of hosts for *Wolbachia*: analysis of recent data
294 suggests that 40% of terrestrial Arthropod species are infected. *PLoS One*, **7**, e38544.

Figure legend

Figure 1. Preferred temperature of uninfected control (orange fill, solid line) and +*Wol* (blue fill, dashed line) *Drosophila melanogaster*. (A) Boxplot of preferred temperature showing median (thick black line), upper and lower quartiles (box), 95% confidence intervals (error bars), and raw data points for control and +*Wol* flies. Each population had $n = 58$ individuals. (B) Density plot showing smoothed distributions of the preferred temperature data for each population.