Assessment of menstrual hygiene management and its determinants among adolescent girls: A cross-sectional study in school adolescent girls in Addis Ababa, Ethiopia.

Ephrem Biruk^{1¶}, Worku Tefera^{2¶}, Nardos Tadesse^{3¶}, Ashagre Sisay^{4¶}

¹Strengthen Ethiopian Urban Health Program, Quality improvement specialist, Addis Ababa, Ethiopia/ SEUHP, Southern Program, Implemented by JSI, Addis Ababa, Ethiopia

²Department of Public Health, Addis Ababa University, Addis Ababa, Ethiopia

³Control of communicable disease prevention SURGE department, HTS coordinator Implemented by Addis Ababa health bureau, Addis Ababa, Ethiopia

⁴Department of urban health extension program, Arada sub city health office, Addis Ababa health bureau Addis Ababa, Ethiopia.

* Corresponding author

E-mail:ephrembiruk@gmail.com (EB)

*Ephrem Biruk

¶These authors contributed equally to this work.

Abstract

Introduction: Managing menstruation is essentially dealing with menstrual flow and also in continuing regular activities like going to school, working etc. However, menstruation can place significant obstacles in girls' access to health, education and future prospects if they are not equipped for effective menstrual hygiene management.

Objective: To assess the menstrual hygiene management and its determinant among school girls in Addis Ababa, Ethiopia.

Methods: Cross-sectional study design with quantitative method was carried out among 770 systematically selected adolescent school girls of Addis Ababa from April 1 to May 5, 2017. A self-administered pre-test close ended Amharic questionnaire at school setting was used for data collection. The coding was done using the original English version and entered to EPI-7 software. The quantitative file exported to statistical package for social science (SPSS) version 25.0 software for analysis. Total mean score was used to categorize individuals as good and poor while AOR; 95% CI with $p < 0.05$ was used to determine factors of menstrual hygiene management practice.

Result: This study had 98% response rate. 530 (70.1%) and 388(51.3%) respondents had good knowledge and practice of menstrual hygiene respectively. The findings also showed a significant positive association between good knowledge of menstruation and girls from mother's whose education were secondary (AOR = 10.012, 95 % CI = 3.628-27.629). Wealth index quantile five (AOR = 9.038, 95 % CI = 3.728-21.909) revealed significant positive association with good practice of menstrual hygiene.

Conclusion and recommendation

Majority of participants had good knowledge and practice of menstrual hygiene and majority of them were from private school. Although knowledge was better than practice, girls should be educated about the process, use of proper pads or absorbents and its proper disposal.

Key Words: practices of menstrual hygiene, Menstrual knowledge, adolescent girl, Sanitary napkins, Menarche, school health.

Introduction

Background

Menarche is an important milestone in a girl's transition to womanhood (1). Around the world women have developed their own personal strategies to cope with menstruation, which vary from country to country and depend on economic status, the individual's personal preferences, local traditions and cultural beliefs and education status. The onset of menstruation presents multiple

challenges for school girls. Many girls lack the knowledge, support and resources to manage menstruation in school (2).

Managing menstruation is essentially dealing with menstrual flow and also in continuing regular activities like going to school, working etc (3). However, menstruation can place significant obstacles in the way of girls' access to health, education and future prospects if they are not equipped for effective menstrual hygiene management (MHM). Good MHM requires access to necessary resources (menstrual materials to absorb or collect menstrual blood, soap and water), facilities (private place to wash, change and dry re-usable menstrual materials, in addition to an adequate disposal system for menstrual materials), and education about MHM (1).

Schools, particularly those in developing countries, often completely lack drinking-water and sanitation and hand washing facilities; even, where such facilities exist they are often inadequate in both quality and quantity. Girls are likely to be affected in different ways from inadequate water, sanitation and hygiene conditions in schools, because the lack of such facilities they cannot attend school during menstruation (4). Girls are particularly vulnerable to dropping out of school, partly because when toilet and washing facilities are not private, not safe or simply not available in schools. Girls who reached puberty and female school staff need gender-related privacy; otherwise they may not use the facilities. This may result in absenteeism rates that can reach 10–20 per cent of school time (5).

Menstrual hygiene and management has not received adequate attention in the health and water, sanitation and hygiene (WASH) sectors in developing countries including Ethiopia and its relationship with and impact on achieving agenda of the sustainable development goal that access to adequate and equitable sanitation and hygiene for all and end open defecation, paying special attention to the needs of women and girls and those in Vulnerable situations (6).

Public investment in institutional sanitation, especially in schools and health facilities in urban areas is limited. The available sanitation facilities in most secondary schools, are poor in construction design; not convenient for sick, disabled, elderly and MHM. This in turn, results in significant unwanted impacts on health, economic activity and education (7).

According to the WHO/UNICEF Joint Monitoring Programme (JMP, 2014), the estimated coverage of urban sanitation indicated as improved, shared and other unimproved facilities have reached 27%, 40% and 26% respectively in 2015 compared to 20%, 30% and 12% respectively in 1990 (7). As a matter of fact, the existing sanitation condition for many of the school in Ethiopia is horrendous. Most school latrines are filthy and unclean, and the poor condition is contributing to high level disease prevalence, creates poor learning environments and especially impacting on girls' education (8).

However, much attention is not given to this problem and studies on menstruation and its hygienic management as well as its influence on girls' education are limited in Ethiopia. This

study therefore, will be conducted with the aim of assessing menstrual hygiene management among primary and secondary school girls at both private and public schools.

Objectives

General objective

To assess the menstrual hygiene management and its determinant among private and public adolescent school girls in Addis Ababa, Ethiopia.

Specific objective

To measure level of knowledge about menstrual hygiene management among adolescent school girls.

To measure level of menstrual hygiene management practice among adolescent school girls.

To determine factors affecting menstrual hygiene management knowledge and practice among adolescent school girls.

Methods

Study area

The study was conducted among primary and secondary school girls in Addis Ababa a capital of Ethiopia, from April 1 to May 15, 2017.

Addis Ababa was established in 1886 by Emperor Taitu and Minilik II and is the capital city of Federal Democratic Republic of Ethiopia. Addis Ababa is located in an area of 540.1 square kilometers and located at 9°2'0" North and 38°42'0" East at range of 2200 – 2800 meters above sea level. Despite its proximity to the equator, its lofty altitude - the third-highest capital in the world - means that it enjoys a mild climate with an average temperature of 16°C (61°F). The hottest, driest months are usually April and May. The city has administrative structures: one city council, 10 sub-cities and 116 woredas. The total population of the city projected for the year 2016, by Population Census Commission, was 3.3 million with male to female ratio of 0.92 (29).

The city has a total of 203 government and 612 private schools. Of the total number of 107,106 students enrolled from grade 7th to 10th education in the year 2016/17 were females (26).

Study design

School based cross-sectional study with a quantitative research methods was employed. The survey was conducted among female adolescent students. The interviews were explored female students' views about menstruation and its hygienic management and availability, accessibility and adequacy improved sanitation and hygiene keeping facilities for menstrual hygiene management.

Population

Source population

The source population of the study was all grade 7th to 10th students in public and private schools of the selected sub cities of Addis Ababa.

Study population

The study population was all grade 7th to 10th students from the selected public and private schools.

Study subjects

The study subjects were 770 randomly selected students and from whom data were collected.

Inclusion and Exclusion criteria

Inclusion criteria:

All students who were from grade 7th to 10th and has monarchy was included in the study.

Exclusion criteria:

All female students who had sight problems and with mental disorders were not be included in this study.

1 Sampling

2 Sample size determination for quantitative study

3 To determine the number of adolescent school girls to be included in the study, a two-population
4 proportion formula were used. Since the specific objectives are three, were calculated a sample size for
5 each in order to take a large sample size.

6 Specific objective 1

7 The sample size of this study was determined using a single proportion formula $n = (Z_{\alpha/2})^2 \frac{P(1-p)}{d^2}$ where
8 $Z_{\alpha/2} = 95\%$ level of confidence (1.96), $p =$ proportion of menstrual hygiene management practice among
9 adolescent school girls in previous study (prevalence of use of sanitary napkins 35.38%) (13) and $d =$
10 margin of error (5%), based on these assumption that sample size found to be 350 and to maximize the
11 response rate of the study sample size population correction were made by multiplying design effect 2 and
12 10% non-response rate. Based on the above assumptions the total sample size for objective one “n” was
13 770 school girls.

14 Specific objective two and three

15 For the second objectives: factors affecting menstrual hygiene management knowledge and practice
16 among adolescent school girls, sample size was calculated using two population proportion formula (13).

$$17 \quad n = \frac{Z_{\alpha/2}^2 (1 + 1/r) p(1-p) - Z_{\beta}^2 \{P_1(1-P_1) + [P_2(1-P_2)/r]\}^2}{(P_1 - P_2)^2}$$

18
19 Where

$Z_{\alpha/2}$: 95% confidence level

P2: the probability of event in the exposed

Z_{β} : power

r: ratio of exposed to unexposed

P1: the probability of event in the unexposed

OR: 1.5

It was calculated using statcalc sample size and power calculation for descriptive study of Epi info
version 7

So, decision was made based on the comparison between the first specific objective (770) and second objectives (434). Finally, due to the issue of representativeness a sample size of 770 were used in the study. By using sample proportional to size, determine sample size to each school. Twenty primary and secondary schools at private and public were selected systematically from the list of schools from the selected sub-cities of Addis Ababa. A total of 770 school girls were randomly selected from students networking list of selected schools based on the proportion to the size of grade seven to ten of each school.

Sampling procedures

The target participants for this study was adolescent school girls from grade 7th to 10th in very selected schools of five sub cities of Addis Ababa. A multi-stage probability sample procedure was used to select participant schools. twenty Primary and secondary schools were selected randomly from the list of schools which have grade seven to ten in the Regional Education bureau. A total of 770 School girls were randomly selected from students networking list of selected schools based on the proportion to the size of grade seven to ten of each school. The reason for the choice of school girls in grade seven to ten was because they start their menarche.

Data collection tool and procedure

A self-administered pre-test close ended Amharic questionnaire at school setting were used. The questionnaire was contained variables related to socio-demographic characteristics, knowledge about menstruation and menstrual hygiene management, practice about menstrual hygiene. Study subjects were invited to take part voluntarily by explaining the purpose of the study and data were collected after obtaining verbal consent. Data was collected by ten female health professional data collectors with health background one supervisor.

Students were instructed on how to fill the questionnaire. Data quality was assured through careful design of the questionnaire. Data collectors and supervisor were received a one-day training on the purpose and procedure of data collection related to this research. During the training, special emphasis was given to establishing trust before asking questions. The training session were also pay attention to careful consideration for sensitive questions, observations where needed, and avoidance of participation bias. Data were checked for completeness and consistency after each day of data collection checking filled questionnaires by supervisors. The overall data collection process was coordinated by the principal investigator.

Variables

Dependent variables

Practice on menstrual hygiene management.

Independent variables: -

Socio-demographic variables

Age

Grade

School type
Religion
Parent's education
Parent's occupation.
Menstrual hygiene related Variables including:
School learning on menstrual hygiene.
Discussion with parents on menstrual hygiene.
Information before menarche on menstrual hygiene.

Data analysis

All responses to the survey questionnaires were coded on pre-arranged coding sheet by the principal investigator to minimize errors. The coding will be using the original English version and were entered to EPI-7 software. The data file will export to statistical package for social studies (SPSS) version 25.0 software for analysis. Descriptive analysis including frequency, proportions, and measures of mean were done. Cross tabulations were made to calculate Crude and adjusted odds ratio. All variables with $p \leq 0.20$ in bivariate analysis were fitted in to the multiple logistic regression model to identify factors associated with menstrual hygienic practice. P value ≤ 0.05 were considered as a level of significance.

Data quality management

The quality of data was assured at the maximum attainable level by using standardized adapted questionnaire and following the necessary procedures in order to get the intended results. To ensure quality of data, pre-test of data collection tools was done on primary school girls in New Era primary school by taking 5% of the total sample size. The data collectors were got orientation. Besides, the questionnaire was checked for completeness and correctness on daily basis by immediate supervisors.

Operational definition

The students' knowledge and practices was scored using a scoring system adapted from a past study. Students' menstrual knowledge score was calculated out of the 12 knowledge specific questions (Table 3). Each correct response earned one point, whereas any wrong or don't know response attracted no mark and thus the sum score of knowledge was calculated (12 points). Accordingly, the mean score of menstrual knowledge (7 ± 1.67) was used to decide the cutoffs of the rank. Good knowledge of menstruation and menstrual hygiene was given to those respondents who scored 7–12 points and Poor Knowledge of menstruation and menstrual hygiene was given to those respondents who scored 0–6 points. Students' practice of menstrual hygiene score was calculated out of the practice specific questions (Table 4). Each correct response earned one point, whereas any wrong or don't know response attracted no mark and thus the sum score of practice was calculated (15 points). And also, the mean score of menstrual practice (8 ± 3.619) was used to decide the cutoffs of the rank. Good practice of menstrual hygiene was given to those respondents who scored 8–15 points and poor knowledge of menstruation and

menstrual hygiene was given to those respondents who scored 0–7 points. Each correct response earned one point, whereas any wrong or don't know response attracted no mark (12).

Hygienic menstrual management practice in school – adolescent school girls using a clean menstrual management material to absorb or collect blood that can be changed in privacy as often as necessary for the duration of the menstruation period, using soap and water for washing the body as required, and having access to facilities to disposed of used menstrual management materials (4).

Secondary school – a high school or a senior high school which provides secondary education, between the ages of 14-19, after primary school and before higher education(26).

Primary school - a primary school or elementary school which provides primary education, between the ages of 6-14, after kinder-garden school and before secondary education(16).

Access to water supply– Sufficient water-collection points and water-use facilities are available in the school to allow convenient access to, and use of, water for drinking, personal hygiene, cleaning and laundry (4).

Address the gender-related needs: The number, location, orientation of school WASH facilities should take into consideration of the gender factor (gender mainstreaming) (8).

Adequacy of safe water- is the availability of 5 liters per person per day for all schoolchildren and staff at day schools(4).

Appropriate designs for different age groups: The detailed design of the facilities provided must also be young child friendly. Steps must be easy to climb. Door handles must be easy to reach. The toilet interior cannot be too dark. Squatting plates must be designed to accommodate a child's feet rather than those of an adult (8).

Improved sanitation - are those more likely to ensure privacy and hygienic use /easily cleanable which is not full, do not have fecal matter in the squat (24)

Physically separate facilities: Physically separated facilities must be provided for girls, spaced sufficiently apart to ensure that girls do not feel embarrassed but secure when approaching and using the facilities. Separate hand-washing areas should also be provided, affording privacy for girls who may need to wash and dry menstrual cloths (8).

Use appropriate orientation of facilities: Specifically, the direction that the toilet entrance faces, must also take into account the perceived security and safety of girls. The orientation of the squatting plate should also take into account cultural and religious norms (8).

Ethical consideration

Ethical clearance was secured from School of Public Health, College of Health Sciences, Addis Ababa University Research Ethics Committee. Approval letter was obtained from Addis Ababa City Administration Education Bureau in the respective schools included in this study. School directors and directresses were briefed on the objectives of the study and permission to conduct the study was obtained from participating schools. The questions from the questionnaire prove not to affect the morale and

personality of study subjects. Informed verbal consent was obtained from each study subject after explanation of the objective of the study. Confidentiality was ensured from all data collectors via using code numbers than names and keeping questionnaires locked. Data collectors also give health education and advice to the subjects during the data collection process.

Result

Socio-demographic characteristics of study population

A total of 756 primary and secondary school girls were participated from twenty primary and secondary schools, with response rate of 98%. Among these participants 38.1 % (288) were from private and the rest 61.9 % were from government schools. Among the total respondent 156 (20.6%), 160 (21.2%), 220 (29.1%) and, 220 (29.1%) were grade seven, eight, nine and ten respectively. The mean age of the study participants was 14.89 with SD $\pm$ 1.285 years, while their age range between 12-20 years. The mean age of menarche of the respondents was 12.84 with SD $\pm$ 0.745 years.

The study also indicated that 267 (35.3%) and 226 (29.9 %) of the respondents' father completed secondary and higher level of education respectively. Regarding respondent's mother occupation, 255 (33.7%) of them were house wife while 156 (20.6%) government employed. The majority 70.2% (531) of the respondents didn't earn pocket money form their families. The quintile division showed that, wealth was almost equally distributed across the five quintiles. 20.5% of respondents were in the lowest quintile whereas 18.7% in the highest quintile (See table 2)

Table 1 Socio-demographic characteristics of primary and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Characteristics of respondents		Frequency	Percentage
Religion	Orthodox	528	69.8
	Muslim	154	20.4
	Protestant	59	7.8
	Catholic	15	2.0
Wealth using the wealth index	Quintile One	155	20.5
	Quintile two	148	19.6
	Quintile three	151	20.0
	Quintile four	161	21.3
	Quintile five	141	18.7
Mothers educational status	Illiterate	206	27.2
	Read & write	84	11.1
	Primary	175	23.1
	Secondary	163	21.6

	College and above	128	16.9
Live with	Both parents	651	86.1
	only mother	59	7.8
	Relatives	24	3.2
	Only father	22	2.9
Fathers occupation	Merchant	204	27.0
	Employed in private organization	197	26.1
	Governmental employee	180	23.8
	Driver	94	12.4
	Daily laborer	81	10.7

153

154 Adolescent school girls' knowledge about menstruation and its hygienic management

155 This study also found that 83 percent of respondents knew about menstruation before it occurred and their
 156 chief source of information was mothers (68.3%) followed by friends (41.08%), elder sister (32.48%) and
 157 school (28.18%). Out of the total respondents, 392 (50.9%) didn't learn about menstrual hygiene in the
 158 school. Two hundred three (26.9%) of the respondents didn't discuss about menstrual hygiene with their
 159 parents and friends.

160 Out of the total, 599 (79.2%) of girls knew that menstruation was a physiological process, whereas 42
 161 (5.6 %) of the them believed that it was a curse from God. Majority of girls (69.3%) correctly responded
 162 hormone as the cause of menstruation. More than half, 64.6% of the respondents knew that uterus is the
 163 source of menstrual blood. (See table 3)

164 Table 2 School girls' Information and knowledge grading on menstruation and its management,
 165 April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Variables	Number	Percentage
Normal menstruation cycle		
Less than 25 days	84	11.1
25 to 28 days	515	68.1
28 to 35 days	104	13.8
More than 35 days	53	7.0

Normal regular menstrual bleeding duration

< 2 days	52	6.9
2 to 7 days	676	89.4
>7	28	3.7

Knew that there is foul smelling during menstruation

No	142	18.8
Yes	614	81.2

Knew that menstrual blood is unhygienic

No	200	26.5
Yes	556	73.5

Knew Pain during menstruation means that's one not sick

No	254	33.6
Yes	502	66.4

Knew menstruation is not harmful for a woman's body if she runs or dances during her period

No	229	30.3
Yes	527	69.7

Hear about menstruation before attaining menarche

No	116	15.3
Yes	640	84.7

Menstruation is not a lifelong process

No	130	17.2
Yes	626	82.8

Knew that a girl should take more nutritious diet during menstruation

No	190	25.1
Yes	566	74.9

Based on the Knowledge summary of the respondent, 70.1% (67.1-73.5) had good knowledge about menstruation and its hygiene while 29.9 % of them had below the mean score of knowledge and categorized as poor knowledge.

Menstrual Hygiene Practice

Out of the total respondents, 52.5 % (48.7-56.1) of the respondents had good practice on menstrual hygiene. Majority 457 (60.4%) of students used sanitary napkins and the rest 295 (39%) and 4 (0.5%) of them used homemade cloth and underwear as menstrual soak-up during their last menstrual period respectively. This study also enlisted the main reason for the non-use of sanitary napkins includes; 149(53.02%) high cost, 99(35.23%) difficulty in disposal, and 33(11.74%) lack of knowledge. Five hundred thirty-one participants change their pads during menstruation at school. Half 384 (59.3 %) of girls change their pads three and above times per day and 129(17.1%) changed their pads once in a day.

Table 3 Practice grading on MHM among private and government school girl's, April 1 to May 15, 2017 Addis Ababa, Ethiopia.

Variables	Number	Percentage
Change pads at school		
No	225	29.8
Yes	531	70.2
Frequency of changing absorbent material per day		
Once	129	17.1
Twice	179	23.7
Three times	399	52.8
More than three times	49	6.5
Clean genitalia		
No	212	28.0
Yes	544	72.0
Material used for genital cleaning		
Soap and water	207	38.05
Water only	307	56.43

Plain paper	30	5.51
Clean external genitalia		
No	254	33.6
Yes	502	66.4
Materials for cleaning external genitalia		
Soap and water	148	29.48
Water only	314	62.55
Plain paper	40	7.97
Hand washing		
No	370	48.9
Yes	386	51.1
Materials for hand washing		
Soap and water	163	42.23
Water only	223	57.77
Shower		
No	238	31.5
Yes	518	68.5
Materials for showering		
Soap and water	506	97.68
Water only	12	2.22
Drying of washed reusable cloths		
In the shade, outside	68	9.0
In the shade, inside	201	26.6
In the sunlight, inside	201	26.6
In the sunlight, outside	131	17.3
Hidden under other clothes	60	7.9

Hidden elsewhere

95

12.6

Six hundred forty (84.7 %) of the respondents disposed their used sanitary pads in the latrine, 68(9%) wrap in paper and put in the bin, and the rest 48(6.3%) threw in the open field. Three hundred thirty-one (43.9%) of the respondents dried their reusable sanitary pads in sunlight. 32.4% of the study participants store their sanitary pads in separated plastic bag for the next use. (see table 4)

This study evidenced that menstrual hygiene practice related school absenteeism was prevalent amongst respondents, 64.3% of whom miss school at least once in a month (mean 1.14, SD 1.132). Out of these respondents, 348(46.03%) of the girls were absent from school during their last menstrual period up to four days. The main reasons for school absenteeism during menstruation were; pain 294 (79.03%), followed by lack of washing facility at school 292 (78.49%), feel uncomfortable or tired 207 (55.65%), no private place to change sanitary pad 197 (48.12%), and didn't have sanitary pad, 98 (26.34%).

Association between overall good knowledge of menstrual hygiene management and socio demographic factors

The crude value in the bivariate analysis, some of socio-demographic characteristics of the respondents were significantly associated with the outcome variable-knowledge of menstrual hygiene. The odds of good knowledge about menstruation and menstrual hygiene were 7.01 times higher for those respondents from private school compared to the government once (COR=7.01%, 95% CI: 4.936-9.952). Whereas the odds of good knowledge about menstruation and menstrual hygiene were 31.493 times higher for respondents' mother education collage and above compared to the illiterate (COR=31.493%, 95% CI: 13.959-71.053). Unlike the crude value, the controlled model found that only the highest categories of wealth were associated with the knowledge. Girls from the wealthiest family were [AOR = 3.199, 95 % CI: 1.082-9.454] more likely to have good knowledge about menstruation and menstrual hygiene when compared to those from non-wealthy. (see table 6)

Table 4 Crosstabulations between socio-demographic variables and knowledge about menstrual hygiene management among primary and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Characteristics	Knowledge			Crude OR
	Poor	Good	Total	95% (CI)

Wealth index				
Quintile One	109	46	155	1
Quintile two	58	96	148	3.68(2.28-5.93) *
Quintile three	35	116	151	7.85(4.71-13.098) *
Quintile four	8	153	161	45.318(20.568-99.85) *
Quintile five	16	125	141	18.52(9.92-34.556) *
School type				
Private	75	213	288	1
Government	151	317	468	0.74(0.533-1.025)
Age at first menarche				
<=13	157	461	618	1
>13	69	69	138	2.94(2.01-4.29) *
Grade				
Primary	166	150	316	1
Secondary	60	380	440	7.01(4.94-9.95) *
Live with				
Both parents	139	512	651	1
Only mother	47	12	59	0.069 (0.036-0.134)
Only father	18	6	24	0.090 (0.035-0.232)
Relatives	22	0	22	0.000 (0.000-0.000)
Mothers educational status				
Illiterate	133	73	206	1
Read & write	27	57	84	3.846(2.24-6.598) *
Primary	36	139	175	7.04(4.42-11.195) *
Secondary	23	140	163	11.09(6.56-18.75) *
College and above	7	121	128	31.49(13.96-71.05) *

Fathers educational status

Illiterate	57	29	86	1
Read & write	42	30	72	1.404 (0.735-2.683)
Primary	30	75	105	4.914(2.66-9.095) *
Secondary	82	185	267	4.434(2.64-7.438) *
College and above	15	211	226	27.65(13.887-55.045) *

Fathers occupation

Merchant	55	149	204	7.74(4.31-13.898) *
Private employee	36	161	197	12.778(6.91-23.62) *
Governmental employee	26	154	180	16.92(8.85-32.35) *
Driver	49	45	94	2.62(1.38-4.98) *
Daily laborer	60	21	81	1

Mothers occupation

Housewife	113	142	255	1.374(0.802-2.357)
Merchant	29	86	115	3.24(1.714-6.138) *
Private employee	41	123	164	3.281(1.809-5.953) *
Governmental employee	8	147	155	20.098(8.52-47.398) *
Daily laborer	35	32	67	1

Earn pocket money

Yes	46	179	225	1.99(1.38-2.89) *
No	180	351	531	1

211 *P<0.05 cutoff point for significance

212 The controlled effect indicated that girls whose mother's education college and above were 6.101 [AOR,
213 95% CI: 1.779-20.919] times more likely to have good knowledge about menstruation and menstrual
214 hygiene than their counterparts. The model also found that girls from secondary schools were [AOR =
215 13.742, 95 % CI: 5.390-35.037] more likely to have good knowledge about menstruation and menstrual
216 hygiene when compared from primary school girls. Respondents whose mothers from government
217 employee were strongly associated with good knowledge about menstruation and menstrual hygiene
218 10.555 (2.047-54.411) [AOR = 10.555, 95 % CI: 2.047-54.411]. This study also found that girls from the

219 wealthiest family were [AOR = 3.199, 95 % CI: 1.082-9.454] more likely to have good knowledge about
220 menstruation and menstrual hygiene when compared to those from lowest quantile. (see table 6)

221 Table 5 Association between socio demographic factors and knowledge about MHM among
222 primary and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Characteristics	Knowledge			Crude OR	Adjusted OR
	Poor	Good	Total	95% (CI)	95% (CI)
Age of the respondents					
<15 years	132	173	305	1	1
>=15 years	94	357	451	2.898(2.102-3.904)	0.819(0.344-1.950)
Wealth index					
Quintile One	109	46	155	1	1
Quintile two	58	96	148	3.677 (2.282-5.925)	2.090 (0.826-5.289)
Quintile three	35	116	151	7.853(4.709-13.098)	2.079 (0.815-5.303)
Quintile four	8	153	161	45.318(20.568-99.85)	4.050(1.265-12.966) *
Quintile five	16	125	141	18.512 (9.917-34.556)	3.199(1.082-9.454) *
Grade					
Primary	166	150	316	1	1
Secondary	60	380	440	7.01(4.936-9.952)	13.742(5.390-35.037) *
Mothers educational status					
Illiterate	133	73	206	1	1
Read & write	27	57	84	3.846 (2.242-6.598)	0.705 (0.257-1.934)
Primary	36	139	175	7.035 (4.420-11.195)	5.718(2.548-12.831) *
Secondary	23	140	163	11.090 (6.558-18.7530)	10.012(3.628-27.629) *
College and above	7	121	128	31.493 (13.959-71.053)	6.101(1.779-20.919) *
Fathers educational status					
Illiterate	57	29	86	1	1
Read & write	42	30	72	1.404 (0.735-2.683)	1.245 (0.376-4.125)

Primary	30	75	105	4.914 (2.655-9.095)	1.284 (0.444-3.709)
Secondary	82	185	267	4.434 (2.644-7.438)	0.796 (0.302-2.098)
College and above	15	211	226	27.648 (13.887-55.045)	2.197 (0.648-7.450)
Fathers occupation					
Merchant	55	149	204	7.740(4.311-13.898)	0.880 (0.284-2.726)
Private employee	36	161	197	12.778 (6.912-23.621)	0.888 (0.274-2.874)
Governmental employee	26	154	180	16.923(8.854-32.346)	2.973(0.863-10.250)
Driver	49	45	94	2.624(1.382-4.981)	1.568 (0.510-4.821)
Daily laborer	60	21	81	1	1
Mothers occupation					
Housewife	113	142	255	1.374(0.802-2.357)	2.301 (0.686-7.721)
Merchant	29	86	115	3.244(1.714-6.138)	1.135 (0.300-4.289)
Private employee	41	123	164	3.281(1.809-5.953)	0.391 (0.103-1.488)
Governmental employee	8	147	155	20.098(8.522-47.398)	10.555(2.047-54.411) *
Daily laborer	35	32	67	1	1
Age at first menarche					
<=13	157	461	618	1	1
>13	69	69	138	2.936 (2.009-4.292)	2.093 (0.914-4.793)
Earn pocket money					
Yes	46	179	225	1.99(1.378-2.890)	0.741(0.392-1.400)
No	180	351	531	1	1

223 *P<0.05 cutoff point for significance

224 Association between overall practice of menstrual hygiene management and socio demographic factors

225 Based on the bivariate analysis, among the ten socio-demographic variables, level of grade (secondary),
226 respondent's mother's educational status (college and above), respondent's father educational status
227 (college and above), both respondent's mother and father occupational status, Age at first menarche
228 (>=13) and pocket money were significantly associated with 95 % CI COR at P<0.05 with overall
229 practice of menstrual hygiene management among respondents.

230 The crude value also showed that the odds of overall practice of menstrual hygiene management among
 231 secondary school girls were 2.364 (95% C.I: 1.759-3.177) times higher compared to primary school girls.
 232 In this study, it was found that pocket money was associated with overall practice of menstrual hygiene
 233 management. The odds of overall practice of menstrual hygiene management among girls 2.177 (95%
 234 C.I: 1.575-3.009) times higher for respondent who had earned pocket money from their families. The
 235 odds of overall practice of menstrual hygiene management among girls 2.330 (95% C.I: 1.720-3.156)
 236 times higher among respondent who were from private school than government school girls. (see table 7)

237 Table 6 Crosstabulations between socio-demographic variables and MHM practice among
 238 primary and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Characteristics	Practice			Crude OR
	Poor	Good	Total	95% (CI)
School type				
Government	259	209	468	1
Private	100	188	288	2.33(1.72-3.16) *
Age of the respondents				
<15 years	180	125	305	1
>=15 years	179	272	451	2.19(1.63-2.94) *
Wealth index				
Quintile One	133	22	152	1
Quintile two	91	57	148	3.79(2.16-6.63) *
Quintile three	74	77	151	6.29(3.620-10.931) *
Quintile four	32	129	161	24.37(13.45-44.16) *
Quintile five	29	112	164	23.35(12.71-42.91) *
Grade				
Primary	189	127	316	1
Secondary	170	270	440	2.36(1.76-3.18) *
Religion				
Orthodox	282	246	528	1
Muslim	40	114	154	0.582(0.204-1.657)

Protestant	31	28	59	1.900(0.636-5.674)
Catholic	6	9	15	0.602(0.19-1.906)
Live with				
Both parents	269	382	651	1
Only mother	44	15	59	0.280(0.131-0.440)
Only father	24	0	24	0.998(0.000-0.000)
Relatives	22	0	22	0.998(0.000-0.000)
Mothers' educational status				
Illiterate	150	56	206	1
Read & write	42	42	84	2.68(1.58-4.54) *
Primary	100	75	175	2.01(1.31-3.08) *
Secondary	48	115	163	6.42(4.07-10.12) *
College and above	19	109	128	15.37(8.64-27.35) *
Fathers' educational status				
Illiterate	64	22	86	1
Read & write	49	23	72	1.365(0.683-2.730)
Primary	39	66	105	4.92(2.634-9.203) *
Secondary	135	132	267	2.84(1.657-4.884) *
College and above	72	154	226	6.22(3.556-10.887) *
Fathers' occupation				
Merchant	74	130	204	9.19(4.757-17.750) *
Private employee	55	142	197	13.51(6.911-26.391) *
Governmental employee	86	94	180	5.717(2.951-11.078) *
Driver	76	18	94	1.239(0.565-2.716)
Daily laborer	68	13	81	1
Mothers' occupation				

Housewife	145	110	255	2.63(1.407-4.92) *
Merchant	41	74	115	6.26(3.140-12.47) *
Private employee	63	101	164	5.56(2.887-10.699) *
Governmental employee	58	97	155	5.798(2.996-11.22) *
Daily laborer	52	15	67	1
Age at first menarche				
<=13	268	350	618	1
>13	91	47	138	2.53(1.718-3.72) *
Earn pocket money				
Yes	77	148	225	2.18(1.575-3.01) *
No	282	249	531	1

239 *P<0.05 cutoff point for significance.

240 After controlling interaction effect of all the variables, it was found that the current age of school
 241 girl students above fifteen years old were 2.283 [AOR (95% C.I: 1.613-4.971)] times more likely
 242 to have good practice than their counterparts. Girls whose mother's education secondary and
 243 above were eight times more likely to have good practice about menstrual hygiene compared to
 244 those from illiterate mothers [AOR = 7.761, 95 % CI: 3.583-16.809)]. Girls whose fathers from
 245 private employee were [AOR (95% C.I): 3.654 (1.215-10.991) times more likely had good
 246 menstrual hygiene management practice than those who were daily laborer family. This study
 247 also found that girls whose age at first menarche greater than thirteen were 2.572 times more
 248 likely to have good practice about menstrual hygiene compared to those who were less than
 249 thirteen years old. [AOR (95% C.I): 2.572 (1.409-4.694)]. Practice of MHM was also distributed
 250 unequally across the wealth quintile. The odd of good practices were 3 times higher in the
 251 highest quintile compared to the 2nd. In addition, the overall knowledge of the respondents was
 252 significantly associated with their practice [AOR (95% C.I): 4.581 (2.462-8.526)] (See table 8)

253 Table 7 Association between socio demographic variables and level of MHM practice among
 254 primary and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n =756)

Characteristics	Practice			Crude OR	Adjusted OR
	Poor	Good	Total	95% (CI)	95% (CI)
School type					
Government	259	209	468	1	1

Private	100	188	288	2.330(1.720-3.156)	1.09(0.64-1.85)
Age of the respondents					
<15 years	180	125	305	1	1
>=15 years	179	272	451	2.188(1.627-2.942)	2.832(1.62-4.97) *
Wealth index					
Quintile One	133	22	152	1	1
Quintile two	91	57	148	3.787 (2.164-6.626)	2.968(1.32-6.67) *
Quintile three	74	77	151	6.291 (3.620-10.931)	3.80(1.730-8.35) *
Quintile four	32	129	161	24.371 (13.45-44.16)	14.776(6.23-35.04) *
Quintile five	29	112	164	23.348 (12.71-42.91)	9.038(3.728-21.91) *
Grade					
Primary	189	127	316	1	1
Secondary	170	270	440	2.364(1.759-3.177)	3.115 (1.606-6.04) *
Mothers' educational status					
Illiterate	150	56	206	1	1
Read & write	42	42	84	2.679(1.582-4.535)	1.588(0.71-3.560)
Primary	100	75	175	2.009(1.308-3.084)	1.929(0.95-3.933)
Secondary	48	115	163	6.417(4.069-10.122)	7.761(3.58-16.81) *
College and above	19	109	128	15.367(8.639-27.352)	17.91(6.65-48.22) *
Fathers' educational status					
Illiterate	64	22	86	1	1
Read & write	49	23	72	1.365(0.683-2.730)	0.482(0.146-1.588)
Primary	39	66	105	4.923(2.634-9.203)	0.552(0.218-1.403)
Secondary	135	132	267	2.844(1.657-4.884)	0.570(0.110-1.663)
College and above	72	154	226	6.222(3.556-10.887)	0.574(0.099-1.761)
Fathers' occupation					
Merchant	74	130	204	9.189(4.757-17.750)	1.933(0.705-5.295)

Private employee	55	142	197	13.505(6.911-26.391)	3.65(1.22-10.991) *
Governmental employee	86	94	180	5.717(2.951-11.078)	1.178(0.413-3.363)
Driver	76	18	94	1.239(0.565-2.716)	0.481(0.166-1.391)
Daily laborer	68	13	81	1	1
Mothers' occupation					
Housewife	145	110	255	2.630(1.407-4.916)	2.125 (0.759-5.952)
Merchant	41	74	115	6.257(3.140-12.470)	1.570 (0.522-4.718)
Private employee	63	101	164	5.558(2.887-10.699)	1.259 (0.448-3.539)
Governmental employee	58	97	155	5.798(2.996-11.219)	0.813 (0.272-2.435)
Daily laborer	52	15	67	1	1
Age at first menarche					
<=13	268	350	618	1	1
>13	91	47	138	2.529(1.718-3.721)	2.57(1.41-4.69) *
Earn pocket money					
Yes	77	148	225	2.177(1.575-3.009)	1.11(0.67-1.81)
No	282	249	531	1	1

255 *P<0.05 cutoff point for significance.

256 Association between overall knowledge and practice of menstrual hygiene management and other
257 menstrual related factors

258 The regression model also evidenced other menstrual related factors like learning and discussing about
259 MHM in the school and with parents and friend and also hearing about it before menarche were
260 significantly associated with both the outcome variables. The odds of good knowledge about menstrual
261 hygiene management among those who learn about menstrual hygiene at school were 3.110 (95% CI:
262 1.569-6.162) times higher than those who didn't learn at their school. (See table 9)

263 Table 8 Association between other related variables and knowledge about MHM among primary
264 and secondary school girls, April 1 to May 15, 2017 Addis Ababa, Ethiopia. (n=756)

Characteristics	Knowledge		Crude OR	Adjusted OR
	Good	Poor	95% (CI)	95% (CI)
Learn about menstrual hygiene in				

the school				
Yes	316	55	4.59(3.24-6.51)	3.11(1.57-6.16) *
No	214	171	1	1
Discuss about menstrual hygiene with their parents				
Yes	481	72	20.99(13.99-31.51)	10.89(5.39-21.98) *
No	49	154	1	1
Heard about menstruation before attaining menarche				
Yes	500	140	10.24(6.49-16.15)	5.03(2.21-11.48) *
No	30	86	1	1

265 *P<0.05 cutoff point for significance.

266 This study found that the odds of good practice about menstrual hygiene management among those who
 267 discuss about menstrual hygiene with their parents were 13.651 (95% CI: 7.087-26.296) times higher than
 268 those who didn't discuss about menstrual hygiene with their parents. (See table 10)

269 Table 9 Association between socio demographic factors and MHM practice among primary and
 270 secondary school girls in Addis Ababa, 2017. (n=756)

Characteristics	Practice		Crude OR	Adjusted OR
	Good	Poor	95% (CI)	95% (CI)
Learn about menstrual hygiene in the school				
Yes	259	112	4.14(3.05-5.61)	2.47(1.52-4.01) *
No	138	247	1	1
Discuss about menstrual hygiene with their parents				
Yes	371	182	13.88(18.86-21.73)	13.65(7.09-26.296) *
No	26	177	1	1
Hear about menstruation before				

attaining menarche

Yes	37	269	4.78(3.003-7.59)	0.779(0.366-1.656)
No	26	92	1	1

*P<0.05 cutoff point for significance.

Discussion

The onset of menstruation is one of the most important changes occurring among the girls during the adolescent years. The bodily changes associated with puberty will have an impact in the girl's physical, psychological and social development (19–21). This study was conducted to identify factors affecting practice of menstrual hygiene among school girls in Addis Ababa.

In this study, the mean age of menarche of the respondents was 12.84 with SD +0.745 years which is similar to studies conducted in Jammu district, India and Argoha village of Haryana with the mean age of menarche 13.43±.83 and 12.76 ± 0.936 years respectively (30,31).

In consistent with study reports from Jimma district, India (66.15%) (30), for about 68% school girls, their mothers were main source of information on menstruation. This could be suggestive of the contribution of mothers for hygienic practice of girls during menarche. In contrast to this finding, school teachers and sisters respectively, were reported to be important source of such information in Amhara North East, Ethiopia (13).

Factors of menstrual hygiene management

Present study also reported that, majority (70.1 %) of the students had good knowledge about menstruation and menstrual hygiene. The finding was similar with the result from studies done in western Ethiopia; 60.9%. This study notified that Five hundred ninety-nine (79.2%) of girls knew that menstruation to be normal a physiological process. Such prevalence found to be consistent with a result of a study done in western Ethiopia (76.9%), Argoha village of Haryana 71.3%, and Central India 89% of school girls knew correctly that menstruation as physiologic process(12,31,32). A possible explanation for this similarity may be that girls had good discussion in families openly. This finding however, was higher than that of those in previous a study done in Northeast Ethiopia; 319 (57.89%) This difference might be due to the study method difference (mixed qualitative and quantitative method for the previous study (12).

Adolescent girls who knew that uterus was the source of blood in menstruation were 64.6% which is similar to studies conducted in Amhara Ethiopia, western Ethiopia, and Central India was found out to be 60.9%, 59.3% and 60% respectively (12,32,33). This study disagrees with results obtained from a study in Argoha village of Haryana 38.7%, possibly due to minimum information provided about menstruation and menstrual hygiene by schools and families (31).

In this study, multivariable analysis showed that girls whose mother's educational status secondary school and above were 10.012 times more likely to had good knowledge about menstruation and menstrual hygiene than their counterparts [AOR = 10.012, 95 % CI: 3.628-27.629]. A similar study done in western

Ethiopia and Jammu District India showed that, parental education was positively associated with girls' menstrual knowledge (12,30). The reason could be that educated mothers may provide information about menstruation and menstrual hygiene to their daughters. Girls from educated families may discuss openly about menstruation.

Unlike a study done in Amhara region, Ethiopia 2016 [AOR = 0.94, 95 % CI: 0.46–1.92], this study found that, the grade level of respondents was positively and significantly [AOR = 13.74, 95 % CI: 5.39–35.04] associated with knowledge about menstruation and menstrual hygiene management. Due to the fact that high school girls might have high possibility for exposure to information regarding menstruation and its hygienic management (11). In another study done in Odisha, India 2015, none of the mother's occupational category were significantly related with knowledge (25). But in this study only government employed mothers were the one associated with knowledge on menstruation.

Regarding hygiene related practices during menstruation, this study found that 9.27% girls took daily bath during menstruation and 29.48% clean their external genital with soap and water during menstruation. A similar study done in Jammu and Kashmir, India indicated that 93.18% had daily bath and 66.67% clean external genital with soap and water (30). The difference might be due to socio cultural, weather condition and economic factors.

In this study, three hundred ninety-seven (52.5%) of the respondents had good practice of menstrual hygiene. The finding of this study was lower than studies conducted in the Amhara region of Ethiopia and Jammu and Kashmir, India which were 84.28 % and 59.09%, respectively (30,33). Comparatively, lower level of practice of menstrual hygiene was recorded from similar study conducted on high school girls in Western Ethiopia, it was indicated that only 39.9 % of the study participants practice good menstrual hygiene (12). Thus, the reason for the observed difference could be due to low awareness and communication of menstrual hygiene by high school Western Ethiopia girls which affects their menstrual hygienic practice.

In this study, it was found that 17.3% of the study participants dried their reusable pads in the sun light outside which is similar with the finding in Amhara, Ethiopia 15.5% of the participants dried in the sun light. In contrast, a study done in central India indicated that 93% of adolescent girls dried reusable pads in the sun light (32,33). This difference might be due to different in the socio-cultural factors. 457(60.4%) of participants use disposable sanitary pads during menses. In a similar study done in Argoha village of Haryana and central India Indicated that, 80.7 % and 98% girls use only napkin during menses respectively (31,32). Comparatively, lower level of use of sanitary pads was recorded from similar study conducted on high school girls in north east Ethiopia, it was indicated that 35.38% of the study participants use disposable sanitary pads during menses. This may be due to differences in socio economic differences. The main reason for not using pads in present study was non-affordability due to high cost (53.02%) followed by non-availability and disposal problems which is similar to study in Jammu district India(78.94%) (30).

The adjusted value of this study found that, both mother's educational category was [AOR = 7.761, 95 % CI: 3.583–16.809] positively associated with practice of menstrual hygiene which agrees with the study done in western Ethiopia, in 2014 and Northeast Ethiopia [AOR = 2.03, 95 % CI: 1.38–2.97] [AOR = 4.26, 95 % CI: 1.61 - 11.28] respectively (11,12). Multivariable analysis showed that girls who learn and discuss about menstrual hygiene in their school and with their parent were 2.472, and 13.651 times more

likely to had good practice about menstruation and menstrual hygiene than their counterparts. A similar study done in northeast Nigeria that learn about menstrual hygiene in the school was positively associated with girl's menstrual practice (34). Possibly due to information provided about menstrual hygiene management at schools. Students whose age above fifteen were 2.832 times more likely to have good practice than age less than fifteen a similar study done in northeast Nigeria. A significant association was also observed between girls whose first menarche were above thirteen (AOR= 2.572) and below with the practice of MHM. Which was also significant in another study done in south India (24).

Conclusion

Seventy percent of the participants had good knowledge of menstruation and menstrual hygiene and it was better among private school girls than the government. Half of the total respondents had good practice of menstrual hygiene among respondents, and alike to that of knowledge, the large proportion of them were from private school.

Good knowledge of menstruation showed a significantly positive association with the level of grade, educational status of the mother, and mother's occupational status. Except for the level of grade, all the above variables plus current age of respondents' and age at first menarche were positively associated with practice of menstrual hygiene management.

All those factors considered in addition to socio demographic variables, which assume to be predictive of the outcome variables including learn about menstrual hygiene in the school, discuss about menstrual hygiene with their parents and friends and hear about menstruation before attaining menarche were positively associated.

Recommendation

Federal level managers are recommended to:

Strengthen the enabling environment through advocacy and policy initiatives for improved WASH and MHM education.

Promote an innovative, intercultural, multi-sectorial and gender approach in all programming, ensuring that MHM aspects are included in planning processes and budget allocation processes by the water and sanitation, health, and education ministries.

Regional Level: in addition to federal recommendations, regional level managers are advised to:

Give technical assistance and advocacy to prioritize budgets and investment in WASH facilities in schools by health, and education bureau.

Strengthen teachers' capacities and equips them with tools to provide in-depth and medically accurate information to students in a safe learning environment by education bureau.

Strengthen school health packages provided by health extension professionals by health bureau.

School level managers suggested to:

Establish coordination between students, teachers and parents to improve MHM conditions at schools.

Complement menstrual hygiene management as part of the school health programs and should also give special attention towards making schools a comfortable place for girl's menstrual hygiene practice by continuous provision of sanitary pad especially for the neediest ones.

Consult parents about the need to support their children with sanitary materials for menstrual hygiene in addition to other basic hygienic products during parent-school teacher meeting.

Educate and counsel girls about the important and the need for good personal hygiene including hand washing practice during menstruation by using peer group discussion which supposedly mediated by female school teachers.

Parents are advised to:

Educate their daughters about the process, good personal hygiene, use of proper pads, and its proper disposal.

Support their children with sanitary materials for menstrual hygiene.

Acknowledgment

It gives me pleasure to express my heartfelt gratitude to Addis Ababa University, School of Public Health librarians for providing me the necessary references. I would like to thank to my advisor Worku Tefera for his unrestricted guidance, envisioning and supporting from the period of instigation to this project.

I am also grateful to thank Addis Ababa education bureau, sub city education offices, school staffs, my office staffs for their cooperation and assistance in giving important information.

I am pleased to extend my thanks to my brother, Mehary Biruk, for his devotion to support me through the academic years.

I would also like to thank all the participants of the study for their patience and collaboration during data collection.

I am exclusively thankful to my friends Ashagre Sisay, Ermias Woldie, Tasew Denbi, Nardos Tadesse and Sintayehu Tadesse for their vital contribution for success of this work.

References

1. Boosey R, Prestwich G, Deave T. Menstrual hygiene management amongst schoolgirls in the Rukungiri district of Uganda and the impact on their education: a cross-sectional study. Pan African Med Journal [Internet]. 2014;8688:1–13. Available from: <http://www.panafrican-med-journal.com>
2. Caruso, A B, Fehr A, Inden K, Sahin M, Anna Ellis, et al. WASH in Schools Empowers Girls' Education in Freetown, Sierra Leone: An Assessment of Menstrual Hygiene Management in Schools. 2013;1–40. Available from: www.unicef.org/wash/schoolsCommentaries
3. Lizette, Burgers, Chiefn Lhalungpa, Samphe. useful information about menstrual health and hygiene. 1st ed. new delhi: govenment of india ministry of rural development department of drinking water supply; 2008. 1-48 p.
4. WHO. Water , Sanitation and Hygiene Standards for Schools in Low-cost Settings [Internet]. Adams J, Bartram J, Chartier Y, Sims J, Adams J, Bartram J, et al., editors. Canberra, Australia: Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland; 2009. 1-41 p. Available from: bookorders@who.int)
5. UNICEF. RAISING CLEAN HANDS Advancing Learning, Health and Participation through WASH in Schools. Sierra Leone; 2010.
6. WaterAid. Is menstrual hygiene and management an issue for adolescent school girls? A comparative study of four schools in different settings of Nepa [Internet]. Nepal; 2009. Available from: www.wateraid.org/nepalWaterAid?s
7. Ehtiopia FDR of, (EPRDE). Integrated Urban Sanitation and Hygiene Strategy (IHSHS). ADDIS ABABA; 2015. p. 1–65.
8. Ethiopia, UNICEF, Health M of, Education M of, Energy M of W and. Design and Construction Manual for Water Supply and Sanitation Facilities in Primary Schools. first edit. Addis Ababa, Ethiopia.: United Nations Children Fund – Ethiopia Program; 2012. 1-115 p.
9. WorldVision, Wash H. Investigating the Perceptions and Barriers to Menstrual Hygiene Management (MHM) in. 2014;
10. Long, Jeanne, A B, Caruso, Lopez D, Vancraeynest K, et al. WASH in Schools Empowers Girls' Education in Rural Cochabamba, Bolivia: An assessment of menstrual hygiene management in schools. 2013;1–52.
11. Gultie T, Hailu D, Workineh Y. Age of menarche and knowledge about menstrual hygiene management among adolescent school girls in amhara province, Ethiopia: Implication to health care workers & school teachers. PLoS One. 2014;9(9):1–9.

- 446 12. Upashe SP, Tekelab T, Mekonnen J. Assessment of knowledge and practice of menstrual hygiene
447 among high school girls in Western Ethiopia. BMC Womens Health [Internet]. 2015;1–8. Available from:
448 <http://dx.doi.org/10.1186/s12905-015-0245-7>
- 449 13. Tegegne TK, Sisay MM. Menstrual hygiene management and school absenteeism among female
450 adolescent students in Northeast Ethiopia. BMC Public Health [Internet]. 2014;14:1118. Available from:
451 <http://www.pubmedcentral.nih.gov/articlerender>.
- 452 14. UNICEF, IRC. A Manual on School Sanitation and Hygiene [Internet]. fifth. New York, USA:
453 UNICEF/Programme Division Water, Environment and Sanitation; 2012. 1-74 p. Available from:
454 www.unicef.org
- 455 15. Ethiopia, UNICEF. School WASH [Internet]. Addis Ababa, Ethiopia.; 2015. Available from:
456 www.unicef.org/ethiopia
- 457 16. IRC, UNICEF. Towards Effective Programming for WASH in Schools, A manual on scaling up
458 programmes for water, sanitation and hygiene in schools [Internet]. Netherland: IRC International Water
459 and Sanitation Centre; 2007. 1-93 p. Available from: <http://www.irc.nl>
- 460 17. Mwanaisha M, UNICEF. School WASH in Tanzania Improving WASH in Schools : Improving
461 the Quality of Education School WASH in Tanzania. Tanzaniya; 2008.
- 462 18. Mooijman, Nnemarieke, Sahin M, Dooley T, UNICEF, Pasic M, et al. Water , Sanitation and
463 Hygiene (WASH) in Schools, A companion to the Child Friendly Schools Manual [Internet]. New York,
464 NY 10017, USA: UNICEF’s Division of Communication.; 2012. 1-48 p. Available from: www.unicef.org
- 465 19. Shanbhag D, Shilpa R, Souza DN, Josephine P, Singh J, Br G. Perceptions regarding
466 menstruation and Practices during menstrual cycles among high school going adolescent girls in resource
467 limited settings around Bangalore city , Karnataka ,. Int J Collab Res Intern Med Public Heal.
468 2012;4(7):1353–62.
- 469 20. McMahon SA, Winch PJ, Caruso BA, Obure AF, Ogutu EA, Ochari IA, et al. The girl with her
470 period is the one to hang her head ’ Reflections on menstrual management among schoolgirls in rural
471 Kenya. McMahon al BMC [Internet]. 2011;1–10. Available from: [http://www.biomedcentral.com/1472-](http://www.biomedcentral.com/1472-698X/11/7)
472 [698X/11/7](http://www.biomedcentral.com/1472-698X/11/7)
- 473 21. Caruso B, Ellis A, Sahin M. WASH in Schools for Girls E-Course Increasing national capacity to
474 conduct research on menstrual hygiene management in schools. 2015 Nov;1–70. Available from:
475 www.unicef.org/wash/schools
- 476 22. Sumpter C, Torondel B. A Systematic Review of the Health and Social Effects of Menstrual
477 Hygiene Management. 2013;8(4).
- 478 23. Santina T, Wehbe N, Ziade FM, Nehme M. Assessment of Beliefs and Practices Relating to
479 Menstrual Hygiene of Adolescent Girls in Lebanon. Int J Heal Sci Res [Internet]. 2013;3(12):75–88.
480 Available from: www.ijhsr.org

24. Omidvar S, Begum K. Factors influencing hygienic practices during menses among girls from south India- A cross sectional study Factors influencing hygienic practices during menses among girls from south India- A cross sectional study. 2010;2(12):411–23.
25. Das P, Kelly BK, Dutta A, Swain T, Sahoo S, Das BS, et al. Menstrual Hygiene Practices , WASH Access and the Risk of Urogenital Infection in Women from Odisha , India. PLoS One. 2015;1–16.
26. AACAREB. First quarter report of 2009. Vol. 1. Addis Ababa, Ethiopia.; 2016.
27. Bartram J, Cairncross S. Hygiene , Sanitation , and Water : Forgotten Foundations of Health. PLoS Med [Internet]. 2010;7(11):1–9. Available from: www.plosmedicine.org
28. UNICEF. Advancing WASH in Schools Monitoring [Internet]. New York, NY 10017 USA; 2015. Available from: www.unicef.org/wash/schoolsAdvancing
29. Commission PC, FDRE. Summary and Statistical Report of the 2007 Population and Housing Census Results 2008. [Internet]. 2008. p. 250. Available from: http://ecastats.uneca.org/aicmd/Portals/0/Cen2007_firstdraft.pdf
30. Channawar K, Prasad V. Menstrual Hygiene: Knowledge and Practice among Adolescent School Girls. Panacea J Med Sci. 2016;6(1):31–3.
31. Sharma S, Choudhary S. Knowledge and Practices About Menstrual Hygiene Among School Adolescent Girls in AGROHA VILLAGE OF HARYANA. 2016;5(9):389–92.
32. Dixit S, Raghunath D, Rokade R, Nawaz SA, Nagdeve T, Goyal I. Awareness about Menstruation and Menstrual Hygiene Practices among Adolescent Girls in Central India. Natl J Community Med. 2016;7(6):468–73.
33. Zeleke, Balew Yosief T. Menstrual hygiene management and associated factors among high school girls: - a case of Bahir Dar city administration. 2016;1–61.
34. Lawan UM, Yusuf NW MA. Menstruation and menstrual hygiene amongst adolescent school girls in Kano, Northwestern Nigeria. Afr J Reprod Heal. 2010;14(3):201–7.