Transcriptional dynamics following freezing stress reveal selection for mechanisms of freeze tolerance at the poleward range margin in the cold water intertidal barnacle *Semibalanus balanoides*

Running title: Transcription following freezing in *S. balanoides*

Katie E. Marshall^a, Eddy J. Dowle^{bc}, Alexandra Petrunina^d, Gregory Kolbasov^d and Benny K. K. Chan^{e*}

^aDepartment of Zoology, University of British Columbia, Vancouver, Canada

^bDepartment of Integrative Biology, University of Colorado, Denver, USA

^cDepartment of Anatomy, University of Otago, Dunedin, New Zealand

^dMoskovskij gosudarstvennyj universitet imeni M V Lomonosova Biologiceskij fakul'tet, Department of Invertebrate Zoology

^eAcademia Sinica, Biodiversity Research Center, Taipei, Taiwan

*Corresponding author: Benny K.K. Chan

chankk@gate.sinica.edu.tw

Abstract

The ability to survive freezing has repeatedly evolved across multiple phyla. This suggests that the mechanisms of freeze tolerance must be readily evolvable from basal physiological traits. While several biochemical correlates to freeze tolerance have been described, the mechanism that confers freeze tolerance is still not well understood. To understand both the basic biochemical mechanisms of freeze tolerance as well as their role in local adaptation at the poleward range edge, we conducted a transcriptomic study on two populations (one from the poleward range margin in the White Sea, Russia, and one from the central coast of British Columbia, Canada) of the cold water acorn barnacle *Semibalanus balanoides* on a time series following a freezing event. We found that the British Columbia population (at the equatorward range margin) was significantly less freeze tolerant than the White Sea population (at the poleward range margin). After assembling and annotating a *de novo* transcriptome for *S. balanoides*, we found that the patterns of differential transcript expression following freezing were almost entirely non-overlapping between the two populations, with the White Sea population expressing a series of heat shock proteins in response to freezing stress as well as several aquaporins, while the British Columbia population expressed a series of proteases instead, indicating severe protein damage. We found strong evidence of purifying selection on the significantly upregulated transcripts in the White Sea population, suggesting local adaptation to freezing threat. Taken together, this shows the importance of freeze tolerance to population survival at the poleward range margin, and highlights the central roles of aquaporins and heat shock proteins to the trait of freeze tolerance across taxa.

Keywords: freeze tolerance, transcriptomics, heat shock proteins, barnacles, local adaptation

Introduction

The occurrence of species range edges is one of the central problems in biogeography (Sexton, McIntyre, Angert, & Rice, 2009). Broadly speaking, it is believed that equatorward range edges are due to biotic interactions, while poleward range edges are due to abiotic effects (Louthan, Doak, & Angert, 2015). Given that minimum temperatures have the steepest and most consistent latitudinal gradient (Sunday, Bates, & Dulvy, 2012), this suggests that low temperature tolerance is central for understanding the presence of poleward range edges. Therefore, understanding the mechanisms and adaptive potential of low temperature tolerance is central for delineating the role of local adaptation in the maintenance of poleward range edges.

Low temperature tolerance includes responses to two overlapping, but distinct, physiological stressors: the direct effects of low temperature on reaction rates, and managing the potential and spread of internal ice. In ectothermic organisms, biochemical reaction rates decrease due to Arrhenius effects as well as by decreased membrane fluidity at low temperatures, which causes a general decrease in rates of physiological functioning (reviewed in R. E. Lee, 2010). At body temperatures below 0 °C, organisms also risk internal ice formation. Surviving these two stressors, while linked by temperature, appear to involve different physiological mechanisms. Some organisms like the eastern spruce budworm (*Choristoneura fumiferana* Clemens) are able to suppress their freezing point well below -30 °C by the production of low molecular weight cryoprotectants and antifreeze proteins, and can survive low temperatures as long as they do not freeze (Han & Baucé, 1995; Katie E. Marshall & Sinclair, 2015; Qin, Doucet, Tyshenko, & Walker, 2007). Others like the woolly bear caterpillar (*Pyrrharctia isabella* Smith) induce and survive freezing at relatively high temperatures (c.a. -10 °C), but are unable to survive

temperatures below -16 °C (Boardman, Terblanche, & Sinclair, 2011; Layne & Kuharsky, 2000). Being able to survive freezing is therefore a physiological trait that is likely mechanistically distinct from being able to survive low temperature (reviewed in Toxopeus & Sinclair, 2018).

The ability to survive internal ice formation is a paraphyletic trait broadly distributed across the animal kingdom, with freeze-tolerant species from multiple phyla including arthropods, molluscs, and vertebrates (Sinclair, Addo-Bediako, & Chown, 2003; Storey & Storey, 1996; Toxopeus & Sinclair, 2018). Freeze tolerance appears to have evolved independently on at least 20 occasions, and this is almost certainly a large underestimate (Sinclair et al., 2003). The repeated emergence of this trait suggests that the underlying mechanisms must be readily evolvable and that freeze tolerance is possible in a variety of underlying physiological conditions. Despite the ubiquity of the trait, and over 200 years of research, the physiological mechanisms underlying animal freeze tolerance remain poorly understood (Lee, 2010; Sømme, 2000; Teets & Denlinger, 2014; Toxopeus & Sinclair, 2018).

Surviving internal ice formation is a unique challenge for animals. They need to be able to manage potential mechanical damage from ice crystal formation, cellular dehydration as water leaves the cell to join the growing extracellular ice lattice, and sustained periods of anoxia since oxygen cannot perfuse easily to frozen tissue (Lee, 2010). Many biochemical correlates to survival of these challenges have been described; most commonly this involves the mass biosynthesis of low molecular weight polyols and sugars that act as osmoprotectants and cryoprotectants that can also decrease the melting (and therefore freezing) point of cells for avoiding intracellular ice formation (Košťál, Korbelová, Poupardin, Moos, & Šimek, 2016; Toxopeus & Sinclair, 2018). However there is no correlation between quantity of low molecular

weight cryoprotectants and degree of freeze tolerance—for instance, woolly bear caterpillars *P. isabella* accumulate circa 2.5 M glycerol as a cryoprotectant but have a lower lethal temperatures of only circa -16 °C (Boardman et al., 2011; K. E. Marshall & Sinclair, 2011), while the freeze tolerant goldenrod gall fly (*Eurosta solidaginis* Fitch) has glycerol content of circa 0.3 M but a lower lethal temperature below -50 °C (Lee, Dommell, Joplin, & Denlinger, 1995). Freeze tolerant intertidal barnacles and mussels appear to not accumulate polyols or sugars at all (Johnston & Clarke, 1990). It therefore appears that while these biochemical correlates might be associated with enhanced survival of freezing, they do not explain the physiological mechanisms of the ability to tolerate internal ice.

More recent work has focused on the role of proteins in freeze tolerance. Antifreeze proteins may protect against intracellular freezing, which is almost always lethal even in very freeze tolerant animals (Davies, 2014; but see review in Sinclair & Renault, 2010). In the extracellular spaces, they may inhibit recrystallization, keeping ice crystals small enough to avoid mechanical damage. However they have been described from only a few freeze tolerant species, and their role in freeze tolerance remains controversial (Duman, Bennett, Sformo, Hochstrasser, & Barnes, 2004; Toxopeus & Sinclair, 2018). Aquaporins facilitate the movement of water out of the cell during freezing and glycerol transporters to facilitate the movement of sugars into the cell have been reasonably well-explored in plant freeze tolerance (Peng, Arora, Li, Wang, & Fessehaie, 2008; Philip, Yi, Elnitsky, & Lee, 2008; Storey & Storey, 2013; Yi et al., 2011). In animals, aquaporins are implicated in the freeze tolerance in *E. solidaginis* and the wood frog (*Lithobates sylvatica* LeConte), but how taxonomically broadly important they are for freeze tolerance is not currently well understood (Philip, Kiss, & Lee, 2011; Philip et al., 2008; Storey

& Storey, 2013; Yi et al., 2011). Finally, heat shock proteins are constitutively upregulated in overwintering insects and have been implicated in plant freeze tolerance, but are not generally thought to be involved in responses to acute freezing stress in animals (King & MacRae, 2015; Rinehart et al., 2006). However, the Antarctic midge (*Belgica antarctica* Jacobs) upregulates Hsp70 in response to repeated freezing episodes, and freeze tolerant drosophilid fly (*Chymomyza costata* Zetterstedt) upregulate small heat shock proteins following freezing suggest, suggesting that heat shock proteins may be more important to freeze tolerance than previously thought (Teets, Kawarasaki, Lee, & Denlinger, 2011).

While the biochemical correlates of freeze tolerance have been explored in broad array of terrestrial organisms, very little is known about the biochemical mechanisms of freeze tolerance in intertidal organisms (Dennis, Loomis, & Hellberg, 2014; Murphy, 1983). In temperate and polar regions, intertidal organisms including barnacles, mussels, and snails are almost without exception freeze tolerant (Murphy, 1983; Waller, Worland, Convey, & Barnes, 2006; but see Sinclair, Marshall, Singh, & Chown, 2004). This is necessary because of the unique nature of the intertidal—at low tide in the winter, the wetness of the habitat and the year-round presence of food in the digestive tract makes freezing a constant threat when animals are emersed. Some efforts have been made to determine the underlying biochemical mechanisms, but they remain elusive (Ansart & Vernon, 2003; Dennis et al., 2014; Murphy, 1983; Storey & Storey, 2013). It has been hypothesized that anaerobic byproducts such as strombine and taurine may be important in freeze tolerance of *Mytilus sp.* mussels (Loomis, Carpenter, & Crowe, 1988), but much remains unknown about how intertidal species survive freezing.

Here we present the first (to our knowledge) transcriptomic study following freezing stress in an intertidal animal. We explore potential mechanisms of freeze tolerance in the acorn barnacle (*Semibalanus balanoides* Linnaeus) from two populations that differ widely in the intensity of freezing stress, one from the central coast of British Columbia, Canada which experience relatively mild winters, and one from the White Sea in Russia which experiences significantly harsher winters. We hypothesize that barnacles from British Columbia will be less freeze tolerant (will have lower survival following freezing) than barnacles from the White Sea. If this is true, we will infer that transcriptional responses to freezing in the White Sea population likely enhance freeze tolerance. In addition, if transcripts upregulated in this population are important for survival in poleward range edge habitats, we predict that we will see evidence of selective sweeps on transcripts identities that are associated with biochemical mechanisms of freeze tolerance and are upregulated following freezing stress. We find strong interpopulation differences in response to freezing stress, and that this includes wholesale transcription of heat shock proteins following freezing stress, as well as the transcription of aquaporins (which show significant evidence of selective sweeps) and potentially new classes of antifreeze proteins. This suggests that these protein-based mechanisms of freeze tolerance may be broadly universal across freeze tolerant phyla, and that they may have selective advantages in intertidal environments.

Methods

Study system and collection

The barnacle *S. balanoides* is a common species in rocky intertidal zones through the Arctic Circle, including northwestern Europe and both coasts of North America. In western North America, it is found as far south as the central coast of British Columbia (Flight & Rand,

2012; Schmidt, Bertness, & Rand, 2000; Wethey, 1983). Because of its polar distribution as well as its intertidal habitat, *S. balanoides* regularly experiences freezing events throughout the winter (Cook & Gabott, 1970; Cook & Lewis, 1971; Dennis J. Crisp & Ritz, 1967). Late-stage larvae (cyprids) of *S. balanoides* have even been found embedded in ice during winter, and successfully settle and metamorphose after spring thaw (Pineda, DiBacco, & Starczak, 2005). Barnacles do not appear to use polyol cryoprotectants for their freeze tolerance, but little is known about the mechanisms of their freeze tolerance (Cook & Gabott, 1970; D.J. Crisp, Davenport, & Gabbott, 1977).

We collected *S. balanoides* from mid intertidal habitats by gathering small stones covered with large groups of barnacles at the White Sea Biological Station (Moscow State University, Russia; 66.5551° N, 34.1976° E) as well as from Calvert Island, British Columbia, Canada (51.5559° N, 128.0365° W) during the spring of 2015. The climate on Calvert Island is moderated by the Alaska Current which brings warm water from the south, and thus winter temperatures rarely drop below -10 °C, while in the White Sea overwinter temperatures regularly drop below -20 °C (Figure S1-S2), thus we expected barnacles from the White Sea be more tolerant of freezing. We have installed robobarnacles (Chan, Lima, Williams, Seabra, & Wang, 2016) in the White Sea to record the estimated body temperature of similarly-shaped *Tetraclita* spp. barnacles during the winter months (Figure S1). Barnacles for transcriptomic experiments were collected in June in both habitats, wrapped in wet newspaper, then were placed in a common seawater aquarium maintained at 10 °C and aerated by an airstone for two weeks before experimentation.

Experimental design and freezing exposures

After acclimation, we measured barnacle test diameter with digital calipers along the longest edge to the nearest mm. To simulate an intertidal freezing event, we cooled barnacles by placing the rocks they had settled on into a Styrofoam box, then placed the box in a freezer set to -20 °C. We continuously monitored barnacle temperature by placing a 36 AWG gauge Type T (copper-constantan) thermocouple through the plates into the barnacle body. The thermocouple was attached to a Picotech TC-08 interface (Pico Technology, Cambridge UK) which relayed temperature data to an attached computer every 0.5 s in real time. We maintained barnacle temperature to within 0.5 °C of the desired temperature by opening and closing the freezer door. Barnacle supercooling points (SCPs) were recorded as the lowest temperature attained before sudden release of heat due to the latent heat of crystallization (Sinclair, Coello Alvarado, & Ferguson, 2015). With this procedure, we achieved cooling rates of circa 1.2 °C/min, which is within the range found in their natural habitat during a low tide (Figure S1). To measure differences in freeze tolerance, we exposed barnacles from both populations to -10 or -6 °C for 4 h, then warmed them by placing them immediately into seawater at 10 °C to mimic inundation during high tide conditions. Survival was monitored for two days following and scored based on the ability to close the top plates when removed from seawater, then compared using generalized linear models with a binomial distribution in R (Team, 2016), while SCPs were compared using ANOVA.

To measure transcriptional changes following freezing, we exposed 48 barnacles from each population to -6 °C for 4 h using the above procedure. We sampled groups from each population at four different time points: before the freezing event took place (“control”), after the freezing event but before thawing in seawater (“before thawing”), 4 h after thawing (“4h

recovery”) and 28 h after thawing (“28h recovery”; experimental design presented in Figure 1). For each population $\times$ timepoint, we rapidly dissected out the soft tissue of 12 barnacles from each population and flash-froze tissue in groups of three individuals in liquid nitrogen for four total biological replicates/population/timepoint.

Transcriptomics

Total RNA was extracted from 32 pools of barnacles using a TRIzol protocol. Sequencing libraries were prepared and sequenced by the Academia Sinica Bioinformatics Center. Libraries were prepared using Illumina’s stranded mRNA TruSeq RNA sample prep v2 kit as per manufacturer’s instructions. Cleaned libraries were sequenced on 2 lanes of 181 bp paired-end Illumina HiSeq 2000 at the Academia Sinica Sequencing Center.

Our general approach for data analysis was informed by best practices for RNA-seq (Conesa et al., 2016). The preliminary data quality checks and analysis were conducted in Galaxy (Afgan et al., 2016). We clipped Illumina adapters from reads and removed low quality and short read sequences using Trimmomatic’s default settings (version 0.36.3; Bolger, Lohse, & Usadel, 2014). We then took the trimmed reads and assembled a transcriptome using Trinity (default settings for paired end reads, version 2.4.0, Grabherr et al., 2011), then clustered redundant sequences using CD-HIT-EST with a sequence identity cut-off of 0.9 (Fu, Niu, Zhu, Wu, & Li, 2012). We then aligned these sequences to the NCBI NR database using Diamond (Buchfink, Xie, & Huson, 2014), then filtered out all reads that did not map to metazoans (viruses, algae, bacteria, etc.) using MEGAN6 (Huson et al., 2016) to produce a *de novo* transcriptome for *S. balanoides*. To annotate the transcriptome, we used Blast2GO (default

settings, version 5.1.13, Gotz et al., 2008) to blastx the transcripts to the NCBI arthropod nr database, then mapped to Interpro IDs and GO terms to produce a final, annotated transcriptome.

To quantify gene expression following a freezing event, we then mapped our trimmed reads to this transcriptome using HISAT2 (D. Kim, Langmead, & Salzberg, 2015), then assembled and quantified transcripts using StringTie (Pertea et al., 2015), gffCompare (Trapnell et al., 2010), and FeatureCounts (Liao, Smyth, & Shi, 2014). Following quantification, we tested for differences in transcript abundance among time points and populations using DESeq2 (Love, Huber, & Anders, 2014) to identify differentially expressed transcripts (adjusted p-value < 0.05) within each population as a function of timepoint, while controlling for multiple comparisons using the Benjamini-Hochberg multiple comparisons procedure.

We then used Blast2GO to test for functional enrichment of the significantly differentially regulated transcripts within each population. We simplified the significantly enriched GO list for plotting using REVIGO to eliminate semantically redundant terms, using the “tiny” setting (Supek, Bošnjak, Škunca, & Šmuc, 2011). We also annotated each sequence with the nearest KEGG orthology term using the KEGG Automatic Annotation Server (Moriya, Itoh, Okuda, Yoshizawa, & Kanehisa, 2007), then tested for KEGG pathway enrichment using the Gage and Pathview (Luo & Brouwer, 2013) packages in R.

Population differentiation

To quantify genetic differences between the two geographic sites, single nucleotide polymorphisms (SNPs) were called using ANGSD 0.916 (Korneliussen, Albrechtsen, & Nielsen, 2014). The ‘before thawing’ samples were removed from the analysis as their transcript

expression profiles and therefore individual transcripts read coverage levels, differed markedly from the other samples. Mapped reads were first filtered to remove PCR duplicates and secondary mapping. Within ANGSD missing values were assigned for base calls of a phred score <30. Loci were limited to those that had a depth of at least 10 reads per sample and an across all samples Minor Allele Frequency (MAF) estimate of >0.05. As each sample is comprised of three individuals (pooled) genotypes could not be called, instead sample MAF were estimated using the allele counts method in ANGSD. Sample MAF estimates were used to conduct a Principle Components Analysis (PCA) within R (prcomp function; PCA presented in Figure S3).

We tested whether there was evidence for selection on the differentially expressed transcripts at either of the two geographic sites. As with the MAF estimates, the ‘before thawing’ samples, PCR duplicates, and secondary mapping were removed. Samples were grouped into two groups by geographic site, with each of the two groups were comprised of 36 individuals total (12 samples of three individuals each). A pileup file was generated and Tajima’s D (Tajima, 1989) estimates were generated for both locations using PoPoolation (Robert Kofler et al., 2011). Within PoPoolation missing values were assigned for base calls of a phred score <20 and a corrected Tajima’s D was calculated for both geographic sites. Tajima’s D estimates were compared between the two sites within R (via a T-test). To further examine genetic differentiation between the two geographic sites. PoPoolation2 (R. Kofler, Pandey, & Schlotterer, 2011) was also used to estimate Fisher Exact Tests (per site) and F_{ST} values (per site and sliding window).

Results

Assessing freeze tolerance

Supercooling point (mean = -3.2 ± 0.51 °C) did not differ between barnacle populations, nor did SCP differ with basal diameter of test ($p > 0.4$ for both terms, $n = 8 - 15$ per population). Barnacles from the White Sea had significantly higher survival at both exposure temperatures (81 – 95% vs. 27 – 78%; Figure 2; deviance explained = 48.4, $df = 1, 18$, $p < 0.001$), and exposure to -10 °C significantly decreased survival in both populations relative to exposure at -6 °C (deviance explained = 74.8, $df = 1, 17$, $p < 0.001$; $n = 92 - 152$ for each combination).

Sequencing and assembly

A total of 776.69 million 181 bp reads were produced from two lanes of Illumina Hiseq 2000, with an average of 24.75 million/library for the White Sea population and 23.79 million/library for the British Columbia population. Of these, an average of 14.2 million reads/library in the White Sea population and 13.1 million reads/library in the British Columbia population passed quality control by Trimmomatic. These reads were then used to assemble a *de novo* transcriptome using Trinity, which was then reduced to 61,881 sequences using CD-HIT-EST. Of these, a total of 61,779 were matched to taxonomic hits by Diamond. We then filtered non-metazoan sequences (RNA viruses, algae, and other microorganisms) out using MEGAN, and were left with a total of 46,535 sequences that were an average of 1277 base pairs long ($N_{50} = 1084$ bp). An average of 4.84 million reads/library in the White Sea population and 3.63 million reads/library in the British Columbia population were mapped to the final transcriptome.

Pattern of transcript expression

We found the patterns of significantly differentially expressed transcripts after a freezing event differed strongly between the two populations (Figure 3, Figure 4). In the more freeze tolerant

population (White Sea), no transcripts were differentially expressed during freezing, while after four hours of recovery, 191 transcripts were differentially expressed (almost evenly split between up and downregulated). By 28 hours after freezing, this had decreased to 185 transcripts, with slightly more upregulated than down. This pattern was dramatically different in the less freeze tolerant population (British Columbia), which had a total of 11,050 transcripts differentially regulated during the freezing event, the majority of which (9432) were significantly upregulated. By four hours of recovery, this number had decreased to 51 (again dominated by 46 upregulated transcripts), and by 28 hours of recovery this had increased to 1739 transcripts (again dominated by 977 significantly upregulated transcripts).

Identities of differentially expressed transcripts

The identities of the top differentially expressed transcripts following freezing were functionally very different between the two populations. The few transcripts that both populations upregulated immediately following freezing were putative transcription factors (annotated by Blast2Go as “proto-oncogenes”; Table 1). In the British Columbia population, the most noticeable pattern is the massive upregulation of transcripts during the freezing event (>9000 transcripts, Figures 3 and 4). This included several ribosomal RNAs, mitochondrially-encoded electron transport chain subunits, and histone methyltransferases and deacetylases (Supplementary Data 1). Among the downregulated transcripts includes several that Blast2Go annotated as “macrophage mannose receptors” which are functionally assigned the “carbohydrate binding” GO term (Supplementary Data 1). At four hours following freezing, this pattern has largely disappeared, with only a total of 46 significantly upregulated transcripts (Figure 3 and 4). These include several transcripts that Blast2Go identified as serine proteases, while the

downregulated transcripts include one identified as “Peter Pan”, a transcript annotated as “macrophage mannose receptor”, and an INO80 complex subunit (Supplementary Data 1). By 28 hours following freezing, differential expression has increased again (Figure 3 and 4), with a total of 50 upregulated transcripts that BLAST to likely proteases and 13 that BLAST to ubiquitin activity, suggesting significant protein scavenging. By contrast, in the White Sea population, a large number of transcripts associated with heat shock proteins were the most highly upregulated transcripts following freezing (Table 2). Two transcripts that were identified as E3 ubiquitin ligases were also upregulated at both 4 and 28 hours following freezing (Supplementary Data 1). Four and 28 hours following freezing, the down-regulated transcripts included a variety of transcripts that perhaps most notably included several mitochondrially-encoded electron transport chain subunits (Supplementary Data 1) including NADH dehydrogenase subunits 1, 2, 4, 5, and 6, cytochrome b, ATP synthase F₀ subunit 6, as well as cytochrome c oxidase subunits 1, 2, and 3 (Supplementary Data 1). No nuclear-encoded electron transport chain subunits were downregulated in our dataset.

Functional enrichment of molecular functions of differentially expressed transcripts

The functions of the transcripts similarly differed significantly between the two populations (Tables 2, 3, Figure 5). In the British Columbia population, we found significant enrichment of 134 GO terms after REVIGO filtering in the downregulated and 80 in the upregulated transcripts during freezing, including several electron transport chain-related terms in both sets (Figure 5). By four hours following freezing there was no significant enrichment of GO terms, and by 28 hours following freezing there was significant enrichment of 56 GO terms in the downregulated transcripts with only 15 in the upregulated transcripts (Figure 5). The most

notable of these enriched terms included “UDP-glucose-6-dehydrogenase activity” and “scavenger receptor activity”. By contrast, in the White Sea population, as we had no significant differential expression of transcripts during freezing, we did not find any significantly enriched GO terms at this timepoint. While we did not find any significant enrichment of GO terms among upregulated transcripts in the White Sea population four hours following recovery, we did find a series of terms associated with the electron transport chain including “oxidoreductase activity”, “heme-copper terminal oxidase activity”, and “hydrogen ion transmembrane activity” that were significantly enriched among transcripts that were significantly downregulated at this timepoint (Figure 5). Interestingly, all of these downregulated terms were associated with transcripts that encoded only mitochondrially-encoded electron chain subunits. By 28 hours of recovery, the pattern had changed significantly with no enriched GO terms among the downregulated transcripts, but among the significantly upregulated transcripts the terms “glycerol transport”, “cellular water homeostasis”, and “water transport” were significantly enriched (which all were annotated as aquaporins).

Using pathview and Gage to examine significantly enriched gene sets based on KEGG orthology we found additional functional enrichment. In the British Columbia population, before thawing we found significant enrichment of the “spliceosome” term and underrepresentation of the “epithelial cell signaling in *Helicobacter pylori*” term. Four hours after thawing, we observed significant overrepresentation of the terms “ribosome biogenesis in eukaryotes” and “spliceosome”, and underrepresentation of “Parkinson’s disease”, “Alzheimer’s disease” and “oxidative phosphorylation”, although we note upon examination that the first two terms all indicated the same underrepresentation of electron transport chain subunits. There was no

functional enrichment 28 hours following freezing in the British Columbia population. Finally, in the White Sea population we observed significant overrepresentation of terms including “ribosome biogenesis in eukaryotes”, “spliceosome”, and “bacterial invasion of epithelial cells” at 28 hours following freezing, but no terms were overrepresented at any other timepoints.

Population differentiation

The estimated F_{ST} value (0.45; 9,164 loci passing filter) suggests little gene flow occurs between the two populations. Across all differentially expressed transcripts Tajima D estimates averaged -1.2 and -1.4 for White Sea and British Columbia respectively, whereas non-differentially expressed transcripts averaged -0.6 (T-test p-value <0.001) and -0.8 (T-test p-value <0.001) for White Sea and British Columbia respectively, suggesting selective sweeps on these differentially-expressed transcripts in each population (Table 4). Tajima’s D estimates for aquaporins in the White Sea population averaged -2.6 compared to the average across differently expressed transcripts of -1.42, indicative of a potential selective sweep in these transcripts (Table 4).

Discussion

Intertidal organisms constantly risk freezing through the winter. We find, consistent with our hypothesis, that the northern White Sea population of *S. balanoides* is significantly more freeze tolerant than the British Columbia population. To our knowledge, there is currently only one genome-sequenced freeze tolerant animal (the Antarctic midge *Belgica antarctica*; Kelley et al., 2014) and no genome-sequenced barnacles, so we assembled a *de novo* transcriptome to study transcriptional responses to freezing stress. In addition, the differing timelines of transcription suggest two very different sets of processes taking place in each population following a freezing

event, which adds to our limited understanding of the molecular mechanisms of freeze tolerance. Finally, the low Tajima's D values on these differentially expressed transcripts suggests that local adaptation to freezing has taken place in the White Sea population. Here we have shown the first study of transcriptional responses following a freezing stress in a non-insect, and demonstrated evidence of selection on transcripts associated with freeze tolerance in a natural population.

One of the most puzzling results we found is the massive upregulation of transcripts during the freezing exposure in the British Columbia population (Figures 3 and 4). There are three possibilities to explain this pattern: this represents genuine upregulation of transcription, degradation of mRNA causing greater numbers of smaller transcripts, or damage to translational machinery causing mRNA to accumulate. The first hypothesis of increased rates of *de novo* mRNA transcription is possible since the cytoplasm of cells generally remains unfrozen during a freezing exposure. However this is difficult to reconcile with the generally-accepted rate of transcription in invertebrates of about 2kb/minute (Ardehali & Lis, 2009), particularly at a subzero temperature where biochemical events would be expected to proceed significantly more slowly. This upregulation is also strikingly absent in the White Sea population, suggesting increased transcription rates during freezing is not a general nor adaptive response to freezing stress. The second hypothesis of degradation of mRNA due to osmotic shock or direct freeze-induced damage is certainly possible, which could increase the total number of detected mRNA transcripts due to cleavage of larger molecules into several smaller ones. However, we note that the average transcript length of these upregulated transcripts is significantly larger (1761 bp vs. 1398 bp for the whole transcriptome), which is inconsistent with the possibility of mRNA degradation. We also note that, at least in *Arabidopsis thaliana*, low temperature tends to

increase mRNA half-life (Chiba et al., 2013). The third hypothesis, cold-induced damage to ribosomal structure or translation initiation machinery, is possible since translation is associated with mRNA degradation (Bicknell & Ricci, 2017). At least in bacteria, the ribosome is particularly sensitive to cold shock and several cellular mechanisms for surviving cold shock are focused on ribosomal proteins (Phadtare, 2004). In addition, we note several ribosomal mRNAs are among the top-upregulated transcripts in this experimental group, as well as several transcripts associated with eukaryotic translation initiation machinery (Supplementary Data 1). Finally, we also note that at 4 hours following freezing, we find that the KEGG term “ribosome biogenesis” is significantly enriched among upregulated transcripts in the British Columbia population, suggesting repair or rebuilding of ribosomal subunits (Table S1). If the ribosomes are damaged and unable to process mRNA, we might expect a buildup of mRNA (Bicknell & Ricci, 2017). If this is true, we would expect that the White Sea population produces ribosomes that are more cold shock resistant. Further functional studies, in addition to further transcriptomic work in freeze tolerant animals, is clearly necessary to differentiate among these potential explanations.

The role(s) of heat shock proteins in freeze tolerance is not well understood, but upregulation of Hsp70 has been documented in response to repeated freezing in *B. antarctica* (Teets et al., 2011). While cold-induced denaturation of proteins has been well studied (Graziano, 2010; Todgham, Hoaglund, & Hofmann, 2007), during survivable freezing significant desiccation of cells is believed to occur (Storey, 1997; Teets & Denlinger, 2014). It is possible that this desiccation leads directly to protein denaturation through the disruption of the hydration shell around proteins during desiccating periods. In the more freeze tolerant White Sea population, we observed significant upregulation of several shock proteins following freezing—it

is unclear whether these are involved with assisting in refolding of damaged proteins or are preparatory for a second stress event. By contrast, the less freeze tolerant population upregulated proteolysis-related transcripts (with significant enrichment of GO terms such as “serine endopeptidase activity” and “scavenger receptor”) following freezing, suggesting significant damage to protein structure has taken place and the only possible solution is to degrade these proteins. Our study points to the importance of maintenance of protein stability at low temperature in an ecological setting.

In addition to direct effects on protein structure, we also saw very different patterns of differential regulation of aquaporins between the two populations following freezing. In the White Sea population, we saw significant enrichment of GO categories associated with aquaporins, which we did not observe in the British Columbia population. This suggests that only the White Sea population is generally upregulating aquaporin transcription following freezing. It is believed that aquaporins aid in water transport out of the cell and that aquaglyceroporins may assist in transport of glycerol into the cell during freezing events (Philip et al., 2008; Storey & Storey, 2013; Yi et al., 2011). While significant glycerol accumulation is not associated with freeze tolerance in marine invertebrates (Storey & Storey, 2013), it is likely that managing water content in the cell is essential for freezing survival. Thus far aquaporins have been associated with freeze tolerance in the wood frog *Rana sylvatica* (Philip et al., 2008) and the goldenrod gall fly *Eurosta solidaginis* (Yi et al., 2011), and with this study we demonstrate their importance in a new class of freeze tolerant organisms, suggesting that this may be a generalized mechanism for freeze tolerance. In addition, we found very low Tajima’s D values in aquaporins that were upregulated following freezing (Table 4), suggesting that selective

sweeps had taken place on the sequence of these putative aquaporins. Follow-up with functional studies, particularly employing the frog oocyte expression system, would provide interesting evidence and further structural information in this poorly understood system (Lind et al., 2017).

The role of antifreeze proteins in freeze tolerance is unclear and little is generally known about invertebrate antifreeze proteins (Davies, 2014; H. J. Kim et al., 2017). We found many transcripts (>200) that were annotated with the term “macrophage mannose receptors” that were upregulated following freezing in both populations. While we cannot definitively assign a function to these based solely on a transcriptomic study, we do note that these are C-type lectins, which are believed to be paralogous to Type II antifreeze proteins in fish (Graham, Loughheed, Ewart, & Davies, 2008; Ng & Hew, 1992). Indeed, when we examined the BLAST results, many of these sequences had high similarity to Type II antifreeze or ice structuring proteins in fish. Several freeze tolerant species are thought to produce antifreeze proteins to inhibit recrystallization of ice at low temperatures (Davies, 2014; Storey & Storey, 2013), although functional studies are necessary for confirming ice recrystallization inhibition activity. Antifreeze proteins are found broadly scattered across the domains of life, and appear to readily evolve (Davies, 2014). In addition, they appear to frequently have been transferred across species through horizontal gene transfer (Davies, 2014; Kiko, 2010). While this study was not designed to differentiate between the hypotheses of similar structure due to convergent evolution or by horizontal gene transfer, further sequencing studies may be able to address this point.

One of the most consistent results we found in both populations is downregulation of transcripts associated with the electron transport chain. There are two potential explanations for this downregulation—either transcription of mitochondrially-encoded transcripts was inhibited,

or the number of mitochondria themselves were downregulated following a freezing stress. This is likely an adaptive response since a degraded or malfunctioning electron transport chain can cause significant oxidative stress, as can a sudden reduction in ATP demand as we might expect at low temperatures (Somero, Lockwood, & Tomanek, 2017). Reducing mitochondrial numbers as a way to reduce metabolic rate is a seasonal adaptation in the overwintering larvae of the moth *Gynaephora groenlandica* (Kukal, Duman, & Serianni, 1989), and this may be a similar phenomena. While we cannot distinguish between these explanations in this study, we hypothesize that reducing mitochondrial activity facilitates freeze tolerance in our barnacles and suggest studies isolating mitochondria to measure mitochondrial respiration and histology to determine mitochondrial numbers following freezing stress.

Conclusions

Here we have presented the first transcriptional time series following freezing in a freeze tolerant barnacle. By comparing two populations that differ strongly in their freeze tolerance, we can control for genetic background and thereby infer the transcripts that are associated with greater freeze tolerance. We find transcriptional evidence for protein damage during a freezing event, and suggest that heat shock protein upregulation appears to be important for recovery from freezing. We also document aquaporin expression associated with freezing in a third freeze tolerant species, and describe a series of new putative antifreeze proteins. These transcripts were associated with selective sweeps in the White Sea population of barnacles, suggesting local adaptation to freezing conditions has occurred. Taken together, we shed new light on the mechanisms of adaptation to the poleward range margin in animals.

Acknowledgements

The authors would like to acknowledge the Company of Biologists for travel funding to KEM, a University of Oklahoma startup grant and Nigel Fish for computing resources for KEM, as well as Taiwan-Russian collaboration funds from MOST, Taiwan (106-2923-B-001 -002 -MY3) and RFBR (grants 17-54-52006 MNT_a, 18-04-00624 A, 15-29-02447 ofi_m) and Academia Sinica Senior Investigator Award (AS-IA-105-L03) to BKKC for funding for sequencing. We would like to thank the Hakai Institute and Mary O'Connor for collecting barnacles in British Columbia, as well as for access to temperature data. We would also like to acknowledge Yao-Fong Tsao and Vanessa Pei-Chen Tsai for laboratory assistance in sample preparation, and Dr. Mei-Yeh Lu for her invaluable advice for library preparation. Finally, we would like to thank Jantina Toxopeus and two anonymous referees for comments that have greatly improved the manuscript.

References

- Afgan, E., Baker, D., van den Beek, M., Blankenberg, D., Bouvier, D., Čech, M., ... Goecks, J. (2016). The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Research*, 44(W1), W3–W10. doi:10.1093/nar/gkw343
- Ansart, A., & Vernon, P. (2003). Cold hardiness in molluscs. *Acta Oecologica*, 24(2), 95–102. doi:10.1016/S1146-609X(03)00045-6
- Ardehali, M. B., & Lis, J. T. (2009). Tracking rates of transcription and splicing in vivo. *Nature Structural & Molecular Biology*, 16(11), 1123–1124. doi:10.1038/nsmb1109-1123
- Bicknell, A. A., & Ricci, E. P. (2017). When mRNA translation meets decay. *Biochemical Society Transactions*, 45(2), 339–351. doi:10.1042/BST20160243
- Boardman, L., Terblanche, J. S., & Sinclair, B. J. (2011). Transmembrane ion distribution during recovery from freezing in the woolly bear caterpillar *Pyrrharctia isabella* (Lepidoptera: Arctiidae). *Journal of Insect Physiology*, 57(8), 1154–1162. doi:10.1016/j.jinsphys.2011.04.022
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, 30(15), 2114–20. doi:10.1093/bioinformatics/btu170
- Buchfink, B., Xie, C., & Huson, D. H. (2014). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, 12(1), 59–60. doi:10.1038/nmeth.3176
- Chan, B. K. K., Lima, F. P., Williams, G. A., Seabra, R., & Wang, H. Y. (2016). A simplified biomimetic temperature logger for recording intertidal barnacle body temperatures. *Limnology and Oceanography: Methods*, 14(7), 448–455. doi:10.1002/lom3.10103
- Chiba, Y., Mineta, K., Hirai, M. Y., Suzuki, Y., Kanaya, S., Takahashi, H., ... Naito, S. (2013). Changes in mRNA Stability Associated with Cold Stress in Arabidopsis Cells. *Plant and Cell Physiology*, 54(2), 180–194. doi:10.1093/pcp/pcs164

- 519 Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A., ...
520 Mortazavi, A. (2016). A survey of best practices for RNA-seq data analysis. *Genome*
521 *Biology*, 17(1), 13. doi:10.1186/s13059-016-0881-8
- 522 Cook, P. A., & Gabott, P. A. (1970). Seasonal changes in the free glycerol level in the body parts
523 of the adult barnacle *Balanus balanoides*. *Marine Biology*, 7(1), 11–13.
524 doi:10.1007/BF00346803
- 525 Cook, P. A., & Lewis, A. H. (1971). Acquisition and loss of cold-tolerance in adult barnacles
526 (*Balanus balanoides*) kept under laboratory conditions. *Marine Biology*, 9(1), 26–30.
527 doi:10.1007/BF00348814
- 528 Crisp, D. J., Davenport, J., & Gabbott, P. A. (1977). Freezing tolerance in *Balanus balanoides*.
529 *Comparative Biochemistry and Physiology Part A: Physiology*, 57(3), 359–361.
530 doi:10.1016/0300-9629(77)90206-7
- 531 Crisp, D. J., & Ritz, D. A. (1967). Changes in temperature tolerance of *Balanus balanoides*
532 during its life-cycle. *Helgoländer Wissenschaftliche Meeresuntersuchungen*, 15(1–4), 98–
533 115. doi:10.1007/BF01618612
- 534 Davies, P. L. (2014). Ice-binding proteins: A remarkable diversity of structures for stopping and
535 starting ice growth. *Trends in Biochemical Sciences*, 39(11), 548–555.
536 doi:10.1016/j.tibs.2014.09.005
- 537 Dennis, A. B., Loomis, S. H., & Hellberg, M. E. (2014). Latitudinal Variation of Freeze
538 Tolerance in Intertidal Marine Snails of the Genus *Melampus* (Gastropoda: Ellobiidae).
539 *Physiological and Biochemical Zoology*, 87(4), 517–526. doi:10.1086/676138
- 540 Duman, J. G., Bennett, V., Sformo, T., Hochstrasser, R., & Barnes, B. M. (2004). Antifreeze
541 proteins in Alaskan insects and spiders. *Journal of Insect Physiology*, 50(4), 259–266.
542 doi:10.1016/J.JINSPHYS.2003.12.003
- 543 Flight, P. A., & Rand, D. M. (2012). Genetic variation in the acorn barnacle from allozymes to
544 population genomics. *Integrative and Comparative Biology*, 52(3), 418–429.
545 doi:10.1093/icb/ics099

- 546 Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: accelerated for clustering the next-
547 generation sequencing data. *Bioinformatics*, 28(23), 3150–3152.
548 doi:10.1093/bioinformatics/bts565
- 549 Gotz, S., Garcia-Gomez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., ...
550 Conesa, A. (2008). High-throughput functional annotation and data mining with the
551 Blast2GO suite. *Nucleic Acids Research*, 36(10), 3420–3435. doi:10.1093/nar/gkn176
- 552 Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., ... Regev, A.
553 (2011). Full-length transcriptome assembly from RNA-Seq data without a reference
554 genome. *Nature Biotechnology*, 29(7), 644–52. doi:10.1038/nbt.1883
- 555 Graham, L. A., Loughheed, S. C., Ewart, K. V., & Davies, P. L. (2008). Lateral transfer of a
556 lectin-like antifreeze protein gene in fishes. *PLoS ONE*, 3(7), e2616.
557 doi:10.1371/journal.pone.0002616
- 558 Graziano, G. (2010). On the molecular origin of cold denaturation of globular proteins. *Physical*
559 *Chemistry Chemical Physics*, 12(42), 14245. doi:10.1039/c0cp00945h
- 560 Han, E. N., & Bause, E. (1995). Non-Freezing Survival of Spruce Budworm Larvae,
561 *Choristoneura-fumiferana*, at Subzero Temperatures during Diapause. *Entomologia*
562 *Experimentalis Et Applicata*, 75(1), 67–74.
- 563 Huson, D. H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., ... Tappu, R. (2016).
564 MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale
565 Microbiome Sequencing Data. *PLOS Computational Biology*, 12(6), e1004957.
566 doi:10.1371/journal.pcbi.1004957
- 567 Johnston, I. A., & Clarke, A. (1990). Cold Adaptation in Marine Organisms [and Discussion].
568 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 326(1237), 655–
569 667. doi:10.1098/rstb.1990.0037
- 570 Kelley, J. L., Peyton, J. T., Fiston-Lavier, A.-S., Teets, N. M., Yee, M.-C., Johnston, J. S., ...
571 Denlinger, D. L. (2014). Compact genome of the Antarctic midge is likely an adaptation to
572 an extreme environment. *Nature Communications*, 5(1), 4611. doi:10.1038/ncomms5611

- 573 Kiko, R. (2010). Acquisition of freeze protection in a sea-ice crustacean through horizontal gene
574 transfer? *Polar Biology*, 33(4), 543–556. doi:10.1007/s00300-009-0732-0
- 575 Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low
576 memory requirements. *Nature Methods*, 12(4), 357–60. doi:10.1038/nmeth.3317
- 577 Kim, H. J., Lee, J. H., Hur, Y. B., Lee, C. W., Park, S.-H., & Koo, B.-W. (2017). Marine
578 Antifreeze Proteins: Structure, Function, and Application to Cryopreservation as a Potential
579 Cryoprotectant. *Marine Drugs*, 15(2). doi:10.3390/md15020027
- 580 King, A. M., & MacRae, T. H. (2015). Insect Heat Shock Proteins During Stress and Diapause.
581 *Annual Review of Entomology*, 60(1), 59–75. doi:10.1146/annurev-ento-011613-162107
- 582 Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., ...
583 Schlötterer, C. (2011). PoPoolation: A Toolbox for Population Genetic Analysis of Next
584 Generation Sequencing Data from Pooled Individuals. *PLoS ONE*, 6(1), e15925.
585 doi:10.1371/journal.pone.0015925
- 586 Kofler, R., Pandey, R. V., & Schlotterer, C. (2011). PoPoolation2: identifying differentiation
587 between populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics*,
588 27(24), 3435–3436. doi:10.1093/bioinformatics/btr589
- 589 Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next
590 Generation Sequencing Data. *BMC Bioinformatics*, 15(1), 356. doi:10.1186/s12859-014-
591 0356-4
- 592 Košťál, V., Korbelová, J., Poupardin, R., Moos, M., & Šimek, P. (2016). Arginine and proline
593 applied as food additives stimulate high freeze tolerance in larvae of *Drosophila*
594 *melanogaster*. *The Journal of Experimental Biology*, 219(15), 2358–2367.
595 doi:10.1242/jeb.142158
- 596 Kukal, O., Duman, J. G., & Serianni, A. S. (1989). Cold-induced mitochondrial degradation and
597 cryoprotectant synthesis in freeze-tolerant arctic caterpillars. *Journal of Comparative*
598 *Physiology B*, 158(6), 661–671. doi:10.1007/BF00693004

- 599 Layne, J. R., & Kuharsky, D. K. (2000). Triggering of cryoprotectant synthesis in the woolly bear
600 caterpillar (*Pyrrharctia isabella* Lepidoptera: Arctiidae). *Journal of Experimental Zoology*,
601 286(4), 367–371.
- 602 Lee, R. E. (2010). A primer on insect cold tolerance. In D. L. Denlinger & R. E. Lee (Eds.), *Low*
603 *Temperature Biology of Insects* (pp. 3–34). Cambridge, UK: Cambridge University Press.
- 604 Lee, R. E., Dommell, R. A., Joplin, K. H., & Denlinger, D. L. (1995). Cryobiology of the freeze-
605 tolerant gall fly *Eurosta solidaginis*: overwintering energetics and heat shock proteins.
606 *Climate Research*, 5, 61–67. doi:10.3354/cr005061
- 607 Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program
608 for assigning sequence reads to genomic features. *Bioinformatics*, 30(7), 923–930.
609 doi:10.1093/bioinformatics/btt656
- 610 Lind, U., Järvå, M., Rosenblad, M. A., Pingitore, P., Karlsson, E., Wrangé, A. L., ... Blomberg,
611 A. (2017). Analysis of aquaporins from the euryhaline barnacle *Balanus improvisus* reveals
612 differential expression in response to changes in salinity. *PLoS ONE*, 12(7), e0181192.
613 doi:10.1371/journal.pone.0181192
- 614 Loomis, S. H., Carpenter, J. F., & Crowe, J. H. (1988). Identification of strombine and taurine as
615 cryoprotectants in the intertidal bivalve *Mytilus edulis*. *Biochimica et Biophysica Acta*,
616 943(2), 113–118. doi:10.1016/0005-2736(88)90542-1
- 617 Louthan, A. M., Doak, D. F., & Angert, A. L. (2015). Where and When do Species Interactions
618 Set Range Limits? doi:10.1016/j.tree.2015.09.011
- 619 Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
620 dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 550.
621 doi:10.1186/s13059-014-0550-8
- 622 Luo, W., & Brouwer, C. (2013). Pathview: an R/Bioconductor package for pathway-based data
623 integration and visualization. *Bioinformatics*, 29(14), 1830–1831.
624 doi:10.1093/bioinformatics/btt285

- Marshall, K. E., & Sinclair, B. J. (2011). The sub-lethal effects of repeated freezing in the woolly bear caterpillar *Pyrrharctia isabella*. *Journal of Experimental Biology*, 214(7), 1205–1212. doi:10.1242/jeb.054569
- Marshall, K. E., & Sinclair, B. J. (2015). The relative importance of number, duration and intensity of cold stress events in determining survival and energetics of an overwintering insect. *Functional Ecology*, 29(3), 357–366. doi:10.1111/1365-2435.12328
- Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C., & Kanehisa, M. (2007). KAAS: An automatic genome annotation and pathway reconstruction server. *Nucleic Acids Research*, 35(SUPPL.2), W182–W185. doi:10.1093/nar/gkm321
- Murphy, D. J. (1983). Freezing resistance in intertidal invertebrates. *Annual Review of Physiology*, 45, 289–299. doi:10.1146/annurev.ph.45.030183.001445
- Ng, N. F. L., & Hew, C. L. (1992). Structure of an antifreeze polypeptide from the sea raven: Disulfide bonds and similarity to lectin-binding proteins. *Journal of Biological Chemistry*, 267(23), 16069–16075. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1644794>
- Peng, Y., Arora, R., Li, G., Wang, X., & Fessehaie, A. (2008). Rhododendron catawbiense plasma membrane intrinsic proteins are aquaporins, and their over-expression compromises constitutive freezing tolerance and cold acclimation ability of transgenic *Arabidopsis* plants. *Plant, Cell and Environment*, 31(9), 1275–1289. doi:10.1111/j.1365-3040.2008.01840.x
- Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*, 33(3), 290–5. doi:10.1038/nbt.3122
- Phadtare, S. (2004). Recent developments in bacterial cold-shock response. *Current Issues in Molecular Biology*, 6, 125–136. Retrieved from <http://www.caister.com/cimb/v/v6/125.pdf>
- Philip, B. N., Kiss, A. J., & Lee, R. E. (2011). The protective role of aquaporins in the freeze-tolerant insect *Eurosta solidaginis*: functional characterization and tissue abundance of EsAQP1. *The Journal of Experimental Biology*, 214(Pt 5), 848–857. doi:10.1242/jeb.051276

- Philip, B. N., Yi, S.-X., Elnitsky, M. a, & Lee, R. E. (2008). Aquaporins play a role in desiccation and freeze tolerance in larvae of the goldenrod gall fly, *Eurosta solidaginis*. *The Journal of Experimental Biology*, 211(Pt 7), 1114–1119. doi:10.1242/jeb.016758
- Pineda, J., DiBacco, C., & Starczak, V. (2005). Barnacle larvae in ice: Survival, reproduction, and time to postsettlement metamorphosis. *Limnology and Oceanography*, 50(5), 1520–1528. doi:10.4319/lo.2005.50.5.1520
- Qin, W., Doucet, D., Tyshenko, M. G., & Walker, V. K. (2007). Transcription of antifreeze protein genes in *Choristoneura fumiferana*. *Insect Molecular Biology*, 16(4), 423–434. doi:10.1111/j.1365-2583.2007.00743.x
- Rinehart, J. P., Hayward, S. A. L., Elnitsky, M. A., Sandro, L. H., Lee, R. E., & Denlinger, D. L. (2006). Continuous up-regulation of heat shock proteins in larvae, but not adults, of a polar insect. *Proceedings of the National Academy of Sciences*, 103(38), 14223–14227. doi:10.1073/pnas.0606840103
- Schmidt, P. S., Bertness, M. D., & Rand, D. M. (2000). Environmental heterogeneity and balancing selection in the acorn barnacle *Semibalanus balanoides*. *Proceedings of the Royal Society B*, 267(1441), 379–384. doi:10.1098/rspb.2000.1012
- Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and Ecology of Species Range Limits. *Annu. Rev. Ecol. Evol. Syst*, 40, 415–36. doi:10.1146/annurev.ecolsys.110308.120317
- Sinclair, B. J., Addo-Bediako, A., & Chown, S. L. (2003). Climatic variability and the evolution of insect freeze tolerance. *Biological Reviews*, 78(2), 181–195. doi:10.1017/s1464793102006024
- Sinclair, B. J., Coello Alvarado, L. E., & Ferguson, L. V. (2015). An invitation to measure insect cold tolerance: Methods, approaches, and workflow. *Journal of Thermal Biology*. doi:10.1016/j.jtherbio.2015.11.003
- Sinclair, B. J., & Renault, D. (2010). Intracellular ice formation in insects: Unresolved after 50 years? *Comparative Biochemistry and Physiology - A Molecular and Integrative*

Physiology, 155(1), 14–18. doi:10.1016/j.cbpa.2009.10.026

Somero, G. N., Lockwood, B. L., & Tomanek, L. (2017). Oxygen and Metabolism. In *Biochemical Adaptation* (pp. 29–138).

Sømme, L. (2000). The history of cold hardiness research in terrestrial arthropods. *Cryo-Letters*, 21(5), 289–296.

Storey, K. B. (1997). Organic Solutes in Freezing Tolerance. *Comparative Biochemistry and Physiology Part A: Physiology*, 117(3), 319–326. doi:10.1016/S0300-9629(96)00270-8

Storey, K. B., & Storey, J. M. (1996). Natural freezing survival in animals. *Annual Review of Ecology and Systematics*, 27, 365–386. doi:10.1146/annurev.ecolsys.27.1.365

Storey, K. B., & Storey, J. M. (2013). Molecular biology of freezing tolerance. *Comprehensive Physiology*, 3(3), 1283–1308. doi:10.1002/cphy.c130007

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, 2(9), 686–690. doi:10.1038/nclimate1539

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. *PLoS ONE*, 6(7), e21800. doi:10.1371/journal.pone.0021800

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123(3), 585–95. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2513255>

Team, R. C. (2016). R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.r-project.org/>

Teets, N. M., & Denlinger, D. L. (2014). Surviving in a frozen desert: environmental stress physiology of terrestrial Antarctic arthropods. *The Journal of Experimental Biology*, 217(Pt 1), 84–93. doi:10.1242/jeb.089490

Teets, N. M., Kawarasaki, Y., Lee, R. E., & Denlinger, D. L. (2011). Survival and energetic costs of repeated cold exposure in the Antarctic midge, *Belgica antarctica*: a comparison between

frozen and supercooled larvae. *The Journal of Experimental Biology*, 214(Pt 5), 806–814.
doi:10.1242/jeb.060624

Todgham, A. E., Hoaglund, E. A., & Hofmann, G. E. (2007). Is cold the new hot? Elevated ubiquitin-conjugated protein levels in tissues of Antarctic fish as evidence for cold-denaturation of proteins in vivo. *Journal of Comparative Physiology B*, 177(8), 857–866.
doi:10.1007/s00360-007-0183-2

Toxopeus, J., & Sinclair, B. J. (2018). Mechanisms underlying insect freeze tolerance. *Biological Reviews*. doi:10.1111/brv.12425

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., ... Pachter, L. (2010). Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature Biotechnology*, 28(5), 511–5. doi:10.1038/nbt.1621

Waller, C. L., Worland, M. R., Convey, P., & Barnes, D. K. a. (2006). Ecophysiological strategies of Antarctic intertidal invertebrates faced with freezing stress. *Polar Biology*, 29(12), 1077–1083. doi:10.1007/s00300-006-0152-3

Wetthey, D. S. (1983). Geographic limits and local zonation: the barnacles *Semibalanus* (*Balanus*) and *Chthamalus* in New England. *The Biological Bulletin*, 165(1), 330–341.
doi:10.2307/1541373

Yi, S. X., Benoit, J. B., Elnitsky, M. A., Kaufmann, N., Brodsky, J. L., Zeidel, M. L., ... Lee, R. E. (2011). Function and immuno-localization of aquaporins in the Antarctic midge *Belgica antarctica*. *Journal of Insect Physiology*, 57(8), 1096–1105.
doi:10.1016/j.jinsphys.2011.02.006

Data accessibility

The original sequencing reads have been archived at the NCBI Short Read Archive under Bioproject number PRJNA496531 and will be made accessible on publication. The assembled and annotated *Semibalanus balanoides* transcriptome has deposited as a Transcriptome Shotgun Assembly at the NCBI database under the same number. Other data (read counts) have been deposited on the Open Science Framework at XXXXXX and will be made accessible upon publication.

Author Contributions

KEM, AP, GK, and BKKC designed the research

BKKC, AP and GK collected the barnacles from the White Sea

KEM, AP, GK, and BKKC performed the laboratory research

KEM and ED analyzed the data

KEM wrote the first draft of the paper, AP, GK, BKKC, and ED contributed to writing

Figure captions and Table captions

Table 1. Transcripts that are significantly differentially regulated in both British Columbia and White Sea populations. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 2. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the White Sea population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 3. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the British Columbia population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Table 4. Tajima D estimates among transcripts. Estimates are averaged across transcripts that either contain or do not contain the search terms. No transcripts annotated as “aquaporins” had sufficient coverage to estimate Tajima D in the British Columbia population. The total number of transcripts passing filter for each test is in brackets. Numbers marked with an asterisk indicate a statistically-significant (t-test, $P < 0.05$) difference between the Tajima D estimates of transcripts containing the search term and those not. Numbers marked with a “^” indicates terms that had insufficient numbers of transcripts to run a t-test.

Figure 1. Outline of experimental design, showing timepoints that barnacles from each population were sampled.

Figure 2. Survival of barnacles from the White Sea and British Columbia populations following a 4 h exposure to either -10 or -6 °C. Asterisk indicates a significant difference in survival between populations in a generalized linear model with binomial error, $n = 92 - 151$ per population/temperature.

Figure 3. Timecourse of the number of differentially expressed transcripts in barnacles from the White Sea (WS) and British Columbia (BC) with expression calculated relative to control (0 h) for each population using DESeq2.

Figure 4. Venn diagrams illustrating the number of differentially expressed (DE) transcripts following a four hour freezing exposure at -6 °C relative to unfrozen control for each population. Up arrows indicate significantly upregulated transcripts, down arrows indicate significantly downregulated transcripts, and a tilde indicates contraregulated transcripts which are upregulated in one group, but downregulated in another. Areas indicate rank order of number of DE transcripts. A) Number of DE transcripts four hours following freezing between the White Sea and British Columbia populations. B) Number of DE transcripts four hours following freezing between the White Sea and British Columbia populations. C) Number of DE transcripts in the White Sea population (there were no DE transcripts before thawing). D) Number of DE transcripts in the British Columbia population.

Figure 5. Significantly enriched GO terms at each timepoint in each population following a four hour freezing exposure at -6 °C relative to unfrozen control for each population as determined by Fisher's exact test and REVIGO. Blue indicates terms enriched in the White Sea population, while red indicates terms enriched in the British Columbia population. Asterisks indicate terms with long names that have been shortened for space. All full terms are available in Supplementary Data 2.

Table 1. Transcripts that are significantly differentially regulated in both British Columbia and White Sea populations. GO terms are listed only for molecular function. Only the top 5 up and down-regulated (shaded) transcripts are given (supplementary data 1 for all).

Timepoint	Description	Log ₂ Fold Change in White Sea	Log ₂ Fold Change in British Columbia	GO Terms
4 hours following freezing	Proto-oncogene c-partial	2.79	6.01	DNA binding; transcription factor activity, sequence-specific DNA binding; sequence-specific DNA binding
	Proto-oncogene c-partial	1.86	2.32	None assigned
	Aromatic-L-amino-acid decarboxylase	1.94	4.12	carboxy-lyase activity
28 hours following freezing	Williams-Beuren syndrome chromosomal region 27	-1.82	-1.17	methyltransferase activity; S-adenosylmethionine-dependent methyltransferase activity
	CAD	-0.92	-1.59	aspartate carbamoyltransferase activity; dihydroorotase activity; carbamoyl-phosphate synthase (ammonia) activity; carbamoyl-phosphate synthase (glutamine-hydrolyzing) activity; ATP

			binding; metal ion binding; amino acid binding
Plasma alpha-L-fucosidase	-1.09	-1.40	alpha-L-fucosidase activity
alpha- partial	1.47	2.44	GTP binding; GTPase activity; structural constituent of cytoskeleton; microtubule-based process
pleckstrin homology domain-containing family F member 2	1.07	0.61	metal ion binding
dystroglycan-like isoform X1	1.04	0.92	calcium ion binding
dystroglycan isoform X1	0.89	0.95	calcium ion binding
Villin quail	0.87	1.79	None assigned

793

794

Table 2. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the White Sea population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated transcripts are given (supplementary data 1 for all).

Description	Log ₂ Fold Change at 4 hours	Log ₂ Fold Change at 28 hours	GO Names
heat shock partial	2.79	6.01	ATP binding
heat shock 68-like	1.86	2.32	ATP binding
heat shock 70 A1-partial	1.94	4.12	ATP binding
heat shock 70 partial	1.82	1.17	ATP binding
heat shock partial	0.92	1.59	ATP binding
gibberellin 20 oxidase	-1.09	-1.40	oxidoreductase activity
sodium-dependent nutrient amino acid transporter	-1.47	-2.44	symporter activity
BAI1-associated 3	-1.07	-0.61	None assigned
Williams-Beuren syndrome chromosomal region 27	-1.04	-0.92	methyltransferase activity; S-adenosylmethionine-dependent methyltransferase activity
None assigned	-0.89	-0.95	None assigned

Table 3. Transcripts that are significantly differentially regulated at both 4 and 28 hours following freezing in the British Columbia population. GO terms are listed only for molecular function. Only the top 5 up and down-regulated (shaded) transcripts are given (supplementary data 1 for all).

Description	Log ₂ Fold Change at 4 hours	Log ₂ Fold Change at 28 hours	GO Names
Cytochrome partial	6.61	5.62	iron ion binding
Proto-oncogene c- partial	6.31	2.52	transcription factor activity, sequence-specific DNA binding
Proto-oncogene c- partial	6.01	2.28	transcription factor activity, sequence-specific DNA binding
techylectin-5A-like isoform X1	3.71	6.85	carbohydrate binding
Proto-oncogene c- partial	2.32	1.77	None assigned
None assigned	-1.02	-1.01	None assigned
INO80 complex subunit	-0.74	-0.59	(Process: DNA repair)
Peter pan	-0.70	-0.49	rRNA binding

Table 4. Tajima D estimates among transcripts. Estimates are averaged across transcripts that either contain or do not contain the search terms. No transcripts annotated as “aquaporins” had sufficient coverage to estimate Tajima D in the British Columbia population. The total number of transcripts passing filter for each test is in brackets. Numbers marked with an asterisk indicate a statistically-significant (t-test, $P < 0.05$) difference between the Tajima D estimates of transcripts containing the search term and those not. Numbers marked with a “^” indicates terms that had insufficient numbers of transcripts to run a t-test.

Search term	British Columbia transcripts not containing search term	British Columbia transcripts containing search term	White Sea transcripts not containing search term	White Sea transcripts containing search term
All Differentially Expressed (DE) Transcripts	-0.64 (13129)	-1.18* (1672)	-0.81 (14765)	-1.42* (875)
Transcripts DE in White Sea	-0.70 (14759)	-1.19* (42)	-0.83 (15393)	-1.44* (247)
“Transcription Factor”	-0.70 (14800)	-1.16^ (1)	-0.84 (15631)	-2.48* (9)
“Aquaporin”	NA	NA	-0.84 (15637)	-2.60^ (3)
“Macrophage Mannose Receptor”	-0.70 (14799)	-1.11^ (2)	-0.84 (15636)	-1.32^ (4)

A

4 hours after freezing

B

28 hours after freezing

C

White Sea

D

British Columbia

Before thawing

4h after freezing

28h after freezing

Differentially expressed transcripts
Total transcripts in category