Muller's Ratchet in Asexual Populations Doomed to Extinction

Logan Chipkin¹, Peter Olofsson^{2,3}, Ryan C. Daileida², Ricardo B. R. Azevedo^{1*}

*For correspondence:
razevedo@uh.edu

¹Department of Biology & Biochemistry, University of Houston, Houston, Texas, U.S.A.;

²Department of Mathematics, Trinity University, San Antonio, Texas, U.S.A.;

³Department of Mathematics, Physics and Chemical Engineering, Jönköping University, Sweden

Abstract Asexual populations are expected to accumulate deleterious mutations through a process known as Muller's Ratchet. Lynch, Gabriel, and colleagues have proposed that the Ratchet eventually results in a vicious cycle of mutation accumulation and population decline that drives populations to extinction. They called this phenomenon mutational meltdown. Here, we analyze the meltdown using a multitype branching process model where, in the presence of mutation, populations are doomed to extinction. We find that extinction occurs more quickly in small populations, experiencing a high deleterious mutation rate, and mutations with more severe deleterious effects. The effects of mutational parameters on extinction time in doomed populations differ from those on the severity of Muller's Ratchet in populations of constant size. We also find that mutational meltdown, although it does occur in our model, does not determine extinction time. Rather, extinction time is determined by the expected impact of deleterious mutations on fitness.

Introduction

"All populations are doomed to eventual extinction." *Lynch and Gabriel (1990)*

In the absence of back mutations, an asexual individual cannot produce offspring carrying fewer deleterious mutations than itself. Indeed, it is always possible that individual offspring will accrue additional deleterious mutations. As a result, the class of individuals with the fewest deleterious mutations may, by chance, disappear irreversibly from the population, a process known as Muller's Ratchet (*Muller, 1964; Felsenstein, 1974; Haigh, 1978*). Successive "clicks" of the Ratchet will cause the fitness of asexual populations to decline. Muller's Ratchet has been invoked to explain the evolution of sex (*Muller, 1964; Felsenstein, 1974; Gordo and Campos, 2008*), the extinction of small populations (*Lynch et al., 1993, 1995a*), the accelerated rate of evolution of endosymbiotic bacteria (*Moran, 1996*), the degeneration of Y-chromosomes (*Charlesworth, 1978; Gordo and Charlesworth, 2000b*), and cancer progression (*McFarland et al., 2013, 2014*).

Haigh (1978) argued that the Ratchet should click at a rate inversely proportional to the size of the least-loaded class in a population. If k is the lowest number of deleterious mutations present in an individual in the population, the size of the least-loaded class at mutation-selection-drift equilibrium is

$$\hat{n}_k = N e^{-U/s}, \quad (1)$$

where N is the size of the population, U is the expected number of deleterious mutations per genome per generation, and s is the deleterious effect of a mutation. Haigh suggested that genetic drift causes the actual value of n_k to deviate stochastically from $\hat{n}_k$. The smaller the value of $\hat{n}_k$, the

greater the probability that n_k will hit zero, causing the Ratchet to click. If $\hat{n}_k > 1$, then after a click of the Ratchet, the size of the new least-loaded class will go to a new equilibrium, $\hat{n}_{k+1}$, equal to $\hat{n}_k$ in **Equation 1**. Haigh concluded that Muller's Ratchet should click faster in small populations, experiencing a high deleterious mutation rate, and mutations with milder deleterious effects (low s). Subsequent work has derived more accurate estimates of the rate of clicking of the Ratchet, both when $\hat{n}_k > 1$ (**Stephan et al., 1993; Gordo and Charlesworth, 2000a,b; Neher and Shraiman, 2012; Metzger and Eule, 2013**) and when $\hat{n}_k < 1$ (**Gessler, 1995; Rouzine et al., 2003, 2008**).

Beginning with Haigh's foundational study, most research on Muller's Ratchet has assumed that the size of a population remains constant as deleterious mutations accumulate (e.g., **Gessler, 1995; Gordo and Charlesworth, 2000a,b; Rouzine et al., 2003; Metzger and Eule, 2013**). This assumption is biologically unrealistic—if true, fitness would decline continuously but the population would be immortal (**Lynch and Gabriel, 1990; Melzer and Koeslag, 1991**). Lynch, Gabriel, and colleagues studied more realistic models where the fitness of an individual influences its fertility and populations experience density-dependent regulation (**Lynch and Gabriel, 1990; Lynch et al., 1993; Gabriel et al., 1993; Lynch et al., 1995a**). They found that Muller's Ratchet causes population size to decline, which accelerates the Ratchet, which further reduces population size. This positive feedback results in a "mutational meltdown" that drives the population to extinction (**Lynch and Gabriel, 1990; Lynch et al., 1993; Gabriel et al., 1993**).

In one model, **Lynch et al. (1993)** considered a population of asexual organisms subject to a carrying capacity of $\hat{N}$ individuals. Each individual produces R offspring. The number of mutations is Poisson distributed with rate U . The offspring then undergo viability selection with a probability of survival

$$w_k = (1 - s)^k, \quad (2)$$

where $k \geq 0$ is the number of deleterious mutations in the individual offspring, and $0 < s < 1$ is the deleterious effect of each mutation. If the number of offspring surviving viability selection is $N' > \hat{N}$, $N' - \hat{N}$ individuals die and $\hat{N}$ individuals survive, independently of their fitness; if $N' \leq \hat{N}$, all N' individuals survive. Reproduction occurs after viability selection and density-dependent regulation. Assuming that initially all individuals in the population are mutation-free and that $NR > \hat{N}$, Muller's Ratchet proceeds in three phases in this model. First, mutations enter the population and accumulate rapidly. As the distribution of mutation numbers approaches mutation-selection-drift equilibrium mutation accumulation slows down. Second, the rate of mutation accumulation settles into a steady rate. This phase proceeds as in the classic constant population size model of Muller's Ratchet (**Haigh, 1978**) and lasts while $NR\bar{w} \gtrsim \hat{N}$. Third, when mean viability reaches $\bar{w} = 1/R$ (i.e., when $NR\bar{w} = \hat{N}$) the population size begins to decline, triggering mutational meltdown. During this phase the population is doomed to extinction.

Lynch et al. (1993) derived some analytical expressions to describe the dynamics of mutation accumulation during the first two phases and the times at which these two phases end. However, they did not present any analytical theory on the dynamics or duration of the third (meltdown) phase itself (see also **Gabriel et al., 1993; Lynch et al., 1995a**). Thus, the validity of the Lynch-Gabriel view of the mutational meltdown is difficult to evaluate. Here we model the mutational meltdown phase of Muller's Ratchet using a multitype branching process. We derive an analytical approximation for the expected time to extinction under this model of populations doomed to extinction. We find that extinction occurs more quickly in small populations, experiencing a high deleterious mutation rate (u), and mutations with *more severe* deleterious effects (high s). Our results differ from predictions on the relationship between the severity of Muller's Ratchet and mutational parameters in populations of constant size. We also find that mutational meltdown, although it does occur in doomed populations, is not an important determinant of time to extinction. Rather, extinction time is approximately inversely proportional to the product us , that is, the expected impact of deleterious mutations on fitness.

Figure 1. Multitype branching process. At each time step, an individual of type k —i.e. with k deleterious mutations (light gray)—can have one of four fates (**A–D**) with different probabilities, P (see **Table 1**). It can either die (**D**) or survive and split into two daughters (**A–C**). The daughters inherit the k mutations from their mother. A daughter can acquire one additional mutation and become a type $k + 1$ individual (dark gray) (**B–C**).

Model

Branching process

A population consists of N_k individuals with $k = 0, 1, 2, \dots$ deleterious mutations. Below, we refer to individuals with k deleterious mutations as belonging to type k .

The size $N = \sum_i N_i$ of the population is allowed to change according to a discrete-time branching process. Each generation, an individual of type k reproduces by splitting into two daughters with probability $w_k/2$ and dies with probability $1 - w_k/2$ (**Figure 1**), where w_k is the expected number of offspring of an individual of type k —i.e., its *absolute fitness* (**Equation 2**). We assume that all mutations have the same deleterious effect s and do not interact epistatically.

Any offspring may acquire one deleterious mutation with probability u . Note that u is defined differently from the mutation rate, U , in Haigh’s model (**Equation 1**). The number of mutant offspring of a surviving individual of any type is binomially distributed with parameters 2 and u (**Figure 1**).

This branching process yields the probability generating function (p.g.f.) of the number of k -type offspring of a k -type individual

$$\varphi_k(x) = 1 - \frac{w_k}{2} (1 - u^2 - 2u(1-u)x - (1-u)^2 x^2) \quad (3)$$

and mean reproduction matrix M with entries

$$\begin{cases} m_{k,k} &= w_k(1-u) \\ m_{k,k+1} &= w_k u \end{cases} \quad (4)$$

where $m_{i,j}$ is the expected number of offspring of type j generated by an individual of type i . All other entries of M are 0.

The mean reproduction matrix for generation t is M^t , the t -th power of M . For any t , all entries of M^t below the diagonal are 0. Assuming the fitness function in **Equation 2**, we can get an explicit form for the entries $m_{i,j}^{(t)}$ of M^t :

$$m_{k,k+j}^{(t)} = (1-u)^{t-j} u^j (1-s)^{tk + \frac{j(j-1)}{2}} \prod_{i=1}^j \frac{1 - (1-s)^{t+1-i}}{1 - (1-s)^i} \quad (5)$$

where $j = 0, 1, 2, \dots$. For a proof, see **Appendix 1**.

Extinction time of individuals of type k

Let τ_k denote the time of extinction of individuals of type k in a population started from N_k ancestors of type k where N_k is a random variable on $\{0, 1, 2, \dots\}$ (if $N_k = 0$, then $\tau_k = 0$). There are no individuals of type $i < k$ in the population. By a standard result from probability theory

$$E[\tau_k] = \sum_{t=0}^{\infty} P(\tau_k > t) \quad .$$

Table 1. Variables and parameters.

Symbol	Description
u	Probability that an individual acquires a deleterious mutation (Figure 1).
s	Deleterious effect of a mutation (Equation 2).
w_k	Fitness of an individual of type k , i.e. with k deleterious mutations (Equation 2).
$m_{i,j}$	Expected number of offspring of type j generated by an individual of type i (Equation 4).
n_0	Initial number of mutation-free individuals in the population.
n_k	Expected number of k -type individuals in the population.
N	Total population size (Equation 19).
t_k	Expected extinction time of k -type individuals in generations, i.e. the k -th click of the Ratchet (Equation 12).
Δt_k	Interval between clicks $k - 1$ and k of the Ratchet (Equation 21).
x_k	Expected number of k -type individuals at the extinction time of type $k - 1$ (Equation 11).
$\mathbf{T}$	Expected extinction time of the entire population in generations (Equation 10).

115 The time of extinction of the entire type- k subpopulation is the time of extinction of the N_k
116 independent subpopulations started from the ancestors. The p.g.f. of the number of k -type indi-
117 viduals in generation t is given by the t -fold composition of φ_k (**Equation 3**) with itself, denoted by
118 $\varphi_k^{(t)}$. We get

$$\tau_k = \max\{\tau_{k,1}, \dots, \tau_{k,N_k}\}$$

119 where $\tau_{k,j}$ is the time of extinction of the subpopulation started from the j th individual, $j = 1, \dots, N_k$.
120 If we let $Z_t^{(k,j)}$ denote the number of type- k individuals in generation t stemming from the j th
121 individual we have the equivalence

$$\tau_{k,j} \leq t \Leftrightarrow Z_t^{(k,j)} = 0$$

122 and get the conditional probability given N_k

$$\begin{aligned} P(\tau_k > t | N_k) &= 1 - P(\tau_k \leq t | N_k) \\ &= 1 - \prod_{j=1}^{N_k} P(\tau_{k,j} \leq t) \\ &= 1 - \left(\varphi_k^{(t)}(0)\right)^{N_k} \end{aligned}$$

123 for $t > 0$ which gives

$$E[\tau_k] = P(\tau_k > 0) + E\left[\sum_{t=1}^{\infty} \left(1 - \left(\varphi_k^{(t)}(0)\right)^{N_k}\right)\right]$$

124 With $n_k = E[N_k]$, a first-order Taylor approximation gives

$$E[\tau_k] \approx P(\tau_k > 0) + \sum_{t=1}^{\infty} \left(1 - \left(\varphi_k^{(t)}(0)\right)^{n_k}\right) \quad (6)$$

125 Note that for $k = 0$ we have $P(\tau_0 > 0) = 1$ because there are always individuals present at time 0.
126 For $k > 0$, however, we have $P(\tau_k > 0) < 1$ because, for example, the entire population may already
127 be extinct in generation 1.

128 In a similar way, we get the variance as

$$\text{Var}[\tau_k] = E[\tau_k(\tau_k - 1)] + E[\tau_k] - E^2[\tau_k] \quad (7)$$

129 where

$$E[\tau_k(\tau_k - 1)] = 2 \sum_{t=1}^{\infty} t P(\tau_k > t) \\ \approx 2 \sum_{t=1}^{\infty} t \left(1 - \left(\varphi_k^{(t)}(0) \right)^{n_k} \right)$$

130 Extinction time of the entire population

131 By well-known results from the theory of branching processes, the extinction time of the entire
132 population has finite mean only in the *subcritical* case, that is, when the mean number of offspring
133 per individual is less than 1. The expected number of offspring of type k produced by an individual
134 of type k is $m_{k,k}$ (**Equation 4**). If $m_{k,k} = 1$ (the *critical* case), the extinction time τ_k is finite but has an
135 infinite mean and if $m_{k,k} > 1$ (the *supercritical* case), τ_k itself may assume the value ∞ .

136 **Equation 4** shows that $m_{k,k} < 1$ (the *subcritical* case) for individuals of any type k provided all
137 mutations are deleterious ($0 < s < 1$) and the mutation rate is nonzero ($u > 0$). Thus, the expected
138 extinction time of every type k is finite. In other words, the population is doomed to eventual
139 extinction.

140 Start with a fixed number n_0 of mutation-free individuals and denote by T_0 the time (generation)
141 of extinction of this class. Conditioned on T_0 , the expected number of individuals in class 1 (those
142 with $k = 1$ mutation) is therefore $m_{0,1}^{(T_0)}$ (see **Equation 5**) which we note is a function of the random
143 variable T_0 . Thus, the expected number of individuals in class 1 at the time of extinction of class
144 0 is obtained by taking the expected value in $m_{0,1}^{(T_0)}$. To this end, recall **Equation 5** and define the
145 function

$$g_1(\cdot) = m_{0,1}^{(\cdot)}$$

146 so that

$$E \left[m_{0,1}^{(T_0)} \right] = E[g_1(T_0)] \\ \approx g_1(E[T_0])$$

147 where we use a first-order Taylor approximation. To generalize the idea, we define the expected
148 number of descendants of type j from a mutation-free individual after t generations

$$g_j(t) = \begin{cases} 0 & , \quad t < j \\ m_{0,j}^{(t)} & , \quad t \geq j \end{cases} \quad (8)$$

149 (see **Equation 5**).

150 Now let T_k be the extinction time for type k and let $t_k = E[T_k]$. Then we have the approximation

$$E \left[m_{0,k+j}^{(T_k)} \right] \approx g_{k+j}(t_k) \quad (9)$$

152 the expected number of individuals of type $k + j$ at the extinction of type k for $j = 1, 2, \dots$ The
153 expected extinction time of the entire population is

$$\mathbf{T} = E[T] = \lim_{k \rightarrow \infty} t_k \quad (10)$$

154 Now let $X_{T_{k-1}}^{(k)}$ be the number of k -type individuals at the extinction time of type $k - 1$ and let

$$x_k = E \left[X_{T_{k-1}}^{(k)} \right] \approx n_0 g_k(t_{k-1}) \quad (11)$$

From **Equation 6**, the expected extinction times of consecutive classes can be computed as

$$t_k \approx t_{k-1} + P(\tau_k > 0) + \sum_{i=1}^{\infty} \left(1 - \left(\varphi_k^{(i)}(0)\right)^{x_k}\right) \quad (12)$$

where n_0 is the initial population size. Note that t_k is the time of the k -th click of the Ratchet. As we noted above, for $k = 0$ we have $P(\tau_0 > 0) = 1$ (**Equation 6**). Thus, if the population is founded by n_0 mutation-free individuals, the time to extinction of the mutation-free class is given exactly by

$$t_0 = 1 + \sum_{i=1}^{\infty} \left(1 - \left(\varphi_k^{(i)}(0)\right)^{n_0}\right) \quad (13)$$

When $k > 0$, **Equation 12** is an approximation (see **Equation 6** and **Equation 11**). In addition, we do not have a closed form expression for $P(\tau_k > 0)$ for $k > 0$. We can, however, place bounds on $P(\tau_k > 0)$ by noting that

$$\tau_k > 0 \Leftrightarrow X_{T_{k-1}}^{(k)} > 0 \quad (14)$$

and that if Y is any random variable on $\{0, 1, 2, \dots\}$ we have

$$\begin{aligned} E[Y] &= E[Y|Y > 0] P(Y > 0) \\ &\geq P(Y > 0) \end{aligned} \quad (15)$$

By **Equation 11**, **Equation 14**, and **Equation 15** we get the bounds

$$0 \leq P(\tau_k > 0) \leq \min(1, x_k) \quad (16)$$

Because extinction of the whole population is irreversible, $P(\tau_k > 0)$ is expected to decline for successive classes:

$$P(\tau_k > 0) \leq P(\tau_{k-1} > 0) \quad .$$

Large initial population size

The expected time to extinction of the mutation-free class, t_0 , is given by **Equation 13**. Following **Jagers et al. (2007)**, there exists a sequence $c(n_0) \rightarrow c$ as $n_0 \rightarrow \infty$ such that

$$t_0 = -\frac{\ln n_0 + c(n_0)}{\ln m_{0,0}} \quad (17)$$

where $m_{0,0} = 1 - u < 1$ (**Equation 4**) is the expected number of mutation-free offspring per mutation-free individual and n_0 is the initial number of mutation-free individuals. Note that the value of c depends on u (e.g., for $u = 0.01$ and 0.02 , numerical estimates using **Equation 13** and **Equation 17** yield $c = 3.3737$ and 2.7058 , respectively).

Equation 17 shows that t_0 grows logarithmically with n_0 with a slope of $-1/\ln m_{0,0}$. If u is small, the slope becomes $\approx 1/u$. Thus, increasing initial population size delays extinction of the mutation-free class more when the mutation rate is low than when it is high.

The value of t_0 is not affected by the effects of mutations, s (**Equation 13** and **Equation 17**), because the rate at which individuals “leave” the mutation-free class is independent of s . The selection coefficient does, however, affect the size of the new least-loaded class (i.e., individuals with $k = 1$ mutation), x_1 (**Equation 11**), and therefore the total time to extinction.

We now investigate the limiting behavior of x_1 as $n_0 \rightarrow \infty$. By **Equation 11** and **Equation 17** we get

$$\begin{aligned} x_1 &\approx n_0 g_1(t_0) \\ &= \frac{C(n_0)u}{s(1-u)} \left(1 - (1-s)^{t_0}\right) \\ &\rightarrow \frac{Cu}{s(1-u)} \end{aligned} \quad (18)$$

Figure 2. Populations are doomed to extinction in our model. **(A)** Dynamics of population size, N , in 40 populations founded by $n_0 = 100$ mutation-free individuals and subject to mutations with deleterious effect $s = 0.01$ and rate $u = 0.01$. Black vertical lines above the time axis indicate extinction times. **(B)** Blue line shows mean N based on stochastic simulations of 10^4 replicate populations like those shown in **(A)**. Light blue region indicates $\bar{N} \pm \text{SD}[N]$ (standard deviation). If $\text{SD}[N] > \bar{N}$, the lower bound of the region was set to zero. Red dashed line indicates expected population size (Equation 19). **(C)** Distribution of extinction times, T , in the stochastic simulations described in **(B)**. Blue line shows mean T based on the 10^4 replicate populations. Red line shows T calculated numerically (see **Materials and Methods**). **(D)** Extinction times of populations with the same mutational parameters as those in **(A)** but with a range of initial populations sizes, n_0 . Blue line shows mean values of T based on stochastic simulations of 10^4 replicate populations for 41 values of n_0 evenly spaced on a log-scale over 4 orders of magnitude. Light blue region indicates $\bar{T} \pm \text{SD}[T]$. If $\text{SD}[T] > \bar{T}$, the lower bound of the region was set to zero. See **Figure 2-Figure Supplement 1** for more on the variability of T . Red line shows T calculated numerically.

Figure 2-Figure supplement 1. Variability of extinction time declines with population size.

Figure 2-source data 1. The code to generate these figures is in the Jupyter notebook <https://github.com/rbazev/doomed/blob/master/python/fig2.ipynb>.

Figure 2-source data 2. The data to generate these figures is at <https://github.com/rbazev/doomed/blob/master/python/data/> (files named fig2*).

as $n_0 \rightarrow \infty$, where $C(n_0) = e^{c(n_0)}$. If u is small and n_0 is large, Equation 18 becomes $x_1 \approx Cu/s$. Interestingly, Equation 18 shows that x_1 approaches a constant as n_0 increases.

Change in population size

If a population is founded by n_0 mutation-free individuals, the expected total population size t generations later is

$$E[N(t)] = n_0 \sum_{j=0}^t g_j(t) \quad (19)$$

(see Equation 8).

Initially, $N(0) = n_0$. Since all individuals have the same fitness, the population size is not expected to change in the following generation: $E[N(1)] = n_0$. One generation later, the population size is expected to decline by $E[N(2)] - E[N(1)] = -n_0us$. In subsequent generations, if mutations have small effects, the population size is expected to continue to decline at approximately the

Figure 3. Mutational parameters have different effects on extinction time in doomed populations and the severity of Muller’s Ratchet in populations of constant size. **(A–B)** Values are mean extinction times, T , in populations founded by $n_0 = 100$ mutation-free individuals but with different mutational parameters. **Figure 3–Figure Supplement 1** shows the variability of T . **(A)** Mutations have deleterious effect $s = 0.01$ and a range of mutation rates, u . **(B)** Mutations occur with $u = 0.01$ and have a range of values of s . Blue lines show mean values of T based on stochastic simulations of 10^4 replicate populations for 21 values of the parameter being manipulated, evenly spaced on a log-scale. Light blue regions indicate $\bar{T} \pm \text{SD}[T]$. If $\text{SD}[T] > \bar{T}$, the lower bound of the region was set to zero. Red lines show T calculated numerically (see **Materials and Methods**) for 41 values of the parameter being manipulated, evenly spaced on a log-scale. **Figure 3–Figure Supplement 2** shows that the theoretical predictions for low s become more accurate with increasing population size. **(C)** Severity of Muller’s Ratchet in populations of constant size and a deleterious mutation rate of $U = 0.01$ per genome per generation. Values are the expected declines in mean fitness per thousand generations, $10^3 \times s/\Delta t$, for 101 values of s evenly spaced on a log-scale. Δt is the time between clicks of the Ratchet calculated using the method of [Gordo and Charlesworth \(2000a,b\)](#). Dashed and solid lines indicate $\hat{n}_k < 10$ and $\hat{n}_k \geq 10$, respectively ([Equation 1](#)). The trend shown in **(C)** was confirmed by simulation (not shown).

Figure 3–Figure supplement 1. Variability of extinction time declines with mutation rate and is approximately invariant with selection coefficient.

Figure 3–Figure supplement 2. Theoretical predictions for low s become more accurate with increasing population size.

Figure 3–source data 1. The code to generate these figures is in the Jupyter notebook <https://github.com/rbazev/doomed/blob/master/python/fig3.ipynb>.

Figure 3–source data 2. The data to generate these figures is at <https://github.com/rbazev/doomed/blob/master/python/data/> (files named fig3*).

192 same rate

$$E[N(t+1)] - E[N(t)] \approx -n_0 u s t \quad (20)$$

193 Results

194 Small population size, high mutation rate, and mutations of large effect accelerate extinction

195
196 In our model, population size, N , can increase as well as decrease from generation to generation. However, all increases are transient and the population will eventually go extinct (**Figure 2A**).
197 The expected value of N can be predicted accurately by [Equation 19](#) (**Figure 2B**). However, the
198 dynamics of the expected value of N are not sufficient to predict the time to extinction. Two pop-
199 ulations with different initial population sizes, n_0 , will be expected to show the same N/n_0 at any
200 time (because they will have the same $g_j(t)$, [Equation 19](#)), but the smaller population is expected
201 to go extinct earlier (**Figure 2C** and **Figure 2D**). [Equation 10](#) provides good estimates of expected
202 extinction time, T , and variability in extinction time, over a broad range of initial population sizes,
203 n_0 (**Figure 2D**, **Figure 2–Figure Supplement 1**, and **Figure 3–Figure Supplement 2**), and mutational
204 parameters (**Figure 3A**, **Figure 3B**, and **Figure 3–Figure Supplement 1**). (See [Appendix 2](#) for an ex-
205 planation of why [Equation 10](#) tends to overestimate the true value of T .) Below, we focus on the
206 numerical calculations of T based on [Equation 10](#).
207

Our model has three parameters— n_0 , u , and s —and all of them influence extinction time. Smaller populations tend to go extinct faster. **Figure 2D** and **Figure 3–Figure Supplement 2A** show that T is approximately proportional to the logarithm of the initial population size, n_0 . When n_0 is large, $t_0 \propto \ln n_0$ (**Equation 17**) and x_1 is approximately constant (**Equation 18**). Thus, $\Delta t_1, \Delta t_2, \dots$ (**Equation 21**) are also expected to approach constant values as $n_0 \rightarrow \infty$. Since t_0 grows logarithmically, this result implies that t_0 represents an increasing fraction of T as n_0 increases. Therefore, we also expect T to grow logarithmically with n_0 . Variability in extinction time declines with n_0 (**Figure 2–Figure Supplement 1**).

At a particular value of n_0 , however, t_0 is not sufficient to explain the variation in total extinction time for different mutational parameters: t_0 dominates T when u/s is low, but not when it is high (**Figure 4B** and **Figure 4–Figure Supplement 1A**). One reason for this pattern is that x_1 increases with u/s (**Equation 18**; **Figure 4C**).

Mutations cause extinction in our model, but how do they influence extinction time? One possibility is that T is determined by the severity of Muller’s Ratchet. High mutation rate and mutations of large effect accelerate extinction in doomed populations (**Figure 3A**, **Figure 3B**, and **Figure 4A**). Mutational parameters act differently on Muller’s Ratchet in populations of constant size: at certain mutation rates the severity of the Ratchet is maximal at intermediate mutational effects (**Gabriel et al., 1993**; **Gordo and Charlesworth, 2000a,b**) (**Figure 3C**). There are two possible explanations for this discrepancy. First, the Ratchet may operate differently in doomed populations and populations of constant size. Second, Muller’s Ratchet may not determine extinction time in doomed populations. We explore each of these possibilities in the next two sections.

Figure 4. High mutation rate and mutations of large effect accelerate extinction in doomed populations. **(A)** Expected extinction time, T , of populations founded by $n_0 = 100$ mutation-free individuals and subject to mutations with deleterious effect s and rate u . T was calculated numerically (see **Materials and Methods**) for $21 \times 21 = 441$ combinations of values of s and u evenly spaced on a log-scale. (See **Figure 4–Figure Supplement 1B** for the variability of T .) **(B)** Expected extinction time of the mutation-free class, t_0 , as a proportion of T for the populations shown in **(A)**. t_0 was calculated numerically (see **Materials and Methods**). **Figure 4–Figure Supplement 1A** shows t_0 . **(C)** Expected number of individuals with $k = 1$ mutation at t_0 , x_1 (**Equation 11**), for the populations shown in **(A)**.

Figure 4–Figure supplement 1. Expected extinction time of the mutation-free class and variability of extinction time.

Figure 4–source data 1. The code to generate these figures is in the Jupyter notebook <https://github.com/rbazev/doomed/blob/master/python/fig4.ipynb>.

Low mutation rate and mutations of large effect accelerate mutational meltdown
Lynch and colleagues have proposed that Muller’s Ratchet accelerates in a doomed population as population size declines, driving the population to extinction—a phenomenon they called mutational meltdown (**Lynch and Gabriel, 1990**; **Lynch et al., 1993**; **Gabriel et al., 1993**). Do our doomed populations experience mutational meltdown?

To answer this question we need a way to quantify the rate of mutational meltdown. Let

$$\Delta t_k = \begin{cases} t_0 & , \quad k = 0 \\ t_k - t_{k-1} & , \quad k > 0 \end{cases} \quad (21)$$

be the interval between clicks $k - 1$ and k of the Ratchet, where t_k denotes the time of the k -th click. In populations of constant size, the Ratchet is expected to click at a steady rate over time, that is, Δt_k is not expected to change with k (Haigh, 1978; Gordo and Charlesworth, 2000a,b). In contrast, under mutational meltdown the Ratchet is expected to accelerate in successive clicks, that is, Δt_k is expected to decline with k . We find that Δt_k declines approximately exponentially with k in doomed populations (Figure 5A). Thus, we use the rate of this decline, β , to measure the rate of mutational meltdown.

Mutational parameters have large effects on the rate of mutational meltdown. Low mutation rate and mutations of large effect accelerate mutational meltdown (Figure 5). This result is consistent with the effects of mutational parameters on Δt_0 and Δt_1 in large populations (Equation 17 and Equation 18). Low values of u increase Δt_0 and decrease x_1 , and therefore Δt_1 , causing mutational meltdown to accelerate. High values of s have no effect on Δt_0 but decrease x_1 , and therefore Δt_1 , also causing the meltdown to accelerate.

Figure 5. Mutational meltdown does not determine extinction time in doomed populations. **(A)** Circles denote time between clicks, Δt_k (Equation 21). Click k indicates the extinction of k -type individuals. Values of t_0 were calculated using Equation 13; values of t_1 , t_2 , and t_3 were calculated using Equation 12 (see Materials and Methods). Note that Δt_k is displayed on a log-scale. Each combination of mutational parameters results in populations having the same expected extinction time ($T = 223.52$). Lines indicate linear regression fits of $\ln \Delta t_k$ on k (meltdown rates: $\beta = -3.20, -1.14$ and -0.32 , respectively). **(B)** Meltdown rates, β , calculated as shown in (A) for $21 \times 21 = 441$ combinations of values of s and u evenly spaced on a log-scale.

Figure 5-source data 1. The code to generate this figure is in the Jupyter notebook <https://github.com/rbazev/doomed/blob/master/python/fig5.ipynb>.

Extinction time is determined by the rate of fitness decline

Does mutational meltdown determine extinction time? Figure 5 shows that doomed populations can experience strong mutational meltdown. However, mutational meltdown does not determine extinction time. Although the three scenarios summarized in Figure 5A have widely different meltdown rates, they have the same expected extinction time, $T = 223.52$. The lack of correlation between T and β is clear when we compare Figure 4A and Figure 5B. Although the meltdown rate does not determine extinction time, for a given extinction time, the meltdown rate is positively correlated with the variability in extinction time (Figure 3-Figure Supplement 1 and Figure 4-Figure Supplement 1B).

The results so far indicate that Muller's Ratchet does not determine extinction time in doomed populations. So what does? Extinction is, trivially, caused by decline in population size. In our model the rate of decline in population size is determined by the product of mutation rate and effect, us (Equation 20). This makes intuitive sense because the rate of decline in population size is

determined by the absolute fitness of individuals and both u and s are inversely related to fitness: increasing s reduces the fitness of an individual directly (*Equation 2*) and increasing u reduces the fitness of an individual's offspring. This explains why time to extinction declines with both u and s (*Figure 4A*).

Discussion

Most models of Muller's Ratchet have assumed that populations maintain a constant size as deleterious mutations accumulate (*Haigh, 1978; Gessler, 1995; Gordo and Charlesworth, 2000a,b; Rouzine et al., 2003; Metzger and Eule, 2013*). This is typically justified as resulting from density-dependent regulation of population size. However, the assumption is unrealistic because it prevents populations from ever going extinct (*Lynch and Gabriel, 1990; Melzer and Koeslag, 1991*). In a series of studies relaxing the assumption of constant population size, Lynch, Gabriel, and colleagues argued that Muller's Ratchet eventually generates a positive feedback where the Ratchet clicks, which causes population size to decline, which strengthens genetic drift relative to natural selection, which in turn accelerates the Ratchet (*Lynch and Gabriel, 1990; Lynch et al., 1993; Gabriel et al., 1993*). They called this vicious cycle mutational meltdown and concluded that it drives populations to extinction. However, the lack of quantitative theory on the mutational meltdown has made it difficult to evaluate the Lynch-Gabriel hypothesis.

Our results challenge the Lynch-Gabriel hypothesis. Although doomed populations can experience mutational meltdown—measured by the acceleration of Muller's Ratchet—the rate of mutational meltdown does not determine extinction time (*Figure 5A*). Therefore, the Lynch-Gabriel mutational meltdown is not a general *cause* of extinction. Rather, our results suggest that extinction time is determined by the expected impact of deleterious mutations on fitness. Interestingly, if we compare populations with the same expected extinction time but different mutational parameters, populations undergoing faster meltdown rate have more variable extinction times than populations undergoing slower meltdown (*Figure 4–Figure Supplement 1*).

The Lynch-Gabriel hypothesis emphasized the role of the change in the strength of genetic drift in causing mutational meltdown (e.g., “we refer to this synergism between mutation accumulation and random genetic drift as a mutational meltdown”; *Lynch et al., 1993*). Our results indicate that extinction in doomed populations is driven by mutation pressure, not genetic drift. *Gessler (1995)* identified a related phenomenon in the operation of Muller's Ratchet under constant population size: if the mutation rate is too high the Ratchet is driven by mutation pressure, not genetic drift. Since the expression “mutational meltdown” is now in common usage (e.g., *Poon and Otto, 2000; Rowe and Beebe, 2003; Allen et al., 2009; McFarland et al., 2014*), we propose that it be revised to refer to extinction caused by mutation pressure.

The extent to which real populations undergo mutational meltdown is unclear. A population must first enter the doomed regime. Models of Muller's Ratchet in populations of constant size have identified three major risk factors that can drive populations into the doomed regime: long-term reductions in population size, increases in mutation rate, and intermediate deleterious effects of mutations (*Lynch and Gabriel, 1990; Lynch et al., 1993; Gabriel et al., 1993; Lynch et al., 1995a; McFarland et al., 2013*). Increased mutation rate can even drive a very large population into the doomed regime—a phenomenon known as lethal mutagenesis (*Bull et al., 2007*). Next, we consider the first two risk factors.

Population size can decline as a result of changes in the environment, such as, climate change, decreased food availability, emergence of infectious diseases, and habitat loss or fragmentation. For example, the emergence of Devil Facial Tumor Disease, a transmissible cancer, has caused the size of the Tasmanian devil population to decline by ~77% within 5 years (*Hawkins et al., 2006; Lazenby et al., 2018*). As a result, the devils are under risk of extinction (*McCallum et al., 2009*). Our results indicate that increasing population size causes relatively small delays in extinction in doomed populations. T is approximately proportional to the logarithm of initial population size

(Figure 2D). Similar results have been obtained in other stochastic models of population dynamics (Lande, 1993; Jagers et al., 2007).

Increases in mutation rate have been observed directly in experimental populations. For example, a population of *Escherichia coli* adapting to a constant environment evolved a mutator mutation after ~25,000 generations that increased mutation rate by ~150-fold (Barrick et al., 2009; Wielgoss et al., 2013). Evolution experiments have revealed that real populations can, indeed, experience increased extinction risk when the mutation rate is high. Zeyl et al. (2001) allowed 12 populations of the yeast *Saccharomyces cerevisiae* with genetically elevated mutation rate to evolve and found that two of them went extinct within 2,900 generations. One of these populations went extinct shortly after a large decline in fitness. Bank et al. (2016) subjected two populations of influenza A virus to gradually increasing concentrations of favipiravir, a drug that increases the mutation rate of the virus, and observed that both populations accumulated mutations rapidly and went extinct. The results from both of these studies are broadly consistent with the occurrence of a mutational meltdown. However, they do not allow us to distinguish between the Lynch-Gabriel model and ours.

The results described in the previous paragraph indicate that mutational meltdown might have clinical applications. Mutagenic agents are being explored as antiviral drugs (Loeb et al., 1999; Crotty et al., 2001; Pariente et al., 2001; Bank et al., 2016). Increased mutation rate in tumor cells has been found to correlate with improved outcomes for some cancers (Silva et al., 2000; Birkbak et al., 2011; Andor et al., 2016). Several inhibitors of key components of the DNA-repair and DNA damage-response machinery (e.g., PARP inhibitors, Lord et al., 2015), are currently being used to treat cancer, or are under preclinical or clinical development (Brown et al., 2017).

The relative theoretical neglect of the evolutionary dynamics in doomed populations is surprising given that it is central to understanding the long-term consequences of both Muller's Ratchet (Lynch and Gabriel, 1990; Lynch et al., 1993; Gabriel et al., 1993; Lynch et al., 1995a) and lethal mutagenesis (Bull et al., 2007; Matuszewski et al., 2017). In both cases, the duration of the meltdown phase was dismissed because it was predicted to be short relative to the time required for the population to become doomed (Lynch et al., 1993, 1995a; Bull et al., 2007). We believe that this neglect of the meltdown phase is misplaced because it is an important phase in the life of a population—the last chance for the population to be rescued by beneficial mutations and avoid extinction. The dynamics and duration of the meltdown phase are expected to be important determinants of the probability of evolutionary rescue. For a given probability of beneficial mutation and selection coefficient of those mutations, populations that decline in size more slowly and retain higher proportions of mutation-free individuals for longer, are more likely to be rescued (Martin et al., 2013). Rescue of doomed populations may play an important role in cancer progression (McFarland et al., 2013, 2014).

A central assumption of our model is that individuals experience hard selection (Wallace, 1975). The expected number of offspring of an individual is its absolute fitness, w_k (Equation 2), and is both density independent and frequency independent. The Lynch-Gabriel models make similar assumptions during the mutational meltdown phase (Lynch et al., 1993, 1995a). In contrast, classic models of Muller's Ratchet typically assume soft selection (Wallace, 1975). Constant population size implies density dependence. Selection is also frequency dependent: the expected number of offspring of an individual depends not only on its fitness, but on the fitness of other individuals in the population. The difference in the mode of selection in these models explains the different effects of the deleterious effect of a mutation, s , on extinction time in our model, and on the severity of Muller's Ratchet in classic models. In doomed populations, increasing s accelerates extinction, albeit with diminishing returns (Figure 2B and Figure 4). In models with soft selection, the Ratchet is most severe at an intermediate value of s (Figure 2C; Lynch et al., 1995a; Gordo and Charlesworth, 2000a,b).

In reality, populations do not necessarily experience either of the extremes of soft or hard selection. Factors such as population structure, resource availability, and the mechanism of competi-

tion can modulate the “softness” of selection in complex ways; different genotypes in a population, and even different genes, can experience different softness of selection (*Laffafian et al., 2010; Ho and Agrawal, 2012*). This raises the interesting question of how mutational meltdown operates in regimes of intermediate softness, or with a smooth transition between soft and hard selection as mutations accumulate.

Our model includes at least three other simplifications that could have important consequences for the evolutionary dynamics of doomed populations. First, the environment, and therefore selection, is constant. *Wardlaw and Agrawal (2012)* showed that temporal variation in selection can accelerate Muller’s Ratchet in populations of constant size. Second, all mutations are deleterious. Beneficial mutations could, potentially, rescue a population from extinction (*Martin et al., 2013*). Furthermore, compensatory mutations might become more common as fitness declines (*Poon and Otto, 2000; Silander et al., 2007*). Third, individuals reproduce asexually. Sex has been shown to slow down Muller’s Ratchet dramatically in populations of constant size (*Pamilo et al., 1987; Charlesworth et al., 1993*), and can delay extinction in large populations (*Lynch et al., 1995b*). We believe that our model provides a promising framework to explore the consequences of relaxing these assumptions for the fate of populations doomed to extinction.

Materials and Methods

Numerical calculations

Expected extinction time of the mutation-free class

We calculated t_0 by computing *Equation 13* until the following criterion was met

$$\left(\varphi_k^{(t)}(0)\right)^{n_0} - \left(\varphi_k^{(t+1)}(0)\right)^{n_0} < 10^{-6}.$$

A similar criterion was applied when evaluating *Equation 7* and *Equation 12*.

Expected extinction time

We calculated $\mathbf{T} = E[T]$ (*Equation 10*) by computing *Equation 12* until the following criterion was met

$$t_k - t_{k-1} < 10^{-6}.$$

We computed $\mathbf{T}$ for both bounds of $P(\tau_k > 0)$ when $k > 0$ (*Equation 16*). We present only results for the lower bound. None of our conclusions would be changed if we used the upper bound results instead (not shown).

Variance in extinction time

We calculated the variance in extinction time by computing

$$\text{Var}[T] = \sum_{k=0}^{\infty} \text{Var}[\tau_k], \quad (22)$$

up to the same value of k used to calculate $\mathbf{T}$ (see *Equation 7*). Again, we only present results for the lower bound of $P(\tau_k > 0)$ when $k > 0$ (*Equation 16*). *Equation 22* assumes that the extinction times of different classes are independent (i.e., we ignore the covariance terms). This assumption was confirmed by simulations (not shown).

Coefficient of variation in extinction time

The coefficient of variation measures the variability of a variable relative to its mean. We calculated the coefficient of variation of extinction time by computing

$$\text{CV}[T] = \frac{\sqrt{\text{Var}[T]}}{\mathbf{T}}. \quad (23)$$

Code availability

Numerical calculations and stochastic simulations of the branching process model were done using software written in Python 2.7 and available at <https://github.com/rbasev/doomed/blob/master/python/doomed.py>.

Acknowledgments

We thank Alex Stewart, Ata Kalirad, Herbert Levine, and Erin Kelleher for helpful discussions.

References

- Allen JM**, Light JE, Perotti AM, Braig HR, Reed DL. Mutational meltdown in primary endosymbionts: selection limits Muller's ratchet. *PLoS One*. 2009; 4(3):e4969. doi: [10.1371/journal.pone.0004969](https://doi.org/10.1371/journal.pone.0004969).
- Andor N**, Graham TA, Jansen M, Xia LC, Aktipis CA, Petritsch C, Ji HP, Maley CC. Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. *Nat Med*. 2016 Jan; 22(1):105–113. doi: [10.1038/nm.3984](https://doi.org/10.1038/nm.3984).
- Bank C**, Renzette N, Liu P, Matuszewski S, Shim H, Foll M, Bolon DNA, Zeldovich KB, Kowalik TF, Finberg RW, et al. An experimental evaluation of drug-induced mutational meltdown as an antiviral treatment strategy. *Evolution*. 2016 Sep; 70(11):2470–2484. doi: [10.1111/evo.13041](https://doi.org/10.1111/evo.13041).
- Barrick JE**, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, Lenski RE, Kim JF. Genome evolution and adaptation in a long-term experiment with *Escherichia coli*. *Nature*. 2009; 461(7268):1243–7. doi: [10.1038/nature08480](https://doi.org/10.1038/nature08480).
- Birkbak NJ**, Eklund AC, Li Q, McClelland SE, Endesfelder D, Tan P, Tan IB, Richardson AL, Szallasi Z, Swanton C. Paradoxical relationship between chromosomal instability and survival outcome in cancer. *Cancer Res*. 2011 May; 71(10):3447–52. doi: [10.1158/0008-5472.CAN-10-3667](https://doi.org/10.1158/0008-5472.CAN-10-3667).
- Brown JS**, O'Carrigan B, Jackson SP, Yap TA. Targeting DNA repair in cancer: beyond PARP inhibitors. *Cancer Discovery*. 2017; 7(1):20–37. doi: [10.1158/2159-8290.CD-16-0860](https://doi.org/10.1158/2159-8290.CD-16-0860).
- Bull J**, Sanjuán R, Wilke C. Theory of lethal mutagenesis for viruses. *J Virol*. 2007; 81(6):2930–2939. doi: [10.1128/JVI.01624-06](https://doi.org/10.1128/JVI.01624-06).
- Charlesworth B**. Model for evolution of Y chromosomes and dosage compensation. *Proc Natl Acad Sci U S A*. 1978; 75(11):5618–5622. doi: [10.1073/pnas.75.11.5618](https://doi.org/10.1073/pnas.75.11.5618).
- Charlesworth D**, Morgan MT, Charlesworth B. Mutation accumulation in finite outbreeding and inbreeding populations. *Genet Res*. 1993; 61(1):39–56. doi: [10.1017/S0016672300031086](https://doi.org/10.1017/S0016672300031086).
- Crotty S**, Cameron CE, Andino R. RNA virus error catastrophe: direct molecular test by using ribavirin. *Proc Natl Acad Sci U S A*. 2001 Jun; 98(12):6895–900. doi: [10.1073/pnas.111085598](https://doi.org/10.1073/pnas.111085598).
- Felsenstein J**. The evolutionary advantage of recombination. *Genetics*. 1974; 78(2):737–756.
- Gabriel W**, Lynch M, Bürger R. Muller's ratchet and mutational meltdowns. *Evolution*. 1993; 47(6):1744–1757. doi: [10.1111/j.1558-5646.1993.tb01266.x](https://doi.org/10.1111/j.1558-5646.1993.tb01266.x).
- Gessler DDG**. The constraints of finite size in asexual populations and the rate of the ratchet. *Genet Res*. 1995; 66(3):241–253. doi: [10.1017/S0016672300034686](https://doi.org/10.1017/S0016672300034686).
- Gordo I**, Charlesworth B. On the speed of Muller's ratchet. *Genetics*. 2000; 156(4):2137–2140.
- Gordo I**, Campos PRA. Sex and deleterious mutations. *Genetics*. 2008; 179(1):621–626. doi: [10.1534/genetics.108.086637](https://doi.org/10.1534/genetics.108.086637).
- Gordo I**, Charlesworth B. The degeneration of asexual haploid populations and the speed of Muller's ratchet. *Genetics*. 2000; 154(3):1379–1387.
- Haigh J**. The accumulation of deleterious genes in a population—Muller's Ratchet. *Theor Popul Biol*. 1978; 14(2):251–267. doi: [10.1016/0040-5809\(78\)90027-8](https://doi.org/10.1016/0040-5809(78)90027-8).
- Hawkins CE**, Baars C, Hesterman H, Hocking GJ, Jones ME, Lazenby B, Mann D, Mooney N, Pemberton D, Pyecroft S, et al. Emerging disease and population decline of an island endemic, the Tasmanian devil *Sarcophilus harrisii*. *Biol Conservat*. 2006 Aug; 131(2):307–324. doi: [10.1016/j.biocon.2006.04.010](https://doi.org/10.1016/j.biocon.2006.04.010).

- 441 **Ho EKH**, Agrawal AF. The effects of competition on the strength and softness of selection. *J Evol Biol.* 2012 Oct;
442 25(12):2537–2546. doi: [10.1111/j.1420-9101.2012.02618.x](https://doi.org/10.1111/j.1420-9101.2012.02618.x).
- 443 **Jagers P**, Klebaner FC, Sagitov S. On the path to extinction. *Proc National Acad Sci.* 2007; 104(15):6107–6111.
444 doi: [10.1073/pnas.0610816104](https://doi.org/10.1073/pnas.0610816104).
- 445 **Laffafian A**, King JD, Agrawal AF. Variation in the strength and softness of selection on deleterious mutations.
446 *Evolution.* 2010; 64(11):3232–3241. doi: [10.1111/j.1558-5646.2010.01062.x](https://doi.org/10.1111/j.1558-5646.2010.01062.x).
- 447 **Lande R**. Risks of population extinction from demographic and environmental stochasticity and random catas-
448 trophes. *Am Nat.* 1993; 142(6):911–927. doi: [10.1086/285580](https://doi.org/10.1086/285580).
- 449 **Lazenby BT**, Tobler MW, Brown WE, Hawkins CE, Hocking GJ, Hume F, Huxtable S, Iles P, Jones ME, Lawrence
450 C, et al. Density trends and demographic signals uncover the long-term impact of transmissible cancer in
451 Tasmanian devils. *J Appl Ecol.* 2018 Feb; 55(3):1368–1379. doi: [10.1111/1365-2664.13088](https://doi.org/10.1111/1365-2664.13088).
- 452 **Loeb LA**, Essigmann JM, Kazazi F, Zhang J, Rose KD, Mullins JI. Lethal mutagenesis of HIV with mutagenic
453 nucleoside analogs. *Proc Natl Acad Sci U S A.* 1999 Feb; 96(4):1492–7.
- 454 **Lord CJ**, Tutt ANJ, Ashworth A. Synthetic lethality and cancer therapy: lessons learned from the development
455 of PARP inhibitors. *Annu Rev Med.* 2015 Jan; 66(1):455–470. doi: [10.1146/annurev-med-050913-022545](https://doi.org/10.1146/annurev-med-050913-022545).
- 456 **Lynch M**, Bürger R, Butcher D, Gabriel W. The mutational meltdown in asexual populations. *J Hered.* 1993 Sep;
457 84(5):339–344.
- 458 **Lynch M**, Conery J, Burger R. Mutation accumulation and the extinction of small populations. *Am Nat.* 1995;
459 146(4):489–518. doi: [10.1086/285812](https://doi.org/10.1086/285812).
- 460 **Lynch M**, Conery J, Burger R. Mutational meltdowns in sexual populations. *Evolution.* 1995; 49(6):1067–1080.
- 461 **Lynch M**, Gabriel W. Mutation load and the survival of small populations. *Evolution.* 1990; 44(7):1725–1737.
462 doi: [10.1111/j.1558-5646.1990.tb05244.x](https://doi.org/10.1111/j.1558-5646.1990.tb05244.x).
- 463 **Martin G**, Aguilée R, Ramsayer J, Kaltz O, Ronce O. The probability of evolutionary rescue: towards
464 a quantitative comparison between theory and evolution experiments. *Phil Trans R Soc B.* 2013 Jan;
465 368(1610):20120088. doi: [10.1098/rstb.2012.0088](https://doi.org/10.1098/rstb.2012.0088).
- 466 **Matuszewski S**, Ormond L, Bank C, Jensen JD. Two sides of the same coin: A population genetics perspective
467 on lethal mutagenesis and mutational meltdown. *Virus Evolution.* 2017 Jan; 3(1). doi: [10.1093/ve/vex004](https://doi.org/10.1093/ve/vex004).
- 468 **McCallum H**, Jones M, Hawkins C, Hamede R, Lachish S, Sinn DL, Beeton N, Lazenby B. Transmission dy-
469 namics of Tasmanian devil facial tumor disease may lead to disease-induced extinction. *Ecology.* 2009 dec;
470 90(12):3379–3392. doi: [10.1890/08-1763.1](https://doi.org/10.1890/08-1763.1).
- 471 **McFarland CD**, Korolev KS, Kryukov GV, Sunyaev SR, Mirny LA. Impact of deleterious passenger mutations on
472 cancer progression. *Proc Natl Acad Sci USA.* 2013; 110(8):2910–2915. doi: [10.1073/pnas.1213968110](https://doi.org/10.1073/pnas.1213968110).
- 473 **McFarland CD**, Mirny LA, Korolev KS. Tug-of-war between driver and passenger mutations in cancer and other
474 adaptive processes. *Proc Natl Acad Sci USA.* 2014; 111(42):15138–15143. doi: [10.1073/pnas.1404341111](https://doi.org/10.1073/pnas.1404341111).
- 475 **Melzer AL**, Koeslag JH. Mutations do not accumulate in asexual isolates capable of growth and extinction—
476 Muller’s ratchet re-examined. *Evolution.* 1991 5; 45(3):649–655. doi: [10.1111/j.1558-5646.1991.tb04335.x](https://doi.org/10.1111/j.1558-5646.1991.tb04335.x).
- 477 **Metzger JJ**, Eule S. Distribution of the fittest individuals and the rate of Muller’s ratchet in a model with over-
478 lapping generations. *PLoS Comput Biol.* 2013; 9(11):e1003303. doi: [10.1371/journal.pcbi.1003303](https://doi.org/10.1371/journal.pcbi.1003303).
- 479 **Moran NA**. Accelerated evolution and Muller’s ratchet in endosymbiotic bacteria. *Proc Natl Acad Sci U S A.* 1996
480 Apr; 93(7):2873–8.
- 481 **Muller HJ**. The relation of recombination to mutational advance. *Mutat Res.* 1964; 1(1):2–9. doi: [10.1016/0027-5107\(64\)90047-8](https://doi.org/10.1016/0027-5107(64)90047-8).
- 482
483 **Neher RA**, Shraiman BI. Fluctuations of fitness distributions and the rate of Muller’s ratchet. *Genetics.* 2012;
484 191(4):1283–1293. doi: [10.1534/genetics.112.141325](https://doi.org/10.1534/genetics.112.141325).
- 485 **Pamilo P**, Nei M, Li W. Accumulation of mutations in sexual and asexual populations. *Genet Res.* 1987;
486 49(02):135–146. doi: [10.1017/S0016672300026938](https://doi.org/10.1017/S0016672300026938).

- 487 **Pariente N**, Sierra S, Lowenstein PR, Domingo E. Efficient virus extinction by combinations of a mutagen and
488 antiviral inhibitors. *J Virol*. 2001 Oct; 75(20):9723–30. doi: [10.1128/JVI.75.20.9723-9730.2001](https://doi.org/10.1128/JVI.75.20.9723-9730.2001).
- 489 **Poon A**, Otto SP. Compensating for our load of mutations: Freezing the meltdown of small populations. *Evolu-*
490 *tion*. 2000; 54(5):1467–1479.
- 491 **Rouzine IM**, Brunet É, Wilke CO. The traveling-wave approach to asexual evolution: Muller’s ratchet and speed
492 of adaptation. *Theor Popul Biol*. 2008; 73(1):24–46. doi: [10.1016/j.tpb.2007.10.004](https://doi.org/10.1016/j.tpb.2007.10.004).
- 493 **Rouzine IM**, Wakeley J, Coffin JM. The solitary wave of asexual evolution. *Proc Natl Acad Sci U S A*. 2003;
494 100(2):587–592. doi: [10.1073/pnas.242719299](https://doi.org/10.1073/pnas.242719299).
- 495 **Rowe G**, Beebe TJ. Population on the verge of a mutational meltdown? Fitness costs of genetic load for an
496 amphibian in the wild. *Evolution*. 2003 Jan; 57(1):177–81.
- 497 **Silander O**, Tenaillon O, Chao L. Understanding the evolutionary fate of finite populations: the dynamics of
498 mutational effects. *PLoS Biol*. 2007; 5(4):e94. doi: [10.1371/journal.pbio.0050094](https://doi.org/10.1371/journal.pbio.0050094).
- 499 **Silva JM**, Garcia JM, Dominguez G, Silva J, Rodriguez R, Portero JL, Corbacho C, Provencio M, España P, Bonilla
500 F. DNA damage after chemotherapy correlates with tumor response and survival in small cell lung cancer
501 patients. *Mutat Res*. 2000 Nov; 456(1-2):65–71.
- 502 **Stephan W**, Chao L, Smale J. The advance of Muller’s ratchet in a haploid asexual population: approximate
503 solutions based on diffusion theory. *Genet Res*. 1993; 61(3):225–231. doi: [10.1017/S0016672300031384](https://doi.org/10.1017/S0016672300031384).
- 504 **Wallace B**. Hard and soft selection revisited. *Evolution*. 1975; 29(3):465–473. doi: [10.2307/2407259](https://doi.org/10.2307/2407259).
- 505 **Wardlaw AM**, Agrawal AF. Temporal variation in selection accelerates mutational decay by Muller’s ratchet.
506 *Genetics*. 2012; 191(3):907–916. doi: [10.1534/genetics.112.140962](https://doi.org/10.1534/genetics.112.140962).
- 507 **Wielgoss S**, Barrick JE, Tenaillon O, Wiser MJ, Dittmar WJ, Cruveiller S, Chane-Woon-Ming B, Médigue C, Lenski
508 RE, Schneider D. Mutation rate dynamics in a bacterial population reflect tension between adaptation and
509 genetic load. *Proc Natl Acad Sci U S A*. 2013 Jan; 110(1):222–227. doi: [10.1073/pnas.1219574110](https://doi.org/10.1073/pnas.1219574110).
- 510 **Zeyl C**, Mizesko M, De Visser JAGM. Mutational meltdown in laboratory yeast populations. *Evolution*. 2001;
511 55(5):909–917. doi: [10.1111/j.0014-3820.2001.tb00608.x](https://doi.org/10.1111/j.0014-3820.2001.tb00608.x).

Appendix 1

Proof of Equation 5

Given real numbers b and x and $n \in \mathbb{N}$ (the nonnegative integers), let A denote the “almost diagonal” $n \times n$ matrix

$$A = \begin{pmatrix} 1 & b & & & \\ & x & bx & & \\ & & x^2 & bx^2 & \\ & & & \ddots & \ddots \\ & & & & x^{n-1} \end{pmatrix},$$

whose i th row is simply x^{i-1} multiplied by $(0 \ 0 \ \dots \ 0 \ 1 \ b \ 0 \ 0 \ 0 \ \dots)$, the 1 occurring in the i th position. Since A is upper triangular, so is its k th power (for $k \in \mathbb{N}$), with diagonal entries

$$A^k(i, i) = x^{k(i-1)}.$$

The superdiagonal entries aren’t quite as simple, but can also be expressed explicitly in terms of x , k and b .

Lemma 1. For $n, k \in \mathbb{N}$, $1 \leq i \leq n-1$ and $1 \leq j \leq n-i$ one has

$$A^k(i, i+j) = b^j x^{\frac{j(j-1)}{2} + k(i-1)} \prod_{\ell=1}^j \frac{x^{k+1-\ell} - 1}{x^\ell - 1},$$

provided that we declare $x^0 = 1$.

Proof. We induct on k . When $k = 1$, for $j = 1$ and any i the given expression becomes

$$bx^{k(i-1)} \frac{x-1}{x-1} = bx^{(i-1)} = A(i, i+1).$$

When $j \geq 2$, then the $\ell = 2$ factor in the product is $\frac{x^{2-2}-1}{x^2-1} = 0$, so that regardless of i the entire expression becomes 0, which again equals $A(i, i+j)$. We conclude that the stated result holds for $k = 1$.

Now assume the result is true for some $k \geq 1$. Since $A^{k+1} = A \cdot A^k$ and $A(i, \ell) = 0$ unless $\ell \in \{i, i+1\}$,

$$\begin{aligned} A^{k+1}(i, i+j) &= \sum_{\ell=1}^n A(i, \ell) A^k(\ell, i+j) \\ &= x^{i-1} A^k(i, i+j) + bx^{i-1} A^k(i+1, i+j) \\ &= x^{i-1} (A^k(i, i+j) + bA^k(i+1, (i+1) + (j-1))). \end{aligned}$$

Using the inductive hypothesis^a we obtain

$$\begin{aligned}
 & A^{k+1}(i, i+j) \\
 &= b^j x^{i-1 + \frac{(j-1)(j-2)}{2} + ki} \left(\prod_{\ell=1}^{j-1} \frac{x^{k+1-\ell} - 1}{x^\ell - 1} \right) \left(x^{j-k-1} \frac{x^{k+1-j}}{x^j - 1} + 1 \right) \\
 &= b^j x^{i-1 + \frac{(j-1)(j-2)}{2} + ki} \left(\prod_{\ell=1}^{j-1} \frac{x^{k+1-\ell} - 1}{x^\ell - 1} \right) \left(\frac{1 - x^{j-k-1} + x^j - 1}{x^j - 1} \right) \\
 &= b^j x^{i-1 + \frac{(j-1)(j-2)}{2} + ki + j - k - 1} \left(\prod_{\ell=1}^{j-1} \frac{x^{k+1-\ell} - 1}{x^\ell - 1} \right) \left(\frac{x^{k+1} - 1}{x^j - 1} \right) \\
 &= b^j x^{\frac{j(j-1)}{2} + (k+1)(i-1)} \left(\prod_{\ell=1}^{j-1} \frac{x^{k+1-\ell} - 1}{x^\ell - 1} \right) \left(\frac{x^{k+1} - 1}{x^j - 1} \right) \\
 &= b^j x^{\frac{j(j-1)}{2} + (k+1)(i-1)} \left(\prod_{\ell=1}^j \frac{1}{x^\ell - 1} \right) \left(\prod_{\ell=0}^{j-1} (x^{k+1-\ell} - 1) \right) \\
 &= b^j x^{\frac{j(j-1)}{2} + (k+1)(i-1)} \left(\prod_{\ell=1}^j \frac{x^{(k+1)+1-\ell} - 1}{x^\ell - 1} \right),
 \end{aligned}$$

which shows that the formula holds for the exponent $k+1$. This concludes the proof. $\square$

^aStrictly speaking, the inductive hypothesis will only apply to the term $A^k(i+1, (i+1) + (j-1))$ when $j \geq 2$. However, if we adopt the convention that any empty product is equal to one, the expression stated in the result agrees with $A^k(i+1, (i+1) + (j-1))$ when $j = 1$ as well.

Appendix 2

Overestimation of the Expected Extinction Time

As shown in the **Model** section, an approximation is made in **Equation 9** in order to derive analytical expressions for click times (t_k) and time to extinction (**T**). **Figure 2D**, **Figure 3A**, **Figure 3B**, and **Figure 3–Figure Supplement 2A** show that the approximation is quite accurate over broad ranges of parameters but consistently overestimates **T**.

Overestimation of **T** is partly explained by the concavity of $E[\tau_k]$ as a function of x_k in Equation 6. The second derivative with respect to x_k is

$$\frac{d^2 E[\tau_k]}{dx_k^2} = - \sum_{i=1}^{\infty} \left(\varphi_k^{(i)}(0) \right)^{x_k} \ln^2 \left(\varphi_k^{(i)}(0) \right)$$

which is negative because both factors of each term in the sum are positive and hence the function is concave. By the reversed Jensen's inequality, for a concave function, $E[f(X)] < f(E[X])$, which implies that $E[\tau_k]$ is consistently overestimated by **Equation 6**. Thus, **Equation 10** overestimates **T**.

Figure 2-Figure supplement 1. Variability of extinction time declines with population size. Blue circles show coefficient of variation of T , $CV[T]$, for the data shown in **Figure 2D**. Red line shows $CV[T]$ calculated numerically using **Equation 23**.

Figure 3-Figure supplement 1. Variability of extinction time declines with mutation rate and is approximately invariant with selection coefficient. Blue circles show coefficient variation of T , $CV[T]$, for the data shown in the corresponding panel of **Figure 3**. Red line shows $CV[T]$ calculated numerically using **Equation 23**.

Figure 3-Figure supplement 2. Theoretical predictions for low s become more accurate with increasing population size. **(A)** Extinction times of populations with $u = 0.01$ and $s = 10^{-3}$ over a range of initial populations sizes, n_0 . Blue line shows mean values of T based on stochastic simulations of 10^4 replicate populations for 41 values of n_0 evenly spaced on a log-scale over 4 orders of magnitude. Light blue region indicates $\bar{T} \pm SD[T]$. If $SD[T] > \bar{T}$, the lower bound of the region was set to zero. **(B)** Variability of extinction times shown in **(A)**. Blue circles show coefficient variation of T , $CV[T]$. Red line shows $CV[T]$ calculated numerically using **Equation 23**.

Figure 4-Figure supplement 1. Expected extinction time of the mutation-free class and variability of extinction time. (A) Expected extinction time of the mutation-free class, t_0 , used in **Figure 4B. (B)** Coefficient of variation of T , $CV[T]$, calculated numerically using **Equation 23**.