Sustaining yam yields amidst climate threat in the forest – savannah transition zone of Ghana

Frimpong F.^{*12}, Owusu Danquah E.², Ennin S. A.², Asumadu H.³, Aidoo A. K.², Maroya N.³

¹Forschungszentrum Jülich, Institute of Bio-Geosciences 2, Plant Sciences, Germany

²Council for Scientific and Industrial Research- Crops Research Institute, P. O. Box 3785 Kumasi, Ghana.

³IITA, Ibadan, Nigeria

*Corresponding Author's Email: f.frimpong@fz-juelich.de

Abstract

With about 70% of yam tuber been water, yield is critically affected during bulking as a result of onset of temporal drought. As a consequence of climate change, farmers who are into *Dioscorea rotundata* (white yam) production for local and international market lose their investments mainly due to erratic precipitation, drought spells culminating into low yields of just 12t/ha compared to the potential of about 22-49t/ha depending on the variety. Innovative land uses technologies with higher and sustained productivity for yam production are imperative. This study verifies improved agronomic package for sustainable yam production in yam growing areas in the forest – savannah transition zone of Ghana during the 2015 and 2016 cropping seasons. The improved agronomic package included use of ridging as seedbed, seed treatment before planting, fertilizer application at a rate of 30:30:36 N:P₂O₅:K₂O kg/ha plus 15 kg/ha Mg and 20 kg/ha S as MgSO₄ and the use of minimum stakes (trellis; 30-50% less number of stakes used by farmers staking). This was compared with farmers' practice which consisted of mounding, no fertilizer application and no seed treatment. The results revealed significant ($P \leq 0.01$) yam yields of more than 60% difference between the improved agronomic practice and farmers' practice from Ejura, Atebubu and Kintampo yam growing communities. Adoption of improved agronomic practices does not only sustain yam production and address deforestation but also provide higher returns on investments promoting climate resilience by small holders.

Key Words

Climate smart agriculture, Fertilizer, Improved technology, Seed treatment, Trellis staking, Yam

Background

In Ghana, crops are already experiencing heat stress, drought spells, several pests and diseases outbreak and shorter growing crop duration as a consequent of the changing climate (1). Thus, a potential catastrophe for smallholder farmers and the millions of people who regularly grow rain-fed full season crops such as yam, cocoyam, rice etc.(2–5). Choices about what to grow are often dictated by the ability of the rainfall regime to support moisture for plant growth (6). One way around this would be to breed for varieties with shorter crop maturity durations or management interventions that build on the resilience of cropping production systems to reduce shocks if the shocks from the climate change cannot be done away with. Evidence suggests that climate smart agriculture can make a contribution to mitigation by supporting more efficient use of fertilizers, weed management and reduced staking options in yam production (1,7,8).

Yam, an important staple food crop across West Africa is a major non-traditional export crop in Ghana contributing to about 16% of the National Agricultural Gross Domestic Product (9,10). However, there are a number of challenges that hamper the production and productivity of yam. Predominantly amongst them are; inadequate rainfall, low soil fertility, weed infestation, pests and diseases in the field (foliar and soil borne) and inadequate storage facilities, attack by organisms such as rodents, access to quality improved seed, implements for mechanization etc.(8,11). Others include shortage of stakes especially from deforested areas and guinea savannah regions. This is as a consequent of clearing new lands year after year popularly known as shifting cultivation and slash and burn agriculture, inadequate labour in view of the labor intensive nature of yam cultivation (e.g. for preparation of mounds, staking, harvesting)(8,12,13). This current yam production system where there is annual shifting of farm to new lands is not sustainable and therefore the urgent need to disseminate an environmentally sound yam production technology that would increase yield and sustain production on continuously cropped fields particularly in the face of climate change (14).

As a follow up on an on-station evaluation conducted in 2013 and 2014 cropping seasons at the CSIR-Crops Research Institute research stations, recommended agronomic package of planting treated yam seeds on ridges with fertilizer rate of 30:30:36 N: P₂O₅:K₂O kg/ha plus 15 kg/ha Mg and 20 kg/ha S as MgSO₄ and trellis staking were verified by comparing it with the farmers practices on farmers' fields at Ejura, Atebubu and Kintampo yam growing communities. The use of ridges and yam seed treatment helps to maintain optimum number of stands per unit area and fertilizer addresses the soil nutrient depletion whiles the trellis staking option uses ropes and few stakes to address the challenge of scarcity of stakes and cutting of more trees/bamboo for staking. The study has a major objective of validating and demonstrating improved yam production technologies (macronutrients (NPK) and micronutrients (Mg & S), minimum staking option and seed treatment) to major yam growing communities in Ghana.

Methodology

Study sites characteristics

The experiments were conducted in 2015 and 2016 cropping seasons at Ejura-Sekyeredumasi (Aframso/Teacherokrom, Ashakoko, Dromankuma & Nkwanta), Atebubu- Amantin (Adom, Dagaati Line, Munchunso & Nwowamu) and Kintampo North (Asantekwa, Suamre, Babaso/Yabroso & Kintampo Magazine) districts of Ghana (Fig 1). These areas lie in the forest-savannah transition agro-ecological zone and amongst the major yam growing areas of Ghana (15). Eight farmers (4 randomly selected for analysis) from each of the three operational areas were selected for the study every season (Table 1). Mean annual rainfall (mm) for 2015 & 2016 across locations are shown in the map (Fig 1). The data were sourced from the local district weather stations which revealed a reduced rainfall amount (mm) in 2016 compared to 2015 with Kintampo communities severely affected (Fig 1). Mean annual rainfall (mm) pairs recorded for 2015 and 2016 were 1256:1034, 929:769, and 863:795 at

Ejura, Atebubu and Kintampo respectively (Fig 1). These locations have bimodal rainfall i.e. major rainy season from March-Mid August and minor rainy season from September-November; peak in October. Temperature ranges from 25-39 °C with soil type of Ferric Acrisol; Asuansi series, upper top soil consist of 5cm grayish brown sandy loam topsoil of dark brown gritty clay loam (16).

Figure 1. Map of Ghana on the (right), zoomed in on Ejura-Sekyeredumasi, Atebubu – Amantin and Kintampo North districts (left) of the forest-savanna transition zone. Farming communities where the studies were undertaken for 2015 and 2016 cropping seasons are illustrated with dots (.) with their names beside. Mean annual rainfall (mm) per location are depicted by bar plots for each season.

Table 1. Farmers and planting dates selected from each of the operational areas for the analysis in 2015 and 2016.

Year	Atebubu		Ejura		Kintampo	
	Farmers and their respective planting dates					
2015	Farmer A	2-Jun	Farmer A	11-Jun	Farmer A	21-Jun
	Farmer B	5-Jun	Farmer B	11-Jun	Farmer B	21-Jun
	Farmer C	6-Jun	Farmer C	11-Jun	Farmer C	21-Jun
	Farmer D	4-Jun	Farmer D	12-Jun	Farmer D	21-Jun
2016	Farmer E	4- Jun	Farmer E	1-Jun	Farmer E	9-Jun
	Farmer F	5-Jun	Farmer F	2-Jun	Farmer F	10-Jun
	Farmer G	5-Jun	Farmer G	3-Jun	Farmer G	10-Jun
	Farmer H	6-Jun	Farmer H	2-Jun	Farmer H	11-Jun

Experimental design

A randomized complete block design with each farmer as a replicate was used for the study. A total of 8 trials (8 replicates) were established in all the operational areas in 2015 and 2016. Local white yam variety “Dente” of *Dioscorea rotundata* species was planted and subjected to two treatment applications from start of planting till harvest. Planting of yam across these locations were started in June and completed by 21-June of each year (Table 1). Harvesting

of yam was completed by the end of December of each year. The treatments were recommended (improved agronomic practice) and local technology/farmers' practice. The recommended practice for yam production included a package of; treating yam seed before planting with fungicide and insecticide, use of ridging as seedbed, fertilization at 30:30:36 N: P₂O₅:K₂O Kg/ha plus 15Kg/ha of Mg and 20Kg/ha S as MgSO₄ and use of trellis for staking while in the farmers' practice, farmers were allowed to use their local technology (planting on mounds), no pre-treatment of seed, no fertilizer application for comparison. Continuously cropped fields that farmers would normally not use for yam production were selected in each operational area for the study. Each improved agronomic field had an area of a quarter of an acre (0.25ac/0.1ha) planted at 1.2m inter row and 0.8m on the ridges (10,416 stands/ha).

The same size (0.25ac/0.1ha) was demarcated for the farmers practice treatment where they mounded sparsely to cover the entire field (3,400 – 4,000 stands/ha). Each farmer field was considered as a replicate and analysis compiled for each district/operational area together. The Fertilizer treatment was applied at 50% split at 5-6 weeks and 11-12 weeks after planting in all the locations. The seed setts (200-250g) of the improved agronomic fields were treated with Dursban (Chlorpyrifos from Dow Agro Sciences; 1.25 l/ha) and Mancozeb (Dithiocarbamate from Ag-Chem Africa 80%; 75 g in 15 l of water) before planting. Farmers' sett sizes used ranged between 350g to 650g depending on each farmer. Emerged weeds in the improved agronomic fields were controlled with glyphosate (2.5 liter per ha) before the sprouting of the yam while farmers only slashed on their fields. There after weeds were manually controlled with cutlass and hoe in either improved agronomic field or farmers' field by hoeing. In 2016, sensory evaluations were conducted with 50 participants (who were mainly farmers from the localities) in all areas after eating boiled yam during the December harvest. Fertilized and unfertilized boiled yam (coded at the blind side of the participants) using one on one questionnaire interviews, farmers scored for taste, texture, aroma and acceptability (Table 3).

Data collection and analysis

During harvest, the tubers were grouped into two; ware yam (tuber sizes of more than 500g) and seed yam (500g and below) and weighed separately for each of the practices. Four replications from each operational area (Ejura, Atebubu and Kintampo) and across the two seasons (2015, 2016) were subjected to statistical analysis. Data on stand harvested, weight of ware yam, weight of seed yam and total yam yield collected were subjected to one way analysis of variance linear model at 5% significant level using 'R' statistical software with the practice treatment as the independent variable. Where treatment means differ, Tukey's HSD test was used to group them and visualized with bar graphs using MS excel 2010. Percentage differences of total yam yield harvested between the improved practice and the farmer's was calculated based on the formula;

$$\text{Percentage difference} = \frac{(\text{improved agronomic practice} - \text{farmers' practice})}{(\text{improved agronomic practice} + \text{farmers' practice}/2)} \times 100$$

Sensory evaluation through one on one interviews upon tasting boiled yam from fertilized and unfertilized samples were calculated with data from 50 participants for each location who were mainly farmers. In order to deduce return on investment after venturing in any of the practices was subsequently calculated using benefit cost ratio for each of the locations using total yam yield and price per kg of yam at the time of harvest.

Results and Discussion

Influence of improved agronomic technology on yam tuber yield

Generally, total yam tuber yields were high in 2015 than in 2016 cropping season irrespective of the practice (Figs 2-4). The improved agronomic practice fields had significantly ($P < 0.01$) higher total tuber yields compared to the farmers' practice fields across the locations despite the season (Figs 2-4). Availability of moisture is critical for yam to sprout and establish during early growth stages and vital to bulk bigger tubers (11,17–20). Higher rainfall during 2015 cropping season (Fig 1) might have ensured yam establishment and increased overall

yields compared to 2016. The use of improved agronomic package of ridging, yam seed pre-treatment, trellis staking and fertilizer rate of 30:30:36 N: P₂O₅:K₂O kg/ha plus 15 kg/ha Mg and 20 kg/ha S as MgSO₄ resulted in total tuber yield percentage difference over farmers' practice of mounding and planting of 68.6%, 78.3%, 80.7% for the 2015 cropping season and 113.6%, 113.9%, 120% for 2016 cropping season in Ejura, Atebubu and Kintampo farming communities (Fig 5) respectively. Stand count/ha on the ridges were significantly (P<0.01) higher on the improved agronomic practice fields than farmers' practice fields for all the locations and across season (Figs 2-4). This we attribute to the use of ridges which made it possible to plant at 1.2m between ridges and 0.8m on the ridges resulting in a planting density of about 10,416 stands per hectare whiles the farmers practice of mounding were relatively sparse (1.5m-2m) resulting in just about 3,400 – 4,000 stands/ha. Thus, ridging helped maintain optimum number of plants and promoted efficient use of fertilizer and conserved moisture than mounding.

It was observed that the improved agronomic fields had better yam canopy as a result of the combined effect of trellis staking and ridging. Revealing similar tuber yield trends on fertilized mounds and unfertilized ridges (8) supports the argument that fertilizer application helps the farmer to achieve value by being more efficient and profitable on ridges than on mounds. Moreover, pre-treating seed yam before planting on the improved agronomic fields resulted in a reduction in yam rot and increased stands/ha than on the farmers' practice fields where seeds were not treated resulting in lower stands/ha. Seed treatment for pest and disease before planting is recommended to promote sprout rate ensuring improvement in final stand density culminating into overall high yam yields (11,12,17). Erratic rainfall and prolong drought require technologies that enable the soil to conserve moisture and promote nutrient use efficiency in order to increase resiliency of any cropping system. An observation we made during the studies was that drought was more pronounced on the farmers' practice which used relatively sparse mounds and vertical staking option than on the improved agronomic

practices which used ridges and trellis staking option. Similar studies suggest that planting on ridges can maintain optimum plant stands and conserve more moisture than mounds resulting in more efficient water used on ridges than on mounds (7,8,17,21). These attributes makes the use of ridges more soil nutrient efficient upon application of fertilizer than on mounds. Furthermore, (8) made a similar observation with planting on ridges increasing yields significantly than on mounds.

It is recommended that adopting and following through the improved agronomic package based on results of 2016 cropping season (Figs 1-5) where rainfall were considerably lower illustrates the resilience ability of it during reduced precipitation. In spite of the gains in yam fertilization, there are perceptions and claims by some consumers and farmers in the public space that fertilizing yam leads to rots and reduces the overall shelf life. We recommend further research into these claims as we settled the dust in this paper (Table 3) on claims that fertilizer affected the taste quality of yam.

Figure 2. Yam tuber yields and stand count as influenced by improved agronomic practice and farmers' practice in the Ejura farming communities for 2015 and 2016 cropping seasons. Mean values and standard errors (n = 4) are plotted. Index letters above the bars indicate significant differences ($P < 0.01$) between media not sharing the same letter by Tukey's HSD test. Asterisks indicate significant differences ($P < 0.01$) between the two practices.

Figure 3. Yam tuber yields and stand count as influenced by improved agronomic practice and farmers' practice in the Atebubu farming communities for 2015 and 2016 cropping seasons. Mean values and standard errors (n = 4) are plotted. Index letters above the bars indicate significant differences ($P < 0.01$) between media not sharing the same letter by Tukey's HSD test. Asterisks indicate significant differences ($P < 0.01$) between the two practices.

Figure 4. Yam tuber yields and stand count as influenced by improved agronomic practice and farmers' practice in the Kintampo farming communities for 2015 and 2016 cropping seasons. Mean values and standard errors ($n = 4$) are plotted. Index letters above the bars indicate significant differences ($P < 0.01$) between media not sharing the same letter by Tukey's HSD test. Asterisks indicate significant differences ($P < 0.01$) between the two practices.

Figure 5. Percentage differences between the two practices across two seasons calculated for each location based on their total yam yield (kg/ha).

Partial budgeting and cost benefit analysis of the two practices

Table 2 presents the partial budgeting and cost benefit analysis of “*Dente*” white yam production under improved agronomic practice and farmers' practice in Ejura, Atebubu and Kintampo operational areas. The results revealed that irrespective of location or the season, yam planted with the improved agronomic package had higher benefit to cost ratio compared to farmers' practice. Benefit to cost ratios of 4.76:3.64, 4.04:2.48 and 2.01:1.08 were achieved for Ejura, Atebubu and Kintampo communities respectively for the 2015 cropping season (**Table 2**) in sequence of improved technology: farmers' practice. Thus, when a farmer invests GhC 1.00 in yam production using the recommended improved technology a profit of GhC 3.76, GhC 3.04 and GhC 1.01 was to be accrued in addition to the GhC 1.00 invested capital at Ejura, Atebubu and Kintampo respectively during the 2015 season. During the 2016 cropping season, drought was more intense particularly for Atebubu and Kintampo areas (Figure 1). This however did not affect benefit to cost ratio for using the improved agronomic package thus achieving 3.76, 3.03 and 1.33 compared to 1.16, 0.75 and -0.55 for Ejura, Atebubu and Kintampo communities respectively (**Table 2**). Thus, a profit of GhC 2.76, GhC 2.03 and GhC 0.33 was to be accrued in addition to the GhC 1.00 invested capital upon the use of improved agronomic practices at Ejura, Atebubu and Kintampo respectively. The use of the farmers' practice resulted in total loss of GhC 1.55 in Kintampo area (**Table 2**). This suggest that the use of the improved agronomic practice would not only increase and sustain

yields on continuously cropped fields but also the it is the best option during drought spells, erratic and reduced rainfall conditions. The improved agronomic package thereby increases farmer's resilience in dealing with such harsh weather conditions with assured returns on their investments.

Table 2. Partial budget and cost benefit analysis of white yam production with improved technology and farmers practices at Ejura, Atebubu and Kintampo farming communities for the 2015 and 2016 cropping seasons.

Location	Ejura operational area				Atebubu operational area				Kintampo operational area			
Practice	Improved agro nom ic prac tice	Farm ers' pra ctic e	Improved agro nom ic prac tice	Farm ers' pra ctic e	Improved agro nom ic prac tices	Farm ers' pra ctic e	Improved agro nom ic prac tices	Farm ers' pra ctic e	Improved agro nom ic prac tices	Farm ers' pra ctic e	Improved agro nom ic prac tices	Farm ers' pra ctic e
Year	2015	2015	2016	2016	2015	2015	2016	2016	2015	2015	2016	2016
Average yield (kg/ha)	1780	870	1420	391	1670	730	1240	340	1060	450	8200	2050
Adjusted yield (kg/ha)	1602	783	1278	351	1503	657	1116	306	9540	405	7380	1845
Farm gate price in December each year (C/kg)	1.4	1.4	1.45	1.4	1.3	1.3	1.4	1.4	1.2	1.2	1.2	1.2
Gross benefit(C/ha)	2242	109	1853	510	1953	854	1562	428	1144	486	8856	2214
Cost of chemical fertilizer, glyphosate, fungicide & pesticide(C)	355	0	355	0	355	0	355	0	355	0	355	0
Labour cost for application of fertilizer & others (C/ha)	153	0	153	0	153	0	153	0	153	0	153	0
Cost of land clearing and stumping (C/ha)	320	0	320	0	290	0	290	0	300	0	300	0
Construction of ridges (C/ha)	300	0	300	0	300	0	300	0	300	0	300	0
Construction of mounds (C/ha)	0	200	0	200	0	250	0	250	0	240	0	240
Cost of seed yam(C)	955	455	955	455	955	455	955	455	955	455	955	455
Labour cost of planting(C/ha)	161	139	161	139	160	140	160	140	160	140	160	140
Cost of stakes(C/ha)	392	282	392	282	392	282	392	282	392	282	392	282
Labour cost of staking(C/ha)	400	460	400	460	415	500	415	500	405	470	405	470

Cost of weeding and reshaping(C/ha)	675	675	675	675	675	675	675	675	600	600	600	600
Harvesting cost(C/ha)	182	153	182	153	182	153	182	153	182	153	182	153
Total cost that vary (C)	3893	2364	3893	2364	3877	2455	3877	2455	3802	2340	3802	2340
Net benefit (C)	18535	8598	14638	2738.55	15662	6086	11747	1829	7646	2520	5054	-126
Benefit cost/Ratio	4.76	3.64	3.76	1.16	4.04	2.48	3.03	0.75	2.01	1.08	1.33	-0.05

Influence of fertilizer on the taste of boiled yam

Dente yam planted in 2016 under the improved agronomic practice (fertilized) and farmers' practice (unfertilized) were boiled after harvest and given to fifty participants each from Atebubu, Ejura and Kintampo who were mainly farmers for sensory evaluation (Table 3). Participants were not previewed as to whether the yam they evaluated at a given time was fertilized or unfertilized as they were coded in order to avoid bias. Participants assessed the various boiled yam on three culinary qualities: 'taste', 'texture' and 'aroma' (Table 3) based on their individual preferences from a scale of 1 up to 5 with 1 been the best score and 5 as the worst. Overall acceptability and STD acceptability on the three traits; taste, texture and aroma was subsequently calculated following the approach of (8). The results was in line with previous results by (8) that, contrary to the view that the use of fertilizer in yam production affects the quality of yam, sensory evaluation showed that the culinary qualities of fertilized yam is good and could even be better than unfertilized yam.

Table 3. Sensory evaluation of fertilized and unfertilized boiled yam involving farmers from Atebubu, Ejura and Kintampo farming communities.

Location	Treatment	Taste	Texture	Aroma	Overall acceptability	STD acceptability
Atebubu	Fertilized yam (30 30 36 N-P ₂ O ₅ -K ₂ O (Kg/ha) + 20kg S & 15kg 15 Mg as MgSO ₄)	2.30	2.5	2	2.20	0.87
	Unfertilized yam (0kg/ha)	2.60	2.5	2.4	2.50	0.86
Ejura	Fertilized yam (30 30 36 N-P ₂ O ₅ -K ₂ O (Kg/ha) + 20kg S & 15kg 15 Mg as MgSO ₄)	2.90	2.9	2.8	2.80	0.91
	Unfertilized yam (0kg/ha)	2.90	2.9	2.8	2.80	0.91
Kintampo	Fertilized yam (30 30 36 N-P ₂ O ₅ -K ₂ O (Kg/ha) + 20kg S & 15kg 15 Mg as MgSO ₄)	2.10	2.2	2.4	2.20	0.81
	Unfertilized yam (0kg/ha)	2.70	3.3	3.1	3.10	0.97

n=50, Score Scale 1-5; 1=best, 5=worst

Conclusion and Policy Implication

The improved agronomic technology has proven to be climate resilient comparing the overall yield (significantly >60% difference) trends of 2015 and 2016 seasons against farmers' practice given the rainfall amounts and pattern in those years. The overarching difference between what farmers do today and the improved agronomic model is the intensification drive and a higher use efficiency (staking, nutrient, soil moisture conservation, improved sprouting) of the technology at a given area as a consequence of the high planting density allowed by the linear arrangement of ridging, trellis staking and seed treatment. It is anticipated that more and

more farmers who have shown keen interest would adopt the technology through the gains made. Claims on yam fertilization affecting taste quality proved otherwise with even a higher overall acceptability. Further research is needed to understand claims and perceptions by farmers on fertilizer affecting yam storage. Up-scaling of the improved agronomic technology to other yam growing communities would further augment the gains already made.

Acknowledgement

We thank the Bill and Melinda Gates Foundation who funded this research work through YIIFSWA project that was coordinated by IITA, Nigeria through CSIR-Crops Research Institute. We are grateful to Mr. Henry Azotibah (Ejura), Mr. Kyeremeh (Atebubu) all of MoFA and Mr. Enoch Bessah of CSIR for their invaluable support and advice during data acquisition and analysis.

Abbreviations

CSIR – Council for Scientific and Industrial Research

FAO – Food and Agriculture Organisation

IITA –International Institute for Tropical Agriculture

MoFA – Ministry of Food and Agriculture

YIIFSWA – Yam Improvement for Income and Food Security in West Africa

References

1. Essegbey GO, Nutsukpo D, Karbo N, Zougmore R. National Climate-Smart Agriculture and Food Security Action Plan of Ghana (2016-2020) [Internet]. Copenhagen, Denmark; 2015. Report No.: 139. Available from: www.ccafs.cgiar.org
2. Maliki R, Toukourou M, Sinsin B, Vernier P. Productivity of yam-based systems with herbaceous legumes and short fallows in the Guinea-Sudan transition zone of Benin. *Nutr Cycl Agroecosystems*. 2011;92(1):9–19.
3. Mignouna BD, Abdoulaye T, Alene AD, Asiedu R, Manyong VM. Characterization of Yam-growing Households in the Project Areas of Ghana. Nigeria; 2014.
4. Srivastava AK, Gaiser T, Cornet D, Ewert F. Estimation of effective fallow availability for the prediction of yam productivity at the regional scale using model-based multiple scenario analysis. *F Crop Res*. 2012;131(November):32–9.
5. Amekudzi L, Yamba E, Preko K, Asare E, Aryee J, Baidu M, et al. Variabilities in Rainfall Onset, Cessation and Length of Rainy Season for the Various Agro-Ecological Zones of Ghana. *Climate* [Internet]. 2015;3(2):416–34. Available from: <http://www.mdpi.com/2225-1154/3/2/416/>
6. Oteng-Darko P, Frimpong F, Sarpong F, Amenorfe LP. Promoting smallholder irrigation for food Security-A review. *African J Food Integr Agric* [Internet]. 2018;2:1–7. Available from: www.asofost.org/journals/index.php/ajfia
7. Owusu Danquah E, Ennin SA, Lamptey JNL, Acheampong PP. Staking Options for Sustainable Yam Production in Ghana. *Sustain Agric Res* [Internet]. 2015;4(1):106–13. Available from: <http://dx.doi.org/10.5539/sar.v4n1p106>
8. Ennin SA, Issaka RN, Acheampong PP, Numafo M, Owusu Danquah E. Mechanization , fertilization and staking options for environmentally sound yam production. *African J Agric Res*. 2014;9(29):2222–30.
9. FAO. The State of Food and Agriculture. Rome, Italy; 2006. (37).
10. Ghana Standard Board. Roots and Tubers –Specification for fresh yams. Accra, Ghana; 2011. (2nd Edition). Report No.: GS 150.
11. Bhattacharjee R, Gedil M, Sartie A, Otoo E, Dumet D, Kikuno H, et al. Dioscorea. In: *Wild Crop Relatives: Genomic and Breeding Resources.Industrial Crops*. C. Kole. Springer-Verlag Berlin Heidelberg; 2011. p. 71–96.
12. Asiedu R, Sart A. Crops that feed the World 1. Yams: Yams for income and food security. *Food Secur Springer*. 2010;2:305–15.

13. Akwaag, A. A., Berchie, J. N., Echavez, M. L., Dapaah, H. K., Kebede, T., Njue, S. W., Paraoam, I., and Labrada HR. Sustainable Farming Practices- Towards Reducing the Expansion of the Forest Savanna Transitional Zone of the Sekyere-West and Ejura-Sekyeredumase Districts of the Ashanti Region of Ghana. International Center for Development Oriented Research in Agricultu. 2000. (86).
14. Otoo E. Yam production in Ghana: A food security enhancer or an environmental degrader. In: Akoroda M, editor. 8th Symposium of the International Society for Tropical Root Crops Africa Branch Triennial Workshop. Ibadan, Nigeria; 2001. p. 387–91.
15. Ministry of Food and Agriculture. Agriculture in Ghana, Facts and figures (2010) [Internet]. Accra, Ghana; 2010. Available from: <http://mofa.gov.gh/site/wp-content/uploads/2011/10/AGRICULTURE-IN-GHANA-FF-2010.pdf>
16. Adu S. V. and Asiamah R. D. Soils of the Ayensu-Densu Basin, Central, Eastern and Greater Accra Regions of Ghana. Council for Scientific and Industrial Research (CSIR) – Soil Research Institute Memoir, Kumasi, Ghana; 1992. Report No.: 9.
17. Ennin S. A., Owusu Danquah E., Frimpong F., Akom M., Osei-Adu J., Lamptey J. N. L., Oteng-Darko P., Osei K., Osuman A. S., Aidoo A. K., Adama I., Appiah-Danquah P. HJ. Yam Production and Marketing Guide. H. Adu-Dapaah, J. N. L. Lamptey, I. S. Banning, B. N. Frimpong JOA and SAE, editor. Kumasi, Ghana: CSIR-Crops Research Institute; 2016.
18. Maroya NG, Balogun M, Asiedu R, Aighewi B, Lava Kumar P, Joao A. Yam Propagation Using ‘ Aeroponics ’ Technology. Annu Res Rev Biol. 2014;4(24):3894–903.
19. Maroya NG, Asiedu R, Kumar PL, Lopez-Montes A, Orchard J, Ndiame F. YIIFSWA Project Description [Internet]. Ibadan, Nigeria; 2014. (YIIFSWA Working Paper). Report No.: 1. Available from: www.iita.org
20. Maroya NG, Balogun M, Asiedu R. Seed yam Production in an Aeroponics System: A Novel Technology [Internet]. Ibadan, Nigeria; 2014. (YIIFSWA Working Paper Series). Report No.: 2. Available from: www.iita.org
21. Ennin SA, Otoo E, Tetteh FM. Ridging, a mechanized alternative to mounding for yam and cassava production. West African J Appl Ecol. 2009;15(1).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5