1

2 **The coast-wide collapse in marine survival of west coast Chinook and steelhead: slow-**
3 **moving catastrophe or deeper failure?**

4

5

6 **Short title:** Coast-wide survival of Chinook and steelhead

7

8 David W. Welch*, Aswea D. Porter, Erin L. Rechisky

9 Kintama Research Services, 4737 Vista View Cr., Nanaimo, B.C. Canada V9V 1N8

10

11 * Corresponding Author

12 E-Mail: david.welch@kintama.com (DWW)

13

14 **Key words:** Chinook salmon; steelhead; Columbia River; Snake River; marine survival;
15 dams; delayed mortality; salmon farming; aquaculture; climate change

16

17 **Abbreviations**

18 BC- British Columbia, Canada

19 FCRPS- Federal Columbia River Power System

20 PSC- Pacific Salmon Commission

21 SAR-Smolt to Adult Return (Survival)

22

23 Abstract

24 Accelerating decreases in survival are evident for northern Hemisphere salmon
25 populations. We collated smolt survival and smolt-to-adult (marine) survival data for
26 all regions of the Pacific coast of North America excluding California to examine the
27 forces shaping salmon returns. A total of 3,055 years of annual survival estimates were
28 available for Chinook (*Oncorhynchus tshawytscha*) and steelhead (*O. mykiss*). This
29 dataset provides a fundamentally different perspective on west coast salmon
30 conservation problems from the previously accepted view. We found that marine
31 survival collapsed over the past half century by a factor of at least 4-5 fold to similar
32 low levels (~1%) for most regions of the west coast. The size of the decline is too large
33 to be compensated by freshwater habitat remediation or cessation of harvest, and too
34 large-scale to be attributable to specific anthropogenic impacts such as dams in the
35 Columbia River or salmon farming in British Columbia. Within the Columbia River,
36 both smolt survivals during downstream migration in freshwater and adult return rates
37 (SARs) of Snake River populations, often singled out as exemplars of poor survival,
38 appear unexceptional and are in fact higher than estimates reported from other regions
39 of the west coast lacking dams. Formal Columbia River rebuilding targets of 2-6%
40 SARs may therefore be unachievable if regions with nearly pristine freshwater
41 conditions also fail to achieve these targets. Finally, we present case studies
42 demonstrating that the historical response to evidence that the salmon problems are
43 primarily ocean-related was to re-emphasize freshwater actions and to stop work on
44 ocean issues. With ocean temperatures forecast to increase far further, the failure of

45 management to identify the drivers of salmon collapse and respond appropriately

46 suggest that the future of most west coast salmon populations is bleak.

47

48 Introduction

49

50 The total abundance of salmon in the North Pacific has now reached record
51 levels [2-4]; however, a dramatic contrast in the winners and losers is obscured by this
52 milestone. Most of the increased abundance is in the lowest valued species (pink
53 *Oncorhynchus gorbuscha* and chum *O. keta* salmon) in far northern regions, at least
54 partly due to major efforts at ocean ranching of these two species [4]. In contrast while
55 essentially all west coast North American Chinook (*O. tshawytscha*) populations
56 (including Alaska) are now performing poorly with dramatically reduced productivity
57 [6]. The situation is similar for most southern populations of steelhead (*O. mykiss*) [7],
58 coho (*O. kisutch*) [8, 9], and sockeye (*O. nerka*) [10-13]. These poorly performing
59 species are of higher economic value and the preferred focus of First Nations, sport,
60 and many commercial fisheries.

61

62 The historical geographic pattern of declines in salmon abundance (greatest
63 problems in the south, least to the north) were originally assumed to reflect a freshwater
64 anthropogenic cause because of the greater degree of terrestrial (i.e., freshwater) habitat
65 modification in the more populous southern regions of the west coast [14, 15]. The
66 growing appreciation of ocean climate change [16-18] has brought a greater awareness

67 of the role of the ocean in influencing salmon survival. As Ryding and Skalski [19]
68 noted almost two decades ago, “*It is becoming increasingly clear that understanding*
69 *the relationship between the marine environment and salmon survival is central to*
70 *better management of our salmonid resources*” (p. 2374).

71

72 Unfortunately, our scientific understanding of the events occurring in the marine
73 phase remains severely limited, so there has been little change in management strategy
74 beyond the essential first step of reducing harvest rates in the face of falling marine
75 survival. The recent recognition of the decline in Chinook returns across essentially all
76 of Alaska [20-22] and the Canadian portion of the Yukon River [23], where
77 anthropogenic freshwater habitat impacts are generally negligible, is another example
78 of how simple explanations looking at freshwater habitat changes are potentially
79 flawed. If freshwater habitat disruption across this vast swathe of relatively pristine
80 territory is severe enough to seriously impact salmon productivity, then there is little
81 hope that freshwater habitat in more southern regions can be “fixed” to support a newly
82 productive environment for salmon.

83

84 The same widespread problem of declining survival is also evident for other
85 diadromous species. Both eulachon [24] and lamprey [25] have undergone sharp
86 unexplained declines along the Pacific west coast of North America. In the Atlantic
87 Ocean, both Atlantic salmon [26] and eels [27, 28] are similarly in sharp decline. In the
88 case of eels, eulachon, and lamprey, the authors attribute the problem to likely marine-

89 related factors, not freshwater. This point is particularly persuasive for eulachon
90 because of the very short freshwater phase [24].

91

92 In this paper, we collate Chinook and steelhead time series for the west coast of
93 North America (excluding California) to look at patterns in survival: (1) freshwater
94 survival of smolts during the downstream migration phase and (2) smolt-to-adult return
95 rates (SARs). The SAR is the three-fold product of freshwater smolt survival during
96 downstream migration multiplied by the marine survival experienced over 2-3 years in
97 the ocean multiplied by freshwater survival during the final upstream migration by the
98 returning adults to the final enumeration point. (Depending upon the specific dataset,
99 adult abundance may be enumerated prior to actual arrival at the spawning grounds; see
100 Methods). In particular, given the very poor perceived returns of salmon to the Snake
101 River, many of our analyses compare regional survival to that of Snake River stocks.
102 We use the term SAR and marine survival interchangeably because, as we will
103 demonstrate, the majority of the SAR is determined in the ocean.

104

105 For the downstream (freshwater) smolt survival analysis, 46 Chinook and 44
106 steelhead time series were collated, comprising 531 annual estimates of survival (see
107 Methods). For the SAR comparison, 101 Chinook time series and 50 steelhead time
108 series were available (Fig. 1) which equate to 1,729 Chinook and 795 annual steelhead
109 SAR estimates. Altogether these datasets total 3,055 years of salmon monitoring—
110 clearly, an enormous effort that likely sums to multiple billions of dollars. As the
111 breakdown by regime periods will demonstrate, the tremendous increase in resources

devoted to survival monitoring as salmon returns have dwindled over time has perhaps provided less actual insight into mechanisms (as opposed to numbers) than might be hoped, a theme we return to in the Discussion.

Figure 1. Map of salmon survival time series used in the analyses. Numbers inside symbols are keyed to the populations in Supplementary Table S2; yellow circles indicate Chinook populations, pink squares indicate steelhead, and blue triangles indicate locations with data for both species. Acronyms: SEAK (SE Alaska/Northern British Columbia Transboundary Rivers); NCBC (North-Central British Columbia); WCVI (West Coast Vancouver Island); WAC (Washington Coastal); ORC (Oregon Coastal); SOG (Strait of Georgia); PS (Puget Sound).

The passive integrated transponder (PIT) tag SAR estimates for Chinook and steelhead are specific to the Columbia River Basin and are reported by the Fish Passage Center, most recently by [5]. Estimates reported in an earlier paper by Raymond [1, 29] which predate PIT tag estimates for Columbia River basin Chinook and steelhead were also included. The primary data source for the coded wire tag (CWT) based SAR time series for Chinook used in this analysis is the official survival estimates submitted by various State and Federal government agencies to the Pacific Salmon Commission under the terms of the US-Canada Salmon Treaty. These data include SAR estimates from OR, WA, BC, and AK. For Washington State steelhead outside the Columbia River basin, SARs were collected and reported by Kendall et al [7] for Puget Sound, as well as a number of locations along the coast of Juan de Fuca Strait and the outer (western) WA coast. In BC, SARs are only available for one steelhead population (Keogh River). We are unaware of additional steelhead SAR data for Alaska or coastal

130 Oregon rivers. Individual time series ranged between 2-39 years for Chinook, and 2-37
131 year for steelhead. (Datasets comprised of only a single year of data were excluded).

132

133 What are acceptable levels of salmon survival? For much of the west coast
134 outside of the Columbia River basin, formal recovery targets (SARs) have not been
135 specified, although it is clear in all regions that historical levels of productivity would
136 be greatly preferred to current return rates. (And, to foretell an underlying theme to the
137 paper, current SAR levels may in fact be much preferred to what climate change has in
138 store for salmon in the future). In the extensively dammed Columbia River basin, the
139 Northwest Power and Conservation Council's Fish and Wildlife Program (NPCC) set
140 rebuilding targets for SARs at 2%-6% ([5], p. 4), roughly the survival observed in the
141 1960s prior to the completion of the 8-dam Federal Columbia River Power System
142 (FCRPS) [29, 30]. The sharp decrease in salmon returns to the Columbia River (and
143 most particularly the Snake River) after the completion of the final Snake River dam in
144 the mid-1970s was widely assumed to be due to the construction of the dams, and great
145 effort has therefore been devoted to improving in-river smolt survival since that time.
146 For this reason, we have chosen to contrast survival in other geographic regions to that
147 of the Snake River as an objective standard of "poor" survival.

148

149 The NPCC SAR objectives did not specify the points in the life cycle where
150 Chinook smolt and adult numbers should be determined. However, one extensive
151 analysis for Snake River spring/summer Chinook was based on SARs calculated as
152 adult and jack returns to the uppermost dam encountered in the migration path [31]:

153 “Median SARs must exceed 4% to achieve complete certainty of meeting the 48-year
154 recovery standard, while ... A median of greater than 6% is needed to meet the 24-year
155 survival standard with certainty” (p. 41). With most current Columbia River basin
156 SARs on the order of ca. 1%, migratory-phase life-cycle survival would have to
157 increase 200%-600% (two- to six-fold) to meet these targets. It is unclear whether this
158 level of rebuilding is actually possible for reasons that we discuss later.

159

160 Unfortunately, as Chinook and steelhead stocks continue to dwindle, progress
161 on addressing and incorporating ocean impacts on salmon dynamics has been slow,
162 perhaps due to a combined lack of understanding about how to address marine survival
163 issues and to pessimism about whether improved understanding of the marine phase
164 can advance conservation. Therefore, lastly we review two case studies which show
165 that even when the overriding role of marine survival is identified, there is still a strong
166 predilection to preferentially identify freshwater factors to study and manipulate. This
167 has resulted in both the failure to directly address the marine survival problem and a
168 rather uncritical approach that too readily identifies widely accepted freshwater
169 stressors as being responsible for the problems evident in specific populations. In our
170 view, a large part of the difficulty lies in some of the fundamental underlying
171 assumptions that the fisheries community makes about the nature of the core problem.
172 Because these assumptions are part of our training and professional ethos, they are
173 difficult to recognize or question. Nevertheless, given the widespread geographic range
174 and magnitude of the collapse in survival that is now evident, we view it as urgent that
175 assumptions about causative agents be carefully assessed.

176

177 Results

178 Freshwater (downstream) smolt survival

179 To separate and assess what are typical freshwater survival levels for smolts
180 migrating downstream, we collated the published studies for west coast North
181 American rivers excluding California (See Methods for a more detailed summary and
182 Table S1 for reported estimates). We used these data to make regional comparisons of
183 smolt survival and survival scaled by distance travelled during the downstream
184 freshwater migration to the sea.

185 Within the Columbia River basin, survival estimates for a range of stocks and
186 river reaches are available, although the majority are for survival through the
187 hydrosystem (dammed segment of the river). For yearling Chinook, smolt survival
188 estimates varied considerably between grouping categories (Fig 2; center column, top

Fig 2. Freshwater smolt survival for west coast North American rivers. A total of N=531 annual survival estimates are included. Top row: smolt survival from release to river mouth (and intermediate locations in the case of the Columbia). Bottom row: survival per 100 km of migration distance. The red horizontal line shows the median value for all Snake River data in a given panel (red coloured bars). Data are shown as a box and whisker plot with associated sample size listed above the appropriate boxes. Abbreviations: LRE, Lower Columbia River and estuary (i.e., the river reach from just below the lowest (Bonneville) dam to the river mouth); Release to BON measures Snake River survival from hatchery release through the Snake River above Lower Granite Dam and down through the 8-dam hydrosystem to the last dam (Bonneville). Full river measures survival from release to the mouth of the Columbia River. Data sources and annual survival estimates are reported in Supplementary Table S1.

189 row); however, when survival is scaled by distance travelled (bottom row), two patterns
190 become apparent. First, regardless of release location or origin (Snake, Upper, or Mid-
191 Columbia), all yearling Chinook from the Columbia River basin have remarkably
192 similar median survival rates of 88% per 100 km of migration distance. Second,
193 survival rates in the dammed and undammed sections of the river (the hydrosystem and
194 LRE) are largely similar.

195 For other populations outside of the Columbia River basin which have
196 published estimates (n=3), survival rates per 100 km varied. Survival rate of the
197 Nimpkish River (B.C.) population was particularly low: estimated survival to the river
198 mouth was 60% but the migration distance was only 8.5 km, resulting in only 0.25%
199 survival per 100 km. Coldwater River (Fraser River/SOG) yearling Chinook survival
200 rate was 63% and 68% per 100 km. Survival of hatchery-reared Chilko River Chinook
201 (a Fraser River/SOG population) was the only population similar to the Columbia River
202 basin; survival was 49% during their 640 km downstream migration in the Fraser River
203 basin, resulting in a survival rate of 89% per 100 km.

204
205 A similar result is evident for Snake River steelhead which had nearly identical
206 median survival rates per 100 km of migration distance (87%) as yearling Chinook
207 irrespective of the section of the Columbia River basin that survival was measured
208 over. Upper Columbia River steelhead tagged and released at Bonneville Dam in the
209 lower Columbia River had survival rates of 70-75% per 100 km in the lower river and
210 estuary, however, steelhead tagged and released at Rock Island Dam (UCOL) had
211 consistently lower median survival rates, only ~41% per 100 km.

212

213 Survival rates per 100 km in the other regions for which we have steelhead data
214 (Keogh River, Strait of Georgia, Puget Sound, and Oregon Coast) were generally lower
215 than either the upper Columbia or Snake River. Keogh River steelhead had particularly
216 low rates; the release site was located only 300 from the river mouth and survival
217 ranged between 72-95%, resulting in an estimated survival rate per 100 km close to
218 zero. Puget Sound and Oregon Coast populations had relatively short migrations to the
219 ocean (0.3-102 km) and highly variable survival rates; these results suggest intense
220 losses concentrated in the lowest reaches of these rivers. The only exception was
221 hatchery-reared Skagit River steelhead which had a survival rate of 90% per 100 km.

222

223 There are no subyearling Chinook survival data available outside of the
224 Columbia River basin, but within the basin, subyearling Chinook had similar median
225 survival rates to yearling Chinook and steelhead in the hydrosystem and in the LRE
226 (~85% per 100 km).

227

228

229 Chinook SARs

230 Coast-wide trends in adult survival (SARs)

231 Adult survival data for Chinook salmon are available for a varying range of years.
232 The most extensive data sets are for the upper Columbia (both subyearling and yearling
233 Chinook) and Snake rivers (yearlings), which extend back to the 1960s (Table S1).

234 Data were available for other regions beginning in the 1970s and for all regions by
235 2001 for yearlings, and 1987 for subyearlings.

236 In essentially all regions where time series extend back to the 1970s or earlier,
237 survival to adult return has substantially decreased with time (Fig 3). A large drop in
238 SARs for yearling Snake River Chinook is evident from the 1960s to approximately the
239 mid-1970s, the time period when Snake River dams were completed [2,28]. Although
240 the timing varies with region, the collapse in survival is also evident in other regions
241 with long time series for both yearling (Upper Columbia River and—notably—Alaskan
242 yearling stocks from SE Alaska), and subyearling Chinook (west coast Vancouver
243 Island, the Strait of Georgia, and Puget Sound). Raymond [1, 29] (and many
244 subsequent authors) ascribed the cause of the drop in survival to dam construction;
245 however, declining SARs are also evident in other regions not affected by the
246 construction of the FCRPS.

Fig 3. Time series of smolt to adult survival (SAR) data for west coast Chinook stocks (excluding California). Left column: subyearlings; Right column: yearlings. Regions are oriented from north (top) to south. Gold dots are SAR measurements based on CWT tags (PSC database), brown dots are SARs reported by Raymond [1], and violet dots are SARs based on PIT tags [5]. A loess curve of survival and associated 95% confidence interval (shaded region) using all available data for each panel is shown as a black line (the smoothing parameter was set to $\alpha=0.75$); the loess curves for Snake River subyearling and yearling survival are overplotted in red to facilitate comparison with other regions. Blank panels indicate regions where the life history type does not occur (for example, Fall (subyearling) Chinook do not occur in Alaska, while Spring (yearling) Chinook do not occur in the low elevation streams on the west coast of Vancouver Island or Oregon coast). The major regime shift years of 1977, 1989, and 1998 are indicated by vertical lines. In this and subsequent figures the pale red band delineates the official Columbia River SAR rebuilding targets of 2-6%.

247 From the time of the major ocean regime shift in 1977 forwards, no substantial
248 recovery in SARs is evident in any region. As more monitoring programs were
249 initiated in the 1980s, SARs for all these regions were either declining or essentially
250 fluctuating around a low mean value closely approximating the Snake River SARs (red
251 lines) in all regions apart from the Oregon Coast; here, SARs were also roughly flat
252 over time but at a persistently higher mean level relative to the Snake.

253

254 Regional survival differences

255 When compared by region (Fig 4), median Snake River yearling (Spring) Chinook
256 SAR (1%) is higher than the regional median SARs for Puget Sound (0.55%) and the
257 Strait of Georgia (0.53%), and is virtually identical to median survival for the Upper
258 (0.96%) and Lower (1.08%) Columbia River populations. Regional yearling SARs are
259 higher than the Snake River values only for three geographic areas: the mid-Columbia
260 River region (1.49%), Northern & Central BC (2.31%), and Alaska (1.88%). Within a

261 few of these geographic regions, striking population-specific differences are also
262 evident which we consider later.

Fig 4. Box and whisker plot of SARs by population (all available years). The black horizontal line within each bar is the median of the SAR data available for each population. Median survival across all available data for each region is shown as a blue line; median Snake River survival for all populations combined is shown as a red line and overplotted on all panels for comparison. The number of years of data is shown to the right. To save space, abbreviated population names are used here along with the map code from Figure 1; full names for the populations are listed in Supplementary Table S2.

263
264 For subyearlings (Fall Chinook), Snake River median SARs (0.81%) are similar to
265 or higher than median survival in other regions of the west coast apart from coastal
266 Oregon (ORC: 2.07%) and the west coast of Vancouver Island (WCVI: 1.34%; Fig 4).
267 As the time series plot (Fig 3) makes clear, the higher median survival evident for
268 WCVI (Robertson Creek) Chinook relative to the Snake River may not actually be due
269 to persistently better SARs, but rather to the longer time series of data for Robertson
270 Creek that extends back to the period of particularly high SARs in the 1970s. Data for
271 this time period are lacking for Snake River subyearling Chinook; we consider this
272 issue further below.

273
274 In addition to the high median SARs for Oregon Coast and WCVI Fall Chinook,
275 two specific subyearling hatchery populations from farther north (University of
276 Washington Accelerated Fall Chinook in Puget Sound (3.96%), and Chilliwack Fall
277 Chinook from the Strait of Georgia (lower Fraser River; 4.56%)) are also of note
278 because of the strikingly large survival difference (~4X) of these stocks relative to the

majority of populations within each region. The higher median SAR for yearling Chinook from the Mid-Columbia region (1.49%) is similarly due to two wild populations (Yakima: 2.21% and John Day: 4.12%) and one hatchery population (Cle Elum: 1.57%) having higher SARs while two other hatchery populations have lower SARs (Carson 0.62% and Warm Springs 0.66%) than both Snake River and Lower Columbia River median SARs (SNAK=1; LCOL=1.08%).

Strikingly, although there are some exceptional populations, no region outside the Columbia River now achieves the Columbia River basin's official SAR recovery targets of 2%-6%. The Alaskan stocks attained these target survival levels in the early 1980s, but since that time Alaskan SARs have fallen below the Columbia River basin rebuilding targets, and in the most recent years have essentially reached the current survival rates of Columbia basin stocks (Fig 2).

3. Relative survival (scaled by Snake River)

The regional-scale aggregation of SAR data provides a useful overview of survival between regions. However, important population-specific differences are potentially obscured because small numerical differences may in fact reflect large differential impacts on survival when SARs are low. For example, when regional SARs are only 1%, a population-specific SAR of 0.5% actually represents a population whose survival rate is only half that of the other populations. In addition, regional

301 comparisons may be distorted because of trends in survival over time, and differing
302 lengths to the various time series.

303 The potential influence of these factors can be reduced by normalizing the SAR
304 estimates. In Fig 5, we divided each annual SAR estimate by the median of all Snake
305 River SAR data available for the same year. This approach removes the potential
306 confounding caused by temporal trends in SAR when time series with different lengths
307 are compared. When SAR data for all available years are normalized in this way, SARs
308 for Snake River yearling Chinook are higher than Puget Sound and Strait of Georgia
309 and virtually indistinguishable from those for the Lower Columbia River (Willamette
310 R) and the Upper Columbia River. Only normalized SARs for mid-Columbia, North &

Fig 5. Normalized Chinook SARs. Values are calculated by dividing individual SAR estimates for each stock and each year by the median Snake River SAR for the same year and aggregating by region. Vertical lines show the median SAR for the Snake River (red) and other regions (blue). Note the logarithmic scale on the x-axis. As in the prior plots, Columbia & Snake River SAR estimates based on PIT tags do not incorporate above-dam survival (or harvest).

311 Central BC, and SE Alaskan populations of Spring Chinook are higher than the Snake
312 River populations.

313 The situation is similar for subyearling Chinook when normalized SARs are
314 compared: Snake River subyearling SARs are either lower (Upper Columbia; Strait of
315 Georgia, Puget Sound), higher (Mid Columbia; Lower Columbia), or closely equivalent
316 (Washington Coast, North-Central BC) to SARs observed for all other regions with
317 data. The only pronounced differences are the nearly 5-fold higher survival of the two
318 Oregon coast stocks and the roughly 2-fold higher SAR for the Robertson Creek
319 population (west coast Vancouver Island).

4. Survival by regime period

Significant changes in ocean productivity are known to impact salmon populations on time scales ranging from decades [16, 18, 32-34] to centuries [35-37]. An alternative approach to comparing survival normalized by year is to break the survival data into recognized ocean regime periods [16-18, 32, 33, 38, 39] and then compare the normalized SARs. We defined four periods based on the year of ocean entry by smolts: 1977 and earlier, 1978-89, 1990-98, and 1999 or later. The results (Fig 6) essentially mirror prior analyses, with the ratio of median Alaskan yearling Chinook survival relative to the Snake River falling from ~19X the Snake River value in the pre-1977 period to ~3X the Snake River value in the next two regime periods and then down to ~2X the Snake River value after the 1990 regime shift. Only the Alaskan, north-central BC, and Mid-Columbia populations remain ~2X higher than the Snake River populations' SARs post-1998, but well below their earlier levels of productivity. (In fact the time series of Alaskan and north-central BC SAR data (Fig 3) show that in the most recent years SARs have fallen to Snake River SAR levels). Upper and Lower Columbia, Puget Sound, and Strait of Georgia populations all have similar or lower survival. An analogous pattern is evident for subyearling Chinook, except here it is only the Oregon Coastal populations that have persistently higher survival. The progressive collapse in survival across regimes is notable for each region.

Fig 6. Comparison of normalized Chinook SARs by regime periods: pre-1977, 1978-1989, 1990-1998, and post 1998. Boxes and whiskers have the conventional interpretation; the horizontal red line shows the Snake R median SAR value for each regime to facilitate comparison (1.0 by definition). Sample sizes are shown above each group (green font) and the ratio of median SARs relative to the Snake River is shown immediately above the upper whiskers (black font).

339

340 Steelhead SARs

341 6. Coast-wide survival

342 Data on steelhead survival (SAR) are more geographically limited than for
343 Chinook (Fig. 1 & Table S2), but share many of the same features (Fig 7). For
344 simplicity, we have included the Keogh R time series from the extreme NE tip of
345 Vancouver Island in the Strait of Georgia/Juan de Fuca Strait (SOG) region, although
346 the population enters Queen Charlotte Strait, not the Strait of Georgia proper.

347 Prior to the 1977 regime shift, data are only available for the Upper Columbia
348 and Snake Rivers (Fig 7). Similar to Snake River yearling Chinook, steelhead SARs in
349 both the Upper Columbia and Snake Rivers declined in the period prior to the mid-
350 1970s (when both FCRPS dam construction was completed and a major marine regime
351 shift occurred). SAR data becomes available for Washington Coast, Puget Sound, and
352 Strait of Georgia regions in the period after the 1977 regime shift. The very high SARs
353 of Keogh R steelhead (northern SOG region) in the early years of the historical record,
354 which exceeded 20% in some years, compresses the SAR differences with other regions
355 (indicated by the LOESS curves), making the differences somewhat difficult to see.

356 However, plotting the data in this way demonstrate that under former climatic
357 conditions, very high SARs were achieved in some regions.

358 Following ocean entry year 1990, further decline is evident in Washington
359 Coast and Strait of Georgia steelhead SARs around ocean entry year 1990 (the time of
360 the subsequent ocean regime shift) as well as a continuing decline in Puget Sound
361 SARs to levels below that of Snake River steelhead. Although SAR data are not
362 available for B.C. steelhead stocks other than the Keogh River (northern Vancouver
363 Island), the pattern of adult returns to other southern B.C. rivers closely matches returns
364 to Keogh River, supporting the view that the Keogh SAR pattern applies more broadly;
365 see [40]. Both Washington outer coast (WAC) and mid-Columbia SARs are
366 substantially higher than those the Snake River (as is Keogh), while Puget Sound SARs
367 drop to lower values after 1990. Upper Columbia River steelhead SARs are closely
368 similar to Snake River values.

Fig 7. Steelhead SARS plotted against ocean entry year. Regions are oriented from north (left) to south (right); the Keogh R (KEOG) is situated on the NE tip of Vancouver Island (BC). Gold dots are SAR measurements based on PIT tags, brown dots are SARs reported by Raymond [1], and violet dots are SARs based on CWT tags. A loess curve of survival and associated 95% confidence interval (shaded region) using all available data for each panel is shown as a black line (the smoothing parameter was set to $\alpha=0.75$); the Snake River loess curve is shown in red and over plotted on all other panels to facilitate comparison. Steelhead survival data are for British Columbia only available for the Keogh River (see Ward et al 2006) for a description of the monitoring program). The major regime shift years of 1977, 1989, and 1998 are indicated by vertical lines.

369

7. Regional survival differences

A few specific steelhead populations are notable for having anomalously high survival (all three mid-Columbia River and three of eight Washington Coast populations; Fig 8). Median SARs for the Snake River region (1.7%) are comparable to the Upper Columbia (1.9%) and the Washington Coast regions (2.3%), but more than double that of Puget Sound steelhead populations (0.8%). Only the median SARs for the mid-Columbia River region (5.5%) and the Strait of Georgia region (Keogh and Snow Creek; 3.3%) are appreciably higher than Snake River survival.

Fig 8. Box and whisker plot of steelhead SARs by population (all available years). Population names are listed in Supplementary Table S1. The black horizontal line within each bar is the median of the SAR data available for that population. Median survival across all available data for each geographic region is shown as a blue line; median Snake River survival for all populations combined is shown as a red line and overplotted on all panels for comparison. The number of years of data is shown to the right.

8. Relative survival (scaled by Snake River)

When annual SAR estimates for individual steelhead stocks are normalized by the Snake River median SAR values in each year, a similar relationship emerges (Fig 9). Median steelhead SARs are either indistinguishable from the Snake River (Upper Columbia River), slightly higher (Washington Coast), or substantially lower (Puget Sound). Only the Mid-Columbia River and Strait of Georgia have substantively higher SARs than the Snake River when compared on a year-for-year basis.

Fig 9. Normalized steelhead SARs, obtained by dividing each individual SAR estimate (i.e., for each stock and each year) by the median SAR calculated across all available Snake River SARs for that year. The median Snake River SAR is overplotted in red. Note the logarithmic scale on the x-axis.

387

388 9. Survival by regime period

389 This pattern becomes particularly clear when normalized steelhead SARs are
390 examined by regime periods (Fig 10). The large drop in Strait of Georgia SARs in the
391 post-1998 regime period is particularly notable, (in absolute terms, the drop in survival
392 corresponds to a change in median SARs from 8.4% in the 1978-88 period to 2.6% in
393 the post-1998 period—a three-fold decline). The second aspect to the steelhead data is
394 the similarity of the other regions. Excluding the Mid-Columbia River, where only data
395 for the post-1998 period are available, most other regions have median SARs roughly
396 similar to the Snake River across all regime periods; only the mid-Columbia and SOG
397 stand out as having consistently higher median SARs, while Puget Sound drops from
398 higher median SARs than the Snake River to substantially lower SARs (less than half)
399 in the post-1998 period.

Fig 10. Comparison of normalized steelhead SARs by regime periods: pre-1977, 1978-1989, 1990-1998, and post 1998. Boxes and whiskers have the conventional interpretation; the horizontal red line shows the Snake R median SAR value for each regime to facilitate comparison (1.0 by definition). Sample sizes are shown above each group (green font) and the ratio of median SARs relative to the Snake River is shown immediately above the upper whiskers (black font).

400

Discussion

Our analysis shows that over time Chinook and steelhead SARs have declined to reach approximately the same low level for almost all measured populations across the entire west coast of North America—with a few important exceptions that we discuss later. Although we do not have direct measurements of survival for Chinook stocks located west of SE Alaska or steelhead for regions north of Vancouver Island, the decrease in the number of adult Chinook returning to the rest of Alaska [20, 21] shows the broad region over which the conservation crisis now extends. We first address juvenile survival during seaward migration as a possible cause of the decline in adult abundance and then demonstrate the importance of the marine habitat.

The freshwater contribution to SARs

Freshwater survival of smolts during downstream migration to the sea has been assessed for a number of river systems only over the past 15 years following the advent of miniaturized acoustic transmitters and the expansion of the PIT tag system within the Columbia River basin. The published studies collated in Table S1 report varying freshwater survival levels lying mostly within the 25-75% range for yearling Chinook. When scaled by migration distance, median survival rates of Columbia River basin yearling Chinook populations are either similar to or better than available populations from outside of the basin per 100 km of migration distance. Snake River steelhead have median survival and median survival rates very similar to Snake River yearling Chinook, and survival rates per 100 km are much better than those of all steelhead populations located outside the Snake River (Fig 2).

423

424 Within the Columbia River basin survival scaled by distance travelled is nearly
425 constant for yearling Chinook irrespective of the source population and migration
426 segment examined. For steelhead, downstream survival rates are lower in the Upper
427 Columbia than the Snake River, but are still higher than values reported for outside of
428 the Columbia River basin.

429 Both observations are at odds with conventional wisdom. Given the enormous
430 focus over the past half century on improving smolt survival within the Columbia River
431 hydrosystem, our interpretation is that these efforts were successful because survival
432 rates are now higher than in undammed river systems. This result extends our earlier
433 finding that Columbia River smolt survival was slightly higher than the adjacent
434 undammed Fraser River [41], particularly for steelhead where estimates are now
435 available for a substantial number of river systems. Thus, significant further
436 improvement is unlikely because the Snake River now boasts the highest measured
437 freshwater survival rates in the Pacific Northwest.

438

439 If survival rates were in fact low in the Columbia River basin, improvements in
440 freshwater survival could potentially increase the SAR. For example, Chinook smolt
441 survival in California ranged from 3-16% for a 516 km migration in the Sacramento
442 River [42] to an astonishingly low 0-2% through the lower 92 km of the San Joaquin
443 River Delta [43]. Such low survival provides substantial scope for potential
444 improvement. This difference is important because the large drop in coast-wide SARs

445 excluding California to around 1% and relatively high freshwater survival isolate the
446 main conservation problem as being in the ocean.

447 Our results also indicate that the river mouth is a perilous location for smolts,
448 something also noted in California [42], because survival rates scaled by distance are
449 extremely low in rivers where post-release distance to the mouth is short, e.g., Keogh
450 River and Big Beef Creek steelhead. Losses (presumably to predators) must be
451 concentrated near the river mouth to result in this pattern, and continued losses from
452 predation may well occur after ocean entry because smolts are still concentrated and the
453 migration timing is predictable, conditions which cause predator aggregation in other
454 situations [44, 45].

455 It is important to outline why past declines in freshwater survival cannot have
456 been the driver of the observed drop in SARs—put simply, currently measured
457 freshwater survival levels are too high. The longer SAR time series indicate at least a
458 4-5 fold decline over time. However, for freshwater survival to be the cause of this
459 decline in SARs, current values of freshwater survival cannot be more than 20-25%.
460 That they are substantially higher for many populations (Fig 2) means that it is
461 mathematically impossible for freshwater survival to have fallen far enough to explain
462 the decline in SARs. For example, even if downstream survival through the dams was
463 originally 80% prior to the 1970s and then fell to 40% this would “only” produce a
464 two-fold decline in SARs, e.g., from 6% to 3%, so the scope for primarily freshwater
465 regulation of SARs is limited.

466

467 The importance of marine habitat

468 Occam's Razor dictates that any coherent theory to explain the large and
469 geographically widespread drop in survival to similar low levels should be applicable to
470 all populations. We are unable to identify a consistent mechanism of action because of
471 the current limits to our understanding of the ocean phase, but some explanations
472 (various forms of anthropogenic freshwater habitat disruption) are clearly less likely as
473 explanations of poor salmon survival than others (climate-related changes in the ocean).

474
475 Salmon, as well as other anadromous fish such as lamprey and eulachon,
476 migrate widely across a complex landscape composed of many successive freshwater
477 and marine habitats; even something as simple as the number of distinct habitats each
478 salmon population occupies over the duration of the marine phase is unknown. The
479 number of returning adults is therefore successively affected by changes in survival in a
480 complex sequence of freshwater and marine habitats, most of which are poorly
481 understood, as the product $SAR = S_1 \cdot S_2 \cdot S_3 \cdot \dots \cdot S_n$. If survival drops to 1/10th of its
482 original value in any one of these habitats, the SAR will also decline equivalently
483 unless density-dependent factors occurring at some later point in the life history buffer
484 the impact on adult returns.

485
486 Despite this, conventional conservation thinking for Pacific salmon primarily
487 focuses on freshwater habitat issues. The rationale for this focus can be traced back to
488 two separate events first occurring in the 1970s. The first was the passage of the U.S.
489 Endangered Species Act (ESA) in 1973, with its strong focus on protecting and

490 preserving habitat as the paramount priority for conservation [46]. Canada’s Species at
 491 Risk Act was enacted in 2003, and was partially modeled on the US ESA. The
 492 Canadian legislation provided a remarkably broad definition of habitat, which
 493 essentially prohibited: “damaging or destroying the residence of one or more
 494 individuals”, with residence defined as “...a dwelling-place such as a den, nest or other
 495 similar area or place, that is occupied or habitually occupied by one or more individuals
 496 during all or part of their life cycles” ([47]; p. 227). Unfortunately, “habitat” in both
 497 countries is ill-defined for migratory animals such as salmon which occupy many
 498 different habitats as they complete their life cycle. The larger question, not discussed in
 499 either country’s legislation, is this: to what degree can (or should) habitat related
 500 declines in some part of the ocean phase be compensated for by remedial action in
 501 some other part of the life history? That is, excluding the direct impacts to habitat
 502 which are obvious candidates for correction, can (and should) ocean impacts be
 503 remediated by intervening in other points in the life history?

504

505 The second event, unappreciated at the time, was a major shift in ocean climate
 506 in 1977 which had impacts on a wide range of marine fish stocks as well as salmon
 507 across the entire west coast of North America [38, 39]. The timing of this regime shift
 508 also nearly coincided with the completion of the final Snake River dam forming the
 509 Federal Columbia River Power System (FCRPS) in 1975. Not surprisingly given the
 510 understanding of salmon dynamics in that era, the ensuing decline in adult salmon
 511 returns a few years later was ascribed purely to poor smolt survival through the dams;

512 however, as we have demonstrated, a similar drop in survival is seen in many other
513 regions after 1977 and in purely marine species as well.

514

515 The decline in marine survival began earliest in the south and then progressively
516 expanded farther north along the coast with time (Figs 3 & 7). Almost none of the
517 rivers outside the Columbia have dams, so the argument that the poor performance of
518 Snake River stocks is primarily due to the completion of the FCRPS is inconsistent
519 with the broader data. (We are not dismissing the argument that extensive past
520 modifications to the FCRPS have improved freshwater survival. Rather, we are making
521 the point that further improvements in freshwater survival will have small or negligible
522 impact on increasing adult returns and that the very large ocean impacts may in fact
523 distort our understanding of how adult returns are related to freshwater modifications).
524 As we will discuss, many other “single factor” reasons for poor salmon survival along
525 the west coast also suffer from the same logical flaw that survival now seems to be poor
526 everywhere.

527 **Overfishing alone can’t explain the decline**

528 Wasser et al [48] cite this blanket statement: “Anadromous salmonids
529 (*Oncorhynchus* sp.), which hatch in fresh water, migrate to the ocean, and then return to
530 their natal waterways to breed, are threatened primarily by habitat loss from dams and
531 overfishing (SOS 2011)” (Lines 98-101 of the SI). The sentiment underpinning this
532 statement is widespread and reflects a fundamental problem with simply making a
533 casual association between the assumed cause (freshwater habitat loss) and the effect
534 (declining salmon stocks). We view the reality as considerably more nuanced: Fall

(ocean-type) Chinook harvest levels of 50%-70% that were formerly sustainable are no longer sustainable because marine survival dropped 4-5 fold over the past few decades. The drop in marine survival is too large (75-80%) to be compensated by even the complete cessation of harvest. The magnitude of the gap is widely unappreciated, and the relatively small percentage difference between the harvest rate (50-70%) and SARs (75-80%) is misleading.

To fully compensate and maintain adult escapements, the initially sustainable harvests of the 1970s would have to be as large as the drop in marine survival has been. Algebraically,

$$E_1 = N \cdot S_1 \cdot (1 - h_1)$$

and

$$E_2 = N \cdot S_2 \cdot (1 - h_2),$$

Here E_t is the escapement at time t , N is the number of smolts beginning migration to the sea, S_t is the SAR, and h_t is the harvest fraction, where $t=1,2$ is the start and end of the time series.

For escapement, E_t , to remain constant in the two time periods implies that

$$\frac{S_2}{S_1} = \frac{(1 - h_1)}{(1 - h_2)}$$

558 or

559

$$h_1 = 1 - \frac{S_2}{S_1}(1 - h_2)$$

560

561 The maximum compensation management can make for declining marine
562 survival occurs when all fisheries are curtailed completely ($h_2=0$). In this case, ceasing
563 or reducing harvest can only fully compensate if the initial rate of sustainable harvest is

564 $h_1 \geq 1 - \frac{S_2}{S_1}$. The key feature of this equation is that it is the ratio of the current to the
565 initial period marine survival that determines how large the initial sustainable harvest
566 rate must have been to allow full compensation by harvest rate reduction. If marine
567 survival drops by almost an order of magnitude, as it has in at least some regions,
568 sustainability can only be maintained if the initial sustainable harvest rate was at least
569 90%.

570

571 Taking the Columbia River basin as a less extreme example, marine survival
572 has dropped from ~6% to 1%, so the initial harvest rate would have to be $h_1 \geq 83\%$ to
573 allow full compensation for changing environmental conditions. Historical harvest
574 rates reported by the PSC [49] suggest that Chinook harvest rates were on the order of
575 50%-60% for many subyearling stocks, implying that complete harvest rate
576 compensation for declining marine survival would only be possible for survival ratios
577 of $S_2/S_1 \sim 0.4-0.5$ (or $\sim 1/2 - 1/2.5$); far less decrease in survival than has actually
578 occurred.

579

580 The same major decline in survival can be seen in British Columbia after the
581 1977 regime shift, the period when the first real measurements of SARs for other west
582 coast regions started. Perhaps the best measurements documenting the magnitude of
583 the drop in British Columbia SARs was reported by Bilton et al [50]. In the early
584 1970s, SARs for Strait of Georgia coho of $\bar{S} = 20.8\%$ (SE: $\pm 0.5\%$) and $S_{\text{median}} = 17.2\%$
585 were obtained in extensive experimental hatchery releases (six replicates of each of
586 three size classes of smolts in each of three months (April, May, and June)). The
587 magnitude of these survival levels (ca. one in five smolts surviving to return as adults)
588 justified Canada's decision to fund the Salmon Enhancement Program (SEP), a major
589 investment in hatcheries. Yet less than two decades after the start of SEP in 1977,
590 average coho SARs for the nearby Big Qualicum hatchery had dropped from 28.6%
591 (1973-77 ocean entry years) to 5.6% (1990-99) and then to 1.5% (2000-2012) (data
592 from [8, 51]). As a result, average survival rates dropped from 1 in 3.5 smolts in the
593 1970s to 1 in 67 smolts—a decrease to 1/20th of the initial value. (See [8] for a
594 detailed description of the decline over time in Strait of Georgia coho SARs).

595

596 To place the magnitude of this change in perspective, by the 2000s coho SARs
597 in the Strait of Georgia were the equivalent to surviving through a sequence of $n =$
598 $\log(S_{2000s})/\log(S_{1970s}) = 3.4$ successive survival periods, with each time period
599 equivalent to the entire survival process experienced by cohorts in 1973-77 (a time
600 when intensive sport and commercial fisheries were operating, unlike recent years).
601 Whatever the change in the environment was, it was the equivalent to the coho now
602 remaining at sea for 60 months (five yr) instead of 1.5 yr while experiencing the overall

603 mortality rates characteristic of the 1970s. As coho harvest rates are near zero in recent
604 years, it is essentially all natural mortality processes that are currently operative.

605

606 Statements about the major role of particular factors in driving salmon declines
607 (dams in the Columbia River or salmon farming in British Columbia, which developed
608 in the 1990s) must therefore be assessed critically because salmon from other regions
609 lacking these specific factors also return from the ocean with very poor marine survival.
610 Thus, dams or salmon aquaculture may contribute as habitat issues to overall losses, but
611 the essential policy debate is (1) whether modifying their operation will materially
612 contribute to improving salmon returns, and (2) whether proposed courses of action are
613 actually credible and cost-effective given the primary influence of ocean conditions.

614

615 The role of dams

616 A wide range of west coast rivers lacking dams have similar or worse reported
617 survival than the Snake River, both in terms of downstream smolt survival and adult
618 return rates. We interpret this as evidence for a fundamental flaw in our biological
619 understanding of the conservation factors controlling salmon productivity.

620

621 Direct Mortality

622 Conventional thinking holds that if average marine survival was 4-6% in
623 regions without dams, then the four- to six-fold lower survival of Columbia River
624 Chinook populations (currently ca. 1%) would be clear evidence that the Columbia
625 River dams were the cause of poor survival. The conclusion would then be that

626 removing or modifying dams lying in the migration path of Snake River basin
627 populations should increase SARs four- to six-fold, thereby achieving rebuilding
628 targets. Yet the same conclusion, which has implicitly guided much conservation
629 thinking, clearly cannot be used in reverse—presumably no one would argue that
630 constructing eight dams in the Fraser River would double salmon returns, raising
631 median Chinook survival in the years since 2000 from a mere 0.53% in the Fraser River
632 to the Snake River’s current 1%. (Median SAR for all other Strait of Georgia yearling
633 Chinook populations is also 0.53%; none have dams in the migration path).

634

635 A similar conclusion is evident when the level of survival through the FCRPS is
636 assessed. Spring Chinook smolt survival through the 8-dam FCRPS ranges from 50-
637 60% (Tables A.1 and A.2 of [5]), so even eliminating all sources of freshwater
638 mortality during hydrosystem migration—direct impacts of the dams on survival,
639 predation, and possible losses from disease—could only increase SARs by a factor of
640 0.5^{-1} - 0.6^{-1} , or 1.7-2%. These levels are well below official rebuilding targets. Further,
641 because a significant fraction of the downstream loss is due to predation by birds [52]
642 and fish [53], unless all predatory wildlife species are eliminated even an increase to
643 1.7-2% SARs is unrealistic.

644

645 Indirect (Delayed) Mortality

646 The mathematical inability of even perfect hydrosystem survival to achieve
647 minimum rebuilding targets likely underlies the theory that delayed mortality caused by
648 multiple dam passage contributes to poor ocean survival [5, 54-64]. Three of five

649 Spring Chinook populations (Fig. 4) and all three steelhead populations (Fig. 8) from
 650 the mid-Columbia region not migrating through the Snake River dams have
 651 substantially higher SARs than Snake River populations, supporting this view;
 652 however, when a broader range of populations is considered the delayed mortality
 653 theory is not supported.. For example, most mid-Columbia stocks of subyearling
 654 Chinook and two of five mid-Columbia yearling Chinook have similar or lower SARs
 655 relative to Snake River populations (Fig. 4). A similar pattern of anomalously high
 656 SARs is also seen for two Washington Coast steelhead populations and one (each)
 657 Strait of Georgia and Puget Sound Fall Chinook populations despite the majority of
 658 nearby populations having SARs consistent with the Snake River median (Figs. 4 & 8).
 659 Thus it is unlikely that greater dam passage causes delayed mortality in the estuary or
 660 ocean both because something unrelated to dam passage also causes a few populations
 661 to have substantially higher survival by the time the adults return from the sea in river
 662 systems lacking dams and because many populations lacking dams in the migration
 663 path now have similarly low levels of survival.

664
 665 **Misplaced efforts: Case studies where the marine**
 666 **environment was implicated, but fresh water research was**
 667 **initiated**

668 The data analyzed in this paper demonstrate both a long term coast-wide decline
 669 in survival for Chinook and steelhead and that the cause of the low SARs must
 670 predominantly be located during the marine phase of the life history. Although

671 managers have moved to reduce Chinook harvest to partially compensate for the drop,
672 relatively little has been done to determine the cause of the decrease in marine survival
673 because much of the focus remains on remediating freshwater habitat.

674

675 Festinger [65] first defined the term “*cognitive dissonance*”. In brief, it can be
676 described as the inability to recognize the true problem, despite the evidence. More
677 formally, in psychology the term has come to mean the process by which an individual
678 manages inconsistent thoughts, beliefs, or attitudes, especially as relating to behavioral
679 decisions and attitude change, by modifying aspects of their cognitive process to
680 achieve internal consistency; for example, discounting or diminishing data inconsistent
681 with the individual’s pre-existing beliefs.

682

683

684 The history of west coast salmon management suggests that cognitive
685 dissonance concerning the marine survival problem is widespread and the reason
686 declining salmon stocks are redressed by addressing primarily freshwater habitat issues.
687 (Interested readers should also consult Janis [66] (especially Chapter 8) for an excellent
688 summary of the sociological factors leading to “*groupthink*” and the poor decision
689 making processes that result). We now review three case studies to illustrate how
690 cognitive dissonance seems to be at play in determining past operational responses to
691 falling marine survival: (i) Rivers-Smith Inlet sockeye (Central B.C.); (ii) Columbia
692 River Chinook and steelhead; and (iii) Upper Fraser steelhead.

693 *Rivers and Smith Inlet Sockeye (B.C.)*

694 The Rivers-Smith Inlet sockeye complex formed the second largest sockeye
695 fishery in British Columbia for much of the last century (the Fraser River being the
696 largest). Adult harvest levels averaged around 1M sockeye for six decades (1910-
697 1970), and escapement (measured from the late 1940s forward) was stable at ca.
698 400,000 adults [67]. The Rivers and Smith Inlet populations are located in adjacent
699 watersheds in the remote central coast of BC where there is little anthropogenic impact.

700 Following 1970, the productivity of both the Rivers and Smith Inlet sockeye
701 populations suddenly collapsed [67-72]. Because escapement remained stable until the
702 1970s [67], recruitment overfishing did not occur during this period. Probably because
703 of the isolated location and the lack of any other nearby significant salmon fisheries,
704 prompt management decisions to reduce harvest to near zero were promptly taken and
705 were maintained. However, despite harvest being curtailed, the stocks did not recover
706 as standard fisheries theory would predict, although escapements remained stable.
707 Following the next ocean regime shift in 1989, escapement levels fell to record lows,
708 from >1 million spawning adults to ca. 9,500 adults by 1999—a collapse to 1/100th of
709 the original stock size in just over two decades. Because the fishery had already been
710 curtailed, no further management action was possible to compensate for the second
711 drop in survival. There was also evidence that additional nearby sockeye stocks were
712 impacted similarly [72]. Thus, the stock collapsed despite prompt and full action by
713 management.

714 A study of the management response [67] to the collapse detailed the reasons
715 for rejecting a freshwater cause (including using data extending back over half a

716 century to demonstrate that pre-smolt abundance in the lake was above the long-term
717 mean). The authors noted that “*Poor marine survival is the most parsimonious*
718 *explanation for the declining fry-to-adult survival in Owikeno Lake, particularly in*
719 *light of coincident declines in sockeye salmon returns per spawner at Long Lake (a*
720 *nearby pristine watershed) and declines in adult sockeye salmon abundance in other*
721 *populations to the north of Rivers Inlet.*”

722

723 The key findings from a joint federal and provincial government technical
724 committee reviewing the collapse are worth quoting verbatim [68, 70]:

725 “(1) *The drastic declines in abundance appear to be due to an extended period*
726 *of poor marine survival that cannot be explained by any one event, such as sea-entry*
727 *during an unusual El Niño year. At least two recent years (1996 and 1997) show signs*
728 *of near-zero marine survival, but the reasons for those low survival rates are not known*
729 *at this time.*

730 (2) *There is little evidence to suggest that logging or other human activity in*
731 *either of the drainage basins has had more than small and localized impacts on sockeye*
732 *spawning and rearing. The simultaneous declines in both basins – i.e., in Owikeno,*
733 *where there has been extensive logging and in Long Lake, where there has been very*
734 *little – is convincing evidence that the cause of the declines does not lie in freshwater*
735 *habitat disturbance”.*

736

737 The Rivers-Smith Inlet study is to our knowledge unique in North America.

738 Not only do the twin conclusions state that the problem lies in the ocean, they also state

739 that freshwater habitat problems were not contributive—something that is generally not
740 possible to rule out with certainty for most salmon populations.

741

742 The joint technical committee then recommended necessary research to clarify
743 the cause of the collapse, and regulatory action that might be taken to improve the
744 situation. Strikingly, despite the conclusions quoted above, marine survival is not cited
745 in any of the research which the various review committees recommended pursuing
746 [68-70]. Instead, the committees recommended three research foci:

747 “(1) *determine absolute escapement levels to Owikeno Lake... in order to*
748 *improve the credibility of stock assessment;*

749 (2) *improve the understanding of habitat use... by sockeye juveniles in Owikeno*
750 *Lake and smolts in the Wannock estuary; and*

751 (3) *investigate the status of ocean-type and lake-spawning sockeye, which are*
752 *less familiar and, although not specifically covered in this plan, may require future*
753 *intervention*”. (The joint committee noted that there was some evidence for an unusual
754 sockeye life history type that went directly to sea without rearing in the lake for a year
755 as pre-smolts (the normal life history pattern) [70]; the other committee reports have
756 similar language).

757

758 No mention is made of addressing the marine survival issue that was at the core
759 of the collapse; the reference to improving the understanding of smolt habitat use in the
760 “Wannock estuary” mentions that “*sockeye smolts do not appear to rear in these*
761 *estuaries for much time*” [69]. The report further mentions that there are numerous

762 estuaries within River and Smith Inlets, with varying sizes and importance to
763 salmonids. It is unclear why the Wannock was identified as particularly worthy of
764 investigation, but the report does note that “*approximately 25% of the Wannock estuary*
765 *was dyked and filled in 1973 for a log dump facility*” (i.e., almost two decades earlier).

766

767 The recommendations under Habitat are even more striking:

768 “5. *Existing conceptual plans for habitat restoration developed by DFO, the provincial*
769 *Watershed Restoration Program, and other stakeholders should be evaluated*
770 *for their potential long term benefits to sockeye, and the feasibility of proposed*
771 *restoration projects should be thoroughly assessed.*

772 6. *Habitat restoration projects could include the reconnection of spawning and early*
773 *rearing habitats along the margins of floodplains and in side-channels that have*
774 *been isolated by road construction or degraded by natural and logging-related*
775 *activities.*

776 7. *Any habitat restoration projects that are undertaken should be monitored to*
777 *determine their benefits for sockeye.*

778 8. *DFO and other agencies and stakeholders should continue to collaborate on*
779 *developing habitat protection strategy during resource development planning*
780 *processes (e.g., CCLCRMP, Forest Development Plans).*

781 9. *The site-specific and cumulative impacts of logging on habitats used by sockeye*
782 *should be more comprehensively evaluated”. (ref. [70]; the other committee*
783 *reports have similar language).*

784

785

786 In other words, despite the reports identifying with high certainty that
787 freshwater habitat issues were not contributory, the committees did not attempt to
788 understand the marine drivers and instead advocated a series of actions in freshwater.

789 ***Columbia River***

790 Two nearly contemporaneous studies identified the importance of either estuary
791 (lower river) or ocean processes in controlling the poor survival of Snake River salmon.
792 First, Kareiva *et al.* [73] applied a matrix life cycle model to demonstrate that recovery
793 of endangered salmon populations in the Columbia River could only be achieved by
794 improving survival in the lower river/estuary or in the coastal ocean and that (similar to
795 our own argument) even raising main stem survival to 100% would not prevent
796 extinction. Second, Marmorek and Peters [74] in a review of the PATH (Plan for
797 Analyzing and Testing Hypotheses) process, stated “*Importantly, we found that the*
798 *different models’ estimate of the survival rate of in-river migrants through the*
799 *hydropower system, a hotly debated value, was NOT an important determinant of*
800 *overall life cycle survival. Rather, the key uncertainties that emerged from these*
801 *sensitivity analyses were related to the cause of mortality in the estuary and ocean*”.
802 (See also [31]).

803

804 Probably owing to the lack of any direct information on juvenile survival in the
805 lower Columbia River and estuary regions, two initiatives were subsequently funded:
806 (a) the development of the JSATS acoustic telemetry system [75], and (b) directed
807 research using commercially available telemetry equipment to formally test the delayed

808 mortality hypothesis in the lower river and coastal ocean [76]. Both approaches
809 established that survival was high in the lower river below Bonneville Dam and lower
810 (but still high) in the estuary/plume region [56, 77-81]. The studies by Rechisky et al
811 extended these results further, showing that survival was even lower in the coastal
812 ocean region extending from the Columbia River plume to the NW tip of Vancouver
813 Island.

814
815 Despite these findings, further work to measure ocean survival and directly
816 address the conclusions of Kareiva *et al.* [73] and Marmorek and Peters [74] was not
817 carried out. After the ocean phase was identified as being the likely cause of poor
818 returns and not the lower river, research shifted to focus exclusively on studying
819 freshwater survival upstream at the hydropower dams. Although several publications
820 subsequently identified the presence of smolts in side channels within the estuary and
821 suggested the potential importance of estuarine wetlands for salmon conservation [82-
822 86], we are unaware of any studies that have actually identified low survival in the
823 estuary or established the period of residency—necessary requisites for improving
824 SARs. In summary, ocean issues remain largely unaddressed by Columbia River basin
825 salmon managers, and it is unclear whether research solely focussing on freshwater or
826 lower river/estuary issues will compensate for poor ocean survival.

827
828 Overall, these studies demonstrate a consistent pattern: a strong proclivity to
829 preferentially identify and work on freshwater habitat, even in cases where marine

830 survival has been identified as either the sole or most serious detriment to population
831 growth.

832

833 We are not arguing that freshwater monitoring should not be conducted;
834 monitoring population trends, and particularly survival, is critical to making informed
835 management decisions. However, monitoring alone is insufficient. As we noted in the
836 Introduction, the survival data used in this paper amount to a total of more than 3,000
837 years of sampling effort. Recent work in BC documented a substantial decline in
838 monitoring effort in north-central BC, and the authors argued that the situation must be
839 improved if salmon conservation efforts are to be effective [87]. While some degree of
840 monitoring is necessary, we note that the previously substantial monitoring effort was
841 insufficient to develop a successful management response. Obviously, if agencies
842 cannot respond effectively to the already available data indicating a widespread
843 collapse in marine survival of salmon populations that has been formally submitted to
844 the PSC on an annual basis, then it is unclear why simply increasing monitoring further
845 will lead to a more effective response. Clearly, greater monitoring alone does not
846 necessarily lead to improved conservation outcomes.

847

848 **Managing salmon research**

849 We are troubled that the increase in monitoring evident as survival has dwindled
850 over time was not matched by an equally intensive analysis to assess whether existing
851 approaches to salmon management are correct. Salmon smolt survival could only be
852 measured in most river systems after the relatively recent development of acoustic

853 telemetry, and PIT tags in the Columbia River Basin. Excluding smolt survival data
854 and focusing only on the adult survival (SAR) data, the number of years of available
855 data for Chinook and steelhead demonstrates a massive increase in monitoring over the
856 decades (pre-1975: 117 yrs; 1976-85: 318 yrs; 1986-95: 456 yrs; 1996-2005: 715 yrs;
857 2006-2014: 918 yrs). Yet, despite a nearly order of magnitude increase in monitoring
858 outputs, the point that basic aspects of this data set are in fundamental disagreement
859 with common assumptions about the cause of the “salmon problem” has gone
860 unrecognized. In brief, a minor industry has developed in salmon monitoring, but the
861 implications remained unappreciated.

862 We view it as critical that the roles of various proposed deleterious impacts on
863 salmon returns be rigorously quantified, rather than simply identified as important
864 without careful thought about other potential contributing factors. As both Lackey [88,
865 89] and Kareiva and Marvier [90] have noted, there is a widespread implicit assumption
866 that ecosystems unaltered by human activity are inherently good, and that restoring
867 anthropogenically altered freshwater ecosystems will help redress the problems (e.g.,
868 [91]).

869

870 Further, competing economic activities may be unfairly blamed for the ongoing
871 collapse of several important salmon species and unrealistic expectations placed on
872 what various recovery options may actually achieve. This is not simply restricted to
873 dam removal in the Columbia River basin or banning open-net salmon aquaculture in
874 British Columbia, two current hot button issues, but extends to impacts of forestry,
875 competing rights to groundwater, or development in general. Policy options for

876 promoting Chinook recovery need to recognize that the wide geographic footprint of
877 poor salmon survival likely implies that efforts focused on “fixing” possible
878 contributing factors specific to some regions are unlikely to be effective. At the very
879 least, these efforts should be held to a significant standard: (a) clearly demonstrating a
880 real and substantive improvement is possible, and (b) demonstrating a clear benefit
881 relative to the proposed costs.

882

883 Refocusing on marine migration pathways

884 The pattern of variation in SARs along the west coast of North America
885 suggests that a progressive worsening of marine survival with time occurred and was
886 accompanied by a geographic expansion northward in the region of poor survival.
887 However, several aspects of this explanation seem to be inconsistent with the roughly
888 similar coast-wide SARs now observed.

889

890 Fall Chinook are believed to remain shelf-resident for their entire marine phase
891 while Spring Chinook migrate north on the shelf before eventually moving off-shelf or
892 into the Bering Sea/Aleutian Islands. Because both groups have poor SARs, this would
893 imply that the area of poor marine survival might be restricted to the coastal shelf off
894 Washington, British Columbia, and SE Alaska; however, the large-scale collapse in
895 adult Spring Chinook returns includes the Yukon and Kuskokwim Rivers (draining into
896 the Bering Sea) and the Kenai River (Cook Inlet, Gulf of Alaska) [20-23, 92, 93]. This
897 suggests that either the area of poor marine survival is now simultaneously large, so
898 that exposure times to regions of poor survival are similar, or that all stocks congregate

899 at some point in the marine phase into a more geographically confined region where
900 their survival is similarly affected.

901

902 We have no evidence for the latter possibility. Fall (subyearling) Chinook
903 stocks only migrate as far north as SE Alaska [94, 95] after one or more years at sea
904 (and at least some Strait of Georgia and Puget Sound Chinook remain resident in
905 southern BC waters for their entire marine lifespan [96-100]). The marine movements
906 of eulachon [24] and lamprey [25], which have also undergone dramatic declines in
907 abundance, are less well-known but are likely similar to Fall Chinook. Thus, the
908 conditions leading to poor marine survival must be geographically widespread because
909 western Alaska Spring Chinook are not known to migrate to the shelf region off SE
910 Alaska or BC.

911

912 A key prediction is that stocks with the lowest SARs should have greater
913 exposure to poor ocean conditions in southern regions. The anomalously high SARs of
914 some specific salmon populations (Fig 4) might provide the basis for an explicit test of
915 this prediction. Although our understanding of population-specific differences in
916 marine migration routes is currently very limited, especially for steelhead [101, 102],
917 there is now some developing evidence for differential salmon survival in the sea; e.g.,
918 [100, 103-105]).

919 Assuming that the region of poor survival progressively expanded northward
920 along the coast at the time of successive regime shifts, there are several testable
921 hypotheses. For example, Strait of Georgia or Puget Sound Chinook populations may

922 have lower survival than adjacent outer coast stocks (west coast Vancouver Island,
923 coastal Washington) because they either remain resident for a longer time period in
924 coastal marine waters with similar survival rates (greater exposure), or because survival
925 rates per unit time are lower in Strait of Georgia waters (greater rates of loss). This
926 could also potentially explain why SE Alaska and north-central BC Chinook stocks in
927 recent years still have SARs ~2X Snake River stocks and ~4X Strait of Georgia stocks
928 (Fig 6)—Strait of Georgia Chinook stocks remain resident in the Strait of Georgia for
929 multiple months after ocean entry [106, 107], while Snake River yearling Chinook
930 juveniles promptly migrate north along the outer shelf to Alaska [55, 56, 108].

931

932 In this context, the consistently low survival of the Dworshak Hatchery
933 yearling Chinook relative to other Snake River Chinook stocks is noteworthy; mean
934 survival from Lower Granite Dam to adult return over the 2000-2015 period was only
935 0.58% for the Dworshak Hatchery stock versus 1.28% for McCall Hatchery and 1.29%
936 for Imnaha Hatchery fish (ref [5], Tables B.16, B.22, & B.24). The Dworshak SAR is
937 thus less than ½ that of the other two populations. All Snake River populations migrate
938 through the same set of dams, so one explanation for the particularly low survival of the
939 Dworshak population could be a differential migration to an area of the North Pacific
940 (or Bering Sea) whose relative survival prospects was only one-half that of other
941 regions (Columbia River Chinook salmon are known to be seasonally present in the
942 Bering Sea and to overwinter in the Gulf of Alaska [109]). Our tenuous understanding
943 of where Chinook and steelhead migrate to in the ocean and how long they remain in

944 various regions (let alone how these patterns differ between populations) clearly needs
945 urgent improvement if these issues are to be resolved.

946

947 One important possibility for establishing the geographic differences in survival
948 is if predators increasingly target returning adult salmon. There is now ample evidence
949 for substantial increases in marine mammal abundance and presumably predation on
950 returning adults [110-115]. Ohlberger et al [116] reviewed the decline in size and age-
951 structure of Chinook across western North America. They noted that consistent with
952 the adult predation hypothesis, the decline was most pronounced in the older age
953 groups in some (but not all) regions of the eastern Pacific. Recent work has also
954 demonstrated that in fish, large females may confer higher fitness on their offspring
955 [117].

956

957 Competition for food may also conceivably play a role. The geographically
958 widespread decline in salmon growth over time seen for multiple species by the mid-
959 1990s, and which was potentially attributed to the growth of hatchery production of
960 pink salmon [118] has apparently continued. Continued increase in pink salmon
961 abundance has been shown to affect plankton populations [119] and reduce survival of
962 at least one marine seabird (shearwaters) [120, 121] as well as some salmon species [4,
963 122]. Thus, geographically variable rates of competition with pink salmon or marine
964 mammal predation at older ages could both contribute to determining differential rates
965 of salmon survival.

966

967 Large differences in SARs point to important directions for future study. A
 968 very few stocks have SARs 3 to 4-fold higher than nearby stocks. At the extreme, the
 969 Chilliwack stock of Fall (subyearling) Chinook has a median SAR of ca. 4%, an order
 970 of magnitude greater than other nearby Strait of Georgia stocks. Oregon Coastal Fall
 971 Chinook also have SARs much higher than any Columbia River basin stocks.
 972 Understanding why only a few populations consistently have high SARs when
 973 returning from the ocean as adults could pay large dividends in understanding what
 974 differences in ocean experience result in those few populations remaining productive
 975 while many others have essentially collapsed. As Peterman and Dorner [13] remarked
 976 for sockeye, “*Further research should focus on mechanisms that operate at large,*
 977 *multiregional spatial scales, and (or) in marine areas where numerous correlated*
 978 *sockeye stocks overlap*”. The markedly higher SARs evident for Oregon coastal
 979 Chinook relative to most other populations (Figs 4 & 5) may provide important
 980 guidance in this context. Riddell et al [123] (p. 580) specifically note the unique
 981 marine distributions of southern Oregon Chinook stocks, which restricts them for their
 982 entire ocean phase to life in the California Current. Nicholas and Hankin [124] (Table
 983 2) report that Fall Chinook from the Salmon and Elk rivers in Oregon are north
 984 migrating stocks and that Oregon coastal stocks show variation in ocean migration
 985 “*with some migrating north, some south, and one stock has a mixed north and south*
 986 *ocean migration*” [14]. Lending credence to the possibility that ocean migration
 987 pathways influence productivity, Nehlsen et al [14] reported that the few “south
 988 migrating” Oregon Fall Chinook stocks were all characterized as having “depressed”

989 runs in 1988 (prior to the 1989 regime shift), whereas the “north migrating” runs all had
990 no or increasing abundance trends.

991

992 It thus seems plausible that specific salmon populations have genetically
993 determined migration behaviours that allow them to home to distinct feeding grounds
994 within the North Pacific, some of which confer better survival [125]. Batten et al [126]
995 identified at least 10 geographically distinct plankton communities evident in a single
996 transect across the North Pacific that were temporally stable across years and
997 demonstrated that geographically distinct seabird assemblages patterned similar to the
998 plankton communities. An analysis of tufted puffin communities [127] found that
999 different forage fish communities also were present in different sub-regions of the
1000 Aleutian Chain. Thus geographically stable and distinct biological communities exist
1001 within the North Pacific Ocean, including the pelagic offshore. Salmon populations
1002 homing to different feeding grounds (or a succession of different feeding grounds)
1003 could therefore have very different fates if these regions develop differently over time,
1004 for which there is at least some experimental evidence [99, 128, 129].

1005 **Columbia River basin policy implications**

1006 A critical policy question for the Columbia River basin concerns whether
1007 recovery of listed fish stocks is limited by the hydropower system as currently operated,
1008 or by ocean conditions [130]. The available evidence indicates that smolt survival
1009 during downstream freshwater migration is not higher in rivers without hydropower
1010 dams (Fig 1 and Table S1) and that a number of much shorter coastal rivers have even

1011 lower smolt survival than is experienced through the Columbia River hydrosystem,
1012 particularly when survival is scaled by distance travelled.

1013

1014 Bisbal and McConnaha [130] suggest several ways in which aspects of the
1015 freshwater habitat might be manipulated to improve ocean survival. However, given
1016 that recovery targets are specified in terms of attained SARs, current evidence indicates
1017 that Snake River SARs are roughly equal to (or better) than those currently achieved in
1018 the nearby Salish Sea region, a region where dams are absent. It therefore seems
1019 unlikely that recovery can be achieved without an improvement in ocean survival.
1020 Unfortunately, current scientific knowledge is simply insufficient to understand how to
1021 promote this.

1022

1023 **The future of Pacific salmon**

1024 Salmon are cold water fish living in a rapidly warming world. There are no
1025 easy answers for maintaining Pacific salmon populations [131] and current problems
1026 are likely to get much worse. At least eight separate ice ages are recorded in the last
1027 800,000 years of the ice core record alone [132] and there were likely more than 50 ice
1028 ages over the past 2.6M year extent of the Quaternary [133]. Climate change is thus the
1029 norm, not the exception. However, projected levels of future climate change are far
1030 outside anything experienced in either the last 150 years of industrialization or the
1031 previous 2.5M years of the Pleistocene Epoch. Recent marine heat waves along the
1032 Pacific Coast [134] are thought to have had significant negative effects on adult salmon
1033 returns [135]. The frequency, duration, and intensity of marine heat waves are all

1034 projected to increase dramatically in future [136], further exacerbating already serious
1035 problems for salmon.

1036

1037 Current CO₂ emission policies are expected to limit warming by 2100 to
1038 approximately 3.0°C [137], or more than four times greater warming than the total
1039 warming experienced over the past 150 years of the instrumental record (~0.7°C).
1040 Even if all countries meet their commitments under the Paris Agreement, these
1041 emissions scenarios are predicted to see global mean temperatures stabilize at 1.5–
1042 2.0°C above pre-industrial levels, or ca. 2-3 times the temperature increase so far—and
1043 an increase achieved in the next 80 years, not 150 years.

1044

1045 Warming rates 4-6 times those experienced in the recent past mean that further
1046 surprises in how much salmon survival drops are inevitable. Given the past slow and
1047 erratic response, the likelihood that the fisheries community will identify the correct
1048 drivers of the problem and then move to successfully address them is not high if current
1049 practice continues. So far, the response has been to re-double efforts on what we know
1050 how to study (freshwater) and to largely avoid what we currently have little ability to
1051 study (the marine phase). There are real economic, social, and biological costs to doing
1052 so, with many groups now identifying various single issue factors as the primary
1053 underlying problem that needs to be fixed (hydropower dams, salmon aquaculture,
1054 forestry, land use practices, water rights). These region-specific issues cannot possibly
1055 be the drivers of the continental-scale response that we document and to further delay
1056 not only increases the threat to salmon, but to those species that rely on them, such as

1057 southern resident killer whales. Wasser et al [48] state that “*Low availability of*
 1058 *Chinook salmon appears to be an important stressor among these fish-eating whales as*
 1059 *well as a significant cause of late pregnancy failure, including unobserved perinatal*
 1060 *loss... Results point to the importance of promoting Chinook salmon recovery to*
 1061 *enhance population growth of Southern Resident killer whales.*” There are many real
 1062 consequences to ineffective policy responses—lost time, inability to boost salmon
 1063 abundances (for both human and non-human salmon predators alike), and elevated
 1064 costs for many other industries.

1065

1066 The history of North American marine research on Pacific salmon has been
 1067 described elsewhere [138-140]. In the last decade, small-scale efforts to describe the
 1068 marine life history of juvenile salmon have developed in specific regions of the
 1069 continental shelf (no small feat in itself; e.g., [141-145]). However, life history
 1070 observation is useful to infer possible mechanisms affecting overall biology, not to test
 1071 and validate the mechanisms driving survival. This means that the rapid learning
 1072 characteristic of physics or chemistry, where hypotheses are explicitly tested and
 1073 important scientific advance occurs when theories are rejected (not merely advanced),
 1074 is unlikely because it is difficult to refute observation-based mechanisms. A key issue
 1075 is that if marine survival problems are widespread along the Pacific Coast, mechanisms
 1076 specific to only some continental shelf regions or river watersheds cannot be the major
 1077 driver unless the movements of all the different salmon populations expose them to
 1078 these stressors. Because poor marine survival is demonstrably widespread, research

1079 and policy predicated on the assumption that the problems are geographically specific
1080 is unlikely to be successful.

1081

1082 Given the massive investment in restoration and monitoring activities for
1083 Pacific salmon, the development of correct conservation analyses and policy planning
1084 is critical. Over \$1 Billion is now spent annually in the continental United States alone
1085 on freshwater habitat restoration [146, 147], and there is great pressure to remove or
1086 modify hydropower dams in the Columbia River basin to rebuild salmon runs to
1087 historical levels of abundance and productivity and more recently to help endangered
1088 orca populations [48]. Within the Columbia River, the total cost of recent conservation
1089 efforts reaches or exceeds ca. 25% of FCRPS annual revenues (including foregone
1090 clean power generation), or >\$0.5 Billion per year [148]. Similarly, in British
1091 Columbia, where dams are not present in the migration paths, much effort has focused
1092 on removing salmon farms to help restore Fraser River salmon populations [149-151].
1093 Clearly, it is important to understand the impact of various anthropogenic impacts on
1094 poor salmon returns, but it is also important that the real prospects for improvement as
1095 a result of these region-specific actions are carefully assessed.

1096

1097 In the novel “The Sun Also Rises”, the character Bill Gorton is asked how he
1098 went bankrupt. He replied, “*Two ways. Gradually, then suddenly.*” [152]. The same
1099 process appears to be playing out in the ways fisheries science has addressed the marine
1100 survival problem for salmon. In west coast salmon management, the first issue was
1101 incorrectly diagnosing the problem (poor and worsening ocean survival) as primarily a

1102 freshwater issue, and the second problem was failing to change behaviour quickly and
 1103 maintaining a focus largely on freshwater issues (which may inflict significant costs on
 1104 other economic activities). As with economic bankruptcy, failing to staunch losses and
 1105 persisting with previously unsuccessful behaviour is a recipe for eventual catastrophic
 1106 loss. Some positive response is certainly evident, in that harvest from Chinook and
 1107 steelhead fisheries was substantially restricted (e.g., [49]). However, harvest rates of
 1108 shelf-resident Fall Chinook were historically in the 50%-60% range. As we have
 1109 demonstrated, even the complete elimination of all harvest can only compensate for a
 1110 roughly two-fold decline in marine survival; for Spring Chinook and steelhead, which
 1111 are much less impacted by saltwater harvest, maximum compensation is far less.

1112

1113 It is not unreasonable to anticipate a further ten-fold decline in the marine
 1114 survival of salmon from climate change in the relatively near future. In fact, the
 1115 survival time series used in this manuscript generally end prior to 2015. The datasets
 1116 therefore do not include more recent years of even worse anticipated survival. An
 1117 overall pattern of low smolt to adult returns of upper Columbia and Snake river Spring
 1118 Chinook salmon and steelhead has been reported for 2015-16 and is considered likely
 1119 to worsen given the apparently poor early ocean survival of juvenile salmon in 2017
 1120 and unprecedented ocean conditions occurring in 2018 in the Northern California
 1121 Current and Gulf of Alaska [153-155].

1122

1123 With the option of reducing harvest rates now almost exhausted, large
 1124 reductions in escapement can now be expected similar to what occurred in the Rivers

Coast-Wide Survival of Chinook and Steelhead **SUBMITTED to PLoS-ONE**

1125 Inlet case study. Without improved understanding of what is happening at sea,
 1126 potentially inappropriate policy recommendations seem likely to continue. As we have
 1127 shown in the case studies, each time salmon research reached the point where it became
 1128 clear that the survival problem was at sea, the ensuing response was to re-focus effort
 1129 on freshwater activities, leaving the marine survival issues unaddressed while often
 1130 increasing potentially costly freshwater interventions. We view this as evidence of
 1131 widespread cognitive dissonance [65] and significant groupthink [66] in salmon
 1132 management. A useful first step towards breaking the current impasse would be to
 1133 determine whether differences in early marine migration pathways and survival of
 1134 geographically close populations cause the strongly disparate SARs that we document
 1135 for some populations.

1136 **Methods**

1137 **Smolt survival data**

1138 To assess freshwater survival levels for Chinook and steelhead smolts migrating
1139 downstream, we collated all published studies for west coast North American rivers
1140 (Table S1), excluding California, and then scaled survival by distance travelled. In a
1141 previous paper [41], we collated data for the Fraser-Thompson and Columbia-Snake
1142 rivers for comparison. Our current paper includes available survival estimates from
1143 coastal Oregon to northern Vancouver Island as well as one additional smolt survival
1144 study for the Fraser River (Chilko Chinook).

1145

1146 Smolt survival during downstream migration was available for several regions,
1147 but the data are most extensive in the Columbia River basin where PIT tag-based
1148 studies have been conducted for over two decades and since the more recent
1149 development of acoustic tags (Juvenile Salmon Acoustic Tracking System (JSATS) or
1150 VEMCO). In other river systems, survival during downstream migration was estimated
1151 using VEMCO acoustic telemetry; there are no published PIT tags survival estimates
1152 outside the Columbia River basin. A total of 531 estimates, representing 73 individual
1153 populations, runs or release sites, and time series between 1-23 years in length were
1154 used in the comparison. All survival estimates were calculated using the Cormack-
1155 Jolly-Seber model or its derivatives. The specific methods can be found in the sources
1156 listed in Table S1.

1157

1158 Within the Columbia River, smolt survival was estimated from various release
1159 points including dams, traps located in rivers, and hatcheries. Downstream census
1160 points were PIT arrays located at dams or acoustic receiver arrays in the lower
1161 Columbia River or estuary. In the Columbia River basin, and particularly the Snake
1162 River, a proportion of smolts are diverted into barges at the dams and then transported
1163 downstream to below the lowest (Bonneville) dam; these fish were not included in the
1164 hydrosystem estimates, but may have been included in lower river and estuary
1165 estimates. Most Columbia River basin smolt survival estimates (N=461) were
1166 calculated by NOAA or the Fish Passage Centre using PIT tag data. Twenty-eight were
1167 from JSATS or VEMCO acoustic tag studies. Smolt survival in the basin was measured
1168 over distances ranging between 144 - 909 km.

1169

1170 In the other regions, smolt survival was estimated from hatcheries or traps, to
1171 acoustic receiver arrays near the river mouths. In some cases, fish were transported
1172 either upstream or downstream of their tagging location, e.g., Chilko Chinook were
1173 reared at a lower Fraser River hatchery but released ~500 km upstream in the Chilko
1174 River. Migration distances to the sea after release were typically much shorter than in
1175 the Columbia or Fraser Rivers (see Table S1). Excluding the Fraser and Columbia
1176 River populations, average smolt migration distance was only 19 km for all other
1177 regions.

1178

1179 To better compare survival across basins we scaled survival measurements by
1180 the migration distance. We used distance because travel time was not reported for all of

1181 the studies. We excluded survival estimates from [41] that were based on populations
1182 where >75% of smolts had fork lengths not meeting current best practices on acceptable
1183 tag burdens [156-161] (<130 mm for VEMCO V7 and < 140 mm for V9). This
1184 resulted in the exclusion of three survival estimates from the Nicola and Spius River
1185 tributaries of the Fraser River because of high tag burden that were included in our
1186 earlier paper [41].

1187 Data sources for Chinook

1188 Most survival rates of Pacific salmon are based on mark-recapture efforts,
1189 where juveniles are “marked” or implanted with a tags--either coded wire tags (CWT)
1190 or passive integrated transponder (PIT) tags, and recaptured in the fishery or upon
1191 return to the river. The basic tag technologies are well described elsewhere [162-166].

1193 CWT technology dates back to the 1960s. A review is provided by [167] and
1194 the application of the methodology to coastal marine migrations of coho and Chinook is
1195 described by [95, 168] and to measuring harvest and survival by [21, 49, 169].

1196 Because the tag is implanted in the nose cartilage of smolts, the fish must be dissected
1197 to recover the tag after capture, ensuring the death of that particular tagged animal and
1198 preventing further study of its movements. CWT technology provides the basis for the
1199 Pacific Salmon Commission’s Chinook survival database used for coast-wide
1200 management of Chinook salmon under the Pacific Salmon Treaty [49]. We used this
1201 database as the source of Chinook survival data for all regions outside the Columbia
1202 River basin and for a few stocks located in the Columbia River basin (Table S2). The

1203 data are contributed by the various governments (provincial, state, and federal agencies)
1204 responsible for conducting the individual monitoring programs.

1205

1206 In contrast, systematic survival data based on PIT tags first came into
1207 widespread use in the Columbia River Basin in 1997 (Table S2). PIT tags are long-
1208 lived but extremely short distance radio frequency tags that can successfully transmit
1209 their unique ID code only when within <0.5 m of a detector. Although there are some
1210 recent exceptions in small rivers, the short detection range essentially limits the use of
1211 PIT tags to the Columbia River dams, which channel sufficient tagged individuals close
1212 to the detectors to generate useful results. Tagging data are contributed to a central
1213 database (PTAGIS- Pit Tag Information System) by the various agencies (state, tribal
1214 and federal) and the SARs are estimated by the Fish Passage Center. All PIT tag-based
1215 SAR estimates reported in this paper are taken from the Fish Passage Center's
1216 Comparative Salmon Survival (CSS) Study (McCann et al [5]) and are listed in our
1217 Table S2.

1218

1219 Earlier survival data for Snake River Chinook populations from the 1960s and
1220 1970s is available from Raymond [1], who noted that "*From the positive relation found*
1221 *between rates of return of adults and survival rates of smolts, it was apparent that*
1222 *mortality of smolts migrating downriver through the dam complex was the main cause*
1223 *of the decline in Snake River salmon and steelhead runs*", a view that has become
1224 commonplace amongst salmon biologists. We have included these data in our analysis
1225 because Raymond's pioneering studies [1, 29, 30] are of unique importance owing to

1226 the documentation of the high SARs occurring in the 1960s and early years of the
1227 1970s, a time period prior to the completion of the Snake River dams and the 1977
1228 marine regime shift, and because they defined the focus for much subsequent work in
1229 the Columbia River basin to improve survival.

1230

1231 The two major tagging technologies available, PIT and CWT, are largely
1232 geographically discrete, with most recent survival data from the Columbia River based
1233 on PIT tag technology and most survival data for other regions based on CWT data.
1234 Although rarely discussed, the differences in the two technologies determine what
1235 aspects of migration-phase survival are estimated. These difference are discussed
1236 below, as is a brief description of Raymond's methods. (Raymond's [1] early survival
1237 analysis was based on direct estimation of the number of smolts migrating downstream
1238 past Snake River dams, and dividing this value into the number of adults returning
1239 several years later; see Raymond [1] for details; as such, comments on the extent of the
1240 migration path monitored also apply to this early study).

1241

1242 ***CWT tags***

1243 The precise technical methods of counting the number of CWT-tagged adults
1244 returning back to each population are not documented in the Pacific Salmon
1245 Commission (PSC) database by the various agencies contributing survival data;
1246 however, an example of the mark-recapture approach used by ADF&G in the
1247 Transboundary Rivers of SE Alaska and Northern British Columbia for wild Chinook
1248 stocks can be found in [21]. Most agencies operate hatcheries or (in a few cases) rotary

1249 screw traps to estimate downstream smolt numbers for wild stocks. In general, CWT-
1250 based survival estimates are calculated for hatcheries by dividing the total number of
1251 maturing adults of various ages estimated to return back to the spawning grounds or
1252 hatchery or caught in the fishery by the number of smolts released in the year of ocean
1253 entry. CWT-based survival estimates (49 time series) are only available for Chinook,
1254 not for steelhead.

1255 The PSC database provides several measures of marine survival. In this
1256 study, we used survival data calculated as the sum of adults returning at all ages or
1257 caught in the fisheries, uninflated for losses to natural mortality for Chinook remaining
1258 at sea for longer than two years because these values are most similar to the CSS PIT-
1259 tag based survival estimates [5]. Survival estimated using this procedure slightly
1260 underestimates true survival to ocean age two because some two year old Chinook
1261 destined to mature at older ages die from natural causes prior to maturing and are
1262 therefore not enumerated. However, in cases where SAR (survival over the migratory
1263 phase of the life history), is 1%, the instantaneous total mortality rate is $M_{\text{Total}}=4.6$.
1264 Ricker [170] suggested that the loss due to natural mortality between age two and older
1265 ages was perhaps $M=0.46 \text{ yr}^{-1}$, or only 10% of M_{Total} . More recent estimates of age-
1266 specific natural mortality for Chinook are even smaller: age 2, 40%; age 3, 30%; age 4,
1267 20%; and age 5 and older 10%; ([49], p. 8). Consequently, not correcting for natural
1268 mortality losses occurring between age 2 and older ages is unlikely to introduce major
1269 errors into the SAR estimates, particularly as the majority of Chinook return at ocean
1270 age two, and especially so in recent years [116]. (The PSC database also includes
1271 survival estimates with age 3+ adults inflated to account for losses at sea; we chose not

1272 to use these estimates because the PIT-tag based survival estimates are not inflated for
1273 mortality at older ages, so for purposes of comparison uninflated values should be more
1274 comparable. We highlight it because of our concern (see Discussion) that fisheries
1275 biologists may be underestimating the magnitude of losses at older ages and thus
1276 incorrectly assuming that the primary survival issue is in the first year after ocean entry.
1277

1278 ***PIT tags***

1279 PIT tag estimates of SARs are taken directly from Appendix B of McCann *et al*
1280 [5], who reports SAR in several ways. We selected the SARs covering the greatest
1281 extent of the migratory life-history (i.e., smolt releases and adult returns to the highest
1282 dam available in the Columbia River basin), and we generally used SAR estimates that
1283 included jacks when available. In the mid-Columbia region, SAR estimates with jacks
1284 were sometimes available only for a shorter migration segment; in these cases we
1285 selected the SAR data sets representing the longer migration segment but excluding
1286 jacks because this was most similar to the CWT survival estimates. The extensive PIT-
1287 tag based SAR estimates for the Columbia River basin total N=45 Chinook SAR time
1288 series and N=22 steelhead SAR time series [5].

1289 Because returning adults must ascend fish ladders with PIT tag detectors, all
1290 PIT tagged adults surviving to return can be censused (ignoring tag shedding).
1291 Dividing adult returns by the estimated number of tagged smolts reaching the most
1292 upstream dam in the year of ocean entry provides an estimate of the SAR. However,
1293 the PIT-tag based SAR estimates for the Columbia River basin differ from CWT-based

1294 estimates in three main ways: (i) they exclude losses to harvest (lowering survival
1295 relative to what is estimated in the PSC database), (ii) they exclude losses occurring
1296 from smolt release to encountering the first dam in the migration path (raising survival),
1297 and (iii) they exclude losses occurring from the time the returning adults migrate past
1298 the last dam until they reach the spawning grounds (raising survival). We review these
1299 differences in the context of the two major life history groups (yearling and
1300 subyearling) for Chinook.

1301 **Data sources for steelhead**

1302 All steelhead data analyzed in this paper were from Kendall et al [7] updated to
1303 incorporate more recent years' data, including new information on the actual age-
1304 structure of the adult returns. Kendall et al [7] should be consulted for a full
1305 description of these data sets and the data are available directly from Dr Kendall (Dr
1306 Neala Kendall, pers. comm. Neala.Kendall@dfw.wa.gov).

1307

1308 **Chinook**

1309 ***Division by life history type***

1310 In general, Chinook salmon display two life history types: subyearling and
1311 yearling. These life history types are identified in our analysis because there are
1312 important ecological differences between them (see reviews by [123, 171], and
1313 references therein) which likely influence survival. Subyearling smolts migrate to the
1314 ocean within a few months of hatching in the spring, while yearlings migrate to sea

1315 after completing one or more full years of life in freshwater, and are thus significantly
1316 larger at ocean entry and (generally) spend one less year in the ocean. The
1317 subyearling/yearling smolt life history types also generally correspond with adult run
1318 timing (“Fall” or “Spring”), but this linkage is somewhat subjective primarily owing to
1319 hatchery rearing practices.

1320

1321 Spring (yearling) populations are largely found in high altitude headwater
1322 tributaries of large river systems penetrating well into the interior of the continent such
1323 as the Columbia and Fraser rivers, and are the only Chinook life history type reported
1324 for Alaskan rivers [172, 173]. In contrast, Fall (subyearling) populations are widely
1325 found in low gradient coastal streams or in the lower mainstem of major rivers but are
1326 absent from Alaska. Early work [174] suggested an ancient genetic divide with
1327 subyearling Chinook smolts primarily produced by adult runs returning to freshwater in
1328 the fall and spawning directly after reaching their natal streams, and yearling smolts
1329 produced mainly by adults that return in the spring and then hold in freshwater without
1330 feeding until spawning in the autumn.

1331

1332 ***Life history & harvest rates***

1333 Spring Chinook are thought to move offshore and become purely open ocean
1334 residents for much of the marine phase, and thus essentially immune to harvest by
1335 fisheries until their return. As a consequence, offshore (pelagic) harvest of Spring
1336 Chinook is likely negligible because a convention banning high seas fishing beyond the
1337 200 mile EEZ of Pacific Rim countries was signed in 1992

1338 (http://www.npafc.org/new/about_convention.html) and enforcement patrols
1339 consistently find few illegal driftnet vessels and only in the far western Pacific, well
1340 beyond the known ocean distribution of North American Chinook stocks [175, 176]
1341 (but possibly not steelhead). However, some incidental harvest of immature and
1342 maturing Chinook occurs in the groundfish fisheries of the Bering Sea, with current
1343 evidence suggesting that Pacific northwest populations form ca. 1/3 of Chinook bycatch
1344 in the Bering Sea/Aleutian Islands region [109]. Unfortunately, owing to a general
1345 inability to use collected Chinook fish scales to determine the duration of the freshwater
1346 period (and thus discriminate yearling from subyearling animals), it is unclear which
1347 life history type the Pacific northwest populations analyzed in [109] represent.

1348

1349 In contrast, Fall Chinook are known to remain as long-term residents of the
1350 continental shelf off the west coast of North America and are thus exposed to
1351 commercial and sport harvest in coastal marine waters over multiple years [171].
1352 Survival of shelf-resident subyearling Fall Chinook populations can therefore be
1353 significantly reduced by coastal fisheries that can harvest these animals over several
1354 years of marine life.

1355

1356 In reality, this relatively simple picture is more complicated. Some hatcheries
1357 hold subyearling (Fall) Chinook for an additional year before releasing them as larger
1358 yearling smolts, and others release Spring run Chinook as subyearlings (e.g., Nooksack
1359 and Skagit-See Table S2). Thus some yearling production is of smolts that
1360 presumably remain shelf-resident for several years because their intrinsic genetic make-

up dictates this behaviour despite their larger (and older) age at release. Sharma and Quinn [171] also document regional differences in migration distribution between lower Columbia River and upper Columbia-Snake River Spring yearling populations which they attribute to possibly greater interbreeding between Spring and Fall run individuals in the lower Columbia River. Clarke et al [177] similarly present evidence from breeding trials that the yearling/subyearling smolting pattern follows simple Mendelian genetic rules in crosses of Fall and Spring adults (with the added twist that the sex of the parent also influences the result)! More recent work by Prince et al [178] has potentially identified a single gene in both Chinook and steelhead that controls early (spring or summer) re-entry of Chinook and steelhead that then mature in freshwater prior to spawning in the autumn; whether and how this gene might also influence marine migration behaviour is unknown.

Very recently, Riddell et al [123] have reviewed the literature and made the argument that repeated parallel evolution of the yearling and subyearling life history types in Chinook may have occurred in different watersheds. If true, this makes Healey's [173] earlier assumption that yearling (Spring) Chinook and subyearling (Fall) Chinook have strongly dichotomous ocean migration pathways untenable unless evolution of age at ocean entry is strongly linked to migratory behaviour in the ocean.

In this paper, we have thus opted to aggregate smolt returns by age at ocean entry (yearling, subyearling) for simplicity, but note that in future it would be very valuable to disentangle the role of age at release from genetically determined differences in marine migration pathways on survival. Unfortunately, a rigorous assessment of the genetic origins of each hatchery program would almost certainly

1384 require a genetic determination of whether each hatchery program was releasing Fall or
1385 Spring Chinook, and would need to take into account whether or not hybrid populations
1386 had been created; it is a fascinating research question whose answer is completely
1387 unclear at the current time to contemplate whether the offspring of an inadvertent
1388 hybridization between a Fall and a Spring Chinook parent would rear offshore or on the
1389 shelf and how it would get there!

1390 The difference in likely marine rearing areas is important because in CWT-
1391 based estimates of survival [49], the commercial and sport harvest of the different age
1392 groups is added to the escapement to generate the reported SAR. In contrast, PIT tag-
1393 based survival estimates for the Columbia River basin do not incorporate losses due to
1394 harvest ([5]; see p. 95). Columbia River survival estimates using PIT tags will
1395 therefore underestimate survival relative to the PSC's CWT-based survival estimates.
1396 For example, the PSC (Table 2.7) reports average annual stock-specific harvest rates of
1397 29-62% for Strait of Georgia Fall (subyearling) Chinook stocks with harvest rates
1398 declining over time [49]. For some Spring (yearling) Chinook, harvest rates are much
1399 lower (at the extreme, Willamette Spring Hatchery Chinook are reported as having only
1400 a 11% mean harvest rate; see Table 2.10 of [49]).

1401

1402 In this report we do not attempt to directly correct for the effects of harvest or
1403 differences in the proportion of the migratory phase survival is measured over because
1404 our most important conclusions seem robust to these differences, but it is important to
1405 recognize that methodological differences exist and influence survival estimates. In a
1406 few situations, we found both CWT and PIT tag-based survival estimates for the same

1407 population and the same release year (Supplementary Info S3). Relative to the 1:1
1408 relationship expected if both methodologies “perfectly” captured the same survival
1409 process, we find ratios of $1.5\text{SAR}_{\text{CWT}}:1\text{SAR}_{\text{PIT}}$ for three subyearling Chinook
1410 populations, consistent with expectation as CWT-based SAR estimates incorporate
1411 harvest, while PIT tag based estimates do not. Unfortunately, we did not find data to
1412 directly compare yearling Chinook survival estimates but provide some indirect
1413 comparisons in Supplementary Info S3.

1414

1415 Summarizing, the PIT tag-based survival estimates for the Columbia River
1416 basin are biased high relative to total migratory phase survival because these estimates
1417 exclude losses in the initial and final phases of the migration period above the dams,
1418 and biased low because they exclude harvest (which varies in potential influence
1419 between large for Fall (subyearling) and low for Spring (yearling) stocks). Finally,
1420 some of the CWT-based survival estimates for wild stocks are also biased low to some
1421 degree because they exclude survival losses occurring in the initial and final phases of
1422 the migration upstream of the enumeration points for smolts and adults. However, at
1423 least for hatchery-reared populations, smolt numbers used in the denominator of the
1424 CWT survival estimate are estimated at the time of release from the hatchery, and
1425 therefore exclude the possibility of migratory losses occurring prior to census.

1426

1427 For these reasons it should be noted that the strongest comparisons are within
1428 individual survival time series (the coast-wide declining trends in survival) which are
1429 based on the most consistent methodologies, while comparison between populations

1430 will be less reliable because of differences in where each populations is censused to
1431 measure survival over the migration phase. However, the coast-wide convergence of
1432 survival in recent years to very low levels at a time when most sport and commercial
1433 harvest has been drastically reduced is strong evidence that a common factor is driving
1434 the collapse in survival. It is unlikely that a single consistent conversion factor between
1435 CWT and PIT tag-based SAR estimates can be derived, because survival losses
1436 incurred upstream of the initial and final census point for calculating SARs can vary
1437 substantially between rivers and between populations within a river system and only
1438 CWT-based methods can account for losses to harvest. Only hatchery releases can
1439 potentially reach this technical standard of measuring survival over the entire migratory
1440 phase of the life history, and only if adult enumeration takes place on the spawning
1441 grounds (or at the hatchery).

1442 **Steelhead**

1443 The migration of steelhead is poorly understood, but it is thought that they may
1444 migrate directly offshore soon after the smolts reach saltwater [102, 179]. Virtually
1445 nothing is known of their marine migration, although the open ocean distribution
1446 extends as a band bounded by specific maximum and minimum sea temperatures across
1447 the North Pacific [180]. This suggests that (similar to Spring Chinook) maturing
1448 steelhead may return directly from the offshore to their natal river and be little exposed
1449 to commercial fisheries operating in continental shelf waters except those lying on the
1450 direct migration path from the offshore. No commercial fisheries target steelhead, so
1451 harvest is limited to freshwater sport fisheries and saltwater bycatch in other fisheries.
1452

1453 Although many steelhead rivers and hatcheries are located in B.C., adult returns
1454 have not been accurately enumerated which prevents direct estimation of survival. As a
1455 result, SAR data for British Columbia is restricted to the Keogh River (Fig 1), where a
1456 weir located within ca. 300 m of the ocean has monitored wild steelhead since 1977
1457 [181]. Despite the lack of SAR data for other populations, it is known that the survival
1458 trends evident for the Keogh River are mirrored in adult returns for the province of B.C.
1459 as a whole, with some differences evident between geographic regions [40, 182, 183] in
1460 more recent regime periods. Importantly, it is broadly recognized that adult steelhead
1461 returns have been falling for decades (e.g., [40, 184]) and are now at record lows; for
1462 example, the Thompson and Chilcotin tributaries of the upper Fraser River now each
1463 have adult steelhead returns of less than 200 adults [185], despite being of roughly
1464 similar size and biogeoclimatic zone to the Snake River.

1465

1466 For Washington State outside the Columbia River basin, steelhead SARs were
1467 assembled and analyzed for Puget Sound (Washington State), as well as a number of
1468 locations along the coasts of Juan de Fuca Strait, and the outer (western) WA coast as
1469 well as Oregon; see [7] for detailed methods. SAR data for the Columbia and Snake
1470 rivers were taken from [5]. We are unaware of additional steelhead SAR data for
1471 Alaskan rivers.

1472

1473 Comparison of relative survival

1474 Several of our analyses are based on comparisons with the SARs of Snake River
1475 populations as these are widely considered to be poor owing to the many dams (8) in

the migration path, and in particular the four Snake River dams. Because the various survival time series vary in length and sampling methodology, and because survival also declines episodically with time, we chose to make the survival analysis as simple and clear cut as possible.

As a result, in each year or regime period, we divided all available individual SAR estimates by the median SAR for all Snake River populations in the same time period. The normalized median SAR for the Snake River region equals one by definition and the frequency distribution of individual normalized estimates allows us to directly judge the similarity of the SAR values between regions in the selected time periods under examination.

1486

1487 Treatment of SAR data

SAR data for salmon are log-normally distributed [186]; i.e., a time series of SAR data, S_t , will have the form $S_t = e^{\mu + \sigma Z_t}$, where μ and σ are respectively the mean and standard deviation of $\log_e(S)$, and Z_t is the standard normal variable $Z \sim N(\mu, \sigma)$. This is important because the log-normal distribution is skewed, exhibiting occasional rare high survivals which increases the expected value above the mean. As a result, the expected value of a log-normally distributed SAR time series is neither the simple mean $\bar{S} = \frac{1}{n} \sum_{t=1}^n S_t$ nor μ , but rather $E(S_t) = e^{\mu + \sigma^2/2}$ (in fact, it is the median value of the log-normal distribution that is related to μ , as $S_{median} = e^{\mu}$). Calculating the average of the untransformed survival data, although often reported, does not have a simple statistical interpretation.

1498

1499 When comparing survival time series between regions, some important but subtle

1500 differences should therefore be kept in mind. We have opted to use the median

1501 (equivalent to the “geometric mean” if the data is truly log-normally distributed,

1502 $\bar{S}_{Geo} = Exp\left[\frac{1}{n} \sum_{t=1}^n \log_e(S_t)\right]$, used in some literature), as well as the simple average $\bar{S}$ of

1503 the untransformed SAR data in a number of key comparisons. The simple average is

1504 what a number of prior studies report, and therefore what most policy makers and

1505 fisheries managers are likely comfortable interpreting. For example, the NWPPC has

1506 set a rebuilding target of 2%-6% for SARs and deemed 1% SARs (roughly the current

1507 average) to be inadequate, but did not define how SAR values should be calculated.

1508

1509 However, when the distribution of SARs are compared between two regions i, j then if

1510 the medians are found to be the similar, the implication is then that $\mu_i = \mu_j$ and that the

1511 simple means of the log-transformed data are also equal; this does not, however, imply

1512 that the expected values $E(S_t) = e^{\mu + \sigma^2/2}$ are equal because this value also depends on the

1513 variance of the time series. For these reasons, we use both measures of central

1514 tendency

1515
$$\bar{S} = \frac{1}{n} \sum_{t=1}^n S_t,$$

$$S_{median} = \bar{S}_{Geo} = e^{\mu}$$

1516 in our analysis, and not the expected mean values of the log-normal distribution

1517 $E(S_t) = e^{\mu + \sigma^2/2}$, owing to the more complex definition and lack of easy interpretation,

1518 which the (simple) mean and the median readily impart.

1519 The precision of survival estimates

1520

1521 The standard error on a binomial proportion reported by the CSS and PSC, survival, is

1522 $SE(S) = \sqrt{S(1-S)/N}$. The precision of a survival estimate, $\phi(S)$, degrades as survival

1523 decreases, because

1524

1525

$$1526 \quad \Phi(S) = \frac{SE(S)}{S} = \sqrt{\frac{1-S}{S \cdot N}}$$

1527 In the limit as survival approaches either 1 or zero,

1528

1529

$$1530 \quad \begin{aligned} &\lim_{S \rightarrow 1} \Phi(S) = 0 \\ &\text{and} \\ &\lim_{S \rightarrow 0} \Phi(S) = \infty \end{aligned}$$

1531 The relative uncertainty in a survival estimate with a given sample size increases

1532 without bound as survival decreases towards zero. With survival values now at 1% or

1533 less, the relative precision of a survival estimate now relative to several decades ago

1534 when survival was in the 5-6% range is

1535

$$\frac{\Phi(S_1 = 0.01)}{\Phi(S_2 = 0.06)} = \sqrt{\frac{(1 - S_1) / (S_1 N)}{(1 - S_2) / (S_2 N)}} = \sqrt{\frac{S_2 (1 - S_1)}{S_1 (1 - S_2)}} \approx \sqrt{\frac{S_2}{S_1}}.$$

1536

1537

1538 In this numeric example, where survival falls from 6% at the start of the record to 1% at
1539 the end, the uncertainty relative to the point estimate increases almost 2.5-fold ($\sqrt{6}$).
1540 (Taking into account that both the number of outgoing smolts and the number of
1541 returning adults is not known without error, as is implicitly assumed in using the
1542 binomial probability distribution, the actual uncertainty will be even larger when these
1543 uncertainties are taken into account).

1544

1545 It is interesting to note that should survival fall from the current ca. 1% level to 0.1%--
1546 a ten-fold further decline—it would in fact be difficult to recognize this massive decline
1547 (a fall as large as the decline from 100% to 10% or 10% to 1% survival) because of the
1548 limited precision with which survival can be measured at such low levels. Thus for
1549 both purely mathematical reasons as well as the methodological differences between
1550 tagging approaches listed in the prior section, it is likely infeasible to obtain a perfect
1551 conversion ratio between survival estimates calculated using different methodologies
1552 (PIT vs CWT).

1553

1554

1555

1556 ORCID

1557 DWW Orcid ID: 0000-0001-8851-5436

1558 ADP ORCID ID: 0000-0002-1258-8265

1559 ELR: ORCID ID: 0000-0002-2811-8399

1560 Acknowledgements

1561 We particularly thank Drs Neala Kendall (WDFW) and Gayle Brown (DFO) for
1562 providing access to steelhead and Chinook SAR data, respectively, and for many
1563 discussions clarifying the interpretation of the data. The Keogh R steelhead survival
1564 project is managed by the Province of British Columbia and is currently primarily
1565 funded by the Habitat Conservation Trust Foundation (HCTF). We have also benefited
1566 from discussions with several of our colleagues, in particular Drs. XXX for early
1567 discussions on this topic. The vast (>3000 years!) of data that this paper relies upon
1568 obviously has involved many more individuals; we collectively thank them all for the
1569 effort required to generate the data used here. All data except the Keogh R steelhead
1570 SAR data are available without restriction at ??? (or from the authors).
1571 Investigators interested in accessing Keogh SAR data should request these data
1572 from YYY.
1573 (Individuals XXX and YYY have been requested to indicate their approval to be
1574 named in the Acknowledgements, as per PLoS ONE stipulation. At time of
1575 submission, formal responses had not been received. Specific names will be
1576 incorporated during the review process).

1577

1578 **CRedit (Contributor Roles Taxonomy)**

1579 Conceptualization, DWW (Lead); Methodology, ADP & DWW; Software, ADP; Validation,
1580 ADP, ELR; Formal Analysis, ADP, DWW; Investigation, DWW; Data Curation, ADP; Writing
1581 – Original Draft Preparation, DWW; Writing – Review & Editing, DWW, ADP, ELR;
1582 Visualization, ADP; Supervision, DWW; Project Administration, DWW, ELR; Funding
1583 Acquisition, DWW, ELR .

1584 **Competing Interests**

1585 DWW is President and owner of Kintama Research Services Ltd., an environmental
1586 consultancy focused on the development of innovative applications of telemetry to improve
1587 fisheries management. ADP and ELR are employed at Kintama. All authors received a
1588 financial benefit in the course of conducting this study and their future salaries depend on their
1589 continued technical and scientific performance, which includes publication of this study.

1590 **Funding**

1591 This study was initially internally funded by Kintama Research as part of a separate research
1592 effort to assess the credibility of the critical period concept in Pacific salmon. In the course of
1593 assembling Strait of Georgia SAR data, we discovered that Chinook survival in many rivers of
1594 the Strait of Georgia region had fallen to levels well below those reported for Snake River
1595 Chinook. A proposal was developed and funding obtained from the US Dept. of Energy,
1596 Bonneville Power Administration, to cover staff time for coast-wide data collation, analysis,
1597 and writing of this paper (Contract # 75025). The funder (BPA) played no role in the design of
1598 the study nor the conclusions reached, and was not provided access to the paper prior to journal
1599 submission.

References

1. Raymond HL. Effects of Hydroelectric Development and Fisheries Enhancement on Spring and Summer Chinook Salmon and Steelhead in the Columbia River Basin. *North American Journal of Fisheries Management*. 1988;8(1):1-24.
2. Schoen ER, Wipfli MS, Trammell EJ, Rinella DJ, Floyd AL, Grunblatt J, et al. Future of Pacific Salmon in the Face of Environmental Change: Lessons from One of the World's Remaining Productive Salmon Regions. *Fisheries*. 2017;42(10):538-53. doi: 10.1080/03632415.2017.1374251.
3. Irvine JR, Fukuwaka M, Kaga T, Park JH, Seong KB, Kang S, et al. Pacific Salmon Status and Abundance Trends. Vancouver, B.C.: NPAFC; 2009. p. 153 pp.
4. Ruggerone GT, Irvine JR. Numbers and Biomass of Natural- and Hatchery-Origin Pink Salmon, Chum Salmon, and Sockeye Salmon in the North Pacific Ocean, 1925–2015. *Marine and Coastal Fisheries*. 2018;10(2):152-68. doi: doi:10.1002/mcf2.10023.
5. McCann J, Chockley B, Cooper E, Hsu B, Schaller H, Haeseker S, et al. Comparative Survival Study of PIT-tagged Spring/Summer/Fall Chinook, Summer Steelhead, and Sockeye. 2017 Annual Report. Portland, Oregon 2017.
6. Dorner B, Catalano MJ, Peterman RM. Spatial and temporal patterns of covariation in productivity of Chinook salmon populations of the Northeastern Pacific. *Can J Fish Aquat Sci*. 2017. doi: 10.1139/cjfas-2017-0197.
7. Kendall NW, Marston GW, Klungle MM. Declining patterns of Pacific Northwest steelhead trout (*Oncorhynchus mykiss*) adult abundance and smolt survival in the ocean. *Can J Fish Aquat Sci*. 2017;74:1275–90. doi: 10.1139/cjfas-2016-0486.
8. Zimmerman MS, Irvine JR, O'Neill M, Anderson JH, Greene CM, Weinheimer J, et al. Spatial and Temporal Patterns in Smolt Survival of Wild and Hatchery Coho Salmon in the Salish Sea. *Marine and Coastal Fisheries*. 2015;7(1):116-34. doi: 10.1080/19425120.2015.1012246.
9. Logerwell EA, Mantua N, Lawson PW, Francis RC, Agostini VN. Tracking environmental processes in the coastal zone for understanding and predicting Oregon coho (*Oncorhynchus kisutch*) marine survival. *Fish Oceanogr*. 2003;12:554-68.
10. Cohen BI. Commission of Inquiry into the Decline of Sockeye Salmon in the Fraser River (Canada). Final Report. Ottawa, Canada: Public Works and Government Services Canada; 2012.
11. COSEWIC. Assessment and Status Report on the Sockeye Salmon *Oncorhynchus nerka*, 24 Designatable Units in the Fraser River Drainage Basin, in Canada. Ottawa: Committee on the Status of Endangered Wildlife in Canada; (In Press) 2017. p. xli + 179 pp.
12. Rand PS, Goslin M, Gross MR, Irvine JR, Augerot X, McHugh PA, et al. Global Assessment of Extinction Risk to Populations of Sockeye Salmon *Oncorhynchus nerka*. *PLoS ONE*. 2012;7(4):e34065. doi: 10.1371/journal.pone.0034065.
13. Peterman Randall M, Dorner B. A widespread decrease in productivity of sockeye salmon (*Oncorhynchus nerka*) populations in western North America. *Can J Fish Aquat Sci*. 2012;1255-60. doi: 10.1139/f2012-063.
14. Nehlsen W, Williams JE, Lichatowich JA. Pacific Salmon at the Crossroads: Stocks at Risk from California, Oregon, Idaho, and Washington. *Fisheries*. 1991;16(2):4-21.
15. Allendorf FW, Bayles D, Bottom DL, Currens KP, Frissell CA, Hankin D, et al. Prioritizing Pacific Salmon Stocks for Conservation. *Conservation Biology*. 1997;11-1:140-52.

- 1645 16. Mantua NJ, Hare SJ, Zhang Y, Wallace JM, Francis RC. A Pacific interdecadal climate
1646 oscillation with impacts on salmon production. Bulletin American Meteorological Society.
1647 1997;78, No. 6, June:1069-79.
- 1648 17. Hare SR, Mantua NJ, Francis RC. Inverse Production Regimes: Alaska and West Coast
1649 Pacific Salmon. Fisheries. 1999;24:6-14.
- 1650 18. Mantua NJ, Hare SR. The Pacific Decadal Oscillation. J Oceanog. 2002;58(1):35-44. doi:
1651 10.1023/A:1015820616384.
- 1652 19. Ryding KE, Skalski JR. Multivariate regression relationships between ocean conditions
1653 and early marine survival of coho salmon (*Oncorhynchus kisutch*). Can J Fish Aquat Sci.
1654 1999;56:2374-84.
- 1655 20. Schindler D, Krueger C, Bisson P, Bradford M, Clark B, Conitz J, et al., editors. Arctic-
1656 Yukon-Kuskokwim Chinook Salmon Research Action Plan: Evidence of Decline of Chinook
1657 Salmon Populations and Recommendations for Future Research 2013: Prepared for the AYK
1658 Sustainable Salmon Initiative (Anchorage, AK). v + 70 pp.
- 1659 21. ADF&G Chinook Salmon Research Team. Chinook Salmon Stock Assessment and
1660 Research Plan, 2013. Anchorage, AK: Alaska Department of Fish & Game; 2013. Available
1661 from:
1662 http://www.adfg.alaska.gov/static/home/news/hottopics/pdfs/chinook_research_plan.pdf.
- 1663 22. Cunningham CJ, Westley PAH, Adkison MD. Signals of large scale climate drivers,
1664 hatchery enhancement, and marine factors in Yukon River Chinook salmon survival revealed
1665 with a Bayesian life history model. Global Change Biology. 2018;0(0). doi:
1666 doi:10.1111/gcb.14315.
- 1667 23. Bradford MJ, von Finster A, Milligan PA. Freshwater life history, habitat, and the
1668 production of Chinook Salmon from the upper Yukon basin. American Fisheries Society
1669 Symposium. 702009. p. 1-20.
- 1670 24. Hay D, McCarter PB. Status of the eulachon *Thaleichthys pacificus* in Canada. CSAS Res
1671 Doc. 2000;2000/145:92.
- 1672 25. Murauskas JG, Orlov AM, Siwicke KA. Relationships between the Abundance of Pacific
1673 Lamprey in the Columbia River and Their Common Hosts in the Marine Environment.
1674 Transactions of the American Fisheries Society. 2013;142(1):143-55.
- 1675 26. Chaput G. Overview of the status of Atlantic salmon (*Salmo salar*) in the North Atlantic
1676 and trends in marine mortality. ICES J Mar Sci. 2012;69(9):1538-48.
- 1677 27. Castonguay M, Hodson PV, Moriarty C, Drinkwater KF, Jessop BM. Is there a role of
1678 ocean environment in American and European eel decline? Fish Oceanogr. 1994;3(3):197-203.
1679 doi: 10.1111/j.1365-2419.1994.tb00097.x.
- 1680 28. Dekker W, Casselman JM. The 2003 Québec Declaration of Concern About Eel
1681 Declines—11 Years Later: Are Eels Climbing Back up the Slippery Slope? Fisheries.
1682 2014;39(12):613-4. doi: 10.1080/03632415.2014.979342.
- 1683 29. Raymond HL. Effects of Dams and Impoundments on Migrations of Juvenile Chinook
1684 Salmon and Steelhead from the Snake River, 1966 to 1975. Transactions of the American
1685 Fisheries Society. 1979;108(6):505-29.
- 1686 30. Raymond HL. Migration Rates of Yearling Chinook Salmon in Relation to Flows and
1687 Impoundments in the Columbia and Snake Rivers. Transactions of the American Fisheries
1688 Society. 1968;97(4):356-9. doi: 10.1577/1548-8659(1968)97[356:MROYCS]2.0.CO;2.
- 1689 31. Marmorek DR, C.N. Peters and I. Parnell (eds.). Path Final Report For Fiscal Year
1690 1998. (Available from Bonneville Power Administration, Portland, Oregon
1691 http://www.efw.bpa.gov/Environment/PATH/reports/ISRP1999CD/PATH%20Reports/WOE_Report/).
1692 1998:263 pp.

- 1693 32. Mantua NJ. Shifting patterns in Pacific climate, West Coast salmon survival rates, and
1694 increased volatility in ecosystem services. *Proceedings of the National Academy of Sciences*.
1695 2015;112(35):10823-4. doi: 10.1073/pnas.1513511112.
- 1696 33. Sharma R, VÉlez-Espino LA, Wertheimer AC, Mantua N, Francis RC. Relating spatial
1697 and temporal scales of climate and ocean variability to survival of Pacific Northwest Chinook
1698 salmon (*Oncorhynchus tshawytscha*). *Fish Oceanogr*. 2013;22(1):14-31. doi:
1699 10.1111/fog.12001.
- 1700 34. Stachura MM, Mantua NJ, Scheuerell MD. Oceanographic influences on patterns in
1701 North Pacific salmon abundance. *Can J Fish Aquat Sci*. 2013;71(2):226-35. doi: 10.1139/cjfas-
1702 2013-0367.
- 1703 35. Finney BP, Gregory-Eaves I, Sweetman J, Douglas MSV, Smol JP. Impacts of Climatic
1704 Change and Fishing on Pacific Salmon Abundance Over the Past 300 Years. *Science*.
1705 2000;290(5492):795-9. doi: 10.1126/science.290.5492.795.
- 1706 36. Rogers LA, Schindler DE, Lisi PJ, Holtgrieve GW, Leavitt PR, Bunting L, et al. Centennial-
1707 scale fluctuations and regional complexity characterize Pacific salmon population dynamics
1708 over the past five centuries. *Proceedings of the National Academy of Sciences*. 2013. doi:
1709 10.1073/pnas.1212858110.
- 1710 37. Finney BP, Gregory-Eaves I, Douglas M, Smol J. Fisheries productivity in the
1711 northeastern Pacific Ocean over the past 2,200 years. *Nature*. 2002;416:729-33.
- 1712 38. Beamish RJ. Climate and exceptional fish production off the west coast of North
1713 America. *Can J Fish Aquat Sci*. 1993;50:2270-91.
- 1714 39. Beamish RJ, Bouillon DR. Pacific salmon production trends in relation to climate. *Can J*
1715 *Fish Aquat Sci*. 1993;50:1002-16.
- 1716 40. Welch DW, Ward BR, Smith BD, Eveson JP. Temporal and spatial responses of British
1717 Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts. *Fish*
1718 *Oceanogr*. 2000;9(1):17-32. doi: 10.1046/j.1365-2419.2000.00119.x
- 1719 41. Welch DW, Rechisky EL, Melnychuk MC, Porter AD, Walters CJ, Clements S, et al.
1720 Survival of Migrating Salmon Smolts in Large Rivers With and Without Dams. *PLoS Biology*.
1721 2008;6(10):2101-8. doi: 10.1371/journal.pbio.0060265.
- 1722 42. Michel CJ, Ammann AJ, Lindley ST, Sandstrom PT, Chapman ED, Thomas MJ, et al.
1723 Chinook salmon outmigration survival in wet and dry years in California's Sacramento River.
1724 *Can J Fish Aquat Sci*. 2015;72(11):1749-59. doi: 10.1139/cjfas-2014-0528.
- 1725 43. Buchanan RA, Brandes PL, Skalski JR. Survival of Juvenile Fall-Run Chinook Salmon
1726 through the San Joaquin River Delta, California, 2010–2015. *North American Journal of*
1727 *Fisheries Management*. 2018 (In Press);0(0):1-17. doi: 10.1002/nafm.10063.
- 1728 44. Lowe CG, Wetherbee BM, Meyer CG. Using acoustic telemetry monitoring techniques
1729 to quantify movement patterns and site fidelity of sharks and giant trevally around French
1730 Frigate Shoals and Midway Atoll. *Atoll Research Bulletin*. 2006;543:281–303.
- 1731 45. Meyer CG, Clark TB, Papastamatiou YP, Whitney NM, Holland KN. Long-term
1732 movement patterns of tiger sharks *Galeocerdo cuvier* in Hawaii. *Marine Ecology Progress*
1733 *Series*. 2009;381:223-35.
- 1734 46. Baur DC, Irvin WR, editors. *Endangered Species Act: Law, Policy, and Perspectives*. 2nd
1735 Edition (November 10, 2009) ed: American Bar Association; 2010.
- 1736 47. Vanderzwaag DL, Hutchings JA. Canada's marine species at risk: science and law at the
1737 helm, but a sea of uncertainties. *Ocean Development & International Law*. 2005;36(3):219-59.
1738 doi: 10.1080/00908320591004333.

- 1739 48. Wasser SK, Lundin JI, Ayres K, Seely E, Giles D, Balcomb K, et al. Population growth is
1740 limited by nutritional impacts on pregnancy success in endangered Southern Resident killer
1741 whales (*Orcinus orca*). PLOS ONE. 2017;12(6):e0179824. doi: 10.1371/journal.pone.0179824.
- 1742 49. CTC (Chinook Technical Committee). 2014 Exploitation Rate Analysis and Model
1743 Calibration. Volume One. TCCHINOOK (15)-1 V. 1. Vancouver, B.C., Canada: Pacific Salmon
1744 Commission, Joint Chinook Technical Committee; 2014.
- 1745 50. Bilton HT, Alderdice DF, Schnute JT. Influence of time and size at release of juvenile
1746 coho salmon (*Oncorhynchus kisutch*) on returns at maturity. Can J Fish Aquat Sci.
1747 1982;39(3):426-47.
- 1748 51. Irvine JR, O'Neill M, Godbout L, Schnute J. Effects of smolt release timing and size on
1749 the survival of hatchery-origin coho salmon in the Strait of Georgia. Prog Oceanogr.
1750 2013;115(0):111-8. doi: <http://dx.doi.org/10.1016/j.pocean.2013.05.014>.
- 1751 52. Hostetter NJ, Evans AF, Cramer BM, Collis K, Lyons DE, Roby DD. Quantifying Avian
1752 Predation on Fish Populations: Integrating Predator-Specific Deposition Probabilities in Tag
1753 Recovery Studies. Transactions of the American Fisheries Society. 2015;144(2):410-22. doi:
1754 10.1080/00028487.2014.988882.
- 1755 53. Ward DL, Petersen JH, Loch JJ. Index of Predation on Juvenile Salmonids by Northern
1756 Squawfish in the Lower and Middle Columbia River and in the Lower Snake River. Transactions
1757 of the American Fisheries Society. 1995;124(3):321-34.
- 1758 54. Gosselin JL, Anderson JJ. Combining Migration History, River Conditions, and Fish
1759 Condition to Examine Cross-Life-Stage Effects on Marine Survival in Chinook Salmon.
1760 Transactions of the American Fisheries Society. 2017;146(3):408-21. doi:
1761 10.1080/00028487.2017.1281166.
- 1762 55. Rechisky EL, Welch DW, Porter AD, Hess JE, Narum SR. Testing for delayed mortality
1763 effects in the early marine life history of Columbia River yearling Chinook salmon. Mar Ecol
1764 Prog Series. 2014;496:159–80 doi: 10.3354/meps10692.
- 1765 56. Rechisky EL, Welch DW, Porter AD, Jacobs-Scott MC, Winchell PM. Influence of
1766 multiple dam passage on survival of juvenile Chinook salmon in the Columbia River estuary
1767 and coastal ocean. Proc Nat Acad Sci USA. 2013. Epub Published online before print April 1,
1768 2013. doi: 10.1073/pnas.1219910110.
- 1769 57. Sandford BP, Zabel RW, Gilbreath LG, Smith SG. Exploring Latent Mortality of Juvenile
1770 Salmonids Related to Migration through the Columbia River Hydropower System. Transactions
1771 of the American Fisheries Society. 2012;141(2):343-52. doi: 10.1080/00028487.2012.664601.
- 1772 58. Schaller H, Petrosky CE. Assessing Hydrosystem Influence on Delayed Mortality of
1773 Snake River Stream-Type Chinook Salmon. North American Journal of Fisheries Management.
1774 2007;27:810–24.
- 1775 59. Muir W, Marsh D, Sanford B, Smith S, Williams J. Post-Hydropower System Delayed
1776 Mortality of Transported Snake River Stream-Type Chinook Salmon: Unraveling the Mystery.
1777 Transactions of the American Fisheries Society. 2006;135:1523–34.
- 1778 60. Budy P, Thiede GP, Bouwes N, Petrosky CE, Schaller H. Evidence linking delayed
1779 mortality of Snake River salmon to their earlier hydrosystem experience. North American
1780 Journal of Fisheries Management. 2002;22:35-51.
- 1781 61. Zabel RW, Williams JG. Comments on "Contrasting patterns of productivity and
1782 survival rates for stream-type chinook salmon (*Oncorhynchus tshawytscha*) populations of the
1783 Snake and Columbia rivers" by Schaller et al (1999). Can J Fish Aquat Sci. 2000;57:1739-41.
- 1784 62. Schaller HA, Petrosky CE, Langness OP. Contrasting patterns of productivity and
1785 survival rates for stream-type chinook salmon (*Oncorhynchus tshawytscha*) populations of the
1786 Snake and Columbia rivers. Can J Fish Aquat Sci. 1999;56(6):1031-45.

63. Schaller HA, Petrosky CE, Tinus ES. Evaluating river management during seaward migration to recover Columbia River stream-type Chinook salmon considering the variation in marine conditions. *Can J Fish Aquat Sci.* 2013;71(2):259-71. doi: 10.1139/cjfas-2013-0226.
64. Independent Scientific Advisory Board (ISAB). Latent Mortality Report. Review of Hypotheses and Causative Factors Contributing to Latent Mortality and their Likely Relevance to the “Below Bonneville” Component of the COMPASS Model. Portland, Oregon. : Northwest Power and Conservation Council., 2007 Document No. ISAB 2007-1.
65. Festinger L. A Theory of Cognitive Dissonance. Stanford, Calif.: Stanford University Press; 1957.
66. Janis IL. Victims of groupthink: a psychological study of foreign-policy decisions and fiascoes. Boston: Houghton Mifflin, 349 pp.; 1972.
67. McKinnell SM, Wood CC, Rutherford DT, Hyatt KD, Welch DW. The demise of Owikeno Lake sockeye salmon. *North American Journal of Fisheries Management.* 2001;21:774-91. doi: 10.1577/1548-8675(2001)021<0774:TDOOLS>2.0.CO;2.
68. Anonymous. Recommendations for a Recovery Plan for the Rivers Inlet and Smith Inlet Sockeye Salmon (Final Draft, V. 8). Fisheries & Oceans Canada. Report prepared by The Technical Coordinating Committee, Rivers and Smith Inlet Recovery Plan Working Group; 2001. p. 112 pp.
69. Rivers and Smith Inlets Salmon Ecosystem Planning Society. Rivers and Smith Inlets Salmon Ecosystem Recovery Plan. Pacific Salmon Endowment Fund Society; 2003.
70. Holtby BC. Recommendations for a Recovery Plan for the Rivers Inlet and Smith Inlet Sockeye Salmon. Prepared by: Technical Coordinating Committee Rivers Inlet and Smith Inlet Recovery Plan Working Group; 2000.
71. English KK, Glova GJ, Blakley AC. An Upstream Battle: Declines in 10 Pacific Salmon Stocks and Solutions for Their Survival. Vancouver: David Suzuki Foundation, 2008.
72. Rutherford D, Wood C. Assessment of Rivers and Smith Inlet Sockeye Salmon, with Commentary on Small Sockeye Salmon Stocks in Statistical Area 8. Canadian Stock Assessment Secretariat research document, 1480-4883 ; 2000/162; 2000. p. 57 p.
73. Kareiva PM, Marvier M, McClure MM. Recovery and management options for spring/summer Chinook salmon in the Columbia River. *Science.* 2000;290:977-9. doi: 10.1126/science.290.5493.977.
74. Marmorek D, Peters C. Finding a PATH toward scientific collaboration: insights from the Columbia River Basin. *Conservation Ecology.* 2001;5(2).
75. McMichael GA, Eppard MB, Carlson TJ, Carter JA, Ebberts BD, Brown RS, et al. The Juvenile Salmon Acoustic Telemetry System: A New Tool. *Fisheries.* 2010;35(1):9-22.
76. Rechisky ER, Welch DW, Porter AD, Jacobs MC, Ladouceur A. Experimental measurement of hydrosystem-induced mortality in juvenile Snake River spring Chinook salmon using a large-scale acoustic array. *Can J Fish Aquat Sci.* 2009;66:1019–24. doi: 10.1139/F09-078.
77. McMichael GA, Hanson AC, Harnish RA, Trott DM. Juvenile salmonid migratory behavior at the mouth of the Columbia River and within the plume. *Animal Biotelemetry.* 2013;1(1). doi: 10.1186/2050-3385-1-14.
78. McMichael GA, Harnish RA, Skalski JR, Deters KA, Ham KD, Townsend RL, et al. Migratory behavior and survival of juvenile salmonids in the lower Columbia River, estuary, and plume in 2010. Richland, Washington.: Pacific Northwest National Laboratory; 2011.
79. Rechisky EL, Welch DW, Porter AD, Jacobs-Scott M, Winchell PM, McKern JL. Estuarine and early-marine survival of transported and in-river migrant Snake River spring Chinook salmon smolts. *Nature Scientific Reports.* 2012;2(448). doi: 10.1038/srep00448.

- 1835 80. Brosnan IG, Welch DW, Rechisky EL, Porter AD. Evaluating the influence of
1836 environmental factors on yearling Chinook salmon survival in the Columbia River plume (USA).
1837 Mar Ecol Prog Ser. 2014;496:181–96. doi: 10.3354/meps10550.
- 1838 81. Brosnan IG, Welch DW, Scott MJ. Survival Rates of Out-Migrating Yearling Chinook
1839 Salmon in the Lower Columbia River and Plume after Exposure to Gas-Supersaturated Water.
1840 Journal of Aquatic Animal Health. 2016;28(4):240-51. doi: 10.1080/08997659.2016.1227398.
- 1841 82. Johnson GE, Ploskey GR, Sather NK, Teel DJ. Residence times of juvenile salmon and
1842 steelhead in off-channel tidal freshwater habitats, Columbia River, USA. Can J Fish Aquat Sci.
1843 2015;1-13. doi: 10.1139/cjfas-2014-0085.
- 1844 83. Harnish RA, Johnson GE, McMichael GA, Hughes MS, Ebberts BD. Effect of Migration
1845 Pathway on Travel Time and Survival of Acoustic-Tagged Juvenile Salmonids in the Columbia
1846 River Estuary. Transactions of the American Fisheries Society. 2012;141(2):507-19. doi:
1847 10.1080/00028487.2012.670576.
- 1848 84. Roegner GC, McNatt R, Teel DJ, Bottom DL. Distribution, Size, and Origin of Juvenile
1849 Chinook Salmon in Shallow-Water Habitats of the Lower Columbia River and Estuary, 2002–
1850 2007. Marine and Coastal Fisheries. 2012;4(1):450-72. doi: 10.1080/19425120.2012.675982.
- 1851 85. McNatt RA, Bottom DL, Hinton SA. Residency and Movement of Juvenile Chinook
1852 Salmon at Multiple Spatial Scales in a Tidal Marsh of the Columbia River Estuary. Transactions
1853 of the American Fisheries Society. 2016;145(4):774-85. doi: 10.1080/00028487.2016.1172509.
- 1854 86. Sather NK, Johnson GE, Teel DJ, Storch AJ, Skalski JR, Cullinan VI. Shallow Tidal
1855 Freshwater Habitats of the Columbia River: Spatial and Temporal Variability of Fish
1856 Communities and Density, Size, and Genetic Stock Composition of Juvenile Chinook Salmon.
1857 Transactions of the American Fisheries Society. 2016;145(4):734-53. doi:
1858 10.1080/00028487.2016.1150878.
- 1859 87. Price MHH, English KK, Rosenberger AG, MacDuffee M, Reynolds JD. Canada's Wild
1860 Salmon Policy: an assessment of conservation progress in British Columbia. Can J Fish Aquat
1861 Sci. 2017;74(10):1507-18. doi: 10.1139/cjfas-2017-0127.
- 1862 88. Lackey RT. Science, scientists, and policy advocacy. Conserv Biol. 2007;21(1):12-7.
- 1863 89. Lackey RT. Is Science Biased Toward Natural? Northwest Science. 2009;83(3):277-9.
- 1864 90. Kareiva P, Marvier M. What Is Conservation Science? BioScience. 2012;62(11):962-9.
1865 doi: 10.1525/bio.2012.62.11.5.
- 1866 91. Lichatowich J. Salmon without Rivers. A History of the Pacific Salmon Crisis.
1867 Washington, D.C.: Island Press; 1999. 317 p.
- 1868 92. Lewis B, Grant WS, Brenner RE, Hamazaki T. Changes in Size and Age of Chinook
1869 Salmon *Oncorhynchus tshawytscha* Returning to Alaska. PLoS ONE. 2015;10(6):e0130184. doi:
1870 10.1371/journal.pone.0130184.
- 1871 93. Orsi J. The Alaska Chinook Salmon Production Enigma... What's Going On?
1872 ONCORHYNCHUS [Internet]. 2013; XXXIII(2):[1-5 pp.]. Available from: [http://www.afs-](http://www.afs-alaska.org/wp-content/uploads/Onco332.pdf)
1873 [alaska.org/wp-content/uploads/Onco332.pdf](http://www.afs-alaska.org/wp-content/uploads/Onco332.pdf).
- 1874 94. Shelton AO, Satterthwaite WH, Ward EJ, Feist BE, Burke B. Using hierarchical models
1875 to estimate stock-specific and seasonal variation in ocean distribution, survivorship, and
1876 aggregate abundance of fall run Chinook. Can J Fish Aquat Sci. 2018 (In Press). doi:
1877 10.1139/cjfas-2017-0204.
- 1878 95. Weitkamp LA. Marine Distributions of Chinook Salmon from the West Coast of North
1879 America Determined by Coded Wire Tag Recoveries. Transactions of the American Fisheries
1880 Society. 2009;139(1):147-70. doi: doi:10.1577/T08-225.1.

- 1881 96. Chamberlin JW, Quinn TP. Effects of natal origin on localized distributions of Chinook
1882 Salmon, *Oncorhynchus tshawytscha*, in the marine waters of Puget Sound, Washington. Fish
1883 Res. 2014;153:113-22.
- 1884 97. O'Neill SM, West JE. Marine Distribution, Life History Traits, and the Accumulation of
1885 Polychlorinated Biphenyls in Chinook Salmon from Puget Sound, Washington. Transactions of
1886 the American Fisheries Society. 2009;138(3):616-32. doi: doi:10.1577/T08-003.1.
- 1887 98. Kagley AN, Smith JM, Fresh KL, Frick KE, Quinn TP. Residency, partial migration, and
1888 late egress of subadult Chinook salmon (*Oncorhynchus tshawytscha*) and coho salmon (*O.*
1889 *kisutch*) in Puget Sound, Washington. Fishery Bulletin. 2017;115(4):544-56.
- 1890 99. Quinn TP, Chamberlain J, Banks E. Experimental evidence of population-specific
1891 marine spatial distributions of Chinook salmon, *Oncorhynchus tshawytscha*. . Environmental
1892 Biology of Fishes. 2011;92(3):313-22. doi: 10.1007/s10641-011-9841-z.
- 1893 100. Arostegui MC, Smith JM, Kagley AN, Spilsbury-Pucci D, Fresh KL, Quinn TP. Spatially
1894 Clustered Movement Patterns and Segregation of Subadult Chinook Salmon within the Salish
1895 Sea. Marine and Coastal Fisheries. 2017;9(1):1-12. doi: 10.1080/19425120.2016.1249580.
- 1896 101. Anonymous. Insights from Genetic Analyses into Stock Specific Early Ocean Migration
1897 Behavior of Juvenile Columbia River Steelhead. Fishery Bulletin. (In Review).
- 1898 102. Hartt AC, Dell MB. Early Oceanic Migrations and Growth of Juvenile Pacific Salmon and
1899 Steelhead Trout. Int North Pacific Fish Comm. 1986;46:1-105.
- 1900 103. Drenner SM, Hinch SG, Furey NB, Clark TD, Li S, Ming T, et al. Transcriptome patterns
1901 and blood physiology associated with homing success of sockeye salmon during their final
1902 stage of marine migration. . Can J Fish Aquat Sci. 2017. doi: 10.1139/cjfas-2017-0391.
- 1903 104. Furey NB, Vincent SP, Hinch SG, Welch DW. Variability in Migration Routes Influences
1904 Early Marine Survival of Juvenile Salmon Smolts. PLoS ONE. 2015;10(10):e0139269. doi:
1905 10.1371/journal.pone.0139269.
- 1906 105. Ruff CP, Anderson JH, Kemp IM, Kendall NW, McHugh PA, Velez-Espino A, et al. Salish
1907 Sea Chinook salmon exhibit weaker coherence in early marine survival trends than coastal
1908 populations. Fish Oceanogr. 2017;26(6):625-37. doi: 10.1111/fog.12222.
- 1909 106. Welch DW, Melnychuk MC, Payne JC, Rechisky EL, Porter AD, Jackson G, et al. *In situ*
1910 Measurement of Coastal Ocean Movements and Survival of Juvenile Pacific Salmon. Proc Nat
1911 Acad Sci USA. 2011;108(21):8708-13 Epub May 10, 2011. doi: 10.1073/pnas.1014044108.
- 1912 107. Neville CM, Beamish RJ, Chittenden CM. Poor Survival of Acoustically-Tagged Juvenile
1913 Chinook Salmon in the Strait of Georgia, British Columbia, Canada. Transactions of the
1914 American Fisheries Society. 2015;144(1):25-33. doi: 10.1080/00028487.2014.954053.
- 1915 108. Fisher JP, Weitkamp LA, Teel DJ, Hinton SA, Orsi JA, Farley EV, et al. Early Ocean
1916 Dispersal Patterns of Columbia River Chinook and Coho Salmon. Transactions of the American
1917 Fisheries Society. 2014;143(1):252-72. doi: 10.1080/00028487.2013.847862.
- 1918 109. Larson WA, Utter FM, Myers KW, Templin WD, Seeb JE, Guthrie Iii CM, et al. Single-
1919 nucleotide polymorphisms reveal distribution and migration of Chinook salmon
1920 (*Oncorhynchus tshawytscha*) in the Bering Sea and North Pacific Ocean. Can J Fish Aquat Sci.
1921 2013;70(1):128-41. doi: 10.1139/cjfas-2012-0233.
- 1922 110. Chasco BE, Kaplan IC, Thomas AC, Acevedo-Gutiérrez A, Noren DP, Ford MJ, et al.
1923 Competing tradeoffs between increasing marine mammal predation and fisheries harvest of
1924 Chinook salmon. Scientific Reports. 2017;7(1):15439. doi: 10.1038/s41598-017-14984-8.
- 1925 111. Chasco B, Kaplan IC, Thomas A, Acevedo-Gutiérrez A, Noren D, Ford MJ, et al.
1926 Estimates of Chinook salmon consumption in Washington State inland waters by four marine
1927 mammal predators from 1970 to 2015. Can J Fish Aquat Sci. 2017;74(8):1173-94. doi:
1928 10.1139/cjfas-2016-0203.

- 1929 112. Thomas AC, Nelson BW, Lance MM, Deagle BE, Trites AW. Harbour seals target
1930 juvenile salmon of conservation concern. Can J Fish Aquat Sci. 2016. doi: 10.1139/cjfas-2015-
1931 0558.
- 1932 113. Nelson BW, Walters CJ, Trites AW, McAllister MK. Wild Chinook salmon productivity is
1933 negatively related to seal density, and not related to hatchery releases in the Pacific
1934 Northwest. Can J Fish Aquat Sci. 2018 (In Press). doi: 10.1139/cjfas-2017-0481.
- 1935 114. Keefer ML, Stansell RJ, Tackley SC, Nagy WT, Gibbons KM, Peery CA, et al. Use of
1936 Radiotelemetry and Direct Observations to Evaluate Sea Lion Predation on Adult Pacific
1937 Salmonids at Bonneville Dam. Transactions of the American Fisheries Society.
1938 2012;141(5):1236-51. doi: 10.1080/00028487.2012.688918.
- 1939 115. Wright BE, Riemer SD, Brown RF, Ougzin AM, Bucklin KA. Assessment of harbor seal
1940 predation on adult salmonids in a Pacific Northwest estuary. Ecol Applic. 2007;17(2):338-51.
1941 doi: 10.1890/05-1941.
- 1942 116. Ohlberger J, Ward EJ, Schindler DE, Lewis B. Demographic changes in Chinook salmon
1943 across the Northeast Pacific Ocean. Fish and Fisheries. 2018;n/a-n/a. doi: 10.1111/faf.12272.
- 1944 117. Barneche DR, Robertson DR, White CR, Marshall DJ. Fish reproductive-energy output
1945 increases disproportionately with body size. Science. 2018;360(6389):642-5. doi:
1946 10.1126/science.aao6868.
- 1947 118. Bigler BS, Welch DW, Helle JH. A review of size trends among north Pacific salmon
1948 (*Oncorhynchus spp.*). Can J Fish Aquat Sci. 1996;53:455-65. doi: 10.1139/f95-181.
- 1949 119. Batten S, Ruggerone G, Ortiz I. Pink salmon induce a trophic cascade in plankton
1950 populations in the southern Bering Sea and around the Aleutian Islands. Fish Oceanogr.
1951 2018;28(1):1-12. doi: 10.1111/fog.12276.
- 1952 120. Springer AM, van Vliet GB, Bool N, Crowley M, Fullagar P, Lea M-A, et al.
1953 Transhemispheric ecosystem disservices of pink salmon in a Pacific Ocean macrosystem.
1954 Proceedings of the National Academy of Sciences. 2018. doi: 10.1073/pnas.1720577115.
- 1955 121. Springer AM, van Vliet GB. Climate change, pink salmon, and the nexus between
1956 bottom-up and top-down forcing in the subarctic Pacific Ocean and Bering Sea. Proceedings of
1957 the National Academy of Sciences. 2014;111(18):E1880-E8. doi: 10.1073/pnas.1319089111.
- 1958 122. Ruggerone GT, Connors BM. Productivity and life history of sockeye salmon in relation
1959 to competition with pink and sockeye salmon in the North Pacific Ocean. Can J Fish Aquat Sci.
1960 2015;72(6):818-33. doi: 10.1139/cjfas-2014-0134.
- 1961 123. Riddell BE, Brodeur RD, Bugaev AV, Moran P, Murphy JM, Orsi JA, et al. Chapter 5:
1962 Ocean Ecology of Chinook Salmon. In: Beamish RJ, editor. The Ocean Ecology Of Pacific Salmon
1963 And Trout. Bethesda. MD.: American Fisheries Society; 2018. p. 555-696.
- 1964 124. Nicholas JW, Hankin DG. Chinook salmon populations in Oregon coastal river basins:
1965 descriptions of life histories and assessment of recent trends in run strengths. In: Oregon
1966 Dept. Fish Wildlife RDS, Information Report No. 88-1, Portland., editor. 1988.
- 1967 125. Welch DW, Boehlert GW, Ward BR. POST—the Pacific Ocean Salmon Tracking Project.
1968 Oceanologica Acta. 2002;25(5):243-53.
- 1969 126. Batten S, Hyrenbach D, Sydeman W, Morgan K, Henry M, Yen P, et al. Characterising
1970 Meso-Marine Ecosystems of the North Pacific. Deep Sea Research. 2006;53 (II):270–90.
- 1971 127. Piatt JF, Arimitsu ML, Sydeman WJ, Thompson SA, Renner H, Zador S, et al.
1972 Biogeography of pelagic food webs in the North Pacific. Fish Oceanogr. 2018;1-18. doi:
1973 10.1111/fog.12258.
- 1974 128. Brodeur RD, Ware DM. Long-term variability in zooplankton biomass in the subarctic
1975 Pacific Ocean. Fish Oceanogr. 1992;1(1):32-8.

- 1976 129. Brodeur RD, Ware DM. Interdecadal variability in distribution and catch rates of
1977 epipelagic nekton in the Northeast Pacific Ocean. *Can Spec Publ Fish Aquat Sci.* 1995;121:329-
1978 56.
- 1979 130. Bisbal G, McConnaha W. Consideration of ocean conditions in the management of
1980 salmon. *Can J Fish Aquat Sci.* 1998;55(9):2178-86. doi: 10.1139/f98-108.
- 1981 131. Lackey RT. Saving Wild Salmon: A 165 Year Policy Conundrum. Dubach Workshop:
1982 Science and Scientists in the Contemporary Policy Process. Portland, Oregon.: Oregon State
1983 University; 2014. p. 1-25.
- 1984 132. Higgins JA, Kurbatov AV, Spaulding NE, Brook E, Introne DS, Chimiak LM, et al.
1985 Atmospheric composition 1 million years ago from blue ice in the Allan Hills, Antarctica.
1986 *Proceedings of the National Academy of Sciences.* 2015;112(22):6887-91. doi:
1987 10.1073/pnas.1420232112.
- 1988 133. Lewis S, Maslin M. The Human Planet: How We Created The Anthropocene. New
1989 Haven & London: Yale University Press; 2018. p. 465 pp.
- 1990 134. Bond NA, Cronin MF, Freeland H, Mantua N. Causes and impacts of the 2014 warm
1991 anomaly in the NE Pacific. *Geophysical Research Letters.* 2015;42(9):3414-20.
- 1992 135. Richerson K, Leonard J, Holland DS. Predicting the economic impacts of the 2017 West
1993 Coast salmon troll ocean fishery closure. *Marine Policy.* 2018 (In Press).
- 1994 136. Frölicher TL, Fischer EM, Gruber N. Marine heatwaves under global warming. *Nature.*
1995 2018;560(7718):360-4. doi: 10.1038/s41586-018-0383-9.
- 1996 137. Sigmond M, Fyfe JC, Swart NC. Ice-free Arctic projections under the Paris Agreement.
1997 *Nature Climate Change.* 2018:1.
- 1998 138. Pearcy W, McKinnell S. The Ocean Ecology of Salmon in the Northeast Pacific Ocean—
1999 An Abridged History. *American Fisheries Society Symposium* 572007. p. 7-30.
- 2000 139. Brodeur RD, Myers KW, Helle JH. Research Conducted by the United States on the
2001 Early Ocean Life History of Pacific Salmon. *N Pac Anadr Fish Comm Bull.* 2003;3:89–131.
- 2002 140. Beamish RJ, Pearsall IA, Healey MC. A history of the research on the early marine life
2003 of Pacific salmon off Canada’s Pacific coast. *N Pac Anadr Fish Comm Bull.* 2003;3:1-40.
- 2004 141. Jacobson K, Peterson B, Trudel M, Ferguson J, Welch D, Baptista A, et al. The Marine
2005 Ecology of Columbia River Basin Salmonids: A Synthesis of Research 1998-2011. Portland, Or:
2006 Northwest Power and Conservation Council; 2012. p. 86 p + appendices.
- 2007 142. Helle JH. Birth of Bering-Aleutian Salmon International Survey (BASIS). *N Pac Anadr*
2008 *Fish Comm Bull.* 2009;5:vii-viii http://www.npafc.org/new/pub_bulletin5.html.
- 2009 143. SSMSWP Workgroup. Research Work Plan: Marine Survival of Puget Sound Steelhead.
2010 2014. p. 117.
- 2011 144. Riddell BE, Pearsall I. Salish Sea Marine Survival Project, Canadian Backgrounder.
2012 Vancouver, B.C.: Pacific Salmon Foundation; 2015. p. 15 pp.
- 2013 145. Riddell B, Pearsall I, Beamish RJ, Devlin B, Farrell AP, McFarlane S, et al. Strait of
2014 Georgia Chinook and Coho Proposal 2009. Available from:
2015 [https://marinesurvivalproject.com/wp-content/uploads/PSF-PROPOSAL-Final-Full-](https://marinesurvivalproject.com/wp-content/uploads/PSF-PROPOSAL-Final-Full-Documents.pdf)
2016 [Document.pdf](https://marinesurvivalproject.com/wp-content/uploads/PSF-PROPOSAL-Final-Full-Documents.pdf).
- 2017 146. Budy P, Schaller H. Evaluating tributary restoration potential for Pacific salmon
2018 recovery. *ecological applications.* 2007;17(4):1068–86.
- 2019 147. Bernhardt ES, Palmer MA, Allan J, Alexander G, Barnas K, Brooks S, et al. Synthesizing
2020 US river restoration efforts. *Science.* 2005;308:636–7. doi: 10.1126/science.1109769.
- 2021 148. NWPCC. 2016 Columbia River Basin Fish and Wildlife Program Costs Report.
2022 Northwest Power and Conservation Council; 2017. p. 23 p.

- 2023 149. Godwin SC, Dill LM, Krkošek M, Price MHH, Reynolds JD. Reduced growth in wild
2024 juvenile sockeye salmon *Oncorhynchus nerka* infected with sea lice. *J Fish Biol.* 2017;1-17. doi:
2025 10.1111/jfb.13325.
- 2026 150. Peacock SJ, Bateman AW, Krkošek M, Connors B, Rogers S, Portner L, et al. Sea -louse
2027 parasites on juvenile wild salmon in the Broughton Archipelago, British Columbia, Canada.
2028 *Ecology.* 2016.
- 2029 151. Morton A, Routledge R. Risk and precaution: Salmon farming. *Marine Policy.*
2030 2016;74:205-12. doi: <http://dx.doi.org/10.1016/j.marpol.2016.09.022>.
- 2031 152. Hemingway E. *The Sun Also Rises*: Charles Scribner & Sons; 1926. 286 pp p.
- 2032 153. Wells BK, Schroeder ID, Bograd SJ, Hazen EL, Jacox MG, Leising A, et al. State of the
2033 California Current 2016–17: Still anything but normal in the north. *CalCOFI Rep.* 2017;58:1-55.
- 2034 154. Brodeur RD, Hunsicker ME, Hann A, Miller TW. Effects of warming ocean conditions on
2035 feeding ecology of small pelagic fishes in a coastal upwelling ecosystem: a shift to gelatinous
2036 food sources. *Mar Ecol Prog Ser.* 2018.
- 2037 155. ISAB. Review of the Comparative Survival Study (CSS) Draft 2018 Annual Report.
2038 Portland, Oregon: Northwest Power and Conservation Council; 2018. p. 27 pp.
- 2039 156. Brown RS, Harnish RA, Carter KM, Boyd JW, Deters KA, Eppard MB. An Evaluation of
2040 the Maximum Tag Burden for Implantation of Acoustic Transmitters in Juvenile Chinook
2041 Salmon. *North American Journal of Fisheries Management.* 2010;30:499–505. doi:
2042 0.1577/M09-038.1.
- 2043 157. Collins AL, Hinch SG, Welch DW, Cooke SJ, Clark TD. Intracoelomic tagging of juvenile
2044 sockeye salmon: swimming performance, growth, survival, and post-surgical wound healing in
2045 freshwater and during a transition to seawater. *Trans Amer Fish Soc.* 2013;142(2):515-23. doi:
2046 10.1080/00028487.2012.743928.
- 2047 158. Smircich M, Kelly J. Extending the 2% rule: the effects of heavy internal tags on stress
2048 physiology, swimming performance, and growth in brook trout. *Animal Biotelemetry.*
2049 2014;2(1):16. doi: 10.1186/2050-3385-2-16. PubMed PMID: doi:10.1186/2050-3385-2-16.
- 2050 159. Morrison PR, Groot EP, Welch DW. The Effect of Short-Duration Seawater Exposure
2051 and Acoustic Tag Implantation on the Swimming Performance and Physiology of Presmolt
2052 Juvenile Coho Salmon. *Transactions of the American Fisheries Society.* 2013;142(3):783-92.
2053 doi: 10.1080/00028487.2013.772536.
- 2054 160. Walker RW, Ashton NK, Brown RS, Liss SA, Colotelo AH, Beirão BV, et al. Effects of a
2055 novel acoustic transmitter on swimming performance and predator avoidance of juvenile
2056 Chinook Salmon: Determination of a size threshold. *Fish Res.* 2016;176:48-54. doi:
2057 <http://dx.doi.org/10.1016/j.fishres.2015.12.007>.
- 2058 161. Chittenden CM, Butterworth KG, Cubitt KF, Jacobs MC, Ladouceur A, Welch DW, et al.
2059 Maximum tag to body size ratios for an endangered coho salmon (*O. kisutch*) stock based on
2060 physiology and performance. . *Environmental Biology of Fishes:* . 2008;84(1):129-40.
- 2061 162. Prentice EF, Flagg TA, McCutcheon. CS. Feasibility of using implantable passive
2062 integrated transponder (PIT) tags in salmonids. *American Fisheries Society Symposium.* 1990a;
2063 7:317– 22.
- 2064 163. Prentice EF, Flagg TA, McCutcheon CS, Brastow DF, Cross DC. Equipment, methods,
2065 and an automated data-entry station for PIT tagging. *American Fisheries Society Symposium.*
2066 1990c;7:335–40.
- 2067 164. Prentice EF, Flagg TA, McCutcheon CS, Brastow DF. PIT-tag monitoring systems for
2068 hydroelectric dams and fish hatcheries. . *American Fisheries Society Symposium.*
2069 1990b;7:323–34.

- 2070 165. Gibbons JW, Andrews KM. PIT tagging: simple technology at its best. Bioscience.
2071 2004;54(5):447-54.
- 2072 166. Jefferts KB, Bergman PK, Fiscus HF. A coded wire identification system for macro-
2073 organisms. Nature. 1963;198:160-2.
- 2074 167. Johnson JK, editor Regional overview of coded wire tagging of anadromous salmon
2075 and steelhead in northwest America. American Fisheries Society Symposium; 1990.
- 2076 168. Weitkamp L, Neely K. Coho salmon (*Oncorhynchus kisutch*) ocean migration patterns:
2077 insight from marine coded-wire tag recoveries. Can J Fish Aquat Sci. 2002;59:1100-115.
- 2078 169. Bernard DR, Clark JE. Estimating salmon harvest with coded-wire tags. Can J Fish
2079 Aquat Sci. 1996;53(10):2323-32.
- 2080 170. Ricker WE. Ocean growth and mortality of pink and chum salmon. J Fish Res Bd
2081 Canada. 1964;21(5):905-6. doi: 10.1139/f64-087.
- 2082 171. Sharma R, Quinn TP. Linkages between life history type and migration pathways in
2083 freshwater and marine environments for Chinook salmon, *Oncorhynchus tshawytscha*. Acta
2084 Oecologica. 2012;41:1-13. doi: <http://dx.doi.org/10.1016/j.actao.2012.03.002>.
- 2085 172. Healey MC. Life history of Chinook salmon (*Oncorhynchus tshawytscha*). In: Groot C,
2086 Margolis L, editors. Pacific Salmon Life Histories. Vancouver: University of British Columbia
2087 Press; 1991. p. 311-94.
- 2088 173. Healey MC. Coastwide Distribution and Ocean Migration Patterns of Stream-and
2089 Ocean-Type Chinook Salmon, *Oncorhynchus tshawytscha*. Canadian Field Naturalist.
2090 1983;97(4):427-33.
- 2091 174. Waples RS, Teel DJ, Myers JM, Marshall AR. Life-history divergence in Chinook salmon:
2092 historic contingency and parallel evolution. Evolution. 2004;58(2):386-403. doi: 10.1554/03-
2093 323.
- 2094 175. Myers KW, Rogers DE. Stock origins of chinook salmon in incidental catches by
2095 groundfish fisheries in the eastern Bering Sea. North American Journal of Fisheries
2096 Management. 1988;8:162-71.
- 2097 176. Myers KW, Haris CK, Knudsen CM, Walker RV, Davis ND, Rogers DE. Stock origins of
2098 chinook salmon in the area of the Japanese mothership salmon fishery. North American
2099 Journal of Fisheries Management. 1987;7 No. 4 (Fall 1987):459-74.
- 2100 177. Clarke WC, Withler RE, Shelbourn JE. Inheritance of smolting phenotypes in
2101 backcrosses of hybrid stream-type x ocean-type chinook salmon (*Oncorhynchus tshawytscha*).
2102 Estuaries. 1994;17:13-25.
- 2103 178. Prince DJ, O'Rourke SM, Thompson TQ, Ali OA, Lyman HS, Saglam IK, et al. The
2104 evolutionary basis of premature migration in Pacific salmon highlights the utility of genomics
2105 for informing conservation. Science Advances. 2017;3(8). doi: 10.1126/sciadv.1603198.
- 2106 179. Anonymous. Insights from Genetic Analyses into Stock Specific Early Ocean Migration
2107 Behavior of Juvenile Columbia River Steelhead. Fisheries Bulletin. Under Review.
- 2108 180. Welch DW, Ishida Y, Nagasawa K, Eveson JP. Thermal limits on the ocean distribution
2109 of steelhead trout (*Oncorhynchus mykiss*). N Pac Anadr Fish Comm Bull. 1998;1:396-404.
- 2110 181. Ward BR, Slaney PA. Life history and smolt-to-adult survival of Keogh River steelhead
2111 trout (*Salmo gairdneri*) and the relationship to smolt size. Can J Fish Aquat Sci.
2112 1988;45(7):1110-22.
- 2113 182. Welch DW, Ward BR, Smith BD, Evenson JP. Influence of the 1990 Ocean Climate Shift
2114 on British Columbia Steelhead (*Oncorhynchus mykiss*) and Coho (*O.kisutch*) Populations.
2115 NPAFC. 1998;310:17p.

Coast-Wide Survival of Chinook and Steelhead **SUBMITTED to PLoS-ONE**

- 2116 183. Smith BD, Ward BR, Welch DW. Trends in British Columbia steelhead (*Oncorhynchus*
2117 *mykiss*) abundance indexed by angler success. Can J Fish Aquat Sci. 2000;57:1-16. doi:
2118 10.1139/f99-254.
- 2119 184. Welch DW, Ward BR, Smith BD, Whitney F. Changes associated with the 1989-90
2120 ocean climate shift, and effects on British Columbia steelhead (*O. mykiss*) populations. In:
2121 Fisheries and Oceans Canada, editor.: Pacific Stock Assessment Review Committee (PSARC)
2122 Paper S97-7; 1997.
- 2123 185. Neilson J, Taylor E. Emergency assessments of the Steelhead Trout (*Oncorhynchus*
2124 *mykiss*): Thompson River and Chilcotin River populations (2018). (Available upon request by
2125 contacting the COSEWIC secretariat at ec.cosepac-cosewic.ec@canada.ca). In: COSEWIC,
2126 editor. Ottawa: Government of Canada, Ministry of Environment and Climate Change; 2018. p.
2127 26 pp.
- 2128 186. Bradford MJ. Comparative review of Pacific salmon survival rates. Can J Fish Aquat Sci.
2129 1995;52(6):1327-38. doi: 10.1139/f95-129.
- 2130

1

Figure 1. Map of salmon survival time series used in the analyses. Numbers inside symbols are keyed to the populations in Supplementary Table S1; yellow circles indicate Chinook populations, pink squares indicate steelhead, and blue triangles indicate a location with data for both species. Acronyms: SEAK (SE Alaska/Northern British Columbia Transboundary Rivers); NCBC (North-Central British Columbia); WCVI (West Coast Vancouver Island); WAC (Washington Coastal); ORC (Oregon Coastal); SOG (Strait of Georgia); PS (Puget Sound).

10

11

Fig 2. Freshwater smolt survival for west coast North American rivers. A total of N=531 annual survival estimates are included. Top row: smolt survival from release to river mouth (and intermediate locations in the case of the Columbia). Bottom row: survival per 100 km of migration distance. The red horizontal line shows the median value for all Snake River data in a given panel (red coloured bars). Data are shown as a box and whisker plot with associated sample size listed above the appropriate boxes. Abbreviations: LRE, Lower Columbia River and estuary (i.e., the river reach from just below the lowest (Bonneville) dam to the river mouth); Release to BON measures Snake River survival from hatchery release through the Snake River above Lower Granite Dam and down through the 8-dam hydrosystem to the last dam (Bonneville). Full river measures survival from release to the mouth of the Columbia River. Data sources and annual survival estimates are reported in Supplementary Table S1.

22

Fig 3. Time series of smolt to adult survival (SAR) data for west coast Chinook stocks (excluding California).

Left column: subyearlings; Right column: yearlings. Regions are oriented from north (top) to south. Gold dots are SAR measurements based on CWT tags (PSC database), brown dots are SARs reported by Raymond [1], and violet dots are SARs based on PIT tags [2]. A loess curve of survival and associated 95% confidence interval (shaded region) using all available data for each panel is shown as a black line (the smoothing parameter was set to $\alpha=0.75$); the loess curves for Snake River subyearling and yearling survival are overplotted in red to facilitate comparison with other regions. Blank panels indicate regions where the life history type does not occur (for example, Fall (subyearling) Chinook do not occur in Alaska, while Spring (yearling) Chinook do not occur in the low elevation streams on the west coast of Vancouver Island or Oregon coast). The major regime shift years of 1977, 1989, and 1998 are indicated by vertical lines. In this and subsequent figures the pale red band delineates the official Columbia River SAR rebuilding targets of 2-6%.

Fig 5. Normalized SARs calculated by dividing individual SAR estimates for each stock and each year by the median Snake River SAR for the same year. Vertical lines show the median SAR for the Snake River (red) and other regions (blue). Note the logarithmic scale on the x-axis. As in the prior plots, Columbia & Snake River SAR estimates based on PIT tags do not incorporate harvest or above-dam survival.

39

40

Fig 6. Comparison of normalized Chinook SARs by regime periods: pre-1977, 1978-1989, 1990-1998, and post 1998. Boxes and whiskers have the conventional interpretation; the horizontal red line shows the Snake River median SAR value for each regime to facilitate comparison (1.0 by definition). Sample sizes are shown above each group (green font) and the ratio of median SARs relative to the Snake River is shown immediately above the upper whiskers (black font).

46

47

Fig 7. Steelhead SARs plotted against ocean entry year. Regions are oriented from north (left) to south (right); the Keogh R (KEOG) is situated on the NE tip of Vancouver Island (BC). Gold dots are SAR measurements based on PIT tags, brown dots are SARs reported by Raymond [1], and violet dots are SARs based on CWT tags. A loess curve of survival and associated 95% confidence interval (shaded region) using all available data for each panel is shown as a black line (the smoothing parameter was set to $\alpha=0.75$); the Snake River loess curve is shown in red and over plotted on all other panels to facilitate comparison. The major regime shift years of 1977, 1989, and 1998 are indicated by vertical lines.

Fig 8. Box and whisker plot of steelhead SARs by population (all available years). Population names are listed in Supplementary Table S1. The black horizontal line within each bar is the median of the SAR data available for that population. Median survival across all available data for each geographic region is shown as a blue line; median Snake River survival for all populations combined is shown as a red line and overplotted on all panels for comparison. The number of years of data is shown to the right.

Fig 9. Normalized steelhead SARs obtained by dividing each individual SAR estimate (i.e., for each stock and each year) by the median SAR calculated across all available Snake River SARs for that year. The median Snake River SAR is overplotted in red. Note the logarithmic scale on the x-axis.

Fig 10. Comparison of normalized steelhead SARs by regime periods: pre-1977, 1978-1989, 1990-1998, and post 1998. Boxes and whiskers have the conventional interpretation; the horizontal red line shows the Snake River median SAR value for each regime to facilitate comparison (1.0 by definition). Sample sizes are shown above each group (green font) and the ratio of median SARs relative to the Snake River is shown immediately above the upper whiskers (black font).

1. Raymond HL. Effects of Hydroelectric Development and Fisheries Enhancement on Spring and Summer Chinook Salmon and Steelhead in the Columbia River Basin. North American Journal of Fisheries Management. 1988;8(1):1-24.
2. McCann J, Chockley B, Cooper E, Hsu B, Schaller H, Haeseker S, et al. Comparative Survival Study of PIT-tagged Spring/Summer/Fall Chinook, Summer Steelhead, and Sockeye. 2017 Annual Report. Portland, Oregon 2017.

DRAFT