1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

Hippocampal connectivity with sensorimotor cortex
during volitional finger movements.
II. Spatial and temporal selectivity

Douglas D. Burman

short title: Spatio-temporal properties of hippocampal-sensorimotor connectivity

Department of Radiology, NorthShore University HealthSystem,
Evanston, Illinois, United States of America

Email: Dburman2@northshore.org

Abstract

Cognitive control refers to brain processes involved in regulating behavior according to internal goals or plans. This study examines whether hippocampal connectivity with sensorimotor cortex during paced movements shows a pattern of spatial and temporal selectivity required for cognitive control. Functional magnetic resonance imaging activity was recorded from thirteen right-handed subjects during a paced, non-mnemonic (repetitive tapping) motor task. Connectivity was examined from psychophysiological interactions in hippocampal activity during two analyses: the first identified motor interactions relative to rest, whereas the second identified differential motor activity between adjacent fingers. Connectivity was observed in both pre- and postcentral gyrus, but only postcentral connectivity was topographical, coincident with finger representations identified in a previous study. Differences in the magnitude of connectivity were observed between finger representations, representing spatial selectivity for the target of movements; the postcentral representation of the moving finger invariably showed greater connectivity than adjacent fingers. Furthermore, the magnitude of connectivity within a pre- or postcentral finger representation was largest when its finger moved, representing temporal selectivity for movement. While the hippocampus is known to be sensitive to spatial and temporal features of the environment, consistent with its role in learning and memory, the pattern of spatial and temporal selectivity of hippocampal connectivity observed in this study occurred during volitional movements in the absence of motor learning or recall. Spatial and temporal selectivity of connectivity during volitional movements meets the criteria for cognitive control adapted from oculomotor studies, suggesting a role for the hippocampus in motor control.

Introduction

To move purposefully, one or more signals within our brain direct the motor system to carry out the intended movement. This is the essence of *cognitive control*, the process by which goals or plans influence behavior. Although various regions of prefrontal cortex have been broadly implicated in cognitive control (1-5), the source of cognitive influence on the motor system is still subject to speculation.

Requirements for cognitive control of movements have been demonstrated in oculomotor studies, particularly those investigating the neural basis of reading (6-8) and other purposive eye movements (9-11). Such studies emphasize three components: 1) neural activity evident during conditions that require cognition; 2) spatial specificity, reflecting the spatial goal of the eye movement (i.e., target location); and 3) temporal specificity, which determines when the movement occurs.

The frontal eye field (FEF) is involved specifically in voluntary saccades, both in human and non-human primates (11, 12). Recording activity from nerve cells is generally avoided in humans, but FEF neurons in non-human primates reflect all three of these properties (13). These three properties are also evident from deficits following FEF lesions; for example, impairments in the accuracy and latency for a variety of volitional, but not reflexive eye movements are observed in both humans (11, 14-20) and non-human primates (21-24). In humans, lesions of the FEF and parietal eye field (PEF) differentially affect the selection of saccade targets and their timing (25); in monkeys, lesions of either area produce modest impairments in the accuracy and initiation of saccades, whereas combined lesions produce profound impairments (26). These studies demonstrate that the FEF, singly or in combination with PEF, is critically involved in the

generation of volitional eye movements, providing the timing of purposive eye movements and their target location.

Analogous to the FEF in the oculomotor system (27), the primary motor cortex is necessary for volitional movements of individual fingers (28, 29), with short fiber tracts that connect postcentral with precentral regions of sensorimotor cortex providing sensory feedback required for accurate performance (30). Neural activity in the FEF and sensorimotor cortex (SMC) both reflect cognitive input. FEF properties that reflect cognitive input includes activity that predicts two sequential saccades (31-33), target selection (34-37), covert attention (38-42), and changes during learning (27, 43). Similarly, cognitive input to SMC is evident from neural changes during motor learning, both in its response properties (44-49) and changes in connectivity (50-52). Although cognitive input to both areas can be inferred, direct evidence for the source (or sources) of this input is sparse. Indirect evidence from neural properties and lesion effects has implicated the supplementary eye field (SEF) (53, 54), basal ganglia (55), and dorsolateral prefrontal cortex (56) in the cognitive control of eye movements, and the basal ganglia (57), cingulate (58), and prefrontal cortex (1, 59-63) for the cognitive control of skeletomuscular movements.

From this, it is tempting to speculate that the prefrontal cortex regulates finger movements. The prefrontal cortex has traditionally been associated with cognitive control, yet its connectivity to the SMC is indirect via the dorsal premotor cortex (2, 64, 65). Prefrontal areas are functionally coupled to premotor areas when learning movement sequences but not during repetitive tapping (66, 67), suggesting prefrontal cortex may be involved in cognitive control when learning complex movement sequences but not during simple movements. In its role, premotor cortex is analogous to the SEF (27), which is specialized for learning eye movement sequences (27, 68-

70) and resolving conflicts between potential saccade targets (71, 72). Because prefrontal connectivity to FEF and SMC is indirect and limited to complex movements, cognitive control of simpler movements mediated specifically through FEF and SMC may arise from a different source. Thus, a relationship between neural activity and task complexity is unlikely when studying cognitive control in SMC, even though such a relationship has been used effectively to study cognitive control elsewhere (4, 73, 74).

The control of finger movements is more complex than eye movements for at least two reasons. First, the spatial reference for finger movements is different. Whereas eye movements are encoded from the current position of the eyes within the orbit (retinocentric space), finger movements are based upon the current position of the body (body-centered space); thus, the combination of finger muscles required to achieve a goal differs with different body positions. Second, the load on eye muscles is constant, whereas muscles moving the fingers may require different forces when moving objects of different resistance. When a task requires the initiation of finger movements from a set position with minimal load, however, requirements for cognitive control are the same as for eye movements: the neural mechanism must be active during volitional movements that require *cognition*, while specifying the timing and goal of the movement. The goal under these conditions reflects the body part that moves (e.g., hand or finger), requiring *cortical specificity* for neural activity within the SMC hand representation.

Whereas Part I of this study examined hippocampal-SMC connectivity within the framework of volitional movements, part II considers whether this connectivity shows spatial and temporal selectivity, necessary to specify which finger moves and when. Connectivity associated with movements of individual fingers is shown to be organized topographically, with connectivity in the representation of the moving finger greater than for adjacent fingers (spatial selectivity);

furthermore, connectivity within each finger representation is greatest when the finger moves, falling when movement stops (temporal selectivity). Combined with Part I, hippocampal-SMC connectivity thus fulfills the three conditions of selectivity required for cognitive control. After considering alternative interpretations, we conclude that hippocampal connectivity is consistent with cognitive control of motor function, and describe experiments that could further elucidate this functionality.

Materials and methods

Subjects

Thirteen right-handed adults from the Chicago metropolitan area participated in the study (ages 24–59, mean=42.3, five females). The nature of experimental procedures were explained to each subject before obtaining written consent; consent procedures complied with the Code of Ethics set forth in the Declaration of Helsinki, and were approved by the Institutional Review Board at the NorthShore University HealthSystem / Research Institute. Consented subjects had no history of the following: a previous concussion, psychiatric illness, learning disability, attention deficit disorder, abnormal physical or cognitive development that would affect educational performance, central neurological disease (or structural pathology), or neurosurgical intervention.

Immediately prior to their fMRI tests, subjects were trained on one cycle of each experimental task.

Experimental tasks

The tasks were described in detail in part I. Briefly, the visual/motor task used in this study comprised 6 cycles of a specified sequence of visual and motor conditions. Motor conditions were preceded by an instruction screen, accompanied by a metronome ticking at 2 beats per second. During the sequence learning task, subjects rehearsed a remembered sequence of button presses, tapping in synchrony with the metronome; this was followed by a visual condition, which required fixation at the center without finger movements while a circular checkerboard flashed at 4Hz. The instruction screen for repetitive tapping followed. For repetitive tapping, subjects repeatedly tapped the same finger on both hands, corresponding to the button identified on-screen; the index finger ('F1') tapped first, and the finger increased incrementally every 4s (eight taps). After tapping movements from all four fingers (F1-F4 during time periods T1-T4, respectively), the visual condition was repeated. Connectivity analysis in this report focused on repetitive tapping.

MRI data acquisition

Images were acquired using a 12-channel head coil in a 3 Tesla Siemens scanner (Verio). Visual stimuli projected onto a screen (Avotec Silent Vision) were viewed via a mirror attached to the head coil, and behavioral responses were recorded by Eprime [Psychology Software Tools, Inc.] from an optical response box (Current Designs, Philadelphia, PA). Blood-oxygen level dependent (BOLD) functional images were acquired using the echo planar imaging (EPI) method, using the following parameters: time of echo (TE) = 25ms, flip angle = 90°, matrix size = 64 x 64, field of view = 22cm, slice thickness = 3mm, number of slices = 32; time of repetition (TR) = 2000ms. The number of repetitions for the visual/motor condition was 182. A structural T1 weighted 3D image (TR = 1600ms, TE = 3.46ms, flip angle = 9°, matrix size = 256 x 256,

field of view = 22cm, slice thickness = 1mm, number of slices = 144) was acquired in the same orientation as the functional images.

fMRI data processing

Data was analyzed using SPM8 software (<http://www.fil.ion.ucl.ac.uk/spm>), using procedures described in Part I. Briefly, images were spatially aligned to the first volume to correct for small movements, sinc interpolation minimized timing-errors between slices, and functional images were coregistered to the anatomical image and normalized to the T1 Montreal Neurological Institute (MNI) template, then smoothed with a 10mm isotropic Gaussian kernel. During analysis, conditions of interest were specified for sequence learning, visual, and each of the four fingers moving during the repetitive tapping block; global normalization scaled the mean intensity of each brain volume to a common value to correct for whole brain differences over time.

Global conjunctive analysis of both motor conditions generated a bilateral activation map for hand movements, used as the ROI for connectivity analysis.

Psychophysiological interactions (PPI)

Preprocessing

Connectivity analysis was carried out using psychophysiological interactions (75), modified to account for individual variability in connectivity (76). As described in Part I, alternate voxels were sampled from the left (13) and right hippocampus (13) of the normalized brain, as delimited by the aal atlas in the WFU PickAtlas toolbox (<http://fmri.wfubmc.edu/software/PickAtlas>).

Two sets of connectivity analyses were employed, differing only in the interaction term used to identify task-specific activity. In the first analysis, an interaction term specified a greater effect of seed activity during movement of a single finger than during the visual condition; this identified the effect of movement relative to rest (e.g., F1>rest). In the second analysis, the interaction term specified a greater effect of seed activity during movement of one in a pair of adjacent fingers (e.g., F1>F2 or F2>F1); this identified preferential activity for one finger. After adjustments for regional differences in timing, a regression analysis identified the magnitude of the BOLD signal in SMC that was correlated with the PPI interaction term.

Seed selection

Two approaches were used to select seeds. The first approach was “structural”; the hippocampus was divided into 8 regions from posterior to anterior, with connectivity evaluated for each region (position-1 through position-8). The second approach was empirical or “functional”; the hippocampal voxel generating the greatest connectivity anywhere within SMC during a task was selected as a seed. Regardless of the method for selection, a voxel seed was selected from both the left and right hippocampus, and a conjunction (global) analysis was used to characterize connectivity.

Finger topography and spatio/temporal selectivity

The topographical arrangement of connectivity associated with individual finger movements was examined during both sets of analysis, exemplified by F1>rest and F1>F2. In the first analysis, the region of connectivity was plotted for each finger. In the second, the region of connectivity for each finger was restricted to the area of overlap between PPI maps for a specified finger; for example, connectivity for F2 was visualized as the overlap in connectivity between F2>F1 and

197 F2>F3.

198 Although differing in voxel resolution, the topographical arrangement of connectivity during
199 these analyses overlapped the finger topography identified in a previous study, which used a
200 variant of our paced repetitive tapping task to identify finger representations through differential
201 activation (77). (See Fig S1 for this comparison.) To investigate patterns of spatial and temporal
202 selectivity, the voxels constituting pre- and postcentral finger representations F1 through F3 from
203 the previous study were identified, and the beta estimate for the magnitude of connectivity was
204 recorded for each voxel.

205 During a specified time period, statistics were derived from voxel-wise comparisons with the
206 mean connectivity in the representation of the moving finger. To minimize confounds due to
207 variations in the baseline, the magnitude of connectivity for the moving finger was operationally
208 defined as the difference in connectivity during movement and the connectivity extremum when
209 the finger was stationery; the magnitude of connectivity for all other fingers (or time periods)
210 was expressed as a percentage change from this. A separate analysis was run for each time
211 period.

212 *Spatial selectivity* compared the magnitude of connectivity between two finger representations
213 during a single time period. Two-tailed paired t-tests identified differences in connectivity
214 between pairs of finger representations; differences between the moving finger and adjacent
215 fingers were particularly relevant. The paired t-test identified patterns of differences between
216 finger representations observed across individuals.

217 *Temporal selectivity* evaluated whether the magnitude of connectivity within each finger
218 representation differed across time periods; it was evaluated separately for each finger
219 representation. Two-tailed paired t-tests identified voxel-wise changes in connectivity between

each pair of time periods; differences in connectivity from the time period when the represented finger was moving were particularly relevant.

Results

Topography of Connectivity

The extent and topography of hippocampal connectivity with SMC derived from interactions between movements of a single finger and rest are shown in Fig 1 (see also Table 1). Functional seed-1 and seed-2 both generated connectivity in left SMC, with seed-2 generating more intense, bilateral connectivity; the expected time course of activity for later fingers was delayed (Fig 1A). By superimposing connectivity maps from each finger, a topographical distribution was evident (Fig 1B). Postcentral connectivity for F4 extended further dorsal than F2 and F1, evident in both the left postcentral gyrus for functional seed-1 (see sagittal section) and bilaterally for functional seed-2 (see coronal sections).

Fig 1. Connectivity and topography for individual fingers. (A) Peak activity predicted for connectivity from the left (solid colored line) and right hippocampal seeds (dashed colored line) were progressively delayed for F1 through F4; the timing predicted for fMRI activation is provided for comparison (black dotted line). Except for F3, connectivity associated with individual finger movements was limited to the left SMC for functional seed-1, but bilateral for functional seed-2. (B) A topography was evident from connectivity in the left postcentral gyrus for seed-1 and bilaterally for seed-2; connectivity for F4 was dorsolateral to the SMC region of connectivity for F2.

Table 1. Finger-specific connectivity in left SMC.

Finger/task interaction	Hemisphere (SMC)	Region	Cluster size	Z-score (peak)	Peak Coordinates
Funct-1 seed					

F1 > visual	left	pre/post	42	3.90	(-34,-24,54)
F2 > visual	left	pre/post	42	3.77	(-34,-24,54)
F3 > visual	left	pre	62	3.93	(-34,-20,58)
		post		3.58	(-38,-36,62)
	right	pre	29	5.06	(34,-12,62)
		post	8	3.80	(38,-36,58)
F4 > visual	left	pre/post	46	3.55	(-34,-24,54)
	right	pre	7	3.24	(42,-12,62)
Funct-2 seed					
F1 > visual	left	pre	102	3.70	(-34,-16,54)
		post		3.66	(-34,-32,62)
		pre		3.17	(-50,-8,42)
	right	pre	154	4.68	(34,-16,58)
		post		3.97	(38,-36,58)
F2 > visual	left	pre	110	3.70	(-34,-36,58)
		post		3.66	(-30,-20,54)
		pre		3.17	(-50,-24,46)
	right	pre	166	5.14	(34,-16,62)
		post		4.80	(48,-24,50)
		post		4.41	(38,-36,58)
F3 > visual	left	pre	124	4.11	(-34,-16,58)
		post		4.10	(-38,-36,62)
		post		3.88	(-46,-24,42)

F4 > visual	right	pre	166	5.48	(34,-16,62)
		post		5.15	(50,-24,54)
		post		4.50	(34,-40,54)
	left	post	154	4.37	(-38,-32,58)
		pre		4.31	(-34,-16,58)
		post		3.64	(-50,-8,42)
	right	post	167	5.53	(46,-20,54)
		pre		5.53	(38,-16,58)
		post		4.24	(38,-40,58)

235 Connectivity clusters in the left SMC from movement of individual fingers, surviving an intensity
236 threshold of $p=0.05$ with a family-wise error correction.

237

238 Fig 2 shows connectivity derived from hippocampal interactions in activity between movements
239 of adjacent fingers (see also Table 2). Arising from differences in the timing of neural activity
240 (Fig 2A, compare left and right columns), the area of cortical connectivity depended on the pair
241 of adjacent fingers used for analysis; connectivity only partially overlapped across pairings with
242 one finger in common (second row for F2 and third row for F3). Regions of overlap designated
243 connectivity for an individual finger. Connectivity showed a similar topographical organization
244 in the right postcentral gyrus (Fig 2B) as shown previously for the left (Fig 1B).

Fig 2. Connectivity and topography identified from psychophysiological interactions between movements of adjacent fingers. (A) The brain location of connectivity and its timing depended on the finger pairing; e.g., the interaction between F2 and F1 shown in the second row arose earlier than the interaction between F2 and F3, and the spatial extent of their connectivity differed. To better define topography, connectivity for F2 was identified from their overlap. (B) Connectivity was organized topographically, from F4 (blue, located dorsal) to F2 (red, intermediate) to F1 (yellow, ventral). For simplicity, connectivity for F3 is not shown.

245

246 **Table 2. Location of finger-specific connectivity in right SMC.**

Finger/task interaction	Hemisphere (SMC)	Region	Cluster size	Z-score (peak)	Peak Coordinates
F1>F2	right	precentral	9	4.36	(38,-12,66)
		postcentral	18	4.05	(38,-32,50)
F2>F1	right	postcentral	22	5.05	(50,-24,50)
		postcentral			(38,-32,62)
		precentral			(50,-16,46)
F2>F3	right	postcentral	22	4.41	(42,-28,54)
		postcentral		4.21	(34,-32,58)
F3>F2	right	postcentral	58	5.11	(50,-24,54)
		precentral		5.03	(34,-12,66)
		precentral		4.69	(46,-16,50)
F3>F4	right	precentral	58	4.01	(34,-16,66)
		postcentral		3.61	(50,-24,54)
		precentral		3.59	(38,-16,46)
F4>F3	right	precentral	71	4.44	(50,-12,46)
		postcentral		4.08	(42,-28,58)
		precentral		4.07	(38,-12,58)

247 Connectivity clusters in the right SMC from movement of individual fingers, surviving an intensity
 248 threshold of $p=0.05$ with a family-wise error correction.

249

Spatial / temporal interactions within SMC finger representations

One or more voxels within each finger representation showed significant connectivity during each time period; spatial and temporal selectivity were identified from differences in the magnitude of connectivity across representations or time periods.

Fig 3 examines spatial and temporal selectivity. Fig 3A illustrates spatial selectivity in the left SMC during the time period for each finger movement, based on the magnitude of connectivity derived from seed-1 interactions between finger movements and rest. Typically, differences >10-20% in connectivity between finger representations were significant (tabulated in Table S1). Differences were invariably observed in connectivity between the postcentral representation of the moving finger and the adjacent finger; differences were also observed during T2 between precentral representations of F2 and adjacent fingers. Connectivity with the postcentral SMC representation of the finger *adjacent* to the moving finger (e.g., F2 during T1) could also differ significantly from the representation of the finger twice removed from movement (in this

Fig 3. Spatial and temporal selectivity of SMC connectivity. (A) Spatial selectivity in left SMC reflected differences in the magnitude of connectivity between finger representations within a time period; except for F2, spatial selectivity was limited to postcentral gyrus. On each timeline, the highlighted period shows when a finger was moving; the colored arcs represent connectivity for those finger representations with elevated connectivity that includes the specified time period (yellow=F1, red=F2, green=F3). (B) Although connectivity in the right postcentral gyrus was identified from seed interactions between movements of adjacent fingers, spatial selectivity was similar to that observed in the left SMC. (C) Temporal selectivity in SMC connectivity reflected significant differences in the magnitude of a single finger representation's connectivity across time periods; systematic differences were not observed between pre- and postcentral gyrus. Connectivity in each finger's representation significantly differed during time periods when the finger was not moving.

Significant differences ($p < 0.05$) are italicized in bold type, with marginally significant results ($p < 0.10$) shown in plain type.

example, F3), suggesting connectivity is also elevated in the representation of the adjacent finger.

Fig 3B illustrates spatial selectivity in the right SMC for each time period, based on the magnitude of connectivity derived from seed-2 interactions between movements of adjacent fingers. Results were similar to those observed in the left SMC, except no connectivity was observed in precentral SMC.

Fig 3C examines temporal selectivity. In each row, the thickened segment highlights the time period serving as the statistical baseline for comparison. Within a finger representation, the magnitude of connectivity was elevated during the time period when the finger was moving, differing significantly from adjacent time periods. Connectivity during an adjacent time period could also be elevated, although modestly, adding a sequential time period without finger movements where differences were observed.

The magnitude of connectivity between and within individual finger representations shows both spatial and temporal selectivity, as summarized in Tables S1 and S2.

Discussion

In two analyses, the topography of hippocampal connectivity with SMC was found to be consistent with that previously demonstrated from fMRI activation (77). The magnitude of connectivity within these finger representations demonstrated both spatial selectivity (differences between finger representations during movement of a single finger) and temporal selectivity (differences across time periods within a finger representation). Because these movements were under volitional control, as demonstrated in Part I, these patterns of connectivity fulfill the

requirements for cognitive control adapted from oculomotor studies. The hippocampus thus augments the role of prefrontal cortex in the cognitive control of volitional finger movements, which is reportedly limited to complex sequences of movements.

Topography of finger representations

Activation studies have shown overlapping activation between nearby fingers, with representations for later fingers (e.g., F4) located dorsal to earlier fingers (77-82). The topographic organization of connectivity associated with individual finger movements followed this pattern in the postcentral gyrus of both hemispheres; no topography was apparent in the precentral gyrus, although precentral connectivity was observed.

Spatial selectivity

Spatial selectivity identified differences in the magnitude of connectivity between finger representations during a time period when a single finger moved. Selectivity was observed during each time period; the magnitude of connectivity was greatest for the moving finger, dropping significantly in the adjacent postcentral finger representation. Elevated connectivity could spill over into an adjacent finger representation, however; significant differences in connectivity were observed during T1, for example, between the representations for F2 and F3, as well as between F1 and F2. A similar spillover effect has been observed for fMRI activation during finger movements, where movement-related activation extends into the representation of adjacent fingers (78-82), perhaps reflecting the fact that adjacent fingers do not move independently

Spatial selectivity was observed bilaterally in postcentral finger representations.

Temporal selectivity within the finger representations

Temporal selectivity identified differences in the magnitude of connectivity at a specific finger representation across time periods. At each finger representation, maximal connectivity occurred when the finger was moving, with reduced connectivity during other time periods. Similar to the spillover into an adjacent finger representation observed for spatial selectivity, however, elevated connectivity could spill over into an adjacent time period.

Temporal selectivity did not differentiate between pre- and postcentral finger representations.

Hippocampal properties are consistent with cognitive control

As noted in the introduction, three features are essential for a role in cognitive control: 1) selective presence during tasks under volitional control; 2) spatial selectivity, reflecting actions directed to specific target locations; and 3) temporal selectivity, reflecting the timing of action. Neural activity consistent with these criteria has previously been reported for the hippocampus.

The hippocampus is selectively active during movements requiring cognitive control. Because the hippocampus is preferentially involved in conscious memories and environmental interactions (83, 84), studies of hippocampal interactions with the motor system have traditionally been focused on its role in explicit learning and memory consolidation.

Hippocampal connectivity with the striatum increases during explicit motor learning and consolidation (85-87); consolidation after rehearsal is further reflected through increased motor activation following sleep in both the primary motor area and the hippocampus (85, 88).

Hippocampal connectivity to SMC occurs during paced repetitive tapping movements as well as sequence learning; subjects' anticipatory responses indicate volitional control of these movements (see Part I of this study). These findings demonstrate hippocampal involvement in movements and motor learning under explicit cognitive control; by contrast, the hippocampus is inactive during implicit learning of motor sequences (89).

The second requirement for cognitive control is spatial selectivity. The hippocampus is sensitive to spatial location and navigation, with spatial functioning localized to the posterior hippocampus (90, 91), including space cells that encode a spatial map (92-96). Because the spatial location of the hand on the response pad was fixed in the current study, the spatial location of a button press corresponded to a SMC finger representation, and posterior structural seeds generated connectivity restricted to the SMC hand representation (see Part I). Furthermore, spatial selectivity was observed between finger representations, as the magnitude of connectivity dropped with distance from the representation of the finger currently moving. Thus, spatial properties of the hippocampus provided the selectivity in motor response required for cognitive control.

The third requirement for cognitive control is temporal selectivity. The hippocampus responds differentially to sequences of events that differ in order (97-100), and to the intervals between stimuli within a sequence (101-103). Cognitive awareness of the temporal intervals between metronome beats provided the basis for anticipatory behavioral responses in the current study. Furthermore, temporal selectivity was observed within each finger representation, such that connectivity was greatest when the represented finger was moving.

Connectivity with SMC during motor tasks is thus consistent with known hippocampal properties and meets the three criteria for the cognitive control of movements. This does not

preclude an additional role in cognitive control of SMC from other areas. Prefrontal cortex has been suggested to play an indirect role through its connectivity with dorsal premotor cortex (2), although functional coupling of prefrontal with premotor areas is limited to movement sequences (66). During paced repetitive tapping, however, hippocampal connectivity provides the best explanation for the cognitive control of SMC.

Cognitive control vs. other theories of hippocampal function

There is currently no consensus about the functional significance of the spatial and temporal properties of the hippocampus. Some have suggested the hippocampus has cognitive functions beyond its traditional roles (104-106), whereas others suggest its diverse properties merely reflect the varied components of episodic and long-term declarative memory (92, 93, 107).

Although hippocampal connectivity with SMC meets the requirements for cognitive control, could the results be better explained through its other known functions?

Memory of the pacing interval, plus associations between the numerical onscreen display and the corresponding fingers, were required for accurate performance during motor tasks in this study. These modest memory requirements might arguably require hippocampal input; neither spatial nor temporal selectivity, however, is required to access this mnemonic information. Spatial selectivity is unnecessary because the mapping between numerals and fingers never changed, whereas temporal selectivity is unnecessary because widespread cortical rhythms (such as theta) could serve to time events. The observed pattern of spatial and temporal selectivity is *required* for cognitive control, however, to specify the finger to be moved and when.

An alternative explanation is that hippocampal connectivity in this study represents a form of sensorimotor processing, facilitating the transformation of sensory signals into motor output

(108-111). This interpretation could explain why its topography and spatial selectivity were most prominent in the postcentral gyrus. Even without hippocampal intervention, however, extensive connections between pre- and postcentral gyrus provides sensory feedback (112), so aside from cognitive control, the role of hippocampal sensorimotor processing in SMC is unclear. Furthermore, facilitation of sensorimotor processing does not require temporal specificity. Given the directional influence of the hippocampus on SMC (implicit in PPI analysis), a hippocampal role in sensorimotor processing begs the question: what directs the hippocampus to provide relevant spatio-temporal information to SMC as needed for task performance? Considering the tight relationship between neural activity in the hippocampal seeds and SMC (shown in Fig 2 of Part I), an additional control signal in SMC seems unlikely. These considerations do not exclude the possibility of other sources of cognitive control under different conditions. By analogy with the oculomotor system, damage to a single structure (e.g., FEF or SMC) can produce deficits without eliminating all function (eye movements or hand movements) due to the preservation of other structures with overlapping functions (SEF / PEF, premotor cortex / basal ganglia). Similarly, executive functions show deficits following damage to prefrontal areas, yet are not eliminated (113-115). Hippocampal dysfunction in Alzheimer's impairs volitional motor control (116), but with overlapping systems for cognitive control, it would not eliminate volitional movements.

Implications and directions for future research

This study provides evidence that the hippocampus is involved in the cognitive control of volitional finger movements. Whether this represents a new function, or represents another

component of a single cognitive function, is unclear. These alternative possibilities suggest two broad experimental questions for further investigation.

First, how extensive is the hippocampal role in cognitive control during volitional movements?

Based on the criteria for cognitive control, hippocampal connectivity should be lost when “mindless” finger movements are generated while simultaneously performing a mentally-challenging task; connectivity also needs to be demonstrated during volitional movements of other body parts. In addition, a fixed spatial relationship between fingers and button presses in the current study allowed cortical specificity for finger movements to provide a measure of spatial selectivity; it’s unclear how the hippocampal pattern of cortical / spatial selectivity would be affected by changes in the spatial target of finger movements.

Second, is the hippocampal role in cognitive control limited to volitional movements, or does it extend to other cognitive activities? The hippocampus is critically involved in the formation and recall of episodic and declarative memories, thus fulfilling the first criterion for cognitive control (preferentially active during volitional states). Whether hippocampal interactions with cortical areas meet the remaining criteria for cognitive control is unclear, although theoretical accounts of memory function suggest they might. The hippocampus has been suggested to index memories, selectively accessing the prefrontal site of memory storage during the time period when memories are recalled (117-122). Such interactions provide cortical and temporal specificity, and may explain how engrams located elsewhere are selected by the hippocampus during recall. Similarly, spatial and temporal selectivity from hippocampal interactions with prefrontal cortex may provide the context for interpreting sensory input (95, 96, 123-126); miscommunication between the hippocampus and prefrontal cortex in schizophrenia would then produce abnormal perceptions (104, 127, 128).

When spatial and temporal selectivity of hippocampal connectivity with cortical areas is necessary to perform cognitive tasks, a role of the hippocampus in cognitive control is implied. The role of the hippocampus in cognitive control may be evaluated through experimental manipulations that affect cognitive awareness and performance, evaluating effects of these variables on the connectivity of the hippocampus with brain areas directly involved in task performance. Comparing effects of these variables on hippocampal and prefrontal connectivity may help identify conditions where these regions are either complementary or synergistic in the cognitive control of behavior.

Conclusions

The hippocampus showed task-specific connectivity with SMC during paced movement tasks, meeting the criteria for cortical control adapted from studies of cognition in the oculomotor system: associated with volitional movements, restricted to the representations of the moving fingers, and selective for the timing of the movements. This pattern of connectivity is consistent with known hippocampal properties, yet cannot be explained by its established functions. A role for the hippocampus in cognitive control is indicated; a collaborative role with prefrontal cortex is suggested from their close functional relationship across tasks and neurological conditions.

Acknowledgments

The author wishes to thank Donald J. Bolger for his suggestions on an early draft of this manuscript, and the Center for Advanced Imaging (CAI) at NorthShore University HealthSystem for its administrative support.

References

1. Cieslik EC, Zilles K, Caspers S, Roski C, Kellermann TS, Jakobs O, et al. Is there "one" DLPFC in cognitive action control? Evidence for heterogeneity from co-activation-based parcellation. *Cerebral cortex* 2012:bhs256.
2. Cole MW, Schneider W. The cognitive control network: integrated cortical regions with dissociable functions. *Neuroimage* 2007;37(1):343-360.
3. Ochsner KN, Ray RD, Cooper JC, Robertson ER, Chopra S, Gabrieli JD, et al. For better or for worse: neural systems supporting the cognitive down- and up-regulation of negative emotion. *Neuroimage* 2004;23(2):483-99.
4. Rajah MN, Ames B, D'Esposito M. Prefrontal contributions to domain-general executive control processes during temporal context retrieval. *Neuropsychologia* 2008;46(4):1088-103.
5. Ridderinkhof KR, Ullsperger M, Crone EA, Nieuwenhuis S. The role of the medial frontal cortex in cognitive control. *Science* 2004;306(5695):443-7.
6. Liversedge SP, Findlay JM. Saccadic eye movements and cognition. *Trends in Cognitive Sciences* 2000;4(1):6-14.
7. Rayner K, Liversedge SP, White SJ, Vergilino-Perez D, Mail E, rayner@psych.umass.edu. Reading disappearing text: Cognitive control of eye movements. *Psychological Science* 2003;14(4) Jul 2003:United Kingdom <http://www>.
8. Pollatsek A, Rayner K. Eye movements and lexical access in reading. In: *Comprehension processes in reading*: Routledge; 2012. p. 165-186.
9. Hayhoe M, Ballard D. Eye movements in natural behavior. *Trends in cognitive sciences* 2005;9(4):188-194.

- 465 10. Hutton SB. Cognitive control of saccadic eye movements. Brain and cognition
466 2008;68(3):327-340.
- 467 11. McDowell JE, Dyckman KA, Austin BP, Clementz BA. Neurophysiology and
468 neuroanatomy of reflexive and volitional saccades: evidence from studies of humans.
469 Brain & Cognition 2008;68(3):255-70.
- 470 12. Vernet M, Quentin R, Chanes L, Mitsumasu A, Valero-Cabré A. Frontal eye field, where
471 art thou? Anatomy, function, and non-invasive manipulation of frontal regions involved
472 in eye movements and associated cognitive operations. Frontiers in integrative
473 neuroscience 2014;8:66.
- 474 13. Bruce CJ, Goldberg ME. Primate frontal eye fields: I. Single neurons discharging before
475 saccades. J. Neurophysiol. 1985;53:603-635.
- 476 14. Guitton D, Buchtel HA, Douglas RM. Frontal lobe lesions in man cause difficulties in
477 suppressing reflexive glances and in generating goal-directed saccades. Exp. Brain Res.
478 1985;58:455-472.
- 479 15. Keating EG. Frontal eye field lesions impair predictive and visually-guided pursuit eye
480 movements. Exp. Brain Res. 1991;86:311-323.
- 481 16. Henik A, Rafal R, Rhodes D. Endogenously generated and visually guided saccades after
482 lesions of the human frontal eye fields. Journal of Cognitive Neuroscience
483 1994;6(4):400-411.
- 484 17. Munoz DP, Everling S. Look away: the anti-saccade task and the voluntary control of eye
485 movement. Nature Reviews Neuroscience 2004;5(3):218.

- 486 18. Rivaud S, Müri RM, Gaymard B, Vermersch AI, Pierrot-Deseilligny C. Eye movement
487 disorders after frontal eye field lesions in humans. *Experimental Brain Research*
488 1994;102(1):110-120.
- 489 19. Stuyven E, Van der Goten K, Vandierendonck A, Claeys K, Crevits L. The effect of
490 cognitive load on saccadic eye movements. *Acta psychologica* 2000;104(1):69-85.
- 491 20. Van der Stigchel S, Van Koningsbruggen M, Nijboer TCW, List A, Rafal RD. The role
492 of the frontal eye fields in the oculomotor inhibition of reflexive saccades: Evidence from
493 lesion patients. *Neuropsychologia* 2012;50(1):198-203.
- 494 21. Gaymard B, Ploner CJ, Rivaud-Pechoux S, Pierrot-Deseilligny C. The frontal eye field is
495 involved in spatial short-term memory but not in reflexive saccade inhibition.
496 *Experimental Brain Research* 1999;129(2):288-301.
- 497 22. Schiller PH, True SD, Conway JL. Deficits in eye movements following frontal eye field
498 and superior colliculus ablations. *J. Neurophysiol.* 1980;44:1175-1189.
- 499 23. Schiller PH, Chou Ih. The effects of frontal eye field and dorsomedial frontal cortex
500 lesions on visually guided eye movements. *Nature neuroscience* 1998;1(3):248.
- 501 24. Tehovnik EJ, Sommer MA, Chou IH, Slocum WM, Schiller PH. Eye fields in the frontal
502 lobes of primates. *Brain Research Reviews* 2000;32(2-3):413-448.
- 503 25. Braun D, Weber H, Mergner TH, Schulte-Mönting J. Saccadic reaction times in patients
504 with frontal and parietal lesions. *Brain* 1992;115(5):1359-1386.
- 505 26. Lynch JC. Saccade initiation and latency deficits after combined lesions of the frontal and
506 posterior eye fields in monkeys. *Journal of Neurophysiology* 1992;68(5):1913-1916.
- 507 27. Chen LL, Wise SP. Supplementary eye field contrasted with the frontal eye field during
508 acquisition of conditional oculomotor associations. *J. Neurophysiol.* 1995;73:1122-1134.

- 509 28. Lang CE, Schieber MH. Differential impairment of individuated finger movements in
510 humans after damage to the motor cortex or the corticospinal tract. *Journal of*
511 *neurophysiology* 2003;90(2):1160-1170.
- 512 29. Lang CE, Schieber MH. Reduced muscle selectivity during individuated finger
513 movements in humans after damage to the motor cortex or corticospinal tract. *Journal of*
514 *neurophysiology* 2004;91(4):1722-1733.
- 515 30. Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, et al. Short frontal lobe
516 connections of the human brain. *Cortex* 2012;48(2):273-291.
- 517 31. Colby CL, Goldberg ME. The updating of the representation of visual space in parietal
518 cortex by intended eye movements. *Science* 1992;255(5040):90-92.
- 519 32. Phillips AN, Segraves MA. Predictive Activity in Macaque Frontal Eye Field Neurons
520 During Natural Scene Searching. *J Neurophysiol* 2009;103:1238-1252.
- 521 33. Umeno MM, Goldberg ME. Spatial processing in the monkey frontal eye field. I.
522 Predictive visual responses. *Journal of neurophysiology* 1997;78(3):1373-1383.
- 523 34. Burman DD, Segraves MA. Primate frontal eye field activity during natural scanning eye
524 movements. *Journal of Neurophysiology* 1994;71(3):1266-1271.
- 525 35. Hanes DP, Patterson Ii WF, Schall JD. Role of Frontal Eye Fields in Countermanding
526 Saccades: Visual, Movement, and Fixation Activity. 1998.
- 527 36. Schall JD, Hanes DP. Neural basis of saccade target selection in frontal eye field during
528 visual search. *Nature* 1993;366(6454):467.
- 529 37. Schall JD, Purcell BA, Heitz RP, Logan GD, Palmeri TJ. Neural mechanisms of saccade
530 target selection: gated accumulator model of the visual-motor cascade. *European Journal*
531 *of Neuroscience* 2011;33(11):1991-2002.

38. Garg A, Schwartz D, Stevens AA. Orienting auditory spatial attention engages frontal eye fields and medial occipital cortex in congenitally blind humans. *Neuropsychologia* 2007;45(10):2307-21.
39. Heitz RP, Cohen JY, Woodman GF, Schall JD. Neural correlates of correct and errant attentional selection revealed through N2pc and frontal eye field activity. *Journal of neurophysiology* 2010;104(5):2433-2441.
40. Moore T, Fallah M. Microstimulation of the frontal eye field and its effects on covert spatial attention. *Journal of neurophysiology* 2004;91(1):152-162.
41. Thompson KG, Bichot NP. A visual salience map in the primate frontal eye field. *Progress in brain research* 2005;147:249-262.
42. Wardak C, Ibos G, Duhamel J-R, Olivier E. Contribution of the monkey frontal eye field to covert visual attention. *Journal of Neuroscience* 2006;26(16):4228-4235.
43. Little DM, Klein R, Shobat DM, McClure ED, Thulborn KR. Changing patterns of brain activation during category learning revealed by functional MRI. *Cognitive Brain Research* 2004;22(1):84-93.
44. Floyer-Lea A, Wylezinska M, Kincses T, Matthews PM. Rapid modulation of GABA concentration in human sensorimotor cortex during motor learning. *Journal of neurophysiology* 2006;95(3):1639-1644.
45. Muellbacher W, Ziemann U, Wissel J, Dang N, Kofler M, Facchini S, et al. Early consolidation in human primary motor cortex. *Nature* 2002;415(6872):640.
46. Rosenkranz K, Kacar A, Rothwell JC. Differential modulation of motor cortical plasticity and excitability in early and late phases of human motor learning. *Journal of Neuroscience* 2007;27(44):12058-12066.

- 555 47. Sanes JN, Donoghue JP. Plasticity and primary motor cortex. Annual Review of
556 Neuroscience 2000;23:393-415.
- 557 48. Pollok B, Latz D, Krause V, Butz M, Schnitzler A. Changes of motor-cortical oscillations
558 associated with motor learning. Neuroscience 2014;275:47-53.
- 559 49. Ostry DJ, Gribble PL. Sensory Plasticity in Human Motor Learning. Trends in
560 Neurosciences 2016;39(2):115.
- 561 50. Ma L, Wang B, Narayana S, Hazeltine E, Chen X, Robin DA, et al. Changes in regional
562 activity are accompanied with changes in inter-regional connectivity during 4 weeks
563 motor learning. Brain research 2010;1318:64-76.
- 564 51. Steele CJ, Penhune VB. Specific increases within global decreases: a functional magnetic
565 resonance imaging investigation of five days of motor sequence learning. The Journal of
566 neuroscience 2010;30(24):8332-8341.
- 567 52. Sun FT, Miller LM, Rao AA, D'Esposito M. Functional connectivity of cortical networks
568 involved in bimanual motor sequence learning. Cerebral cortex 2006;17(5):1227-1234.
- 569 53. Stuphorn V, Schall JD. Executive control of countermanding saccades by the
570 supplementary eye field. Nature neuroscience 2006;9(7):925.
- 571 54. Stuphorn V, Brown JW, Schall JD. Role of supplementary eye field in saccade initiation:
572 executive, not direct, control. Journal of neurophysiology 2009;103(2):801-816.
- 573 55. Hikosaka O, Takikawa Y, Kawagoe R. Role of the basal ganglia in the control of
574 purposive saccadic eye movements. Physiological reviews 2000;80(3):953-978.
- 575 56. Pierrot-Deseilligny C, Milea D, Muri RM. Eye movement control by the cerebral cortex.
576 Current Opinion in Neurology 2004;17:17-25.

57. Graybiel AM. The basal ganglia: learning new tricks and loving it. *Current opinion in neurobiology* 2005;15(6):638-644.
58. Hoffstaedter F, Grefkes C, Caspers S, Roski C, Palomero-Gallagher N, Laird AR, et al. The role of anterior midcingulate cortex in cognitive motor control. *Human brain mapping* 2014;35(6):2741-2753.
59. Azuar C, Reyes P, Slachevsky A, Volle E, Kinkingnehun S, Kouneiher Fdr, et al. Testing the model of caudo-rostral organization of cognitive control in the human with frontal lesions. *Neuroimage* 2014;84:1053-1060.
60. Badre D. Cognitive control, hierarchy, and the rostro-caudal organization of the frontal lobes. *Trends in cognitive sciences* 2008;12(5):193-200.
61. Cole MW, Yarkoni T, Repovš G, Anticevic A, Braver TS. Global connectivity of prefrontal cortex predicts cognitive control and intelligence. *The Journal of Neuroscience* 2012;32(26):8988-8999.
62. Koechlin E, Ody C, Kouneiher Fdr. The architecture of cognitive control in the human prefrontal cortex. *Science* 2003;302(5648):1181-1185.
63. Ridderinkhof KR, Van Den Wildenberg WPM, Segalowitz SJ, Carter CS. Neurocognitive mechanisms of cognitive control: the role of prefrontal cortex in action selection, response inhibition, performance monitoring, and reward-based learning. *Brain and cognition* 2004;56(2):129-140.
64. Chouinard PA, Paus TÅ. The primary motor and premotor areas of the human cerebral cortex. *The neuroscientist* 2006;12(2):143-152.

- 598 65. Koch G, Franca M, Del Olmo MF, Cheeran B, Milton R, Saucó MA, et al. Time course
599 of functional connectivity between dorsal premotor and contralateral motor cortex during
600 movement selection. *The Journal of neuroscience* 2006;26(28):7452-7459.
- 601 66. Abe M, Hanakawa T. Functional coupling underlying motor and cognitive functions of
602 the dorsal premotor cortex. *Behavioural Brain Research* 2009;198(1):13-23.
- 603 67. Kantak SS, Stinear JW, Buch ER, Cohen LG. Rewiring the brain: potential role of the
604 premotor cortex in motor control, learning, and recovery of function following brain
605 injury. *Neurorehabilitation and neural repair* 2012;26(3):282-292.
- 606 68. Chen LL, Wise SP. Neuronal activity in the supplementary eye field during acquisition of
607 conditional oculomotor associations. *J. Neurophysiol.* 1995;73:1101-1121.
- 608 69. Gaymard B, Pierrot-Deseilligny C, Rivaud S. Impairment of sequences of memory-
609 guided saccades after supplementary motor area lesions. *Annals of Neurology* 1990.
- 610 70. Isoda M, Tanji J. Cellular activity in the supplementary eye field during sequential
611 performance of multiple saccades. *Journal of neurophysiology* 2002;88(6):3541-3545.
- 612 71. Husain M, Parton A, Hodgson TL, Mort D, Rees G. Self-control during response conflict
613 by human supplementary eye field. *Nature neuroscience* 2003;6(2):117.
- 614 72. Parton A, Nachev P, Hodgson TL, Mort D, Thomas D, Ordidge R, et al. Role of the
615 human supplementary eye field in the control of saccadic eye movements.
616 *Neuropsychologia* 2007;45(5):997-1008.
- 617 73. Harvey P-O, Fossati P, Pochon J-B, Levy R, LeBastard G, Lehericy S, et al. Cognitive
618 control and brain resources in major depression: an fMRI study using the n-back task.
619 *Neuroimage* 2005;26(3):860-869.

74. Lavie N. Attention, distraction, and cognitive control under load. *Current Directions in Psychological Science* 2010;19(3):143-148.
75. Gitelman DR, Penny WD, Ashburner J, Friston KJ. Modeling regional and psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution. *Neuroimage* 2003;19(1):200-207.
76. Burman DD, Minas T, Bolger DJ, Booth JR. Age, sex, and verbal abilities affect location of linguistic connectivity in ventral visual pathway. *Brain and language* 2013;124(2):184-193.
77. Burman DD, Lie-Nemeth T, Brandfonbrener AG, Parisi T, Meyer JR. Altered Finger Representations in Sensorimotor Cortex of Musicians with Focal Dystonia: Precentral Cortex. *Brain Imaging and Behavior* 2009;3(1):10-23.
78. Beisteiner R, Windischberger C, Lanzenberger R, Edward V, Cunnington R, Erdler M, et al. Finger somatotopy in human motor cortex.[see comment]. *Neuroimage* 2001;13(6 Pt 1):1016-26.
79. Dechent P, Frahm J. Functional somatotopy of finger representations in human primary motor cortex. *Human Brain Mapping* 2003;18(4):272-83.
80. Hlustik P, Solodkin A, Gullapalli RP, Noll DC, Small SL. Somatotopy in human primary motor and somatosensory hand representations revisited. *Cerebral Cortex* 2001;11(4):312-21.
81. Kleinschmidt A, Nitschke MF, Frahm J. Somatotopy in the human motor cortex hand area. A high-resolution functional MRI study. *European Journal of Neuroscience* 1997;9(10):2178-86.

- 642 82. Volkmann J, Schnitzler A, Witte OW, Freund H. Handedness and asymmetry of hand
643 representation in human motor cortex. *Journal of Neurophysiology* 1998;79(4):2149-54.
- 644 83. Behrendt R-P. Hippocampus and consciousness. *Reviews in the Neurosciences*
645 2013;24(3):239-266.
- 646 84. Huijgen J, Samson S. The hippocampus: a central node in a large-scale brain network for
647 memory. *Revue neurologique* 2015;171(3):204-216.
- 648 85. Albouy G, King BR, Maquet P, Doyon J. Hippocampus and striatum: Dynamics and
649 interaction during acquisition and sleep-related motor sequence memory consolidation.
650 *Hippocampus* 2013;23(11):985-1004.
- 651 86. Albouy G, Fogel S, King BR, Laventure S, Benali H, Karni A, et al. Maintaining vs.
652 enhancing motor sequence memories: Respective roles of striatal and hippocampal
653 systems. *Neuroimage* 2015;108:423-434.
- 654 87. Fernandez-Seara MA, Aznarez-Sanado M, Mengual E, Loayza FR, Pastor MA.
655 Continuous performance of a novel motor sequence leads to highly correlated striatal and
656 hippocampal perfusion increases. *Neuroimage* 2009;47(4):1797-1808.
- 657 88. Walker MP, Stickgold R, Alsop D, Gaab N, Schlaug G. Sleep-dependent motor memory
658 plasticity in the human brain. *Neuroscience* 2005;133(4):911-917.
- 659 89. Rose M, Haider H, Salari N, Buchel C. Functional dissociation of hippocampal
660 mechanism during implicit learning based on the domain of associations. *J. Neurosci.*
661 2011;31:13739-13745.
- 662 90. Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the hippocampal
663 longitudinal axis. *Nature Reviews Neuroscience* 2014;15(10):655.

- 664 91. Travis SG, Huang Y, Fujiwara E, Radomski A, Olsen F, Carter R, et al. High field
665 structural MRI reveals specific episodic memory correlates in the subfields of the
666 hippocampus. *Neuropsychologia* 2014;53:233-245.
- 667 92. Eichenbaum H. On the Integration of Space, Time, and Memory. *Neuron*
668 2017;95(5):1007.
- 669 93. Eichenbaum H. The role of the hippocampus in navigation is memory. *Journal of*
670 *neurophysiology* 2017;117(4):1785.
- 671 94. Ekstrom AD, Ranganath C. Space, time, and episodic memory: the hippocampus is all
672 over the cognitive map. *Hippocampus* 2017.
- 673 95. Libby LA, Hannula DE, Ranganath C. Medial temporal lobe coding of item and spatial
674 information during relational binding in working memory. *J. Neurosci.* 2014;34:14233-
675 14242.
- 676 96. Rolls ET. Pattern separation, completion, and categorisation in the hippocampus and
677 neocortex. *Neurobiology of learning and memory* 2016;129:4-28.
- 678 97. Davachi L, DuBrow S. How the hippocampus preserves order: the role of prediction and
679 context. *Trends in cognitive sciences* 2015;19(2):92-99.
- 680 98. Ekstrom AD, Copara MS, Isham EA, Wang W-c, Yonelinas AP. Dissociable networks
681 involved in spatial and temporal order source retrieval. *Neuroimage* 2011;56(3):1803-
682 1813.
- 683 99. Paz R, Gelbard-Sagiv H, Mukamel R, Harel M, Malach R, Fried I. A neural substrate in
684 the human hippocampus for linking successive events. *Proceedings of the National*
685 *Academy of Sciences* 2010;107(13):6046-6051.

100. Warburton EC, Brown MW. Findings from animals concerning when interactions
between perirhinal cortex, hippocampus and medial prefrontal cortex are necessary for
recognition memory. *Neuropsychologia* 2010;48(8):2262-2272.
101. Barnett AJ, O'neil EB, Watson HC, Lee ACH. The human hippocampus is sensitive to
the durations of events and intervals within a sequence. *Neuropsychologia* 2014;64:1-12.
102. MacDonald CJ, Fortin NJ, Sakata S, Meck WH. Retrospective and prospective views on
the role of the hippocampus in interval timing and memory for elapsed time. *Timing &
Time Perception* 2014;2(1):51-61.
103. MacDonald CJ. Prospective and retrospective duration memory in the hippocampus: is
time in the foreground or background? *Phil. Trans. R. Soc. B* 2014;369(1637):20120463.
104. Behrendt R-P. Consciousness, memory, and hallucinations. In: *Hallucinations*: Springer;
2013. p. 17-31.
105. Olsen RK, Moses SN, Riggs L, Ryan JD. The hippocampus supports multiple cognitive
processes through relational binding and comparison. *Frontiers in human neuroscience*
2012;6:146.
106. Poch C, Campo P. Neocortical-hippocampal dynamics of working memory in healthy
and diseased brain states based on functional connectivity. *Frontiers in human
neuroscience* 2012;6.
107. Moscovitch M, Cabeza R, Winocur G, Nadel L. Episodic memory and beyond: the
hippocampus and neocortex in transformation. *Annual review of psychology*
2016;67:105-134.
108. Bast T, Feldon J. Hippocampal modulation of sensorimotor processes. *Progress in
neurobiology* 2003;70(4):319-345.

709 109. Cruikshank LC, Singhal A, Caplan JB. Theta oscillations reflect a putative neural
710 mechanism for human sensorimotor integration. *Journal of Neurophysiology* 2012;jn.
711 00893.2010.

712 110. Dypvik AT, Bland BH. Functional connectivity between the red nucleus and the
713 hippocampus supports the role of hippocampal formation in sensorimotor integration.
714 *Journal of neurophysiology* 2004;92(4):2040-2050.

715 111. Scheidt RA, Zimbelman JL, Salowitz NMG, Suminski AJ, Mosier KM, Houk J, et al.
716 Remembering forward: neural correlates of memory and prediction in human motor
717 adaptation. *Neuroimage* 2012;59(1):582-600.

718 112. Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, et al. Short frontal lobe
719 connections of the human brain. *cortex* 2012;48(2):273-291.

720 113. Alvarez JA, Emory E. Executive function and the frontal lobes: a meta-analytic review.
721 *Neuropsychology review* 2006;16(1):17-42.

722 114. Barbey AK, Colom R, Solomon J, Krueger F, Forbes C, Grafman J. An integrative
723 architecture for general intelligence and executive function revealed by lesion mapping.
724 *Brain* 2012;135(4):1154-1164.

725 115. Leh SE, Petrides M, Strafella AP. The neural circuitry of executive functions in healthy
726 subjects and Parkinson's disease. *Neuropsychopharmacology* 2010;35(1):70.

727 116. Buchman AS, Bennett DA. Loss of motor function in preclinical Alzheimer's disease.
728 *Expert review of neurotherapeutics* 2011;11(5):665-676.

729 117. Eichenbaum H. Prefrontal-hippocampal interactions in episodic memory. *Nature reviews.*
730 *Neuroscience* 2017;18(9):547-558.

- 731 118. Eichenbaum H. Memory: Organization and Control. Annual review of psychology
732 2017;68:19.
- 733 119. Preston AR, Eichenbaum H. Interplay of Hippocampus and Prefrontal Cortex in Memory.
734 Current Biology 2013;23(17):R764-R773.
- 735 120. Rosenbaum RS, Winocur G, Moscovitch M. New views on old memories: re-evaluating
736 the role of the hippocampal complex. Behavioural Brain Research 2001;127(1-2):183-97.
- 737 121. Teyler TJ, DiScenna P. The hippocampal memory indexing theory. Behavioral
738 neuroscience 1986;100(2):147.
- 739 122. Teyler TJ, Rudy JW. The hippocampal indexing theory and episodic memory: updating
740 the index. Hippocampus 2007;17(12):1158-1169.
- 741 123. Burgess N, Becker S, King JA, O'Keefe J. Memory for events and their spatial context:
742 models and experiments. Philosophical Transactions of the Royal Society of London -
743 Series B: Biological Sciences 2001;356(1413):1493-503.
- 744 124. Burgess N, Maguire EA, Spiers HJ, O'Keefe J. A temporoparietal and prefrontal network
745 for retrieving the spatial context of lifelike events. Neuroimage 2001;14(2):439-53.
- 746 125. Jenkins LJ, Ranganath C. Prefrontal and medial temporal lobe activity at encoding
747 predicts temporal context memory. Journal of Neuroscience 2010;30(46):15558-15565.
- 748 126. Hsieh L-T, Gruber MJ, Jenkins LJ, Ranganath C. Hippocampal activity patterns carry
749 information about objects in temporal context. Neuron 2014;81(5):1165-1178.
- 750 127. Behrendt R-P. Contribution of hippocampal region CA3 to consciousness and
751 schizophrenic hallucinations. Neuroscience & Biobehavioral Reviews 2010;34(8):1121-
752 1136.

- 753 128. Behrendt R-P. Consciousness, memory, and hallucinations. In: Hallucinations: Springer;
754 2012. p. 17-31.

Fig S1. SMC topography from fMRI activation and PPI connectivity during paced tapping. The fMRI topography was demonstrated in the left SMC by contrasting activation by each finger in the right hand with all other fingers, applied to images acquired with 1mm in-plane resolution and 3mm slice thickness (Burman et al., 2009); the PPI topography in the right SMC of the current study was demonstrated through global connectivity from left and right hippocampal seeds reflecting interactions between adjacent fingers, applied to 4mm isotropic images. Despite differences in voxel size and hemisphere, the two methods provide good correspondence in topographical organization.

Fig 1

Fig 2

bioRxiv preprint doi: <https://doi.org/10.1101/479436>; this version posted November 26, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Fig 3