1 **Individual Behavioral and Physiological Responses During Different** 2 **Experimental Situations – Consistency over Time and Effects of** 3 **Context**

4 Alexandra Safryghin, Denise V. Hebesberger & Claudia A.F. Wascher*

5 Behavioural Ecology Research Group, School of Life Sciences, Anglia Ruskin University,
6 United Kingdom

7

8 *Corresponding author at: Claudia A.F. Wascher

9 Email address: claudia.wascher@anglia.ac.uk

10

11

12

Abstract

In a number of species, consistent behavioral differences between individuals have been described in standardized tests, *e.g.* novel object exploration, open field test. Different behavioral expressions are reflective of different coping strategies of individuals in stressful situations. A causal link between behavioral responses and the activation of the physiological stress response is assumed but not thoroughly studied. Also, most standard paradigms investigating individual behavioral differences, are framed in a fearful context, therefore the present study aimed to add a test in a more positive context, the feeding context. We assessed individual differences in physiological (heart rate, HR) and behavioral responses (presence or absence of pawing, startle response, defecation, snorting) of twenty domestic horses (*Equus caballus*) in two behavioral experiments, a novel object presentation and a pre-feeding excitement test. Experiments were conducted twice, in summer and autumn. Both experiments caused higher mean HR in the first ten seconds after stimulus presentation compared to a control condition, but mean HR did not differ between the experimental conditions. Interestingly, in the novel object experiment, horses displaying stress-related behaviors during the experiments also showed a significantly higher HR increase compared to horses which did not display any stress-related behaviors, reflecting a correlation between behavioral and physiological responses to the novel object. On the contrary, in the pre-feeding experiments, horses that showed fewer behavioral responses had a greater HR increase, indicating the physiological response being due to emotional arousal and not behavioral activity. Moreover, HR response to experimental situations varied significantly between individuals, and although we found HR to be significantly repeatable across experiments, repeatability indices were low. In conclusion, our findings show that horses' behavioral and physiological responses differed between test situations and that high

37 emotional reactivity, shown via mean HR and HR increase, is not always displayed
38 behaviorally.

39

40 Key words: individual variation, heart rate, novel object, pre-feeding excitement, domestic
41 horses, *Equus caballus*

42

43

44 **Introduction**

45 The perception of a potential threat to homeostasis, caused by extrinsic or intrinsic
46 stimuli (stressors) results in the activation of the physiological stress response in animals
47 (Chrousus and Gold 1992; Moberg 1985). Regardless of the intensity of the stressor,
48 individuals respond both behaviorally and physiologically, and these responses aim to
49 counteract the effects of the stimuli and to re-establish homeostasis (Chrousus and Gold
50 1992). Physiological activation in response to a stressor causes a fast release of
51 norepinephrine in the brain, triggering the activation of the sympatho-adreno-medullary
52 (SAM) axis, and the hypothalamic-pituitary-adrenal (HPA) axis (Goldstein 2010). As a
53 consequence, heart rate (HR) and glucocorticoid levels increase (Elia et al. 2010; McGreevy
54 et al. 2005).

55
56 The way an individual perceives a stressor can significantly influence its
57 responsiveness (Moberg 1985). Indeed, both endogenous and exogenous factors have been
58 shown to modulate an animal's sensitivity to a perceived threat, which results in individual
59 differences in responsiveness within the same species (Gosling 2001). For example, the
60 presence of constant environmental stressors can lead to a desensitization of the threat-
61 perception, or learned helplessness, which can cause a reduction of individual responses to a
62 stressor (Ellis et al. 2014). Furthermore, early differences in environmental stimuli, such as
63 differences in maternal investment, may cause individuals to follow diverse trajectories,
64 shaping their behavioral repertoire and physiological responsiveness in different manners
65 (Claessens et al. 2011; Stamps 2003). For example, Meany (2001) showed how rat pups
66 raised by low-licking mothers are more fearful and more sensitive to stress in adulthood
67 compared to pups reared by high-licking mothers. Moreover, animals can be bred to either
68 show high or low responsiveness to stressors, indicating that individual differences in stress-

responsiveness are heritable (Carere et al. 2003; Flaherty and Rowan 1989). Studies on great tits (*Parus major*) have shown individual physiological stress responses to be related to differences in exploration strategies (Carere and van Oers 2004) and heritable throughout four generations (Drent et al. 2003). If individual differences in physiological and/or behavioral responses are found to be relatively stable across time and contexts, they are regarded as temperamental traits (Cockrem 2007; Goldsmith et al. 1987; Lansade et al. 2008; Le Scolan et al. 1997).

Temperamental traits have been found in both vertebrate and invertebrate taxa (Carere et al. 2003; Kralj-Fišer et al. 2010; Momozawa et al. 2003; Riechert and Hedrick 1993; Verbeek et al. 1994; While et al. 2009; Øverli et al. 2006). Studies have shown that individual differences along the shy-bold axis are linked with aggressiveness (Kralj-Fišer et al. 2010), exploratory behavior (Sibbald et al. 2009), neophobia (Momozawa et al. 2003), and physiological responsiveness, such as hormonal (Carere et al. 2003), and HR modulation (Kralj-Fišer et al. 2010). Heart rate presents a valid indicator of emotional arousal, defined as an internal state, which is triggered by specific extrinsic or intrinsic stimuli (Anderson and Adolphs 2014). Arousal can range between the subject being calm – low arousal, and excited – high arousal, as well as the experience being of positive or negative valence (Russell 1980). Briefer et al. (2015) have studied the effect of emotional arousal and valence on the physiological and behavioral response in goats, showing both, positive (feeding) as well as negative (frustration, isolation) emotional context to cause a significant physiological response compared to a control situation.

An evaluation of individual differences in emotional arousal in response to situations of positive emotional valence or the relationship of activation of the physiological stress

94 response in situations of different emotional valence is still lacking. In the present study, we
 95 aim at investigating individual differences in emotional arousal in response to two
 96 experimental paradigms, a novel object exposure and a test of pre-feeding excitement. So far,
 97 most studies investigating temperamental traits in non-human animals have tested those in
 98 fearful contexts, *e.g.* novel object exploration or open field tasks (Dall and Griffith 2014;
 99 Smith and Blumstein 2008). Thereby we aimed to investigate physiological and behavioral
 100 responses in a fearful (novel object) and anticipatory (pre-feeding excitement) context.

101

102 In horses (*Equus caballus*), diverse factors influencing individual differences in
 103 behavior and physiology have been identified in terms of experience, such as habituation
 104 (Leiner and Fendt 2011), diet (Bulmer et al. 2015), handling (Visser et al. 2002), and
 105 maternal behavior (Haupt and Hintz 1983). Similarly, breed has also been found to strongly
 106 influence individual reactivity, suggesting a relationship between individual responsiveness
 107 and the heritability of traits (Hausberger et al. 2004; Lloyd et al. 2008). Further studies on
 108 equine temperament have focused on the assessment of horse responsiveness to different
 109 stimuli such as diverse environmental conditions (McCall et al. 2006; Schmidt et al. 2010a;
 110 Schmidt et al. 2010b), novel situations (Ellis et al. 2014; Fureix et al. 2009; Leiner and Fendt
 111 2011; Visser et al. 2001; Visser et al. 2002) human interactions in terms of both handling
 112 (Ellis et al. 2014; Fureix et al. 2009; König von Brostel et al. 2011) and riding (Visser et al.
 113 2008) and have shown how an individual's response to threat – or 'fearfulness' – is stable
 114 across time (Lansade et al. 2008; Visser et al. 2001; Visser et al. 2003). However, contrasting
 115 results have been found according to the relationship between HR and behavioral parameters
 116 (Christensen et al. 2005; Lansade et al. 2008; Momozawa et al. 2003). For instance, only
 117 Momozawa et al. (2003) found behavioral correlations with HR in their fear-inducing

118 experiments; results which were not supported in subsequent research (Christensen et al.
119 2005; Lansade et al. 2008).

120

121 In the present study, we aim at gaining further understanding of the relationship
122 between individual differences in behavioral and physiological reactivity in horses. Heart rate
123 represents a standardized, objective and non-invasive measure to infer an individual's
124 emotional arousal level, readily available to inexperienced behavioral observers (Lansade et
125 al. 2008; Visser et al. 2003). We question whether physiological responses during a novel
126 object presentation and pre-feeding test are caused by behavioral changes or purely emotional
127 arousal, in the absence of behavioral activity. Furthermore, we ask whether behavioral and
128 physiological responses are stable across time and contexts. We expect that horses show a
129 greater physiological reaction to a fear-inducing situation such as being exposed to a novel
130 object, compared to the anticipatory pre-feeding exposure.

Methods

Animals and housing

The study was conducted at the equine yard of the College of West Anglia (United Kingdom) between July and November 2017. The research was conducted on 20 horses which were individually stabled in loose boxes. Five of the 20 horses were tested only in one of the two experimental conditions, two solely for pre-feeding excitement and three only for novel object test, due to their lack of availability during testing periods (Table 1). The sample included 14 geldings (Age: mean 11.8 ± 3.8 years (yrs), range 6 - 18 yrs) and 6 mares (Age: mean 11.8 ± 2.6 yrs, range 10-17 yrs) of diverse breeds, use, and training experiences (Table 1). The horses were fed twice a day: once in the morning (0800-0830) and once in the afternoon (1500-1600). Water was available *ad libitum*, and feces were removed from the stables after the horses were fed. In the late afternoon (1600-1700), some of the horses were turned out in the paddock for the night.

Experimental design

We conducted two experimental tests – the pre-feeding excitement and the novel object test, presented in random order, with one trial per horse and repeated in summer (July/August) and in autumn (October/November). Behavior was recorded by video camera (Canon Legria HF R56), and HR was recorded using a Polar V800 system belt, placed around the chest of the horses. The belt consists of an electrode belt with a built-in transmitter, connected via Bluetooth to a wristwatch (receiver). To optimize the contact between the belt and the skin, both the coat of the horse in the interested area and the belt were wetted. The receiver was placed at the stable entrance or inside the stable. Prior to each test, an adjustment period of 5 minutes was allowed to let the horse habituate to the belt.

The pre-feeding excitement test was conducted during morning feeds (0800-0830)

and started with the horses being shown a bucket containing their individual mix of hard feed on the floor outside the stable, while the other horses were fed. The horses' physiological and behavioral responses were recorded for five minutes. Thereafter, the horses were fed by placing the feeding bucket inside the box and their behavioral and physiological responses were measured for the following ten minutes.

In the novel object test, the horses were exposed to two of three different objects in their stable. The first object was formed of a main cylindrical hard body (approximately 30 cm in length) filled with gravel which was fixed to a soft foam rubber ball (about 15 cm in diameter) and covered in fabric. The second object was formed of two cylindrical plastic tubes fixed together to form an 'x'. Similar to the first object, the cylinders were approximately 30cm in length, filled with gravel and covered with fabric. In addition, twelve tennis balls of different colors and materials were pierced and attached to 4 strings (3 balls per string) of approximately 50 cm in length. These were then tied to the main body of the object and left hanging. The third object was an inflatable guitar of approximately one meter in length. All objects were attached to a string, around four-meters-long, to allow their retrieval from the stable.

The object assigned to each individual was randomized for each season as well as the order of the horses tested. To avoid the horses seeing the object before testing, the objects were covered from sight when carried around the yard. The novel object tests took place between the hours of 0900 and 1300 and between 1500 and 1800 when the yard was quiet, and the horses were fed. The testing procedure was based on the one used by Dai et al. (2015) and Górecka-Bruzda et al. (2011) and adapted for the present experiment. A novel object was placed over the box entrance, with the cord hanging over the stable door to keep the object at the height of approximately one meter. The object was kept in this position for the following

five minutes and was then dropped to the floor (the objects filled with gravel created a muffled noise). The horse reaction was recorded for the following five minutes. Thereafter, the object was removed from the stable, whilst behavioral monitoring and HR measurement continued for another 15 minutes.

Ethical statement

All applied methods were non-invasive, and the experimental procedure was approved by Anglia Ruskin University's Departmental Research Ethics Panel.

Data processing

Raw HR data were purged with a moving average filter to remove biologically implausible outlier values. The following HR variables were calculated: (1) mean HR in beats per minute (bpm) for the ten seconds preceding and following the food presentation, as well as preceding (control) and following the presentation, drop and removal of the object; (2) HR increase in bpm following the food presentation and novel object presentation, drop, and removal, calculated as difference between maximum value and three seconds average HR before the presentation of the stimuli.

Behavioral responses of the horses were analyzed from videos using SOLOMON Coder v. beta 17.03.22 (András Péter, www.solomoncoder.com). The behavior of the individuals was analyzed for the five minutes prior to the presentation of the hard feed for the pre-feeding excitement. For the novel object task, the five minutes following the presentation and drop of the object and the two minutes following its removal were analyzed. Behavior was recorded as continuous variables, *e.g.*, feeding as duration of behavior in s per observation period, or frequency of behavior per observation period, *e.g.*, snorting. The classification of the individual in high and low behavioral respondents was based on the

frequency and duration of vocalizations, pawing behavior, startle response and defecation that the individuals performed during the experimental settings (Table 2).

Statistical Analysis

All data were analyzed using R version 3.4.3 (R Core Team, 2017; RStudio Team 2016). Generalized linear mixed models (GLMM) were conducted with the additional packages ‘glmmADMB’ (Skaug et al. 2016). For the comparison of HR, two generalized linear models were calculated. Experiment (pre-feeding excitement versus novel object), test period and behavioral category (high versus low respondents) were considered as fixed factors in both models, together with the interaction between behavioral category and experimental condition. For the purpose of the analysis, the different conditions of each experiment were individually included in the dataset, resulting in horses having multiple values per each experimental condition. The response variable was assigned to the ten second average HR (GLMM1) and the HR increase (GLMM2). Tukey test for multiple comparisons was chosen to gain further understanding of the effect of the categories of the fixed factors in the models. Specifically, the ‘multcomp’ (Hothorn et al. 2008) package was used to conduct the post hoc analysis. Moreover, we used a likelihood ratio test to compare models fit according to presence or absence of the individual random effect. Also, individual repeatability of the physiological and behavioral measurements was assessed using the ‘rptR’ package (Stoffel et al. 2017). In particular, we assessed the repeatability of the ten second average HR and HR increase with 1000 permutations for the physiological reactivity data collected for the control, novel object and pre-feeding conditions. The significance level was set at $\alpha = 0.05$.

Results

Seasonal variation and behavioral categorization

Horses had a significantly higher mean HR during summer compared to autumn (GLMM1: $z = 2.713$, $p = 0.007$), but HR increase was not significantly different between the seasons (GLMM2: $z = -0.94$, $p = 0.348$). Out of the 20 horses tested, only four individuals were consistently categorized as high behavioral responders and one individual as low behavioral responder, in both experiments (novel object, pre-feeding excitement) and seasons (summer and autumn; see Table 1 for details).

Physiological and behavioral responses to experimental situations

Average HR of the horses was significantly higher during the novel object experiment compared to the control period (Tukey: $z = 5.205$, $p < 0.001$; Figure 1A, Figure 2) and tended to be higher during the pre-feeding excitement compared to the control period (Tukey: $z = 2.197$, $p = 0.0662$; Figure 1A, Figure 2). Heart rate between novel object and pre-feeding excitement was not significantly different (Tukey: mean: $z = -1.887$, $p = 0.133$; Figure 1A, Figure 2), despite horses showing a significantly lower HR increase in the pre-feeding condition (GLMM2: $z = -3.59$, $p < 0.001$; Figure 1B). We found a significant interaction between behavior and experiment affecting HR. Mean HR during the novel object experiment was significantly higher in the group of horses showing a high behavioral response compared to horses showing a low behavioral response (GLMM1: $z = -3.139$, $p = 0.002$; Figure 1A). This pattern reversed regarding HR increase in the pre-feeding experiment, with the horses showing a low behavioral response having a higher HR increased compared to individuals with a high behavioral response (GLMM2: $z = 3.56$, $p < 0.001$; Figure 1B). Finally, the models including individual identity as random factor had a significantly better fit compared to the models without the random effect (ANOVA: mean

251 HR: Deviance = 14.632, df = 1, p = 0.001; HR increase: Deviance = 7.984, df = 1, p = 0.005),
 252 indicating that the random effect also significantly variance in the data.

253

254 *Repeatability*

255 Individual horses' HR was significantly repeatable across experiments. However, the
 256 repeatability indices were low for both parameters tested, with the highest repeatability score
 257 recorded for the HR increase ($R = 0.265$, CI 95% [0.07, 0.442], $p < 0.001$); followed by mean
 258 HR ($R = 0.201$, CI 95% [0.06, 0.343], $p < 0.001$).

259

260

Discussion

In the present study, we investigated individual behavioral and physiological responses of horses during two experimental procedures, a novel object experiment and a pre-feeding excitement test. Surprisingly, we found rather little behavioral consistency across tests and only a limited number of individuals responded similarly across contexts and seasons, which would have been expected when behavioral responses in experimental tests are indicative of temperamental traits. In contrast to behavior, heart rate (HR) was significantly repeatable across experiments, although it has to be pointed out that the repeatability indices were low. This indicates that physiological responses to experiments were consistent over time and contexts, as expected for temperamental traits. Repeatable individual variation in behavior, *i.e.* personality has been very much in focus of scientific research in recent years and was described in a vast variety of species, including horses (Grajfoner et al. 2010; Momozawa et al. 2005), however most of these studies are focusing on behavior rather than underlying physiological processes. For example, Grajfoner et al. (2010) compared individual horse behavioral ratings between high and low performers, showing how the combination of multiple traits, such as a horse being “nice”, “patient”, “easy to handle”, shape the perceived personality of horses. On the contrary, only a few studies investigated individual differences in physiological responses and their relationships with behavioral reactivity in non-human animals (Briefer et al. 2015; Ellis et al. 2014).

We described context dependent links between behavioral and physiological response. The two conducted experiments differed regarding their context, with the novel object experiment being considered as a fear-inducing situation (Dalmau et al. 2009; Lansade et al. 2008; Leiner and Fendt 2011), whereas the pre-feeding excitement test presents an anticipatory context (Mahnhardt et al. 2014). In the novel object experiment, a classic

temperament test in non-human animals' personality research (Carter et al. 2013), individuals who were behaviorally active also showed a higher average HR. This is especially relevant as the presentation of a novel object can be seen as a fear-inducing situation (Leiner and Fendt 2011). The results of the present study confirm that increased physical activity resulting from the performance of stress-related behaviors (*e.g.*, startle response) correlates with higher mean HR compared to individuals not responding behaviourally in the experiment. It is difficult to conclude whether the physiological activation is caused by increased emotional arousal, *i.e.* the perception of fear or the increased physical activity, *e.g.* locomotion (von Borell et al. 2007; Visser et al. 2002).

A different pattern became apparent during the pre-feeding excitement task. Here, individuals lacking a behavioral response during the experiment showed a higher HR increase. This indicates that emotional arousal, but not physical activity, accounted for the increase in HR. Horses that were classified as highly behaviorally responsive during the pre-feeding experiments had an undetectable increase in HR, reflecting that physical activity may have increased their HRs before the presentation of the feed. Once the feed was presented their arousal level may not have changed in amplitude, as no significant HR increase was recorded, but possibly in valence, from negative to positive. Effects of emotional arousal on physiological responses have already been identified in other non-human animals. For example, in the study by Wascher et al. (2008), immobile greylag geese (*Anser anser*) watching aggressive interaction between conspecifics showed a significantly higher increase in HR compared to geese watching non-social interactions. Moreover, an increase in physiological reactivity resulting solely from emotional arousal was also identified in guide dogs (Fallani et al. 2007).

Our findings show that HR in the control situations did not necessarily predict HR during the experiments and horses differed in the strength of their responses, with some even decreasing HR. Such variation, together with the repeatability of the physiological reactivity, reflected how behavioral responses of horses do not necessarily predict physiological reactions during a novel object and pre-feeding excitement test. This is in line with previous studies on horses describing that individual classified as calmer had higher HR compared to more excited individuals despite showing less behavioral signs of stress when tested for pre-feeding reactivity (Ellis et al. 2014), as well as with other studies on cattle (*e.g.*, Christensen et al. 2005; Jezierski et al. 1999; Lansade et al. 2008; Welp et al. 2004).

Classical models regarding individual differences in behavior and physiology, assume them to be associated with each other to form different coping styles (Koolhaas et al. 1999). However, evidence for independent modulation of the hypothalamic-pituitary-adrenal axis (Boulton et al. 2015; Dosmann et al. 2015; Ferrari et al. 2013), the sympatho-adreno-medullary axis (Qu et al. 2018), and behavioral traits have been recently accumulating. Our study provides further evidence that HR and behavior might be regulated independently.

To conclude, our study illustrates the importance of accounting for context when studying individual differences in non-human animals, and we suggest that the inclusion of tests in non-fear related contexts is desirable. Further, we point at the importance of studying interconnectedness and independencies between physiological and behavioral reactions.

Acknowledgements

We would like to thank Elizabeth Ragg and Nikki Bentley from the College of West Anglia for granting us access to their facilities and horses to conduct our experiments. Furthermore,

we would like to thank all members of staff for helping us to facilitate our research on site.

Finally, we would also like to thank Irene Susini for discussion and comments on the

manuscript.

Author Contribution statement

A.S., D.V.H. & C.A.F.W. designed the experiments

A.S. & D.V.H. conducted the experiments

A.S. & C.A.F.W. analyzed the data

A.S., D.V.H. & C.A.F.W. wrote the paper

Conflict of interest

The authors declare that they have no competing interests.

References

Anderson, D. J. and Adolphs, R., 2014. A framework for studying emotions across species.

Cell, 157(1), pp. 187-200. doi: 10.1016/j.cell.2014.03.003

Boulton, K., Couto, E., Grimmer, A. J., Earley, R. L., Canario, A. V. M., Wilson, A. J., et al.,

2015. How integrated are behavioral and endocrine stress response traits? A repeated

measures approach to testing the stress-coping style model. *Ecology and Evolution*,

5(3), pp. 618–633. doi: 10.1002/ece3.1395

- 355 Briefer, E. F., Tettamanti, F. and McElligott, A. G., 2015. Emotions in goats: Mapping
356 physiological, behavioural and vocal profiles. *Animal Behaviour*, 99, pp. 131-143.
357 doi: 10.1016/j.anbehav.2014.11.002
- 358 Bulmer, L., McBride, S., Williams, K. and Murray, J., 2015. The effects of a high-starch or
359 high-fibre diet on equine reactivity and handling behaviour. *Applied Animal*
360 *Behaviour Science*, 165, pp. 95-102. doi: 10.1016/j.applanim.2015.01.008
- 361 Carere, C., Groothuis, T. G., Möstl, E., Daan, S. and Koolhaas, J. M., 2003. Fecal
362 corticosteroids in a territorial bird selected for different personalities: daily rhythm
363 and the response to social stress. *Hormones and Behavior*, 43(5), pp. 540-548. doi:
364 10.1016/S0018-506X(03)00065-5
- 365 Carere, C. and van Oers, K., 2004. Shy and bold great tits (*Parus major*): body temperature
366 and breath rate in response to handling stress. *Psychology & Behavior*, 82, pp. 905-
367 912. doi: 10.1016/j.physbeh.2004.07.009
- 368 Carter, A. J., Feeney, W. E., Marshall, H. H., Cowlshaw, G., and Heinsohn, R., 2013.
369 Animal personality: What are behavioural ecologists measuring? *Biological Reviews*,
370 88(2), pp.465–475. doi: 10.1111/brv.12007
- 371 Christensen, J. W., Keeling, L. J. and Nielsen, B. L., 2005. Responses of horses to novel
372 visual, olfactory and auditory stimuli. *Applied Animal Behaviour Science*, 93(1-2), pp.
373 53-65. doi: 10.1016/j.applanim.2005.06.017
- 374 Chrousos, G. P. and Gold, P. W., 1992. The concepts of stress and stress system disorders.
375 Overview of physical and behavioral homeostasis. *Journal of the American Medical*
376 *Association*, 267(9), pp. 1244-52. doi: 10.1001/jama.1992.03480090092034

- 377 Claessens, S. E., Daskalakis, N.P., van der Veen, R., Oitzl, M. S., de Kloet, E. R. and
378 Champagne, D. L., 2011. Development of individual differences in stress
379 responsiveness: an overview of factors mediating the outcome of early life
380 experiences. *Psychopharmacology*, 214(1), pp. 141-154. doi: 10.1007/s00213-010-
381 2118-y
- 382 Cockrem, J. F., 2007. Stress, corticosterone responses and avian personalities. *Journal of*
383 *Ornithology*, 148(2), pp. 169-178. doi: 10.1007/s10336-007-0175-8
- 384 Dai, F., Cogi, N. H., Heinzl, E. U. L., Dalla Costa, E., Canali, E. and Minero, M., 2015.
385 Validation of a fear test in sport horses using infrared thermography. *Journal of*
386 *Veterinary Behavior-clinical Applications and Research*, 10(2), pp. 128-136. doi:
387 10.1016/j.jveb.2014.12.001
- 388 Dall, S. R. X. and Griffith, S. C., 2014. An empiricist guide to animal personality variation in
389 ecology and evolution. *Frontiers in Ecology and Evolution*, 2(3). doi:
390 10.3389/fevo.2014.00003
- 391 Dalmau, A., Fabrega, E., and Velarde, A., 2009. Fear assessment in pigs exposed to a novel
392 object test. *Applied Animal Behaviour Science*, 117(3–4), pp.173–180. doi:
393 10.1016/j.applanim.2008.12.014
- 394 Dosmann, A. J., Brooks, K. C., and Mateo, J. M., 2015. Within-individual correlations reveal
395 link between a behavioral syndrome, condition and cortisol in free-ranging Belding's
396 ground squirrels. *Ethology*, 121(2), pp.125–134. doi: 10.1111/eth.12320
- 397 Drent, P. J., van Oers, K. and van Noordwijk, A. J., 2003. Realized heritability of
398 personalities in the great tit (*Parus major*). *Proceedings of the Royal Society of*

- 399 *London. Series B: Biological Sciences*, 270(1510), pp. 45-51. doi:
400 10.1098/rspb.2002.2168
- 401 Elia, J. B., Erb, H. N. and Houpt, K. A., 2010. Motivation for hay: Effects of a pelleted diet
402 on behavior and physiology of horses. *Physiology & Behavior*, 101(5), pp. 623-627.
403 doi: 10.1016/j.physbeh.2010.09.010
- 404 Ellis, A. D., Stephenson, M., Preece, M. and Harris, P., 2014. A novel approach to
405 systematically compare behavioural patterns between and within groups of horses.
406 *Applied Animal Behaviour Science*, 161, pp. 60-74. doi:
407 10.1016/j.applanim.2014.09.017
- 408 Fallani, G., Previde, E. P. and Valsecchi, P., 2007. Behavioral and physiological responses of
409 guide dogs to a situation of emotional distress. *Physiology & Behavior*, 90(4), pp.
410 648-655. doi: 10.1016/j.physbeh.2006.12.001
- 411 Ferrari, C., Pasquaretta, C., Carere, C., Cavallone, E., von Hardenberg, A., and Réale, D.,
412 2013. Testing for the presence of coping styles in a wild mammal. *Animal Behaviour*,
413 85(6), pp.1385–1396. doi: 10.1016/j.anbehav.2013.03.030
- 414 Flaherty, C. and Rowan, G., 1989. Rats (*Rattus norvegicus*) selectively bred to differ in
415 avoidance behavior also differ in response to novelty stress, in glycemic conditioning,
416 and in reward contrast. *Behavioral and Neural Biology*, 51(2), pp. 145-169. doi:
417 10.1016/S0163-1047(89)90782-6
- 418 Fureix, C., Pagès, M., Bon, R., Lassalle, J. M., Kuntz, P. and Gonzalez, G., 2009. A
419 preliminary study of the effects of handling type on horses' emotional reactivity and
420 the human-horse relationship. *Behavioural Processes*, 82(2), pp. 202-210. doi:
421 10.1016/j.beproc.2009.06.012

- 422 Goldsmith, H., Buss, A., Plomin, R., Rothbart, M., Thomas, A., Chess, S., et al., 1987.
423 Roundtable: What is temperament? Four approaches. *Child Development*, 58(2), p.
424 505-529. doi: 10.2307/1130527
- 425 Goldstein, D. S., 2010. Adrenal responses to stress. *Cellular and Molecular Neurobiology*,
426 30(8), pp. 1433-1440. doi: 10.1007/s10571-010-9606-9
- 427 Gosling, S. D., 2001. From mice to men: what can we learn about personality from animal
428 research? *Psychological Bulletin*, 127(1), pp. 45–86. doi: 10.1037/0033-
429 2909.127.1.45
- 430 Grajfoner, D. D., Austin, E. J. and Wemelsfelder, F., 2010. Horse personality profiles and
431 performance. *Journal of Veterinary Behavior-clinical Applications and Research*,
432 5(1), pp. 26-27. doi: 10.1016/j.jveb.2009.10.035
- 433 Górecka-Bruzda, A., Jastrzębska, E., Sosnowska, Z., Jaworski, Z., Jezierski, T. and
434 Chruszczewski, M. H., 2011. Reactivity to humans and fearfulness tests: Field
435 validation in Polish cold blood horses. *Applied Animal Behaviour Science*, 133(3), pp.
436 207-215. doi: 10.1016/j.applanim.2011.05.011
- 437 Hausberger, M., Bruderer, C., Le Scolan, N. and Pierre, J. S., 2004. Interplay between
438 environmental and genetic factors in temperament/personality traits in horses (*Equus*
439 *caballus*). *Journal of Comparative Psychology*, 118(4), pp. 434-446. doi:
440 10.1037/0735-7036.118.4.434
- 441 Hothorn, T., Bretz, F. and Westfall, P., 2008. Simultaneous inference in general parametric
442 models. *Biometrical Journal*, 50(3), pp.346-363. doi: 10.1002/bimj.200810425

- 443 Houpt, K. A. and Hintz, H., 1983. Some effects of maternal deprivation on maintenance
444 behavior, spatial relationships and responses to environmental novelty in foals.
445 *Applied Animal Ethology*, 76(2), pp. 297-307. doi: 10.1016/0304-3762(83)90002-0
- 446 Jezierski, T., Jaworski, Z. and Górecka- Bruzda, A., 1999. Effects of handling on behaviour
447 and heart rate in Konik horses: comparison of stable and forest reared youngstock,
448 *Applied Animal Behaviour Science*, 62(1), pp.1-11. doi: 10.1016/S0168-
449 1591(98)00209-3
- 450 König von Brostel, U., Euent, S., Graf, P., König, S. and Gauly, M., 2011. Equine behaviour
451 and heart rate in temperament tests with or without rider or handler. *Physiology &*
452 *Behavior*, 104(3), pp. 454-463. doi: 10.1016/j.physbeh.2011.05.010
- 453 Koolhaas, J. M., Korte, S. M., De Boer, S. F., Van Der Vegt, B. J., Van Reenen, C. G.,
454 Hopster, H. et al., 1999. Coping styles in animals: Current status in behavior and
455 stress- physiology. *Neuroscience and Biobehavioral Reviews*, 23(7), pp.925–935. doi:
456 10.1016/S0149-7634(99)00026-3
- 457 Kralj-Fišer, S., Weiß, B. M. and Kotrschal, K., 2010. Behavioural and physiological
458 correlates of personality in greylag geese (*Anser anser*). *Journal of Ethology*, 28(2),
459 pp. 363-370. doi: 10.1007/s10164-009-0197-1
- 460 Lansade, L., Bouissou, M. and Erhard, H. W., 2008. Fearfulness in horses: A temperament
461 trait stable across time and situations. *Applied Animal Behaviour Science*, 115(3), pp.
462 182-200. doi: 10.1016/j.applanim.2008.06.011
- 463 Le Scolan, N., Hausberger, M. and Wolff, A., 1997. Stability over situations in
464 temperamental traits of horses as revealed by experimental and scoring approaches.
465 *Behavioural Processes*, 41(3), pp. 257-266. doi: 10.1016/S0376-6357(97)00052-1

- 466 Leiner, L. and Fendt, M., 2011. Behavioural fear and heart rate responses of horses after
467 exposure to novel objects: Effects of habituation. *Applied Animal Behaviour*
468 *Science*, 131(3), pp. 104-109. doi: 10.1016/j.applanim.2011.02.004
- 469 Lloyd, A. S., Martin, J. E., Bornett-Gauci, H. L. and Wilkinson, R. G., 2008. Horse
470 personality: Variation between breeds. *Applied Animal Behaviour Science*, 112(3), pp.
471 369-383. doi: 10.1016/j.applanim.2007.08.010
- 472 Mahnhardt, S., Brietzke, J., Kanitz, E., Schön, P. C., Tuchscherer, A., Gimsa, U. et al., 2014.
473 Anticipation and frequency of feeding affect heart reactions in domestic pigs. *Journal*
474 *of Animal Science*, 92(11), pp.4878–4887. doi: 10.2527/jas.2014-7752
- 475 McCall, C. A., Hall, S., McElhenney, W. H. and Cummins, K. A., 2006. Evaluation and
476 comparison of four methods of ranking horses based on reactivity. *Applied Animal*
477 *Behaviour Science*, 96(1-2), pp. 115-127. doi: 10.1016/j.applanim.2005.04.021
- 478 McGreevy, P. D., Righetti, J. and Thomson, P. C., 2005. The reinforcing value of physical
479 contact and the effect on canine heart rate of grooming in different anatomical areas.
480 *Anthrozoos*, 18(3), pp. 236-244.
- 481 Meaney, M. J., 2001. Maternal care, gene expression, and the transmission of individual
482 differences in stress reactivity across generations. *Annual Review of Neuroscience*. 24.
483 pp. 1161-1192. doi: 10.1146/annurev.neuro.24.1.1161.
- 484 Moberg, G. P., 1985. Biological response to stress: key to assessment of animal well-being?
485 In: G. P. Moberg, ed. *Animal Stress*. New York: Springer.
- 486 Momozawa, Y., Ono, K., Sato, F., Kikusui, T., Takeuchi, Y., Mori, Y. et al., 2003.
487 Assessment of equine temperament by a questionnaire survey to caretakers and

- 488 evaluation of its reliability by simultaneous behavior test. *Applied Animal Behaviour*
489 *Science*, 84(2), pp. 127-138. doi: 10.1016/j.applanim.2003.08.001
- 490 Momozawa, Y., Kusunose, R., Kikusui, T., Takeuchi, Y. and Mori, Y., 2005. Assessment of
491 equine temperament questionnaire by comparing factor structure between two
492 separate surveys. *Applied Animal Behaviour Science*, 92(1-2), pp. 77-84. doi:
493 10.1016/j.applanim.2004.11.006
- 494 Qu, J., Fletcher, Q. E., Réale, D., Li, W., and Zhang, Y., 2018. Independence between coping
495 style and stress reactivity in plateau pika. *Physiology and Behavior*, 197, pp.1–8. doi:
496 doi: 10.1016/j.physbeh.2018.09.007
- 497 Riechert, S. E. and Hedrick, A. V., 1993. A test for correlations among fitness-linked
498 behavioural traits in the spider *Agelenopsis aperta* (*Araneae, agelenidae*). *Animal*
499 *Behaviour*, 46(4), pp. 669-675. doi: 10.1006/anbe.1993.1243
- 500 Russell, J. A., 1980. A circumplex model of affect. *Journal of Personality and Social*
501 *Psychology*, 39(6), pp. 1161-1178. doi: 10.1037/h0077714
- 502 Schmidt, A., Biau, S., Möstl, E., Becker-Birck, M., Morillon, B., Aurich, J. et al. 2010a.
503 Changes in cortisol release and heart rate variability in sport horses during long-
504 distance road transport. *Domestic Animal Endocrinology*, 38(3), pp. 179-189. doi:
505 10.1016/j.domaniend.2009.10.002
- 506 Schmidt, A., Möstl, E., Wehnert, C., Aurich, J., Müller, J. and Aurich, C., 2010b. Cortisol
507 release and heart rate variability in horses during road transport. *Hormones and*
508 *Behavior*, 57(2), pp. 209-215. doi: 10.1016/j.yhbeh.2009.11.003

- 509 Sibbald, A. M., Erhard, H. W., McLeod, J. E. and Hooper, R. J., 2009. Individual personality
510 and the spatial distribution of groups of grazing animals: an example with sheep.
511 *Behavioural Processes*, 82(3), pp. 319-326. doi: 10.1016/j.beproc.2009.07.011
- 512 Skaug H., Fournier D., Bolker B., Magnusson, A. and Nielsen, A., 2016. _Generalized Linear
513 Mixed Models using 'AD Model Builder'_. R package version 0.8.3.3.
- 514 Smith, B. R. and Blumstein, D. T., 2008. Fitness consequences of personality: a meta-
515 analysis. *Behavioural Ecology*, 19(2), pp. 448-455. doi: 10.1093/beheco/arm144
- 516 Stamps, J. A., 2003. Behavioural processes affecting development: Tinbergen's fourth
517 question comes of age. *Animal Behaviour*, 66(1), pp. 1-13. doi:
518 10.1006/anbe.2003.2180
- 519 Stoffel, M. A., Nakagawa, S. and Schielzeth, H., 2017. rptR: repeatability estimation and
520 variance decomposition by generalized linear mixed-effects models. *Methods in*
521 *Ecology and Evolution*, 8(11), pp. 1639-1644. doi: 10.1111/2041-210X.12797
- 522 R Core Team, 2017. R: A language and environment for statistical computing. R Foundation
523 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 524 RStudio Team, 2016. RStudio: Integrated Development for R. RStudio, Inc., Boston, MA
525 URL <http://www.rstudio.com/>.
- 526 Verbeek, M. E., Drent, P. J. and Wiepkema, P., 1994. Consistent individual differences in
527 early exploratory behaviour of male great tits. *Animal Behaviour*, 48(5), pp. 1113-
528 1121. doi: 10.1006/anbe.1994.1344

- 529 Visser, E. K., Ellis, A. D. and van Reenen, C. G., 2008. The effect of two different housing
530 conditions on the welfare of young horses stabled for the first time. *Applied Animal*
531 *Behaviour Science*, 114(3), pp. 521-533. doi: 10.1016/j.applanim.2008.03.003
- 532 Visser, E. K., van Reenen, C. G., Hopster, H., Schilder, M. B. H., Knaap, J., Barneveld, A. et
533 al., 2001. Quantifying aspects of young horses' temperament: consistency of
534 behavioural variables. *Applied Animal Behaviour Science*, 74(4), pp. 241-258. doi:
535 10.1016/S0168-1591(01)00177-0
- 536 Visser, E. K., van Reenen, C. G., Schilder, M. B. H., Barneveld, A. and Blokhuis, H. J., 2003.
537 Learning performances in young horses using two different learning tests. *Applied*
538 *Animal Behaviour Science*, 80(4), pp. 311-326. doi: 10.1016/S0168-1591(02)00235-6
- 539 Visser, E. K., van Reenen, C. G., van der Werf, J., Schilder, M. B. H., Knaap, J., Barneveld,
540 A. et al., 2002. Heart rate and heart rate variability during a novel object test and a
541 handling test in young horses. *Physiology & Behavior*, 76(2), pp. 289-296. doi:
542 10.1016/S0031-9384(02)00698-4
- 543 von Borell, E., Langbein, J., Després, G., Hansen, S., Leterrier, C., Marchant-Forde, J. et al.,
544 2007. Heart rate variability as a measure of autonomic regulation of cardiac activity
545 for assessing stress and welfare in farm animals: A review. *Physiology & Behavior*,
546 92(3), pp. 293-316. doi: 10.1016/j.physbeh.2007.01.007
- 547 Wascher, C. A. F., Scheiber, I. B. R., and Kotrschal, K., 2008. Heart rate modulation in
548 bystanding geese watching social and non-social events. *Proceedings of the Royal*
549 *Society B: Biological Sciences*, 275(1643), pp.1653-1659. doi:
550 10.1098/rspb.2008.0146

- 551 Welp, T., Rushen, J., & Kramer, D. L., Festa-Bianchet, M. and de Passille, A. M., 2004.
 552 Vigilance as a measure of fear in dairy cattle. *Applied Animal Behaviour Science*.
 553 87(1-2), pp. 1-13. doi: 10.1016/j.applanim.2003.12.013
- 554 While, G. M., Sinn, D. L. and Wapstra, E., 2009. Female aggression predicts mode of
 555 paternity acquisition in a social lizard. *Proceedings of the Royal Society B: Biological*
 556 *Sciences*, 276(1664), pp. 2021-2029. doi: 10.1098/rspb.2008.1926
- 557 Øverli, Ø., Sørensen, C. and Nilsson, G., 2006. Behavioral indicators of stress-coping style in
 558 rainbow trout: Do males and females react differently to novelty? *Physiology &*
 559 *Behavior*, 87(3), pp. 506-512. doi: 10.1016/j.physbeh.2005.11.012

Table 1. Age in years, sex and breed of the 20 horses tested for this study. Mean heart rate and standard deviation during the different experimental situations as well as whether the horses were categorized as high (orange) or low (blue) behavioral responders.

ID	Age	Sex	Breed	NO 1	NO2	PF1	PF2
1	13	Gelding	Cob	NA	NA	48.29 ± 9.89	57.14 ± 12.43
2	14	Gelding	Welsh Pony	51.47 ± 8.87	51.21 ± 5.66	67.27 ± 12.41	54.42 ± 8.01
3	17	Mare	Gypsy Cob	54.77 ± 9.71	45.16 ± 3.44	NA	NA
4	8	Gelding	Cob	44.54 ± 1.56	38.37 ± 0.40	41.08 ± 5.62	33.69 ± 2.48
5	14	Gelding	Cob	46.82 ± 2.69	54.67 ± 3.85	51.29 ± 9.89	55.47 ± 10.65
6	9	Gelding	KWPN	43.87 ± 6.93	45.93 ± 12.30	39.01 ± 2.80	39.95 ± 7.07
7	10	Mare	Welsh Crossbred	62.40 ± 12.40	43.16 ± 1.92	48.06 ± 5.75	65.76 ± 14.01
8	16	Gelding	Welsh Crossbred	71.47 ± 18.82	44.41 ± 2.40	68.63 ± 9.19	53.98 ± 13.29
9	13	Gelding	Cob	NA	NA	55.57 ± 5.06	50.73 ± 1.53
10	12	Gelding	KWPN	54.30 ± 5.87	40.24 ± 8.05	NA	NA
11	16	Gelding	Shire X Warmblood	83.26 ± 20.00	80.03 ± 29.02	60.78 ± 7.80	46.88 ± 6.05
12	11	Mare	Appaloosa X Cob	42.84 ± 1.54	41.92 ± 8.30	60.89 ± 3.36	48.30 ± 12.39
13	11	Gelding	Welsh Section A	62.73 ± 20.70	55.29 ± 25.91	NA	NA
14	12	Mare	Thoroughbred X Cob	43.93 ± 3.47	47.27 ± 4.77	61.14 ± 7.16	46.12 ± 3.40
15	18	Gelding	Thoroughbred	44.43 ± 8.47	36.94 ± 4.96	41.58 ± 1.78	33.59 ± 1.90
16	10	Mare	Cob	44.28 ± 12.98	66.93 ± 15.66	42.84 ± 4.98	44.89 ± 7.01
17	6	Gelding	KWPN	36.67 ± 2.59	31.41 ± 1.52	42.10 ± 4.30	38.41 ± 2.57
18	9	Gelding	Irish Sports Pony	69.63 ± 15.71	62.12 ± 8.10	64.30 ± 12.74	36.77 ± 3.04
19	11	Mare	Warmblood	40.25 ± 4.11	42.52 ± 6.27	50.69 ± 17.53	38.71 ± 4.06
20	6	Gelding	Cob	44.22 ± 9.89	39.63 ± 4.86	53.74 ± 8.18	37.48 ± 3.31

Table 2. Description of behavioral categories and number of individuals per category, per season.

Condition	Value	Description	n	n
			(summer)	(autumn)
Pre-Feeding	Low	Absence or occasional vocalization (snort, neigh), no pawing behavior.	8	10
	High	Repetitive vocalization and/or pawing behavior.	9	7
Novel Object	Low	Total absence or occurrence of just one of the following behaviors: pawing at object, startle response, defecation, snorting.	9	14
	High	Presence of at least two of the following behaviors: pawing, startle response, defecation, snorting.	9	4

565

Figure 1. Effect of behavioral categorization on the mean heart rate (HR) (A) and HR increase (B) of the horses recorded during the study. Boxplots represent the median (black bar), the interquartile range – IQR (boxes), maximum and minimum values excluding outliers (whiskers) and outliers (black dots).

Figure 2. Individual average heart rate of horses across experiments. Points represent a value per individual horse and the lines connect individual horses over different experiments.