1 **Response of soil microbial communities to alpine meadow** 2 **degradation severity levels in the Qinghai-Tibet Plateau**

3

4 Wenjuan Zhang^{1,2*}, Xian Xue ¹, Fei Peng^{1,3}, Quangang You¹, Jing Pan ^{1,2},
5 Chengyang Li^{1,2}, Chimin Lai ⁴

6

7 ¹ *Key Laboratory of Desert and Desertification, Chinese Academy of Sciences,*
8 *Northwest Institute of Eco-Environment and Resources, Chinese Academy of*
9 *Sciences, 320 West Donggang Road, Lanzhou, China 730000*

10 ² *University of Chinese Academy of Sciences, Beijing 100049, China*

11 ³ *International platform for dryland research and education, arid land research*
12 *center, Tottori University*

13 ⁴ *College of Forestry, Fujian Agriculture and Forestry University, Fuzhou 350000,*
14 *Chinafujian*

15

16

17 *Corresponding author: Wenjuan Zhang (Tel) 86-931-4967567;

18

(Fax) 86-931-827389

19 E-mai:zhangwenjuan@lzb.ac.cn

20

21 Type of Paper: Original research

22

23

Abstract

Soil microbial community structure is an effective indicator to reflect changes in soil quality. Little is known about the effect of alpine meadow degradation on the soil bacterial and fungal community. In this study, we used the Illumina MiSeq sequencing method to analyze the microbial community structure of alpine meadow soil in five different degradation levels (i.e., non-degraded (ND), slightly degraded (LD), moderately degraded (MD), severely degraded (SD), and very severely degraded (VD)) in the Qinghai-Tibet Plateau. *Proteobacteria*, *Actinobacteria*, and *Acidobacteria* were the mainly bacterial phyla in meadow soil across all five degradation levels investigated. *Basidiomycota* was the mainly fungal phylum in ND; however, we found a shift from *Basidiomycota* to *Ascomycota* with an increase (severity) in degradation level. The overall proportion of *Cortinariaceae* exhibited high fungal variability, and reads were highest in ND (62.80%). Heatmaps of bacterial genera and fungal families showed a two-cluster sample division on a genus/family level: (1) an ND and LD group and (2) an SD, VD, and MD group. Redundancy analysis (RDA) showed that 79.7% and 71.3% of the variance in bacterial and fungal composition, respectively, could be explained by soil nutrient conditions (soil organic carbon, total nitrogen, and moisture) and plant properties (below-ground biomass). Our results indicate that meadow degradation affects both plant and soil properties and consequently drives changes in soil microbial community structure.

Introduction

The Qinghai-Tibet Plateau (QTP) has an area over 250×10^4 km² within China and is the highest plateau in the world with an average elevation of ~4500 m a.s.l. [1]. Alpine meadows comprise approximately 38% of all grassland area in the QTP and are the primary ecosystem utilized by the Tibetan people and their livestock [2,3]. Accordingly, the alpine meadow is considered to be one of the most critically important ecosystems in the QTP. The ecological functions of alpine meadow ecosystems in the QTP are also important, which include water storage [4],

biodiversity maintenance [5], and soil carbon (C) sequestration [6].

Furthermore, the lower air temperature and higher altitude make alpine meadows more sensitive to global warming. Thus, these ecosystems are considered as good indicators of environmental change [2]. In recent decades, frequent reports on the degradation of alpine meadows in the source regions of the Yangtze and Yellow rivers have been attributed to climate warming and anthropogenic activities [7,8]. The alpine grassland degradation has led to a variety of ecological consequences, including alterations plant community composition, decreased plant species richness and biomass [9], and accelerated soil erosion [10].

Soil bacteria and fungi play crucial roles in soil nutrient supplies and element cycling in terrestrial ecosystems [11], and their composition and diversity are sensitive to disturbances [12,13]. Different microbes exhibit various ability to efficiently utilize soil organic matter (SOM) and the composition of microbial decomposers directly influence a variety of ecosystem processes, such as CO₂ flux and litter decomposition. Previous studies have shown that soil microbial communities are affected by plant characteristics and soil properties [14,15]. For example, vegetation type has a strong effect on soil microbial communities in determining the physical soil environment and the availability of nutrients [16]. Soil substrate availability and heterogeneity are important factors responsible for changes in microbial communities [17,18]. Potential changes in soil nutrient availability [19], soil moisture [20], and plant composition and biomass during processes of grassland degradation [21] would inevitably alter the composition and diversity of soil microbial communities. The current literature on alpine meadow degradation mainly focuses on plant and soil characteristics; however, knowledge regarding the effects of meadow degradation on soil microbial communities and their diversity remains largely inadequate [22]. Therefore, it is essential to understand how the composition and diversity of microbial communities respond to alpine meadow degradation in the QTP and to consider the key influencing factors and to provide important insights for the alpine meadow health assessment and management.

In this study, we selected alpine meadows under different levels of degradation

severity (i.e., non-degraded (ND), slightly degraded (LD), moderately degraded (MD), severely degraded (SD), and very severely degraded (VD)), applying the space-for-time substitution method [9]. We used the Illumina MiSeq sequencing method (Illumina, Inc., USA) to determine soil bacterial and fungal composition and their diversity with changes in edaphic and plant properties under degradation. The specific aims of this study were: 1) to investigate how soil bacterial and fungal communities vary with changes in explicit soil and plant properties in the QTP, and 2) to discern which factors under conditions of alpine meadow degradation significantly influence associative microbial community structure.

Materials and methods

Study area

The study area was located at the source of the Yangtze River on the QTP (34°49'N, 92°55'E; 4635 m a.s.l.) (Fig 1). The annual average mean air temperatures are -3.8°C. Mean annual precipitation is 285 mm with greater than 93% falling during the warm growing season (April–October) [2]. The site is used for grazing during the summer. The typical vegetation is alpine meadow dominated by *Kobresia capillifolia*, *Kobresia pygmaea*, and *Carex moorcroftii* [3].

The ND, LD, MD, SD, and VD alpine meadow sites were chosen based on vegetation coverage as described by Liu et al. (2018) [23]. Vegetation coverage percentages are as follows: 80%~90% for ND, 70%~80% for LD, 50%~70% for MD, 30%~50% for SD, and <30% for VD.

Plant measurement and soil sampling procedures

Five quadrats (5 m × 5 m) at a spatial distance of greater than 50 m were randomly selected at each of the five alpine meadow sites. Three plots were sampled from each quadrat. Plant cover, height, and composition of each quadrat were measured using point-intercept sampling, employing a 30 cm × 30 cm square frame, with 100 sampling points spaced equidistantly within the frame [24]. Plant height was estimated by randomly measuring 10 individuals in each quadrat. Following this, plants were clipped at the soil surface, and harvested plants were categorized into

different functional groups: graminoids, sedges, and forbs. Aboveground and belowground plant biomass (abbreviated as AGB and BGB in this study, respectively) were separated, dried in an oven at 65°C for 48 h before being weighed.

On July 3, 2017, three 50 mm diameter surface soil cores (from the top 10 cm soil layer) were randomly collected from each quadrat. These soil cores were then mixed, homogenized, and sieved (<2 mm) to remove roots and other plant material [25]. Each soil sample was placed in a sterile centrifuge tube before being immediately transported to the laboratory and stored at -80°C for total DNA extraction and molecular analyses.

From each quadrat, three soil profiles were randomly collected for soil sampling (0 to 10 cm). The soil samples were measured for soil bulk density (BD), soil moisture (SM), and soil nutrient content piling at each of its four corners.

Soil characterization

Soil BD was measured using 100 cm³ rings and calculated from dry soil matter. At the same time, SM was measured by drying soil samples taken from the rings at 105°C for 48 h. Soil organic carbon (SOC) was determined using the potassium dichromate oxidation titration method [26]. Total nitrogen (TN) in soil was determined using the semimicro Kjeldahl digestion procedure [2]. Ammonium-nitrogen (NH₄⁺-N) and nitrate nitrogen (NO₃⁻-N) content in soil were measured using the UV-3300 Spectrophotometer (Mapada Instruments Co., Ltd., Shanghai, China).

DNA extraction, Polymerase chain reaction and Illumina sequencing

DNA was extracted from 0.5 g soil samples using the PowerSoil DNA Isolation Kit (Mo Bio Laboratories, Carlsbad, CA, USA) according to the manufacturer's instructions. The universal primer pair 338F/806R [27] for bacteria and ITS1 [28] for fungi were used for amplification and MiSeq sequencing of the polymerase chain reaction (PCR) products. PCR amplification was conducted in a 25 µL reaction system using TransGen AP221-02 (TransGen Biotech, Beijing, China) and carried out in an ABI GeneAmp 9700 (Applied Biosystems, Inc., Carlsbad, USA). After purified

and quantitated, the PCR products were sequenced using the Illumina Miseq platform.

The 16s and ITS1 rRNA gene sequences associated this study were submitted to the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) (accession no. PRJNA490659).

Data analysis

The QIIME (version 1.8.0) was used to estimate α -diversity (i.e. Chao1, Shannon, and Good's coverage) for each sample [29]. The changes of species composition under different degradation stages, principal component analysis (PCA), and heatmaps were conducted using the R language (3.2.0). One-way analysis of variance (ANOVA) tests was used to evaluate significant levels for all factors analyzed herein. The SPSS 17.0 software package (SPSS Inc., Chicago, IL, USA) was used to calculations and analyses. The CANOCO software (version 4.5) was used to run redundancy analysis (RDA) [30].

Results

Response of plant and soil factors to alpine meadow degradation

Plant coverage in MD, SD, and VD decreased significantly ($P < 0.05$) by 32.48%, 17.52%, and 84.67% compared to ND and by 38.13%, 24.42%, and 85.95%, respectively, compared to LD (Table 1). Plant AGB in MD, SD, and VD significantly ($P < 0.05$) decreased by 47.09%, 51.60%, and 84.14%, respectively, compared to LD (Table 1). Plant BGB in SD and VD also significantly decreased by 60.43% and 67.12% compared to ND and by 59.67% and 66.49% in SD and compared to LD, respectively (Table 1). The ND and LD sites not exhibited a significant differ in plant coverage, AGB, or BGB (Table 1). There was no significant change in species richness (SR) and plant height throughout the whole degradation process (Table 1). For plant group, sedge biomass in SD and VD significantly decreased compared to LD, but we found no significant change in the biomass/proportion of graminoids and forbs as degradation severity increased (Table 2).

Soil TN, SOC, NO₃⁻-N, NH₄⁺-N, and SM content decreased significantly in MD, SD, and VD compared to ND, but BD exhibited the opposite response pattern (Table 3). In MD, there were significant decreases in soil TN, SOC, NO₃⁻-N, NH₄⁺-N, and SM by 62.20%, 67.75%, 48.40%, 47.94%, and 43.57%, respectively, compared to ND (Table 3); in SD, the content of soil TN, SOC, NO₃⁻-N, NH₄⁺-N, and SM decreased significantly by 65.35%, 67.11%, 60.34%, 50.09%, and 65.66%, respectively, compared to ND (Table 3); in VD, there were significant decreases in soil TN, SOC, NO₃⁻-N, NH₄⁺-N, and SM by 70.08%, 76.16%, 57.64%, 43.25%, and 66.72%, respectively, compared to ND (Table 3). However, soil BD increased significantly ($P < 0.05$) by 37.14%, 28.57%, and 47.14% in MD, SD, and VD, respectively, compared to ND (Table 3). We found no significant difference in SOC, TN, NO₃⁻-N, NH₄⁺-N, SM, and BD between ND and LD (Table 3).

α -diversity indices based on MiSeq sequencing

For bacteria, MD yielded the highest richness value (Chao1 = 1187.75), while SD yielded the lowest (Chao1 = 1086.23). The Shannon index not only provides simple SR (the number of species present) but also the level of abundance of each species (species evenness) as distributed among all species in a community. ND yielded the highest diversity value (Shannon = 8.46) among the five samples, and SD yielded the lowest diversity value (Shannon = 8). From 962 to 1104 OTUs were detected in total for all samples at a 3% genetic distance. The progression of rarefaction curves (99.01%–99.26%; Good's coverage) was very close for all samples (S1 Table).

For fungi, LD also yielded the highest richness value (Chao1 = 645.85), followed by VD (Chao1 = 575.29), and ND yielded the lowest richness value (Chao1 = 517.94). VD yielded the highest diversity value (Shannon = 6.38), however, ND yielded the lowest diversity value (Shannon = 3.13). The number of OTUs ranged from 423 to 564 in the samples, for which LD had the highest one and ND the lowest (S1 Table).

Taxonomic composition based on MiSeq sequencing

Bacterial OTUs could be assigned to 28 phyla, 156 families, and 170 genera. In total, 20 different phyla (*Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Bacteroidetes*,

Chloroflexi, *Gemmatimonadetes*, *Verrucomicrobia*, etc.) out of the 28 total bacterial phylotypes were common to the five libraries (Fig 2A), contributing from 99.84%, 99.62%, 99.72%, 99.91%, and 99.74% of the total reads in the ND, LD, MD, SD, and VD libraries, respectively. *Proteobacteria* was the most abundant division, comprising 26.48% (350) of the OTUs and 44.18% (10 078) of the reads across all samples. *Acidobacteria*, the second most abundant phylum, comprised 14.83% (196) of the OTUs and 18.44% (4213) of the reads in all libraries. These two phyla collectively accounted for 60.88%, 66.83%, 64.31%, 63.65%, and 57.42% of the total reads in the ND, LD, MD, SD, and VD libraries, respectively (Fig 2A). However, the proportion of *Proteobacteria* exhibited low variability in the different samples, namely, ND (46.63%; 10 224 reads), LD (46.80%, 11,442 reads), MD (39.08%, 8,069 reads), SD (49.68%, 10,637 reads) and VD (38.71%, 10,021reads). Additionally, reads from *Acidobacteria* fluctuated in the different samples, for which the proportion of reads was highest in MD (25.23%; 5210 reads).

Although members of the α -*proteobacteria* class dominated the *proteobacteria* phylum (10.82% and 143 OTUs) in all libraries, they accounted for 30.34% of the total reads (Fig 2B). The subdivision of β -*proteobacteria*, δ -*proteobacteria*, and γ -*proteobacteria* classes comprised of 7.16%, 4.42%, and 2.21% of the total reads, respectively.

The fungal communities were assigned to 8 phyla, 78 families, and 138 genera. *Ascomycota* was the most dominant division, comprising 34.52% (339) of the OTUs and 44.43% of the total reads (Fig 3A). *Basidiomycota* was the second largest division, with 12.63% (124) of the OTUs and 29.92% of the total reads. These two phyla collectively accounted for 95.88%, 72.04%, 64.78%, 74.50%, and 64.55% of the total reads in ND, LD, MD, SD, and VD, respectively (Fig 3A). However, *Ascomycota* exhibited high variability in read abundance of the different samples; namely, in increasing order of abundance, ND (14.47%; 5641 reads), LD (37.90%; 13 649 reads), MD (48.06%; 20 251 reads), SD (63.35%; 24 980 reads), and VD (58.38%; 23 434 reads). In contrast, *Basidiomycota* reads decreased with an increase in degradation severity; accordingly, ND had the highest read value (81.40%; 31 729 reads). Reads

from *Zygomycota* and *Glomeromycota* fluctuated in the different samples; namely, VD had the highest read value (Fig 3A).

On a family level, we found abundant unidentified fungal sequences in each sample. The total relative abundance of unidentified fungi in ND, LD, MD, SD, and VD were 9.65%, 40.06%, 46.40%, 43.89%, and 37.19%, respectively (Fig 3B). *Cortinariaceae* members dominated the Basidiomycota phylum, and the proportion of reads exhibited high variability in read abundance of the different samples; namely, ND (62.80%; 24 476 reads), LD (0.42%; 150 reads), MD (0.23%; 98 reads), SD (0.26%; 101 reads), and VD (0.22%; 90 reads). *Incertae-sedis*, the third most abundant group (7.84%; 77 OTUs), comprised 10.78% (4208) of the reads in all libraries. These three groups combined, namely, unidentified, *Cortinariaceae*, and *incertae-sedis*, comprised of greater than 53.21% of the total sequences from all five sample sites.

Taxonomic composition based on MiSeq sequencing

In order to analyze microbial community similarity among the five mixed samples, we generated heatmaps applying hierarchical cluster analysis. For bacteria, the heatmap (Fig 4A) was based on the top 20 abundant bacterial genera. The heatmap showed a two-sample cluster division. The first cluster was the ND and LD group, and the second cluster was the SD and VD group, which first clustered together before clustering with MD, resulting in the second SD, VD, and MD group. Results from PCA also showed that the bacterial communities of SD and VD grouped to the left of the graph along the PC1 axis, accounting for 41.08% of total variation, whereas ND and LD grouped along the PC2 axis, with a total variance of 31.92% (Fig 5A).

For fungi, the heatmap (Fig 4B) was based on the top 20 fungal families. The figure shows a two cluster sample division on a family rank level: ND and LD into one group, and SD, VD, and MD into another. The PCA score plot (Fig 5B) was in agreement with the heatmap, indicating high fungal community similarity between ND and MD and between MD, SD, and VD. ND and LD grouped to the right, and MD, SD, and VD grouped to the left of the graph along the PC1 axis, with a total

variance of 30.93%. The PC2 axis accounted for 26.42% of the total variances.

Correlation between community structure and environmental factors

The Monte Carlo test showed that the total explanatory powers of measure variables explained 44.5% and 35.2% of the total variation in bacterial communities on the first and second axis, respectively (Fig 6A). Results showed that soil bacteria under different degradation stages was significantly correlated to soil NO_3^- -N ($P < 0.01$), SOC ($P < 0.05$), TN ($P < 0.05$), SM ($P < 0.05$), and BGB ($P < 0.05$), whereas soil NH_4^+ -N, BD, AGB, etc. ($P > 0.05$) had no obvious effect on bacteria (S2 Table).

For fungi, the Monte Carlo test showed that the total explanatory powers of the measured variables explained 44.3% and 27.0% of total variation in fungal communities on the first and second axis, respectively (Fig 6B). Results showed that soil fungi were significantly correlated to soil TN ($P < 0.05$), SOC ($P < 0.05$), SM ($P < 0.05$), BD ($P < 0.05$), BGB ($P < 0.05$), and coverage ($P < 0.05$) whereas soil NO_3^- -N, NH_4^+ -N, AGB, etc. ($P > 0.05$) had no obvious effect on bacteria (S2 Table).

Discussion

This study investigated how soil bacterial and fungal communities vary with changes in specific soil and plant properties in the QTP while discerning which factors significantly influence associative microbial community structure with alpine meadow degradation.

Soil microbial community characteristics and its relationship with biotic and abiotic factors

Consistent with previous studies [19,21], increasing meadow degradation in our study significantly decreased the soil nutrient status (Table 3). This was due to a decrease in plant coverage and biomass, a reduction in SM, and an increase in soil BD in conjunction with degradation severity (Tables 1 and 3), which could potentially lead to nutrient leaching and contribute to nutrient loss [31]. In our study, both the bacteria and fungi genera heatmaps showed a two-cluster sample division. The first cluster

was the ND and LD group, and the second cluster was the SD and VD group, which first clustered together before clustering with MD, resulting in the second SD, VD, and MD group. These results could be explained by a significant change in plant (i.e., coverage and BGB) and soil (i.e., SOC, TN, NO₃-N, NH₄-N, SM, and BD) factors in MD, SD, and VD compared to ND, while ND and LD exhibited no significant difference in the above factors (Tables 1 and 3).

About 79.7% and 71.3% of variation in bacterial and fungal composition could be explained by soil physicochemical properties and plant characteristics, respectively (Fig 6A, B), indicating that soil nutrient (i.e., SOC and TN) and moisture content, plant coverage, and BGB could potentially be key factors in defining changes in microbial composition under conditions of meadow degradation. Similar to our results, previous studies have demonstrated that the factors listed above affect soil microbial communities [32–34]. Variability in SM may reduce soil water availability averages [35], leading to reduced soil microbial community C use efficiency [36] and ultimately shifting the biomass or ratios of fungi and bacteria in soil [37]. At present, a number of studies have focused on the effect of nitrogen (N) gradients on soil microbial communities. This is because ecosystems around the world are being subjected to elevated levels of N [38,39]. For example, Ramirez et al. (2012) [40] found that N fertilization significantly affects total bacterial community composition. Conversely, increased N fertilizer dosages could potentially have a negative impact on C cycling in soil while, at the same time, promoting fungal genera with known pathogenic traits [38]. Moreover, the significant ($P < 0.05$) correlation found between bacterial and fungal communities and plant BGB (S2 Table) demonstrates that degradation-induced changes in plant species composition and biomass also exert strong effects on microbial communities. Wallenstein et al. (2007) [32] also suggested that plants strongly regulate microbial communities through the role they play in substrate supplies (e.g., litter, root turnover, and exudates) and by changing the physical environment in the active soil layer.

Bacterial and fungal community response to meadow

degradation

The bacterial phyla in meadow soil investigated in this study exhibited low variability in the different samples (Fig 2A); however, the composition of fungal phyla significantly changed under conditions of degradation severity (Fig 3A), indicating that fungal communities are more sensitive to degradation than bacterial communities. It was also reported that biotic and abiotic factors have a greater influence on fungi than bacteria [41]. The higher sensitivity of fungi that have been reported by some recent studies suggests that soil fungal communities are highly responsive to changes in SM and soil nutrient limitations than bacteria [42–44].

If fungal groups differ in their preference to substrate utilization processes [45], degradation-induced effects on biogeochemical properties will cause marked changes in specific species [46]. In our study, we consistently found that the dominate *Ascomycota* community structure significantly increased with increasing levels of meadow degradation severity (Fig 3A). Growth rates of *ascomycetes* were correlated to N availability, while their activity may dramatically accelerate C decomposition [47]. Additionally, the abundance of *Basidiomycota* significantly decreased in response to meadow degradation (Fig 3A). *Basidiomycetes* are widely recognized as lignin decomposers [48], and its capacity to utilize this recalcitrant substrate will likely hinder the development of this fungal group given that we detected a reduction in plant litter (biomass) with an increase in meadow degradation severity (Tables 1 and 2). Specifically, members of the *Cortinariaceae* family dominated the Basidiomycota phylum, and the proportion of reads exhibited high variability in abundance in the different samples; namely, in decreasing order of abundance, ND (62.80%), LD (0.42%), MD (0.23%), SD (0.26%), and VD (0.22%). Therefore, the association between fungi and plants may be responsible for the higher sensitivity of fungal composition to degradation severity compared to the bacterial composition.

For the bacterial groups, the overall proportion of *Proteobacteria* generally decreased while *Actinobacteria* generally increased in MD and VD compared to ND and LD (Fig 2A). This could have been because many members of *Proteobacteria*

(particularly *α-proteobacteria*; Fig 2B) prefer nutrient-rich environments, whereas *Actinobacteria* are well adapted to oligotrophic environments [49,50]. Therefore, observed shifts are consistent with a decrease in the status of soil nutrients (i.e., C and N content) and the degradation of permafrost. Thus, degradation-induced environmental changes exert effects on microbial communities.

Conclusions

In conclusion, this study, using the high throughput Illumina MiSeq sequencing method, provided a detailed description of variation in alpine meadow bacterial and fungal communities under different levels of degradation severity. For bacterial communities, these differences likely resulted from combined differences in soil properties and plant characteristics (most closely associated with SOC, TN, SM, NO₃⁻-N, and BGB) rather than to single biotic or abiotic factors that combined to create the unique characteristics representative of each site. On the other hand, changes in soil fungal communities were mainly attributed to variations in SOC, TN, SM, BD, BGB, and plant coverage. Although this study will aid in our understanding of changes in soil microbial communities throughout the whole alpine meadow degradation process, further research into the mechanisms that underlie our findings would also be of great interest.

Acknowledgements

The study was financially supported by the National Key Research and Development Program of China (2016YFC0501803) and the National Science Foundation of China (41771229, 41771233).

References

1. Qiu J. Climate change is coming fast and furious to the Tibetan plateau. 2008;454: 4.
2. You Q, Xue X, Peng F, Xu M, Duan H, Dong S. Comparison of ecosystem characteristics between degraded and intact alpine meadow in the Qinghai-Tibetan Plateau, China. Ecological Engineering. 2014;71: 133–143.

- doi:10.1016/j.ecoleng.2014.07.022
3. Xue X, You Q, Peng F, Dong S, Duan H. Experimental Warming Aggravates Degradation-Induced Topsoil Drought in Alpine Meadows of the Qinghai-Tibetan Plateau: Experimental Warming Aggravates Degradation-Induced Topsoil Drought. *Land Degradation & Development*. 2017; doi:10.1002/ldr.2763
4. Zhang L, Su F, Yang D, Hao Z, Tong K. Discharge regime and simulation for the upstream of major rivers over Tibetan Plateau. *Journal of Geophysical Research: Atmospheres*. 2013;118: 8500–8518. doi:10.1002/jgrd.50665
5. López-Pujol J, Zhang F-M, Sun H-Q, Ying T-S, Ge S. Centres of plant endemism in China: places for survival or for speciation? *Journal of Biogeography*. 2011;38: 1267–1280. doi:10.1111/j.1365-2699.2011.02504.x
6. Piao S, Fang J, Ciais P, Peylin P, Huang Y, Sitch S, et al. The carbon balance of terrestrial ecosystems in China. *Nature*. 2009;458: 1009–1013. doi:10.1038/nature07944
7. Wang Z, Zhang Y, Yang Y, Zhou W, Gang C, Zhang Y, et al. Quantitative assess the driving forces on the grassland degradation in the Qinghai–Tibet Plateau, in China. *Ecological Informatics*. 2016;33: 32–44. doi:10.1016/j.ecoinf.2016.03.006
8. Ren X, Dong Z, Hu G, Zhang D, Li Q. A GIS-Based Assessment of Vulnerability to Aeolian Desertification in the Source Areas of the Yangtze and Yellow Rivers. *Remote Sensing*. 2016;8: 626. doi:10.3390/rs8080626
9. Wang CT, Long RJ, Wang QL, Jing ZC, Shi JJ. Changes in plant diversity, biomass and soil C, in alpine meadows at different degradation stages in the headwater region of three rivers, China. *Land Degradation & Development*. 2009;20: 187–198. doi:10.1002/ldr.879
10. Xue X, Guo J, Han B, Sun Q, Liu L. The effect of climate warming and permafrost thaw on desertification in the Qinghai–Tibetan Plateau. *Geomorphology*. 2009;108: 182–190. doi:10.1016/j.geomorph.2009.01.004
11. Gessner MO, Swan CM, Dang CK, McKie BG, Bardgett RD, Wall DH, et al.

- 402 Diversity meets decomposition. *Trends in Ecology & Evolution*. 2010;25:
403 372–380. doi:10.1016/j.tree.2010.01.010
- 404 12. Allison SD, Martiny JBH. Resistance, resilience, and redundancy in microbial
405 communities. *Proceedings of the National Academy of Sciences*. 2008;105:
406 11512–11519. doi:10.1073/pnas.0801925105
- 407 13. Zhang Y, Dong S, Gao Q, Liu S, Zhou H, Ganjurjav H, et al. Climate change
408 and human activities altered the diversity and composition of soil microbial
409 community in alpine grasslands of the Qinghai-Tibetan Plateau. *Science of The
410 Total Environment*. 2016;562: 353–363. doi:10.1016/j.scitotenv.2016.03.221
- 411 14. Chen S, Liu W, Qin X, Liu Y, Zhang T, Chen K, et al. Response characteristics
412 of vegetation and soil environment to permafrost degradation in the upstream
413 regions of the Shule River Basin. *Environmental Research Letters*. 2012;7:
414 045406. doi:10.1088/1748-9326/7/4/045406
- 415 15. Kumar M, Männistö MK, van Elsas JD, Nissinen RM. Plants impact structure
416 and function of bacterial communities in Arctic soils. *Plant and Soil*. 2016;399:
417 319–332. doi:10.1007/s11104-015-2702-3
- 418 16. Chu H, Neufeld JD, Walker VK, Grogan P. The Influence of Vegetation Type
419 on the Dominant Soil Bacteria, Archaea, and Fungi in a Low Arctic Tundra
420 Landscape. *Soil Science Society of America Journal*. 2011;75: 1756.
421 doi:10.2136/sssaj2011.0057
- 422 17. Myers RT, Zak DR, White DC, Peacock A. Landscape-Level Patterns of
423 Microbial Community Composition and Substrate Use in Upland Forest
424 Ecosystems. *Soil Science Society of America Journal*. 2001;65: 359.
425 doi:10.2136/sssaj2001.652359x
- 426 18. Ranjard L, Richaume A. Quantitative and qualitative microscale distribution of
427 bacteria in soil. *Research in Microbiology*. 2001;152: 707–716.
428 doi:10.1016/S0923-2508(01)01251-7
- 429 19. Dong SK, Wen L, Li YY, Wang XX, Zhu L, Li XY. Soil-Quality Effects of
430 Grassland Degradation and Restoration on the Qinghai-Tibetan Plateau. *Soil
431 Science Society of America Journal*. 2012;76: 2256.

- doi:10.2136/sssaj2012.0092
20. Zhang X, Xu S, Li C, Zhao L, Feng H, Yue G, et al. The soil carbon/nitrogen ratio and moisture affect microbial community structures in alkaline permafrost-affected soils with different vegetation types on the Tibetan plateau. *Research in Microbiology*. 2014;165: 128–139. doi:10.1016/j.resmic.2014.01.002
21. Wu GL, Ren GH, Dong QM, Shi JJ, Wang YL. Above- and belowground response along degradation gradient in an alpine grassland of the Qinghai-Tibetan Plateau. *CLEAN - Soil, Air, Water*. 2014;42: 319–323. doi:10.1002/clen.201200084
22. Stark S, Männistö MK, Eskelinen A. When do grazers accelerate or decelerate soil carbon and nitrogen cycling in tundra? A test of theory on grazing effects in fertile and infertile habitats. *Oikos*. 2015;124: 593–602. doi:10.1111/oik.01355
23. Liu S, Zamanian K, Schleuss P-M, Zarebanadkouki M, Kuzyakov Y. Degradation of Tibetan grasslands: Consequences for carbon and nutrient cycles. *Agriculture, Ecosystems & Environment*. 2018;252: 93–104. doi:10.1016/j.agee.2017.10.011
24. Wang S, Duan J, Xu G, Wang Y, Zhang Z, Rui Y, et al. Effects of warming and grazing on soil N availability, species composition, and ANPP in an alpine meadow. *Ecology*. 2012;93: 2365–2376. doi:10.1890/11-1408.1
25. Duan Y, Wu F, Wang W, He D, Gu J-D, Feng H, et al. The microbial community characteristics of ancient painted sculptures in Maijishan Grottoes, China. Chaturvedi V, editor. *PLOS ONE*. 2017;12: e0179718. doi:10.1371/journal.pone.0179718
26. Zeng C, Zhang F, Wang Q, Chen Y, Joswiak DR. Impact of alpine meadow degradation on soil hydraulic properties over the Qinghai-Tibetan Plateau. *Journal of Hydrology*. 2013;478: 148–156. doi:10.1016/j.jhydrol.2012.11.058
27. Dennis KL, Wang Y, Blatner NR, Wang S, Saadalla A, Trudeau E, et al. Adenomatous polyps are driven by microbe-instigated focal inflammation and are controlled by IL-10-Producing T cells. *Cancer Research*. 2013;73:

- 462 5905–5913. doi:10.1158/0008-5472.CAN-13-1511
- 463 28. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of
464 fungal ribosomal RNA genes for phylogenetics, in: PCR Protocols. Elsevier;
465 1990. pp. 315–322. doi:10.1016/B978-0-12-372180-8.50042-1
- 466 29. Schloss PD, Gevers D, Westcott SL. Reducing the Effects of PCR Amplification
467 and Sequencing Artifacts on 16S rRNA-Based Studies. Gilbert JA, editor. PLoS
468 ONE. 2011;6: e27310. doi:10.1371/journal.pone.0027310
- 469 30. Giaramida L, Manage PM, Edwards C, Singh BK, Lawton LA. Bacterial
470 communities' response to microcystins exposure and nutrient availability:
471 Linking degradation capacity to community structure. International
472 Biodeterioration & Biodegradation. 2013;84: 111–117.
473 doi:10.1016/j.ibiod.2012.05.036
- 474 31. Mchunu C, Chaplot V. Land degradation impact on soil carbon losses through
475 water erosion and CO₂ emissions. Geoderma. 2012;177–178: 72–79.
476 doi:10.1016/j.geoderma.2012.01.038
- 477 32. Wallenstein MD, McMahon S, Schimel J. Bacterial and fungal community
478 structure in Arctic tundra tussock and shrub soils. FEMS Microbiology Ecology.
479 2007;59: 428–435. doi:10.1111/j.1574-6941.2006.00260.x
- 480 33. He Z, Piceno Y, Deng Y, Xu M, Lu Z, DeSantis T, et al. The phylogenetic
481 composition and structure of soil microbial communities shifts in response to
482 elevated carbon dioxide. The ISME Journal. 2012;6: 259–272.
483 doi:10.1038/ismej.2011.99
- 484 34. Jumpponen A, Jones KL. Tallgrass prairie soil fungal communities are resilient
485 to climate change. Fungal Ecology. 2014;10: 44–57.
486 doi:10.1016/j.funeco.2013.11.003
- 487 35. Fay PA, Blair JM, Smith MD, Nippert JB, Carlisle JD, Knapp AK. Relative
488 effects of precipitation variability and warming on tallgrass prairie ecosystem
489 function. Biogeosciences. 2011;8: 3053–3068. doi:10.5194/bg-8-3053-2011
- 490 36. Zeglin LH, Bottomley PJ, Jumpponen A, Rice CW, Arango M, Lindsley A, et al.
491 Altered precipitation regime affects the function and composition of soil

492 microbial communities on multiple time scales. *Ecology*. 2013;94: 2334–2345.
 493 doi:10.1890/12-2018.1

494 37. Cregger MA, Schadt CW, McDowell NG, Pockman WT, Classen AT. Response
 495 of the Soil Microbial Community to Changes in Precipitation in a Semiarid
 496 Ecosystem. *Applied and Environmental Microbiology*. 2012;78: 8587–8594.
 497 doi:10.1128/AEM.02050-12

498 38. Paungfoo-Lonhienne C, Yeoh YK, Kasinadhuni NRP, Lonhienne TGA,
 499 Robinson N, Hugenholtz P, et al. Nitrogen fertilizer dose alters fungal
 500 communities in sugarcane soil and rhizosphere. *Scientific Reports*. 2015;5.
 501 doi:10.1038/srep08678

502 39. Leff JW, Jones SE, Prober SM, Barberán A, Borer ET, Firn JL, et al. Consistent
 503 responses of soil microbial communities to elevated nutrient inputs in
 504 grasslands across the globe. *Proceedings of the National Academy of Sciences*.
 505 2015;112: 10967–10972. doi:10.1073/pnas.1508382112

506 40. Ramirez KS, Craine JM, Fierer N. Consistent effects of nitrogen amendments on
 507 soil microbial communities and processes across biomes. *Global Change*
 508 *Biology*. 2012;18: 1918–1927. doi:10.1111/j.1365-2486.2012.02639.x

509 41. Nielsen UN, Osler GHR, Campbell CD, Burslem DFRP, Van Der Wal R. The
 510 influence of vegetation type, soil properties and precipitation on the
 511 composition of soil mite and microbial communities at the landscape scale:
 512 Biogeography of mites and microbes. *Journal of Biogeography*. 2010;37:
 513 1317–1328. doi:10.1111/j.1365-2699.2010.02281.x

514 42. Hawkes CV, Hartley IP, Ineson P, Fitter AH. Soil temperature affects carbon
 515 allocation within arbuscular mycorrhizal networks and carbon transport from
 516 plant to fungus. *Global Change Biology*. 2008;14: 1181–1190.
 517 doi:10.1111/j.1365-2486.2007.01535.x

518 43. Lauber CL, Strickland MS, Bradford MA, Fierer N. The influence of soil
 519 properties on the structure of bacterial and fungal communities across land-use
 520 types. *Soil Biology and Biochemistry*. 2008;40: 2407–2415.
 521 doi:10.1016/j.soilbio.2008.05.021

44. Dooley SR, Treseder KK. The effect of fire on microbial biomass: a meta-analysis of field studies. *Biogeochemistry*. 2012;109: 49–61. doi:10.1007/s10533-011-9633-8
45. McGuire KL, Bent E, Borneman J, Majumder A, Allison SD, Treseder KK. Functional diversity in resource use by fungi. *Ecology*. 2010;91: 2324–2332. doi:10.1890/09-0654.1
46. Allison SD, Treseder KK. Warming and drying suppress microbial activity and carbon cycling in boreal forest soils. *Global Change Biology*. 2008;14: 2898–2909. doi:10.1111/j.1365-2486.2008.01716.x
47. Fontaine S, Henault C, Aamor A, Bdioui N, Bloor JMG, Maire V, et al. Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming effect. *Soil Biology and Biochemistry*. 2011;43: 86–96. doi:10.1016/j.soilbio.2010.09.017
48. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 2010;26: 2460–2461. doi:10.1093/bioinformatics/btq461
49. Saul DJ, Aislabie JM, Brown CE, Harris L, Foght JM. Hydrocarbon contamination changes the bacterial diversity of soil from around Scott Base, Antarctica. *FEMS Microbiology Ecology*. 2005;53: 141–155. doi:10.1016/j.femsec.2004.11.007
50. Yergeau E, Bokhorst S, Kang S, Zhou J, Greer CW, Aerts R, et al. Shifts in soil microorganisms in response to warming are consistent across a range of Antarctic environments. *The ISME Journal*. 2012;6: 692–702. doi:10.1038/ismej.2011.124

Legends

Fig 1. The location of the sampling site.

Table 1. Properties of plant community at different degraded alpine meadow (n=3).

Table 2. Properties of plant AGB and proportion at different degraded alpine

551 meadow (n=3).

552 Table 3. Properties of soil at different degraded alpine meadows (n=3,
553 depth=0-10cm).

554 Fig 2. Relative abundance of the bacterial community at the phylum (A) level
555 and class distribution of proteobacteria (B).

556 Fig 3. Relative abundance of the fungal community at the phylum (A) and
557 family(B) level.

558 Fig 4. Heatmaps representation and cluster analysis of the microbial community
559 among five samples.

560 Fig 5. The results of bacterial (A) and fungal (B) communities for the five mixed
561 samples according to the principal components analysis (PCA).

562 Fig 6. Relationships between soil and plant properties (arrows) and bacterial (A)
563 and fungal (B) community structure according to the redundancy analysis
564 (RDA). (A red solid arrow line means $P < 0.05$; A black dashed line means $P > 0.05$).

565

566 Supporting information

567 S1 Table. The results of MiSeq sequencing and α -diversity estimates of the five
568 mixed samples.

569 S2 Table. Correlations between environmental factors with the ordination axis and
570 significance test of single variables for the bacteria and fungi community structure
571 obtained from the RDA results.

Fig 1. The location of the sampling site

Fig 2A. Relative abundance of the bacterial community at the ph

Fig 2B. Relative abundance of proteobacteria at the class (B)level

Fig 3A. Relative abundance of the fungal community at the phylum

Fig 3B. Relative abundance of the fungal community at the family

Fig 4A. Heatmaps representation and cluster analysis of the mic

Fig 4B. Heatmaps representation and cluster analysis of the fungal community

Fig 5A. The results of bacterial (A) communities for the five mixe

Fig 5B. The results of fungal (B) communities for the five mixed s

Fig 6A. Relationships between soil and plant properties (arrows)

Fig 6B. Relationships between soil and plant properties (arrow)

Table 1. Properties of plant community at different degraded alpine meadow (n=3)

Items	ND	LD	MD	SD	VD
Plant coverage (%)	91.33±1.78ab	99.67±0.27a	61.67±5.04c	75.33±6.15bc	14±4.49d
Plant height (cm)	1.63±0.27a	2.47±0.08a	1.66±0.06a	2.32±0.33a	1.89±0.17a
Species richness (No./m²)	6.67±0.72a	7±0.82a	6±0.82a	6±0.47a	7±0.47a
Above-ground biomass (g/m²)	241.63±1.81ab	383.85±43.19a	203.11±4.94bc	185.78±50.39bc	60.86±15.39c
Blow-ground biomass (g/m²)	1918.45±48.36a	1882.14±310.92ab	861.30±205.47abc	759.05±25.54bc	630.79±237.42c

Note: The data at the table are showed by average value ± standard error. Data with the different letter in same row are significantly different (P<0.05). The same below.

Table 2. Properties of plant AGB and proportion at different degraded alpine meadow (n=3)

Degradation gradients	Above-ground biomass (g/m ²)	graminoids		sedges		forbs	
		biomass (g/m ²)	proportion (%)	biomass (g/m ²)	proportion (%)	biomass (g/m ²)	proportion (%)
ND	241.63±1.81ab	-	-	190.22±5.35ab	78.71±1.87a	51.41±4.36a	21.29±1.87a
LD	383.85±43.20a	24.59±20.08a	16.16±4.40a	261.93±38.21a	67.45±3.96a	97.33±6.81a	27.16±5.33a
MD	203.11±4.94bc	-	-	159.26±7.21ab	78.29±1.75a	43.85±2.46a	21.71±1.75a
SD	185.78±50.40bc	21.96±9.98a	26.36±3.73a	87.19±32.50bc	49.79±10.67 a	76.63±27.80a	41.4±10.04a
VD	60.85±15.39c	3.11±2.54a	19.76±5.38a	28.52±3.97c	50.76±5.39a	29.22±12.52a	42.65±7.83a

Table 3. Properties of soil at different degraded alpine meadows (n=3, depth=0-10cm)

Items	ND	LD	MD	SD	VD
Organic carbon (SOC, g/kg)	15.69±1.10a	17.95±0.42a	5.06±0.62b	5.16±0.29b	3.74±0.49b
Total nitrogen (TN, g/kg)	1.27±0.06a	1.4±0.03a	0.48±0.03b	0.44±0.01b	0.38±0.03b
Nitrate nitrogen (mg/kg)	14.07±0.23b	16.37±0.56a	7.26±0.55c	5.58±0.12c	5.96±0.18c
Ammonium nitrogen (mg/kg)	16.81±0.64a	14.73±0.22a	8.75±0.60b	8.39±0.23b	9.54±0.78b
Soil moisture content (%)	27.61±0.69a	23.46±0.42a	15.58±0.84b	9.48±0.79c	9.19±1.00c
Soil bulk density (g/cm ³)	0.70±0.01b	0.67±0.02b	0.96±0.03a	0.90±0.04a	1.03±0.02a