1 The algae community in taxon Haptophyceae at the early bloom stage of 2 *Phaeocystis globosa* in Northern Beibu Gulf in winter

3

4 Bin Gong^{1,2*}, Haiping Wu^{*2}, Jixian Ma⁴, Meimiao Luo², Xin Li²

5

6 1. Institute of Marine Drugs, Guangxi university of chinese medicine, Nanning, Guangxi, China,
7 530200

8 2. Guangxi Key Laboratory of Beibu Gulf Marine Biodiversity Conservation, Qinzhou University,
9 Qinzhou, China, 535000

10 3. Guangxi Key Laboratory of Marine Disaster in the Beibu Gulf, Qinzhou University, Qinzhou,
11 China, 535000

12 4. Marine Environment Monitoring Center, Qinzhou Oceanic administration, Qinzhou, China,
13 535000

14

15 *Correspondent author

16 Haiping Wu: wuhaipingsky@163.com

17 Bin Gong: 342965691@aliyun.com

18

19 **Abstract:**

20 *Phaeocystis globosa* (Order Phaeocystales, family Phaeocystaceae) caused significant impact on
21 aquaculture farming, global climate change and industry. Since the year 2010, intensive red tides
22 of *P. globosa* began to appear in Beibu Gulf, where previously free of harmful algal blooms, and
23 posed great threats to the cooling systems of a nuclear power plant in 2014 and 2015. In order to
24 discover the bloom mechanism, the community structure of marine microalgae, with a focus on
25 Haptophyceae taxa, in winter in the northern Beibu Gulf near the Qinzhou Bay, Sanniang Bay
26 (SNB) and Dafenjiang River Estuary (DRE), were explored via 18S ribosomal DNA analysis of the
27 V4 region using the Illumina-Based Sequencing platform. The correlation between the relative
28 abundance of five kinds of Haptophyceae algae and environmental factors of seawater were

analyzed. The most abundant Haptophyceae-related OTU in terms of number of reads was identified as *Phaeocystis* and *Chrysochromulina*. The abundance for other Haptophyceae class was relatively low, such as *Haptolina*, *Prymnesium* and *Isochrysis*. *Phaeocystis* was present in all samples sites except S6, S11, S12, S14 and S15, and particularly abundant at S8, nearly 29 times more than the second most abundant site. Most notably, the results showed that *Phaeocystis* displayed highly positive linear correlation with the concentration of NO_3^- -N (Pearson $r=0.856$, $p<0.01$). Linear regression analysis indicated that *Phaeocystis* was significantly linearly related to the NO_3^- -N ($R^2=0.732$; $Y=-0.005 + 0.410 \cdot X$, Y is the relative abundance of *P.globosa*, X is the concentration of NO_3^- -N; $F=38.227$, $P<0.05$) and NO_3^- -N has a significant positive effect on *P.globosa* (regression coefficient is 0.410, $P=0.000$). Moreover, the relative abundance of *Phaeocystis* was significant related to temperature of sea water (Pearson $r=-0.882$, $p<0.01$). Water temperature can explain the 77.8% change reason for the *P.globosa* ($R^2=0.778$), and has a significant effect on *P. globosa* ($Y=0.169-0.009 \cdot X$, $F=49.031$, $P<0.05$), and the regression coefficient is -0.009 ($P=0.000$) which indicated a significant negative impact relationship between them. Our high throughput sequencing (HTS) based research illustrated how the *P. globosa* bloom generated and its relationship with NO_3^- -N and temperature of sea water in northern Beibu Gulf for the first time, and bringing hope for solving this big problem.

Key Words:

Phaeocystis globosa, high throughput sequencing, nitrate, temperature, Haptophyceae, Beibu Gulf

Introduction:

Haptophyceae (or Prymnesiophyceae), a class of the phylum Chrysophyta, contained the Order Prymnesiales, Discoasterales, Phaeocystales, Isochrysidales. The bloom of some Haptophyceae algae had occurred frequently worldwide and led to great ecological disaster and substantial economic losses. For instance, *Prymnesium parvum*, a species of Haptophyceae algae (Order

Prymnesiales, family Prymnesiaceae), is capable of producing a toxin, prymnesin, and kills fish [1]. Similarly, large area bloom of the *Chrysochromulina polylepis* (Order Prymnesiales, family Prymnesiaceae) have resulted in mortality of trout and salmon in Scandinavian waters during Spring 1988 [2]. Further more, the bloom of *Phaeocystis globosa* (Order Phaeocystales, family Phaeocystaceae) caused mortality of cultured fish [3], Mussel Mortalities [4], higher concentration of DMS [5] and Clogging of Cooling System of Power Plant [6], so both significantly impact aquaculture farming, global climate change and industry.

Beibu Gulf is an important habitat for protection and propagation of marine organisms, known for Indo-Pacific humpback dolphins [7], horseshoe crabs [8], and also an ecologically sensitive region [9]. Recently, rapid economic development and human activities had already resulted in great degradation of marine environment [10,11]. Most noticeably, human-induced nutrient enrichment is becoming a serious problem for coastal marine areas of Beibu Gulf [9,12]. Since the year 2010, intensive red tides of *P. globosa* began to appear in Beibu Gulf, where previously free of harmful algal blooms, and posed great threats to the cooling systems of a nuclear power plant in 2014 and 2015 [6]. Up to now, knowledge on where the *P. globosa* originated and the bloom mechanism is still quite limited. In our opinion, in order to solve the great confusion above, special attention should be paid to the early bloom stage and the environmental characteristics leading to the bloom. For this purpose, many methods had been utilized, such as automatic monitoring buoy [13], satellite remote sensing [14], “molecular probes” [15], methods to detect and quantify toxins [16] and a combination of methods above [17].

Environmental DNA-based Techniques (EDT) was utilized in monitoring Prokaryote and Eukaryotes in water environments for many years and significantly gained impetus over traditional approaches presently [18]. Recently, there are many EDT methods advented for this object; for instance, DNA metabarcoding [19,20], Microsatellite DNA marker [21], and high throughput sequencing (HTS) techniques based metagenomics [22,23] and amplicons [24,25]. In recent years, the HTS techniques has been widely used and remarkably promoted the ecological studies of bacteria [26], fungi [27], algae [28] and animals [29]. However, to date, molecular diversity of microeukaryotes, including algae, in the Beibu Gulf marine region, remains

unexplored. Thus, the recent study about the community and diversity structure of marine microalgae, with a focus on Haptophyceae taxa in winter in the northern Beibu Gulf near the Qinzhou Bay, Sanniang Bay (SNB) and Dafenjiang River Estuary (DRE), were explored via 18S ribosomal DNA analysis of the V4 region using the Illumina-Based Sequencing platform. This study provides a valid taxonomic reference dataset for future microeukaryotic community structure and diversity studies, aimed at monitoring environmental change in the northern Beibu Gulf.

Materials and Methods

Sample Collection

The samples were collected on 27 Dec 2017, when large *Phaeocystis* bloom was observed between eastern Qinzhou Bay (EQB) and Dafenjiang River Estuary (DRE), during the middle several days of Jan to the end of Mar in 2018. Seawater was collected from surface, middle and deep using a CTD Rosette Water Sampling System (Sea-Bird Electronics, USA) between eastern Qinzhou Bay (EQB) and Dafenjiang River Estuary (DRE) (21°32'58"N, 108°39'56"E-21°37'39"N, 108°55'57"E) and mixed evenly. A total of 16 seawater samples (S1-S16, Figure 1) were collected in 3L sterile polyethylene bottles, kept in the dark at 4-8 °C, and filtered at laboratory within 4h. Each 3000 ml seawater sample was filtered through a 1.2 µm mixed cellulose membrane filter (Advantec, Japan) and filtrates were frozen immediately at -20°C for subsequent molecular analyses.

Figure 1. Locations of the the 16 sites (S1-S16) where seawater samples were collected on 27 Dec 2017, including six sites near Sandun Dock (S1-S6), six sites near Sanniang Bay (SNB, S7-S12) and four sites in the Dafengjiang River Estuary (DRE,S13-S16).

DNA Extraction, PCR and illumina sequencing

The PowerWater DNA isolation kit (MoBio Laboratories Inc., CA, USA) was used to extract the DNA of the total organisms on the 1.2 μ m filters following the manufacturer's protocol. The DNA samples were detected by 1% agarose gels and NanoDrop One spectrophotometer (Thermo Fischer Scientific Inc., USA), and then were amplified using the primers 528F and 706R [30,31] which was designed to amplify the hypervariable region V4 of eukaryote 18S rRNA gene. Illumina sequencing was carried out by the Novogene Company (Beijing, China). Sequencing libraries were generated using TruSeq® DNA PCR-Free Sample Preparation Kit (Illumina, USA) following manufacturer's recommendations and index codes were added. The library quality was assessed on the Qubit@ 2.0 Fluorometer (Thermo Scientific) and Agilent Bioanalyzer 2100 system. At last, the library was sequenced on an Illumina HiSeq2500 platform and 250 bp paired-end reads were generated. The raw data sequences were assigned to samples by their unique barcodes. The 18S rDNA primers and barcodes were cut off to generate pair-end (PE) reads. Paired-end reads were merged using FLASH (V1.2.7) [32]; the raw tags were filtered to obtain the high-quality clean tags using QIIME software package (V1.7.0) [33]. Sequences analysis were performed by Uparse software (Uparse v7.0.1001) [34]. Sequences with $\geq 97\%$ similarity were assigned to the same

OTUs. For each representative sequence, the GreenGene Database [35] was used to annotate taxonomic information.

Physical and chemical analyses of seawater characteristics

The seawater temperature ($^{\circ}\text{C}$), salinity were measured using the SBE 911 plus CTD (Sea-Bird Electronics, USA). Dissolved oxygen (DO) concentrations (ml/l) were measured using the SeaBird 43 (Sea-Bird Electronics, USA). The pH values were measured by pH meter (METTLER, FE38-Meter, 30254110). Chlorophyll-a fluorescence was measured using the WETStar (WET Labs, USA). Inorganic nutrient concentrations (nitrate [NO_3^- -N], ammonia [NH_4^+ -N], nitrite [NO_2^-], and phosphate [PO_4^{3-}]) were determined from 100 mL samples with an Alliance Integral Futura Autoanalyzer II [36,37]. The TOC content was determined with TOC analyzer (Multi N/C 3100, Analytik Jena AG, Jena, Germany) according to the procedure explained by Ali [38]. Spatial distribution of seawater characteristics was interpolated by the measurements of the 16 sampling sites using Kriging method [39]. Interpolations outside the sampling area and over the terrestrial landscape were subtracted.

Statistical analysis

Data were compiled and transformed in Microsoft Excel. Correlation between variables were made using a linear Pearson's r coefficient. Linear regression analysis was facilitated and conducted between closely related. Statistics were generated using the SigmaStat version 2.01 software package (SPSS, Inc., Chicago, Ill). All comparisons were performed at the 95% confidence level.

Result

OTUs profile based on 18s RNA amplicon analysis using high throughput sequencing (HTS)

The eukaryote communities in the seawater were represented by 1,594 OTUs identified in our

present study. A good coverage of the eukaryotic diversity in all samples were illustrated by rarefaction curves, which were calculated and reached a plateau in all cases. All OTUs were classified into 44 main Phylum, and the most abundant were Arthropoda, Ascomycota, Chlorophyta, Diatomea, Basidiomycota, Protalveolata and Picozoa (Fig 2). Among all the OTUs, our most concerned OTU was that identified as class Haptophyceae, which some Harmful algae (such as *P.globosa* and *Prymnesium*) causing serious ecological disasters represented in this group. In our study, 19 OTUs were classified into Haptophyceae, belonging to the groups of *Prymnesium*, *Haptolina*, *Isochrysis*, *Chrysochromulina* and *Phaeocystis* respectively (Fig 3). *Phaeocystis* was represented by OTU 174 and OTU 1030, with OTU 174 identified as *P. globosa* (100% similarity). *Prymnesium* was represented only by one OTU (OTU 1269), which closely related to *Prymnesium zebrinum* (98% similarity) and *Prymnesium pienzaarii* (97% similarity). The most biggest taxonomic group in Haptophyceae was identified as genus *Chrysochromulina* which including 4 OTUs. The other taxonomic groups belonging to Haptophyceae were *Haptolina fragaria* (100% similarity, OTU370), *Isochrysis galbana* (100% similarity, OTU1449), *Haptophyceae* sp. (96% similarity, OTU1155) (Fig 3).

165

166 Figure 2. The top ten OTUs were identified based on the hypervariable region V4 of eukaryote
167 18S rRNA gene and sequences with $\geq 97\%$ similarity were assigned to the same OTUs.

Figure 3. The taxonomic relationship of 18S rDNA phylotypes of class Haptophyceae from the northern coast of Beibu Gulf. Kimura two-parameter model and midpoint rooting options were used to reconstruct phylogenetic relationships. Numbers above branches indicate bootstraps for NJ analysis (> 50).

173

174 The most abundant Haptophyceae-related OTU in terms of number of reads was identified
 175 as *Phaeocystis* and *Chrysochromulina*. *Chrysochromulina* was the most abundant taxa containing
 176 5353 reads, while *Phaeocystis* displayed a relatively lower number of 2298 reads. The abundance
 177 for other Haptophyceae class was relatively low, such as *Haptolina* (228 reads), *Prymnesium* (9
 178 reads), *Isochrysis* (6 reads) (Fig 4). The relative abundance of the five kinds of Haptophyceae at
 179 different sampling sites showed different characteristics. The spatial distribution of *Phaeocystis*
 180 and *Prymnesium* were similar. *Phaeocystis* was present in all samples sites except S6, S11, S12,
 181 S14 and S15, and particularly abundant at S8 (2150 reads, 3.4%), nearly 29 times more than the
 182 second most abundant site S2 (73 reads). *Prymnesium* showed the same pattern, recording the
 183 highest percentage at site S8 (7 reads, 0.01%) and haven't appeared on the other sites. While
 184 *Isochrysis* indicated very different property, with major peaks occurred at the sites S3 (5 reads)
 185 and none at the other sites. Most of algal species *Chrysochromulina* were abundant and fairly
 186 distributed at different stations with reads ranged from 158 to 899, except highest percentage
 187 (2.05 %, Site 8) and relatively lower at S6 (0%), S7 (19 reads, 0.03%), S9 (46 reads, 0.07%), S11 (30
 188 reads, 0.04%), S12 (0%) and S16 (54 reads, 0.08%). The relative abundance of *Haptolina* among
 189 different sites displayed similar pattern with *Chrysochromulina*, and the major peaks occurred at
 190 sites S8 (Fig 5).

Sampling sites

191

— ● — ● — ● — ● — ●

Figure 4. The number of reads detected in 16 sampling sites using 18s RNA amplicon analysis.

Figure 5. Spatial variations of relative abundance (%) of five Haptophyceae algae

(Prymnesium, Haptolina, Isochrysis, Chrysochromulina and Phaeocystis) at the different

stations in the northern coast of Beibu Gulf in Dec 2017.

Physical and chemical analyses of seawater characteristics

Physical and chemical characteristics of the seawater in the 16 sampling sites, including nutrient concentrations (TOC, $\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, $\text{NO}_2^-\text{-N}$, and phosphate [PO_4^{3-}]) and other environmental condition (chlorophyll-a, DO, salinity, pH, temperature of seawater), were detected. Just as our above discovery that relative abundance of some Haptophyceae algae were obviously higher at site S8 than others, some environmental factors at S8 were apparently special as well. Seawater temperature during the period of the study ranged from 15.4 to 18.5 °C and the lowest temperature appeared at site S8, while the salinity ranged from 25.7 to 37.3 ppt and S8 contained the lowest value 25.69 ppt. The highest value recorded for DO was 9.32 mg/L at site S8 and lowest was 8.33 mg/L at site S13. The highest value of pH was 8.71 at site 12 and lowest was 8.22 at S8. For the nutrient concentrations, the site S8 possessed the highest concentration of $\text{NO}_3^-\text{-N}$ (0.066 mg/L). Spatial distribution of seawater characteristics over the study region, including DO, $\text{NO}_3^-\text{-N}$, temperature of seawater and pH, were simply illustrated on the map by using blue and red colour (Figure 6).

237

238

239

240

241

242 Figure 6. Spatial distribution of seawater characteristics over the study region, including DO (a),
243 NO_3^- -N (b), T-temperature of seawater (c) and pH (d).

244

245 **The Correlation between environmental factor and Haptophyceae algae**

246 The correlation between the relative abundance of five kinds of Haptophyceae algae and
247 environmental factors of seawater (temperature, pH, salinity, DO, NO_3^- -N, NO_2^- -N, NH_4^+ -N,
248 Chlorophyll a and TOC) were analyzed. The results showed that Prymnesium, Phaeocystis and
249 Haptolina displayed highly positive linear correlation with the concentration of NO_3^- -N (Pearson
250 $r=0.85\sim 0.92$, $p<0.01$). Isochrysis and Chrysochromulina indicated high negative correlation with
251 temperature of sea water (Pearson $r=-0.88\sim -0.89$, $p<0.01$). Except for this, there was a significant
252 negative correlation just for Prymnesium, Haptolina, Chrysochromulina and Phaeocystis with pH
253 and salinity (Pearson $r=-0.52\sim -0.78$, $p<0.01$). Chrysochromulina revealed significant negative
254 correlation with seawater temperature (Pearson $r=-0.787$, $p<0.01$). Haptolina and
255 Chrysochromulina also exhibited significant negative correlation with Chlorophyll-a (Pearson
256 $r=-0.52\sim -0.53$, $p<0.01$). There was a significant positive correlation of Prymnesium and
257 Phaeocystis with DO (Pearson $r=0.534$, $p<0.01$), and also the same relationship of
258 Chrysochromulina with NO_3^- -N (Pearson $r=0.69$, $p<0.01$). There was no other obvious correlation
259 between other environmental factors and those five kinds of Haptophyceae algae (Fig 7).

260

Figure 7. The correlation between the relative abundance of five kinds of Haptophyceae algae and environmental factor (NO₃⁻-N, temprature, H, salinity, DO, NO₂⁻-N, NH₄⁺-N, Chlorophyll a and TOC) was indicated by Correlation matrix (Pearson's product moment correlation coefficient). The more the colure of shaded cells was close to red, the more significantly the correlation was positively related (P < 0.05); inversely, the more the colure of shaded cells was close to blue, the more significantly the correlation was negatively related (P < 0.05).

Chrysochromulina was the most abundant taxon of Haptophyceae algae in this study. The environmental factors NO₃⁻-N, pH and seawater temperature on the relative abundance of Chrysochromulina were studied by regression analysis. The concentration of NO₃⁻-N possessed significant positive impact on the relative abundance of Chrysochromulina (Pearson $r=0.692$, $p<0.01$; $R^2=0.479$; $Y=-0.001 + 0.281 \cdot X$, Y is the relative abundance of Chrysochromulina, X is the concentration of NO₃⁻-N; $F=12.859$, $P<0.05$; regression coefficient is 0.281, $P=0.003$). Conversely, the temperature of seawater indicated obvious negative influence on the relative abundance of Chrysochromulina (Pearson $r=-0.733$, $p<0.01$; $R^2=0.537$; $Y=0.123-0.007 \cdot X$, Y is the relative abundance of Chrysochromulina, X is the value of seawater temperature; $F=16.266$, $P<0.05$; regression coefficient is -0.007, $P=0.001$), and pH showed the same features on it (Pearson $r=-0.653$, $p<0.01$; $R^2=0.426$; $Y=0.282-0.032 \cdot X$, Y is the relative abundance of Chrysochromulina, X is the value of seawater pH; $F=10.384$, $P<0.05$; regression coefficient is -0.032, $P=0.006$).

There is consistent relation between the relative abundance of Phaeocystis and environmental factor NO₃⁻-N, DO, pH and temperature of sea water. The relative abundance of

Phaeocystis has obvious correlation with NO_3^- -N (Pearson $r=0.856$, $p<0.01$). Linear regression analysis indicated that Phaeocystis was significantly linearly related to the NO_3^- -N ($R^2=0.732$; $Y=-0.005 + 0.410 \cdot X$, Y is the relative abundance of *P.globosa*, X is the concentration of NO_3^- -N; $F=38.227$, $P<0.05$) and NO_3^- -N has a significant positive effect on *P.globosa* (regression coefficient is 0.410, $P=0.000$). The relative abundance of Phaeocystis had a positive correlation with DO (Pearson $r=0.535$, $p<0.01$; $R^2=0.286$; $Y=-0.140 + 0.016 \cdot \text{DO}$, $F=5.616$, $P<0.05$; regression coefficient is 0.016, $P<0.05$). The relative abundance of Phaeocystis was significant related to temperature of sea water (Pearson $r=-0.882$, $p<0.01$). Water temperature can explain the 77.8% change reason for the *P.globosa* ($R^2=0.778$), and has a significant effect on *P. globosa* ($Y=0.169-0.009 \cdot X$, $F=49.031$, $P<0.05$), and the regression coefficient is -0.009 ($P=0.000$) which indicated a significant negative impact relationship between them. The relative abundance of Phaeocystis also has negative correlation with pH of sea water (Pearson $r=-0.643$, $p<0.01$).

Prymnesium exhibited very similar condition with Phaeocystis, concerning relationship between their relative abundance and environmental characteristics. For example, the relative abundance of Prymnesium showed highly correlation with NO_3^- -N (Pearson $r=0.858$, $p<0.01$) and NO_3^- -N had a significant positive impact on the relative abundance of Prymnesium ($R^2=0.736$; $Y=-0.000 + 0.002 \cdot X$, Y is the relative abundance of Prymnesium, X is the concentration of NO_3^- -N; $F=39.079$, $P<0.05$; regression coefficient is 0.002, $P=0.000$). In contrast, the temprature of seawater exhibited obvious negative effect on Prymnesium ($R^2=0.776$; $Y=0.001 - 0.000 \cdot X$, Y is the relative abundance of Prymnesium, X is the value of temperature of seawater; $F=48.579$, $P<0.05$; regression coefficient is -0.000, $P=0.000$).

Discussion

NGS: a promising approach to study the community of algae in marine environments

P.globosa bloom occurred in winter from the year of 2014-2015 near the coast of Beibu Gulf annually. In order to discover the bloom mechanism, *P. globosa* was monitored between eastern Qinzhou Bay (EQB) and Dafenjiang River Estuary (DRE) during Nov 15th 2017 to Feb 15th 2018.

The beginning of the *P. globosa* bloom appeared from the middle several days of Jan in 2018, and disappearing occurred at the end of Mar in 2018. All the above findings were based on microscopy observations and cell counts, and it seems that the bloom appeared suddenly and instantly. We believed that finding out what happened before the emergence of *P. globosa*, not only by observations of microscopy and naked eyes, was very significant for our understanding of bloom mechanism of *P. globosa* in Beibu Gulf.

Classically, the algae ecological value is weighted based on the relative abundance of morphologically identified species. This traditional method is costly, time-consuming, and requires excellent taxonomic expertise, which is not always available [40]. Comparatively speaking, the eDNA and NGS approach for identification and quantification of algae open a new avenues for assessing and monitoring of aquatic ecosystems [41]. In our present work, by utilizing NGS based eDNA detection, not only the *P. globosa*, but also all the algae species belonging to Haptophyceae, taxonomically including *P. globosa*, were discovered and analyzed. The two most abundant OTUs in taxon Haptophyceae were affiliated with Chrysochromulina (5353 reads) and Phaeocystis (2298 reads), significantly more reads than Haptolina (228 reads), Prymnesium (9 reads) and Isochrysis (6 reads). The genus Chrysochromulina include two species *Chrysochromulina scutellum* and *Chrysochromulina* sp., both not very clear about their significance in ecosystem. But for the genus Phaeocystis, *P. globosa* was obviously detected among the samples, and this was in accord with our traditional monitoring approaches for the findings of *P.globosa* bloom in Beibu Gulf. Our result convincingly implicated that algae community reflected by eDNA and 18S ribosomal DNA analysis of the V4 region using the Illumina-Based Sequencing platform was suitable for monitoring the harmful algae in the early bloom stage. Many scientist had utilized HTS or NGS to detect algae in aquatic ecosystems and got some attractive results; for instance, NGS had been employed to microalgal diversity in the lichen Ramalina farinacea [42], Diatom resting stages in surface sediments [43], Diatom biomonitoring [40] and detection of harmful algal bloom species [44]. All the results show that eDNA and HTS sequencing is a promising approach to explore the community of algae in aquatic environments [45].

Hypothesis about the bloom mechanism of *P.globosa* in Beibu Gulf

Some mechanisms about the bloom of *P. globosa* could be proposed from our present work.

The first and most interesting finding was the bloom mode of *P. globosa*. In this study, at the early bloom stage, *P. globosa* was only obviously detected at site S8 with relatively much higher reads (2150) than other sixteen sites (148 in all). Therefore, the bloom of *P. globosa* may originate from a point of site (S8) and then spread to other regions, generally speaking, just as “diffusion from point to face”. Several evidences supported our opinion. Firstly, the marine environment in the Beibu Gulf was protected relatively better than other coastal zones in China [46]. Large area pollution, especially excessive nutrient concentration, had never appeared and been reported previously. Inversely, the probability of point source pollution in the coastal zone was even greater. The first appearance and flourish of *P. globosa* only at site S8 was probably because of its special condition, such as aquaculture farming nearby and consequent eutrophication phenomenon, rich nutrients brought by bottom-up stream and terrestrial drainage. Then the Phaeocystis was carried by transport of some water and drifted along the coastline under the influence of stream in Beibu Gulf. Some previous research could support our opinion. For instance, during the Phaeocystis bloom of the year 1957 around the coast of North Wale, Phaeocystis was only able to proliferate in Liverpool Bay, and then spread to other regions [47]; when Phaeocystis bloom in the coastal of north-western English Channel in 1990, Phaeocystis bloom emanated from near-shore, then spread towards the south-east in accord with the wind direction [48]. If our hypothesis is correct, releasing effective bio-agents and environmental controls on certain region to inhibit harmful algal blooms (HABs) *P. globosa* will be promising [49-51]. In this way, we only need to paid more attention to the early stage of *P. globosa* bloom, not after outbreaks in large scale, and this should be extremely effective.

The second interesting finding about the bloom of *P. globosa* was in relevant to the concentration of NO_3^- -N, pH, temperature and DO of seawater. More specifically, the bloom of *P. globosa* has a significant positive correlation with NO_3^- -N and negatively related to temperature of seawater. In the Beibu Gulf, the average temperature of seawater was above 15-16 °C and the most suitable temperature for growth of *P. globosa* is 15-16 °C [52], so the most relative abundance of *P. globosa* appeared first at site S8 may originate from its appropriate temperature of seawater. Nutrient elements, especially nitrogen and phosphorus, have obvious influence on

Phaeocystis blooms [52,53]. However, the impact of different forms of N-sources to the bloom of Phaeocystis had never been discovered previously. In our present work, it was the first time that the great correlation between NO_3^- -N and Phaeocystis bloom was illustrated. This result provided a helpful reference to our government on how to manage marine environment and control the *P.globosa* bloom. They should pay more attention to reduce the emission of nitrogen, especially NO_3^- -N, to the coastal zone of northern Beibu Gulf.

Acknowledgments

Supported by the National Natural Science Foundation of China (31570875); Guangxi Natural Science Foundation (2014GXNSFBA118135); Foundation of Guangxi Key Laboratory of Marine Disaster in the Beibu Gulf, Qinzhou University (No.2017TS03); Foundation of the Key Laboratory of Coastal Science and Engineering, Beibu Gulf, Guangxi (No.2016ZYB07).

Author Contributions

Conceptualization: Bin Gong

Funding acquisition: Bin Gong, Haiping Wu

Formal analysis: Bin Gong

Project administration: Bin Gong, Haiping Wu

Supervision: Haiping Wu

Validation: Jixian Ma, Meimiao Luo, Xin Li

Writing-original draft: Bin Gong

Writing-review & editing: Haiping Wu

Reference:

1. Roelke D L, Gable G M, Valenti T W, Grover J P, Brooks BW, Pinckney J L. Hydraulic flushing as a *Prymnesium parvum* bloom-terminating mechanism in a subtropical lake.[J]. Harmful Algae, 2010, 9(3):323-332.
2. Dahl E, Lindahl O, Paasche E, Throndsen J. The *Chrysochromulina polylepis* Bloom in Scandinavian Waters During Spring 1988[J]. 1989, 35:383-405.
3. Jiang T, Jiang T J, Zeng M, Peng L, Tong M M. Toxicity of red tide alga *Phaeocystis globosa* scherffel from Zhanjiang against young *Penaeus vannamei* boone and fish fry[J]. Journal of Tropical & Subtropical Botany, 2006, 14.
4. Peperzak L, Poelman M. Mass mussel mortality in The Netherlands after a bloom of *Phaeocystis globosa* (prymnesiophyceae)[J]. Journal of Sea Research, 2008, 60(3):220-222.
5. Shen P P, Qi Y Z, Wang Y, Huang L M. *Phaeocystis globosa* Scherffel,a harmful microalga,and its production of dimethylsulfoniopropionate[J]. Chinese Journal of Oceanology and Limnology, 2011, 29(4):869-873.
6. Yu R C, Lü S H, Liang Y B. Harmful algal blooms in the coastal waters of China[J]. Global Ecology and Oceanography of Harmful Algal Blooms, 2018: 309-316.
7. Wu H , Xu Y , Peng C , Liao Y, Wang X, & Jefferson T A. Long-term habitat loss in a lightly-disturbed population of the Indo-Pacific humpback dolphin, *Sousa chinensis*[J]. Aquatic Conservation Marine & Freshwater Ecosystems, 2017, 27(Supplement 1):1-11.
8. Liao Y, Hsieh H L, Xu S, Zhong Q, Lei J, Liang M. Wisdom of Crowds reveals decline of Asian horseshoe crabs in Beibu Gulf, China[J]. Oryx, 2017: 1-8.

- 413 9. Lai J, Jiang F, Ke K, Xu M, Lei F, Chen B. Nutrients distribution and trophic status assessment
414 in the northern Beibu Gulf, China[J]. Chinese journal of oceanology and limnology, 2014,
415 32(5): 1128-1144.
- 416 10. Wang Y J, Liao R Q, Liu W L, Kannan K, Ohura T, Wu M H. Chlorinated polycyclic aromatic
417 hydrocarbons in surface sediment from Maowei Sea, Guangxi, China: occurrence,
418 distribution, and source apportionment.[J]. Environmental Science & Pollution Research,
419 2017, 24(1):1-12.
- 420 11. Han Q Y, Huang X P, Xing Q G, Shi P. A review of environment problems in the coastal sea of
421 South China[J]. Aquatic Ecosystem Health & Management, 2012, 15(2):108-117.
- 422 12. Yang B, Kang Z J, Lu D L, Dan S F, Ning Z M, Lan W L, Zhong Q P. Spatial Variations in the
423 Abundance and Chemical Speciation of Phosphorus across the River-Sea Interface in the
424 Northern Beibu Gulf[J]. Water, 2018, 10(8): 1103.
- 425 13. Rouen M A, George D G, Hewitt D P. Using an automatic monitoring station to assess the
426 impact of episodic mixing on the seasonal succession of phytoplankton[J]. Internationale
427 Vereinigung für theoretische und angewandte Limnologie: Verhandlungen, 2000, 27(5):
428 2972-2976.
- 429 14. Kutser T, Metsamaa L, Strömbeck N, Vahtmäe E. Monitoring cyanobacterial blooms by
430 satellite remote sensing[J]. Estuarine Coastal & Shelf Science, 2006, 67(1):303-312.
- 431 15. Antonella P, Luca G. The quantitative real-time PCR applications in the monitoring of
432 marine
433 harmful algal bloom (HAB) species[J]. Environmental Science & Pollution Research
434 International, 2013, 20(10):6851-6862.
- 435 16. Mackenzie L A. In situ passive solid-phase adsorption of micro-algal biotoxins as a monitoring

436 tool.[J]. Curr Opin Biotechnol, 2010, 21(3):326-331.

437 17. Pang S J, Liu F, Shan T F, Xu N, Zhang Z H, Gao S Q. Tracking the algal origin of the bloom in
438 the Yellow Sea by a combination of molecular, morphological and physiological analyses[J].
439 Marine Environmental Research, 2010, 69(4):207-215.

440 18. Creer S, Deiner K, Frey S, Porazinska D, Taberlet P, Thomas W K, Potter C, Bik H M. The
441 ecologist's field guide to sequence-based identification of biodiversity[J]. Methods in
442 Ecology and Evolution, 2016, 7(9): 1008-1018.

443 19. Elbrecht V, Peinert B, Leese F. Sorting things out: Assessing effects of unequal specimen
444 biomass on DNA metabarcoding.[J]. Ecology & Evolution, 2017, 7(17):6918-6926.

445 20. Lanzén A, Lekang K, Jonassen I, Thompson E M, Troedsson C. High-throughput
446 metabarcoding of eukaryotic diversity for environmental monitoring of offshore oil drilling
447 activities[J]. Molecular Ecology, 2016, 25(17):4392-4406.

448 21. Divu D. Microsatellite DNA marker based molecular ecological evaluation studies in
449 Macrobrachium species[D]. Karnataka Veterinary, Animal and Fisheries Sciences University,
450 Bidar, 2009.

451 22. Cuvelier M L, Allen A E, Monier A, Mccrow J P, Monique M, Tringe S G. Targeted
452 metagenomics and ecology of globally important uncultured eukaryotic phytoplankton[J].
453 Proceedings of the National Academy of Sciences, 2010, 107(33): 14679-14684.

454 23. Steele H L, Streit W R. Metagenomics: advances in ecology and biotechnology[J]. FEMS
455 microbiology letters, 2005, 247(2): 105-111.

456 24. Połka J, Rebecchi A, Pisacane V, Morelli L, Puglisi E. Bacterial diversity in typical Italian salami
457 at different ripening stages as revealed by high-throughput sequencing of 16S rRNA

- 458 amplicons[J]. Food Microbiology, 2015, 46: 342-356.
- 459 25. Tedersoo L, Anslan S, Bahram M, Pölme S, Riit T, Liiv I, Kõljalg U, Kisand V, Nilsson H,
460 Hildebrand F, Bork P, Abarenkov K. Shotgun metagenomes and multiple primer
461 pair-barcode combinations of amplicons reveal biases in metabarcoding analyses of fungi[J].
462 MycoKeys, 2015 (10): 1-43.
- 463 26. Mortazavi R, Attiya S, Ariya P A. Arctic microbial and next-generation sequencing approach
464 for bacteria in snow and frost flowers: selected identification, abundance and freezing
465 nucleation[J]. Atmospheric Chemistry and Physics, 2015, 15(11): 6183-6204.
- 466 27. Kristiansen H B. Characterization of marine fungal communities using next generation
467 sequencing techniques[D]. 2014.
- 468 28. Medlin L K, Orozco J. Molecular techniques for the detection of organisms in aquatic
469 environments, with emphasis on harmful algal bloom species[J]. Sensors, 2017, 17(5): 1184.
- 470 29. Rees H C, Maddison B C, Middleditch D J, Patmore J R M, Gough K C, Crispo E. REVIEW: The
471 detection of aquatic animal species using environmental DNA-a review of eDNA as a survey
472 tool in ecology[J]. Journal of Applied Ecology, 2015, 51(5):1450-1459.
- 473 30. Elwood H J, Olsen G J, Sogin M L. The small-subunit ribosomal RNA gene sequences from the
474 hypotrichous ciliates *Oxytricha nova* and *Stylonychia pustulata*[J]. Molecular biology and
475 evolution, 1985, 2(5): 399-410.
- 476 31. Cheung, M. K., Au, C. H., Chu, K. H., Kwan, H. S. Wong, C. K. Composition and genetic
477 diversity of picoeukaryotes in subtropical coastal waters as revealed by 454 pyrosequencing.
478 ISME J. 4, 1053-1059 (2010).
- 479 32. Magoč T, Salzberg S L. FLASH: fast length adjustment of short reads to improve genome

480 assemblies[J]. Bioinformatics, 2011, 27(21): 2957-2963.

481 33. Caporaso J G, Kuczynski J, Stombaugh J, Bittinger K, Bushman F D, Costello E K, Fierer N, Peña
482 A G, Goodrich J K, Gordon J I, Huttley G A, Kelley S T, Knights D, Koenig J E, Ley R E, Lozupone
483 C A, McDonald D, Muegge B D, Pirrung M, Reeder J, Sevinsky J R, Turnbaugh P J, Walters W
484 A, Widmann J, Yatsunenko T, Zaneveld J, Knight R. QIIME allows analysis of high-throughput
485 community sequencing data[J]. Nature methods, 2010, 7(5): 335.

486 34. Edgar R C. UPARSE: highly accurate OTU sequences from microbial amplicon reads[J]. Nature
487 methods, 2013, 10(10): 996.

488 35. DeSantis T Z, Hugenholtz P, Larsen N, Rojas M, Brodie E L, Keller K, Huber T, Dalevi D, Hu P,
489 Andersen G L. Greengenes, a chimera-checked 16S rRNA gene database and workbench
490 compatible with ARB[J]. Applied and environmental microbiology, 2006, 72(7): 5069-5072.

491 36. Strickland J D H, Parsons T R. A practical handbook of seawater analysis[J]. 1972.

492 37. Aminot A, Krouel R. Dissolved organic carbon, nitrogen and phosphorus in the NE Atlantic
493 and the NW Mediterranean with particular reference to non-refractory fractions and
494 degradation[J]. Deep Sea Research Part I: Oceanographic Research Papers, 2004, 51(12):
495 1975-1999.

496 38. Ali U, Mahmood A, Syed J H, Li J, Zhang G, Katsoyiannis A, Jones K C, Malik R N. Assessing the
497 combined influence of TOC and black carbon in soil-air partitioning of PBDEs and DPs from
498 the Indus River Basin, Pakistan[J]. Environmental pollution, 2015, 201: 131-140.

499 39. Elbasiouny H, Abowaly M, Abu_Alkheir A, Gad A. Spatial variation of soil carbon and nitrogen
500 pools by using ordinary Kriging method in an area of north Nile Delta, Egypt[J]. Catena,
501 2014, 113: 70-78.

502 40. Vasselon V, Domaizon I, Rimet F, Kahlert M, Agns B. Application of high-throughput

503 sequencing (HTS) metabarcoding to diatom biomonitoring: Do DNA extraction methods
504 matter?[J]. Freshwater Science, 2017, 36(1):162-177.

505 41. Visco JA, Apothéloz-Perret-Gentil L, Cordonier A, Esling P, Pillet L, Pawlowski J.
506 Environmental Monitoring: Inferring the Diatom Index from Next-Generation Sequencing
507 Data[J]. Environmental Science & Technology, 2015, 49(13):7597-605.

508 42. Moya P, Molins A, Martínez-Alberola F, Muggia L, Barreno E. Unexpected associated
509 microalgal diversity in the lichen Ramalina farinacea is uncovered by pyrosequencing
510 analyses[J]. Plos One, 2017, 12(4):e0175091.

511 43. Piredda R, Sarno D, Lange C B, Tomasino M P, Zingone A, Montresor M. Diatom Resting
512 Stages in Surface Sediments: A Pilot Study Comparing Next Generation Sequencing and
513 Serial Dilution Cultures[J]. Cryptogamie Algologie, 2013, 38(1):31-46.

514 44. Linda K. Medlin, Jahir Orozco. Molecular Techniques for the Detection of Organisms in
515 Aquatic Environments, with Emphasis on Harmful Algal Bloom Species[J]. Sensors (Basel,
516 Switzerland), 2017, 17(5):1184.

517 45. Oliveira M C, Repetti S I, Iha C, Jackson C J, Pilar D T, Valéria C. High-throughput sequencing
518 for algal systematics[J]. European Journal of Phycology, 2018, 53(3):1-17.

519 46. Li J, Gao J, Wang Y, Feng X L, Fang X S, Cao P, Sun X Q, Ye W X, Khokiattiwong S,
520 Kornkanitnan N. Distribution and dispersal pattern of clay minerals in surface sediments,
521 eastern Beibu Gulf, South China Sea[J]. Acta Oceanologica Sinica, 2012, 31(2): 78-87.

522 47. Jones P G W, Haq S M. The distribution of Phaeocystis in the eastern Irish Sea[J]. ICES Journal
523 of Marine Science, 1963, 28(1): 8-20.

524 48. Davies A G, De Madariaga I, Bautista B, Fernández E, Tranter P R G. The ecology of a coastal

525 Phaeocystis bloom in the north-western English Channel in 1990[J]. Journal of the Marine
526 Biological Association of the United Kingdom, 1992, 72(3): 691-708.

527 49. Brussaard C P D. Viral Control of Phytoplankton Populations-a Review 1[J]. Journal of
528 Eukaryotic Microbiology, 2004, 51(2): 125-138.

529 50. Weston K, Greenwood N, Fernand L, Pearce D J, Sivyer D B. Environmental controls on
530 phytoplankton community composition in the Thames plume, U.K.[J]. Journal of Sea
531 Research, 2008, 60(4):246-254.

532 51. Guan C, Guo X, Cai G, Zhang H J, Li Y, Zheng W, Zheng T L. Novel algicidal evidence of a
533 bacterium *Bacillus* sp. LP-10 killing *Phaeocystis globosa*, a harmful algal bloom causing
534 species[J]. Biological Control, 2014, 76(3):79-86.

535 52. Schoemann V, Becquevort S, Stefels J, Lancelot C. Phaeocystis, blooms in the global ocean
536 and their controlling mechanisms: a review [J]. Journal of Sea Research, 2005, 53(1):43-66.

537 53. Riegman R. Nutrient-related selection mechanisms in marine phytoplankton communities
538 and the impact of eutrophication on the planktonic food web[J]. Wat.sci.technol, 1995,
539 32(4):63-75.

540

541

542

543

544