Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

1 **Title: Paternal grandparental exposure to crop failure or surfeit during a childhood slow**
2 **growth period: Epigenetic marks on grandchildren's growth, glucoregulatory and stress**
3 **genes**

4 **Authors:** Lars Olov Bygren^{1,2*}, Patrick Müller¹, David Brodin¹, Gunnar Kaati¹, Jan-Åke
5 Gustafsson¹, John G. Kral³

6 **Affiliations:**

7 ¹ Department of Biosciences and Nutrition, Karolinska Institutet, SE_14183 Huddinge,
8 Sweden.

9 ² Departments of Community Medicine and Rehabilitation, Umeå University, SE_90187,
10 Umeå, Sweden.

11 ³ Departments of Surgery, Medicine, Cell Biology, SUNY Downstate Medical Center, 450
12 Clarkson Avenue, Box 40, Brooklyn, NY 11203, USA.

13 * Correspondence to: lars.olov.bygren@ki.se

14 **ABSTRACT**

15 This latest in our series of papers describes transgenerational methylation related to mid-
16 childhood food availability in 19th century Överkalix, Sweden. Failed vs. bountiful crops
17 differentially influenced methylation in grandchildren of paternal grandparents exposed to
18 feast or famine during their Slow Growth Period (SGP), a sensitive period preceding the pre-

Bygren & al. Early life Feast and Famine: The Methyome of disparate stressors inherited

19 pubertal growth spurt. In this case-study of 8 tracked 75-year old progeny with differential
20 ancestral exposure, we found, in 40 posited gene ontology pathways, 39 differentially
21 methylated CpG regions (DMRs) related to famine, excess food and food-insecurity stress, 9
22 of which with DMRs above 5%. Three gene ontology terms (GOs) “insulin processing”,
23 “adipose development” and “hypothalamus development” were key, with DMRs >14%.
24 An unbiased test of known pathways revealed four nuclear transcription factors upstream of
25 promoters repressing the pathway following paternal grandparental famine experience, as
26 well as 4 upregulated GOs with average DMRs >20%. We conclude that this is the first
27 demonstration of human transgenerational inheritance of epigenetic marks following
28 ancestral childhood exposure to variable food availability inducing early developmental
29 origins of adult disease.

30 Paternal grandparent food supply, preceding their prepubertal growth spurt, induced
31 epigenetic marks in three gene pathways reflecting famine, excess food and food-insecurity
32 stress. Our epidemiological findings of adverse transgenerational effects of ancestral
33 overnutrition and conversely, beneficial effects of famine during SGP¹⁻⁶ prompted us to
34 study individuals whose paternal grandparents were randomly exposed during their Slow
35 Growth Period (SGP) - a sensitive period preceding the pre-pubertal growth spurt - to failed
36 or bumper crops potentially engendering epigenetic marks on mechanistic molecular gene
37 pathways.

38 For detailed analyses of specific gene ontology (GO) pathways, we tested three posited
39 structural classes of genes associated with each of 40 preselected GO terms
40 (<http://amigo.geneontology.org>) for differentially methylated CpG regions (DMRs) between
41 the experimental groups. We selected GO pathways with average methylation differing >
42 5%. The posited pathways comprised glucoregulatory genes interpreted as immediate
43 sensors of decreased glycogen stores, whereas those of lipids and ketones were posited to
44 reflect the duration and magnitude of the famine or alternatively, responses to
45 overnutrition. Food insecurity stress was assumed to have activated the HPA-axis. We
46 focused on two putative response lines in direct descendants from a) grandfather to son
47 with grandson and b) paternal grandmother to son with granddaughter, adjusted for crop
48 size during famine or surfeit. We found 39 DMRs in grandchildren after grandparental

Bygren & al. Early life Feast and Famine: The Methyome of disparate stressors inherited

differential exposure to food availability in GO-pathways associated with famine, excess food and environmental stressors. Nine of them were large, three GOs exhibiting remarkable DMRs in grandchildren (Table 1). An unbiased analysis detected transcriptional pathways, such as CCAAT-box (AKA C/EBPbeta) affecting adipogenesis, laying down fat, the primal tissue for energy storage.

RESULTS

Exploiting an “experiment” of nature, we studied DNA methylation in grandsons and granddaughters in the paternal lineage. We found 39 DMRs among 40 posited pathways in *islands*, North and South *shores* and North and South *shelves* related to famine, nutrient excess and stress, among which 9 pathways had DMRs above 5%, and 3 above 14%. The other 31 pathways had 0-5% DMRs. An unbiased test of pathways in *islands* revealed 2 pathways with DMRs 40% and 2 with DMRs 22 and 21% (Table 1b).

Paternal grandparental exposure vs. methylated genes in adult grandchildren.

The main metabolic pathways implicated in grandpaternal SGP exposure to food-insecurity stress transferred to grandsons were “appetite”, “insulin processing”, “ketone body catabolic process” and “hypothalamic development”. Differential grandmaternal exposure to feast or famine was associated with marks in “adipogenesis”, “insulin-like growth factor receptor (IGF-R) binding”, “ketone body catabolic process” and “adrenal gland development” (Table 1). The probands’ 4 ketone body catabolism genes (GO 30070) were the same for grandfathers’ as for grandmothers’ exposures, supporting the findings. The directions of the differences were intuitive: famine decreased energy-demanding pathways whereas overnutrition enhanced them. Altogether three pathways were key.

Three key gene ontology terms implying transgenerational epigenetic inheritance.

GO 0030070: “insulin processing” modulates insulin production downstream from proteolysis of the precursor preproinsulin via C-peptide. Proinsulin is cleaved to release C peptide. This pathway had a DMR in the male line in the Southern Shore involving 2 genes: PCSK2, proprotein convertase (also called NE2 HUMAN) in chromosome 20 and CPE Carboxypeptidase E (also called CPBE HUMAN) in chromosome 2. This pathway is related to GO 1901142: “insulin metabolic process”.

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

78 GO 1904179: "adipose development" includes gene NR1H4 (also called BAR;FXR;HRR1;
79 RIP14) and the nuclear hormone receptor pthr 24082, "bile acid receptor". Other genes are
80 SIRT1 (also called SIR2L1) and chromatin regulatory protein sir2, pthr 11085.

81 GO 0021854: "hypothalamus development" involves GSX1 (GShomeobox1), PTX2 (Pituitary
82 homeobox2), UBB (Polyubiquitin-B) and CRH (Corticoliberin; Corticotropin-releasing
83 hormone).

84 *Unbiased correlations (Table 1b).*

85 Regardless of sex, the following gene ontology terms were highly correlated with famine:

86 GO 0016602: A transcription complex that binds to the CCAAT-box upstream of promoters
87 typically consisting of three subunits, i.e. NFYA-c. Four genes were repressed: NFYA on
88 Chromosome 6, NFYB on chromosome 12, NFYC on chromosome 1 and ING2 on
89 chromosome 4.

90 GO 0045540: "Regulation of cholesterol biosynthesis" (formation of cholesterol) was
91 repressed in 13 genes, such as ERLIN1, chromosome 8 and 10; SEC14L2, chromosome 22 and
92 SODI, chromosome 21.

93 Several cell differentiation pathways correlated with food surplus, notably the following two:

94 GO 0030854: "Positive regulation of granulocyte differentiation" available for expression of
95 its 9 genes, e.g. HAXI on chromosome 1, OGT on the X chromosome, ILS on chromosome 5
96 and HCSL1 on chromosome 3.

97 GO 0045616: "Regulation of keratinocyte differentiation" with 16 genes, such as HAXI, OGT,
98 ILS and C1QC on chromosome 1.

99 *Parental age data and selection*

100 Average paternal ages at birth of first child were 32 , 31 and 27 years in the three
101 generations with grandparents exposed to famine and 34 , 27 and 27 years with
102 grandparents exposed to feast. Seven index persons were alive at 74, the eighth having died
103 at 63 years. For grandparents and parents exposed to famine during SGP, the average ages
104 of death were 61 and 78 years respectively whereas after feast they were 87 and 71 years

Bygren & al. Early life Feast and Famine: The Methylome of disparate stressors inherited

(Table 2). Sequential genetics findings contributing to differential methylation were not taken into account.

DISCUSSION

We propose that known adaptations to energy deficiency or excess in cells, organs or organisms are expressed transgenerationally, appearing as epigenetic marks in humans. Famine might induce marks around promoters in at least two types of pathways: one general or nutrient-specific, related to *diminished* substrate stores of glycogen, lipid or protein, or *excess*, nutritoxic exposure (cf. eutrophication in plants), the other to environmental stresses of famine through activation of the hypothalamo-pituitary-adrenal (HPA) axis, as in food insecurity.

In earlier papers we described a unique cohort of descendants of grandparents exposed randomly to these weather-dependent variations in food availability. Data emanated from detailed archival records enabling correlations between demographic and crop yield statistics¹⁻⁶. Our finding of SGP sensitivity findings have been replicated in Germany and Sweden^{7,8}. Our current paper adds mechanistic information based on blood sampling of seven 75-year-old grandchildren and one 63 years old, enabling determination of candidate epigenetic markers reflecting differential methylation in gene pathways posited to be affected by exposure of their grandparents to crop failure versus bountiful harvest during the grandparental childhood SGP preceding the prepubertal peak stature growth. Changes in DNA methylation levels of cytosine and guanine (CpGs) differentially methylated in one generation could explain transgenerational epigenetic inheritance in the third generation⁹.

A 2015 review concluded that evidence suggesting that acquired epigenetic marks pass to the next generation was limited; many others have noted the absence of any mechanism by which gene-regulatory information is transferred from somatic cells to germ cells in the study of transgenerational epigenetic inheritance (review: Nagy and Turecki¹⁰). Our epidemiological and epigenetic results demonstrate sex differences, e.g. in development of preproinsulin as has been seen in human pancreatic islets¹¹. The sex-bound epigenetic inheritance has been interpreted simply as epigenetic actions and disease manifestations being sex specific, as is the case with many conventional genetic variants¹². The transcriptions are also not fully understood. Most confusion emanates from not knowing

Bygren & al. Early life Feast and Famine: The Methylome of disparate stressors inherited

how exposed somatic cells can communicate their exposure to the germline which induces changes lasting for generations.

It is difficult to understand or accept epigenetic inheritance if there is a real rather than just a theoretical barrier between somatic and germline cells. One route to overcome this barrier might be the proposal that extracellular nano-vesicles, neighboring germ cells, shed by somatic cells contain the material required for transcription, able to bypass the barrier¹³. In the SGP the male primordial germ cells form active spermatozoa whereas human ovarian stem cells that can be modified in the SGP probably are required.

Presently the preponderance of evidence suggests transgenerational cumulative effects of exposures: diet, behavior, environmental chemicals, activity and the microbiome. For reviews, see Sales et al.⁹ and Vaiserman et al.¹⁴.

We have focused on the paternal line to discern transgenerational epigenetic inheritance disentangled from maternal-fetal and maternal-infant influences. The pathways in the paternal line are probably similar in the maternal line. We found three interesting gene pathways influenced by paternal grandparents' exposure resulting in grandchildren's DMRs, pathways induced by famine, overnutrition and food insecurity stress. When the paternal grandfather had been exposed to famine, the grandson exhibited DMRs of insulin processing. When the paternal grandmother had been exposed the granddaughter had DMRs of GOs related to environmental stress such as "hypothalamus development"(influencing the HPA-axis) related to increased female susceptibility to stress, yet explaining the benefit to mental health recently found in a different setting⁷.

A strength of our study is that it exploits a natural secular phenomenon viz. variable food security, the effects of which are documented in a homogeneous well-documented 19th century population, allowing exceptional tracking of ancestors. Few founders colonized the area 500 years earlier whereby the area metaphorically became an island of native speakers of Swedish surrounded by Sami- and Finnish-speaking neighbors. Furthermore the index persons, the grandchildren, are dispersed all over Sweden⁵ and have neither been exposed to famine nor, being born before WW2, experienced current food excesses during their childhood, preceding the current obesity epidemic. Differences in blood tissue composition

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

or heterogeneity of blood probably were of minor importance in this very homogenous population.

A second strength is the focus on two lines of inheritance that emerged in the epidemiological studies, the line from the paternal grandmother's son and his daughter and the line from the paternal grandfather with son and grandson. Thirdly our *a priori* selection of gene ontologies, GOs, based on our earlier epidemiological findings of cardiovascular morbidity related to food security reduces the risk of spurious correlations. A confound of multiple testing could be suspected having 40 GOs but the ontologies were chosen for their evidence-based responses to only three potential pathways: feast, famine and stress. In addition, our unbiased test identified related transcriptional genes.

Our earlier published studies in the community and correlation of age at the birth of the first child, number of children and variance of survival over three generations ruled out that variable availability of food during SGP was caused selectively². In human studies, measures are very crude but the figures for the eight subjects do not demonstrate any biased selection owing to ancestors' random exposure during the SGP. The main weakness is having only 8 index cases, with low statistical power. Furthermore, seven were survivors up to age 75 years (one to 63) during which their own exposures might have induced methylations confounding those inherited from the grandparents and parents. On the other hand, epimutations analogous to mutation most often do not appear on the clinical horizon until late in life. The subjects were sufficiently healthy to participate and travel to the nearest health facility to give blood.

Confounding could have occurred through differences in exposure to the feast and famine between index cases owing to social circumstances such as belonging to a family with better food resources, or to other mechanisms differentially affecting allele-specific or other variables' methylation¹². However, the natural experiment exposed all families with little variation in poverty in the 19th century mitigating this confound, as supported by our earlier research in the community⁴. The isolation of this community during the centuries up to the present might have resulted in a relatively inbred cohort and diminished assortative mating, both benefitting the epigenetic analysis. We present a case series of epigenomes of 8 adult grandchildren with archival data exhibiting their paternal grand-paternal or -maternal

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

exposure to crop failure or large crops during their SGP linked to differential methylation of genes associated with energy balance and hypothalamic development. The salient novel finding is that grandparental childhood exposure was reflected in grandchild profiles of CpG methylation of key genes. Our epidemiological data from the cohort from which these strategic cases were drawn strengthens the molecular findings and indicates possible mechanisms of inheritance that might be important for the field. The presence of epigenetic changes attributable to environmental influences potentially engendering several prevalent multifactorial chronic diseases and conditions. These findings replicated in other settings may provide guidance for preventive and therapeutic interventions targeting methylation.

METHODS

Classification of exposure to food availability in the environment

A seven point scale for estimating annual crop sizes was employed during 1816-1849 [Statistics Sweden (Tabellverket)]¹⁵. It spanned from “total failure” to “abundant” crops and has been used since in demographic research. For the years 1865-1902, birth years of the 8 grandparents, statistical tables of crop yields were the primary source, validated against food price statistics¹⁶ and qualitative reports from a gubernatorial *Crops and National Aid Office* (Hushållningssällskapet). Using these sources, we created the ordinal seven-item “Hellstenius scale”. Over the years the criteria for the items of the scale have changed apace with changes in the Swedish language. The current translation is as follows: Total failure (0), Sparse general growth (1), Weak or Small (2), Below average (3), Average or Mediocre (4), Above average (5) and Good or Abundant (6). We defined 0-2 as famine, 3-4 as mediocre, and 5-6 as excess food availability. We used May 1 of the year following crop failure as the time of least food availability in this area around the 66th latitude and determined November 1 of the year following good harvest with pig and cattle slaughter when the meat could be frozen, as the time of greatest food availability (excess, eutrophic). Grandparent age on those dates was used to relate food availability to his or her SGP.

SGP, the slow growth period

Two developmental periods in our first analysis were posited to differ in sensitivity to food supply affecting transgenerational responses: the prepubertal growth peak and the

Bygren & al. Early life Feast and Famine: The Methylome of disparate stressors inherited

preceding slow growth period (SGP). The growth peak was seen as a sensitive period for a child to be exposed to famine. The slowest growth period before the peak was posited to be less prone to energy variation. These two periods were set in relation to puberty, taking into account individual variation, nutrient availability shifting the timeframes, and realizing that the age distribution of puberty was skewed to the right during the 19th century compared to later. The two periods were derived by superimposing the stature growth velocity curves of Prader et al.¹⁷ upon the ages of 19th century pubarche described by Tanner¹⁸. The periods for ancestors in the 19th century were set at the ages 8-10 years for female ancestors and at 9-12 years of age for male ancestors. Excess food was not understood to be a problem in the 19th century in Överkalix but to analyze potential transgenerational responses we were obliged to focus on the SGP where we indeed found mismatches reflecting transgenerational responses not only to famine but also to excess food availability. We introduced the concept of the “slow growth period”¹, the causes and mechanisms of which we discuss here in depth.

Samples and pedigrees

Eight subjects, seven of which 75 years of age and one 63, consented to blood sampling. They were selected from the 1935 birth cohort in an original epidemiologic study of transgenerational responses to variable availability of food during ancestors’ SGP in Överkalix, Sweden, described in Tingshög et al.⁵ During SGP two paternal grandmothers were exposed to famine (grade 0-2) in 1867 and 1900 respectively, and two to surfeit (grade 5-6) in 1871 and 1879. Two paternal grandfathers were exposed to famine (grade 0-2) in 1867 and 1877 respectively and two were exposed to surfeit (grade 5-6) in 1887 and 1881. These subjects represented four pairs of grandchildren of paternal grandparents exposed to feast or famine (Fig1). Individuals inheriting grandparents who had been exposed to medium crops (grade 3-4) were not included in this strategic sample.

Whole blood samples in EDTA were drawn at county health service units close to the subjects’ residence, frozen at 70° C and sent to the project center. Pedigrees were originally tracked at parish offices in Sweden and Finland and presently in the National archive, the Regional archive of north Sweden, the Research archive of Umeå University, the National tax bureau and digitized genealogical net-based registries. Detailed possible confounders were not identified in this natural experiment: probands, the grandchildren, did not experience

Bygren & al. Early life Feast and Famine: The Methyome of disparate stressors inherited

any lifetime exposure to famine. The modern food availability though was there from their young adulthood.

Metabolic pathways posited

The gene ontologies chosen *a priori* described processes related to cardiovascular risks related to hunger, excess food and reactions to environmental stress (<http://amigo.geneontology.org>). Posited processes during *famine* were “glucose homeostasis” and “glucose transport”, “ketone body biosynthesis and catabolic processes”, “cellular ketone body metabolic pathways”, and “cholesterol homeostasis”. In *hunger and excess* “insulin processing and binding”, “insulin-like growth factor”, “insulin receptor signaling” and “autophagy pathways” were posited. Individuals’ emotional reactions to the stress of famine were assumed to influence hypothalamic and adrenal gland development. Availability and absorption of folate providing methyl donors was of obvious interest, though not the focus of this report. Other posited GOs include appetite, ghrelin, leptin, calcium signaling, eating behavior, adipogenesis, insulin secretion, and lipids.

Altogether, 40 pathways were tested, many of them overlapping and covering three kinds of response pathways reflecting the 3 posited mechanisms.

Pathways undergoing unbiased testing

We tested all gene ontology terms in a public database, (UniProt-GOA) including about 40 000 gene ontologies, for grandchildren’s responses to paternal grandparental exposure to poor *versus* excess food.

Genome-wide methylation

Genomic DNA extraction from whole-blood (buffy coat in one) used the GeneCatcher Blood kit (Invitrogen, Carlsbad, CA, USA). 500 ng of genomic DNA was bisulfite converted with EZ-96 DNA Methylation kit (Zymo Research, Irvine, CA, USA) and genome wide DNA methylation analysis was carried out using the Infinium MethylationEPIC Bead Chip (Illumina, San Diego, CA, USA). The Laboratory procedures followed the manufacturers’ protocol.

The array was designed for genome wide methylation analysis with coverage across gene regions with sites in the promoter region, 5UTR, enhancer and gene body, and interrogating

Bygren & al. Early life Feast and Famine: The Methyome of disparate stressors inherited

more than 850 000 methylation sites at single nucleotide resolution. This technique uses two different probe types (Infinium 1 and 2) with different characteristics, thus requiring normalization to reduce technical bias. GenomeStudio software, version 2011.1 (Illumina Inc.), was used for analysis, visualization and extraction of methylation data.

Methylation levels (beta values) were estimated as the ratio of signal intensity of the methylated alleles to the sum of methylated and unmethylated intensity signals. The beta-values vary from 0 (no methylation) to 1 (100% methylation).

The chip covers CpG islands, shores, shelves and the promoted genes. GOs containing genes and biological pathways were assigned from the literature, primarily from the Gene Ontology project (geneontology.org) chosen for their known intra-generational relation to excess and lack of food.

The assay protocol combined bisulfite conversion of genomic DNA and whole-genome amplification with array-based capture and scoring of the CpG loci. Signal intensity was measured by scanner to generate beta values, the degree of methylation at a locus. Allele-specific single base extension of the probes incorporated a biotin nucleotide or a dinitrophenyl labeled nucleotide. Signal amplification of the incorporated label further improved the overall signal-to-noise ratio of the assay.

Human genomes punctuated by DNA sequences with high frequencies of CpG sites, >200 bases, with CpG content of 50% were termed “*islands*” (CGIs). Differential methylation induced by environment visible in CGIs or in “northern- and southern-*shores*”, within 2 kilo base-pairs (<2kbp) of the islands or in “northern- and southern-*shelves*” within another 2 kilo base pairs from the shores (<2kbp) were studied in CGIs in grandchildren of grandparents exposed to extremes of food availability during their childhood SGP. They are presented as DMRs expressed as percent.

Statistical analysis

Associations between genes and GO terms were retrieved from the Gene Ontology Association (UniProt-GOA) Database, <http://www.ebi.ac.uk/GOA>.

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

Number of genes associated with a GO term and represented with at least one probe on the EPIC chip and number of probes associated with the genes were analyzed among index cases, the grandchildren.

Differentially methylated female index cases whose paternal grandmothers were exposed to bumper harvests in their mid-childhood SGP and to crop failure, respectively, were recorded as well as male index cases whose paternal grandfather had the same extreme exposures.

The criteria for differential methylation was a p-value < 0.05 and an average difference in beta-value between the group averages >0.1. No adjustment for multiple comparisons was carried out. The decreased power of genome-wide analysis, by having negative controls in the remainder of the genome, was compensated by the strict criteria chosen.

Tables were prepared after beta-mixture quantile normalization (BMIQ) correcting probe design bias and with BMIQ-normalized beta values and annotations. The analyses were carried out in Bioconductor-package ChAMP package default probe filtering based on detection p-value, minimum bead count and probes possibly confounded by SNPs whereupon cross hybridization (<https://www.ncbi.nlm.nih.gov/pubmed/24063430>) left 813007 probes for further analysis.

REFERENCES

1. Bygren LO, Kaati G, Edvinsson E. Longevity determined by paternal ancestors' nutrition during their slow growth period. *Acta Biotheoretica* 2001;49:53-59.
2. Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality determined by nutrition during parents' and grandparents' slow growth period. *Eur J Hum Genet* 2002;10:682-8.
3. Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K et al. Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 2006;14:159-166
4. Kaati G, Bygren LO, Pembrey M, Sjöström M. Transgenerational response to nutrition, early life circumstances and longevity. *Eur J Hum Genet* 2007;15:784-90.
5. Tingshög P, Carstensen J, Kaati G, Edvinsson S, Sjöström M, Bygren LO. Migration and mortality trajectories: a study of individuals born in the rural community of Överkalix. *Soc Sci Med*. 2011;73:744-51.
6. Bygren LO, Tingshög P, Carstensen J, Edvinsson S, Kaati G, Pembrey ME,

Bygren & al. Early life Feast and Famine: The Methylome of disparate stressors inherited

- 343 Sjöström M. Change in paternal grandmothers' early food supply influenced
344 cardiovascular mortality of the female grandchildren. BMC Genet 2014;15:12.
- 345 7. Van den Berg GJ, Pinger PR. Transgenerational effects of childhood conditions on third
346 generation health and educational outcomes. EconHumBiol 2016;23:103-120.
- 347 8. Vågerö D, Pinger PR, Aronsson V, van den Berg G. Paternal grandfather's access to food
348 predicts all-cause and cancer mortality in grandsons. Nat Comm
- 349 9. Sales VM, Ferguson-Smith AC, Patti ME. Epigenetic mechanisms of transmission of
350 metabolic disease across generations. Cell Met 2017;25:559-571.
351
- 352 10. Nagy C, Turecki G. Transgenerational epigenetic inheritance: an open discussion.
353 Epigenomics 2015;7:781-790.
- 354 11. Hall E, Volkov P, Dayeh T, Esguerra JLS, Salö S et al. Sex difference in the genome-wide DN
355 A methylation pattern and impact on gene expression, microRNA levels and insulin secretion
356 in human pancreatic islets. Genome Biology 2014;15:522-544.
357
- 358 12. Guo YT, LUO HV, Wu YM, Magdalou J, Chen LB, Wang H. Influencing factors, underlying
359 mechanisms and interactions affecting hypercholesterolemia in adult offspring with caffeine
360 exposure during pregnancy, Reprod Toxicol 2018;79:47-56
- 361 13. Earon SA, Jayasooriah N, Buckland ME, Buckland ME, Martin DI, Cropley JE, Suter CM. Roll
362 over Weismann: extracellular vesicles in the transgenerational transmission of environmental
363 effects Epigenomics 2015; 7:1165-71.
- 364 14. Vaiserman AM, Koliada AK, Jirtles RL. Non-genomic transmission of longevity between
365 generations: potential mechanisms and evidence across species. Epigenetics Chromatin
366 2017;DOI 10.1186/s 13072-017-0145-1
- 367 15. Hellstenius, J. Skördarna i Sverige och deras verkningar. [Harvests in Sweden and their
368 repercussions]. Statistisk Tidskrift 77-119. Stockholm, Sweden 1871 (sic).
369
- 370 16. Jörberg, L. (1972). A history of prices in Sweden 1732–1914. CWK Gleerup, Lund,
371 Sweden.
- 372 17. Tanner, J.M. A History of the Study of Human Growth. Cambridge University Press,
373 Cambridge 1981
- 374 18. Prader, A., R.H. Largo, L. Molinari and C. Issler (1989). Physical growth of Swiss children
375 from birth to 20 years of age. First Zurich Longitudinal Study of Growth and Development.
376 Helvetica Paediatrica Acta 43:Suppl 52: 1-125

377 Figure 1. Flowchart

Bygren & al. Early life Feast and Famine: The Methylome of disparate stressors inherited

Table 1. Gene ontology pathways related to hunger, excess food and stress associated with paternal grandparental exposure to variable food availability with DMRs in grandchildren.

Gene Ontology pathway	Genes	CpGs	MethylC pG	DMR %	Shore, Shelf ³
Paternal grandfather with son and grandson					
32100 Positively regulating appetite	3	163	10	6%	Northern shore
30070 Insulin processing ¹	2	4	1	25%	Southern shore
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
21854 Hypothalamus development ²	6	33	5	15%	Southern shelf
Paternal grandmother with son and granddaughter					
1904179 Positive adipose development	2	11	2	18%	Northern shore
70341 Fat cell proliferation ⁴	5	16	3	18%	Southern shelf
51599 IGF receptor binding ⁵	9	94	12	13%	Northern shelf
46952 Ketone body catabolic process	4	16	1	6%	Southern shore
30325 Adrenal gland development	12	57	4	7%	Southern shelf

¹Mature insulin formed by proteolysis of the precursor pre-proinsulin. The sequence starts with pre-proinsulin; cleavage of proinsulin leads to release of C-peptide, leaving the A and B chain of mature insulin.

²Evolution of the hypothalamic region of the forebrain from its initial formation to its mature state.

³CpG sites with >200 bases with CpG content of 50% are termed “islands” (CGIs). Differential methylation induced by environment is most often seen in “northern- and southern-shores” <2kb from the islands, and “northern- and southern shelves” <2kb apart from the shores.

⁴Multiplication or recruitment of fat cells leading to expansion of their population (adipogenesis).

⁵Interacting selectively and non-covalently with insulin-like growth factor receptor.

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

Table 1 b

Gene ontology (GO) pathways in 8 grandchildren (4 men, 4 women) whose paternal grandparents were exposed to famine or food excess.

Gene ontology pathway %	GO name	DMR
<u>Exposure</u>		
GO:0016602 ^a	Famine	CCAAT binding complex 40
GO:0045540 ^a	Famine	Regulation of cholesterol biosynthetic process 40
GO:0030854 ^a	Excess	Positive regulation of granulocyte differentiation 22
GO:0030852 ^a	Excess	Regulation of keratinocyte differentiation 21

Beta cutoff 0.02, Methylation difference >15%.

		CpGs	Methylated CpGs	p-value,	exponent
		observed	expected		
GO: 0016602 ^a	47	5	0.255	6.13	-6
GO: 0045540 ^b	407	5	0.221	3.12	-6
GO: 0030854 ^c	75	6	0.017	1.53	-14
GO: 0030852 ^d	260	7	0.059	1.95	-13

Numbers of genes in GO pathways, their ID, chromosome #:

^a n= 4 NFYA, 6; NFYB, 12; NFYC, 1; ING2, 4.

^b n=13 ERLIN1, 10; ERLIN2, 8; SEC14L2, 22; SOD1, 21; APOE, 19; APOB, 2; FGF1, 5; POR, 7; SREBF1, 17; ABCG1, 21; SCAP, 3; PRKAA1, 5; DHCR7, 11.

^c n= 9 HAX1, 1; OGT, X; IL-5, 5; HCLS1, 3; RUNX1, 21; KMT2E, 7; TESC, 12; TRIB1, 8; LEF1, 4.

^d n=16 HAX1, 1 ; OGT, X ; C1QC, 1 ; IL-5, 5 ; RARA, 17 ; CEACAM1, 19 ; HCLS1, 3 ; RUNX1, 21 ; CUL4A, 13 ; ADIPOQ, 3 ; ZBTB46, 20 ; KMT2E, 7 ; INPP5D, CHR_HG2232_PATCH; TESC, 12; TRIB1, 8; LEF1, 4.

Bygren & al. Early life Feast and Famine: The Methyloome of disparate stressors inherited

425 Table 2. Reproductive fitness following grandparental famine or feast in the Slow Growth
426 Period

Index person	Age at first child's birth			Number of children		Longevity		
Exposure	Grandparent	Father	Grandchild	Father	Grandchild	Grandparent	Father	Grandchild
Paternal grandfathers' exposure								
Famine								
Index person No 15	39	38	NA	4	NA	60	69	74+
Index person No 53	27	31	27	7	1	49	98	74+
Feast								
Index person No 14	28	23	30	6	3	80	70	74+
Index person No 23	33	26	NA	11	NA	94	83	63
Paternal grandmothers' exposure								
Famine								
Index person No 59	39	38	34	3	2	80	83	74+
Index person No 86	24	20	22	6	2	55	64	74+
Feast								
Index person No 3	36	29	23	10	4	86	65	74+
Index person No 22	35	31	29	4	2	89	68	74+

NA. Not available