Novel, economically important semi-dwarf and early mutants: Selection and development from Improved White Ponni Rice (*Oryza sativa* L.)

***S. Ramchander*¹, *Andrew Peter Leon*², *Jesima Khan Yasin*³, *KK. Vinod*⁴ and *M. Arumugam Pillai*^{2*}**

¹ *Visiting Scientist (SERB –National Post-Doctoral Fellow),*

IRRI-South Asia Hub, ICRISAT,

Patancheru, Hyderabad, India

²*Department of Plant Breeding and Genetics,*

Agricultural College and Research Institute, Killikulam

Tamil Nadu Agricultural University, India

³ *Scientist, Division of Genomic Resources,*

ICAR-National Bureau of Plant Genetic Resources,

and Faculty, ICAR-IASRI, New Delhi, India

⁴*Principal Scientist*

Rice Breeding and genetics research Centre,

Aduthurai 612 101Tamil Nadu, India

**Corresponding author E-mail: mapillai1@hotmail.com*

Abstract

Rice variety, Improved White Ponni is a medium duration crop, but highly susceptible to lodging impacting maximum yield losses. The present investigation aimed to identify early and early semi-dwarf mutants in Improved White Ponni by inducing variations using gamma rays without altering its native grain quality traits. Seeds were treated with various doses of after fixation of the LD₅₀ value of gamma radiation and reported that most of the traits exhibit variations in the mutants at various levels of irradiation. The selection for earliness and dwarf plant height was imposed in M₂ and it was confirmed by evaluation of M₃ generation. Apart from semi-dwarf early mutants, high tillering habit, narrow rolled leaf, upper albino leaf and grassy stunt and extreme dwarf mutants were also identified. Characterization of mutants using already reported genic and linked microsatellite markers associated with semi-dwarfism and earliness resulted that PIC value ranges between 0.037 and 0.949 with an average of 0.382. Single marker analysis revealed that RM302 and RM310 on chromosome 1 and 8 had exhibited an association with the traits plant height, culm and internodal lengths. Of these gene-specific markers, GA20Oxi_1 and GA20Oxi_2 have shown polymorphism among mutants. GA20Oxi_2 showed null alleles in the dwarf mutants and this clearly emphasized that there are some base deletions exists in the region of exon 2 of *sd1* region. GA₃ response study shown that identified mutants were GA₃ responsive except IWP 11-2, IWP 48-2, IWP 50-11 and IWP 33-2 which showed very low responsive. Agar plate assay revealed that, IWP 50-11, IWP 33-2, IWP 43-1, IWP 47-2 and IWP 18-1 had low production of α - amylase. Scanning electron microscope examination on confirmed mutants exhibited larger cell size and a lesser number of cells per unit area than the wild-type which shows that these mutants are defective in GA mediated pathway.

Keywords: Early maturing, irradiation, LD₅₀, lodging resistance, semi-dwarf, Single Marker

Introduction

Rice (*Oryza sativa* L.) improvements especially in case of quantity and quality enhance the rice global production and play a vital role to overcome food shortage, enlighten the local consumption and export. Introducing new varieties of rice characterized by early heading, short stature, lodging resistance, blast resistance and improved grain quality characters are the main objectives for a quantum increase of grain yield of rice. Semi-dwarfism, an important trait in cereals and governs by green revolution gene *semi-dwarf 1 (sd-1)*, which have an impact of short and thick culms, imparts lodging resistance and nitrogen responsiveness and it has initially derived from Dee-Gee-Woo-Gen and responsible for higher yield without affecting the native grain quality parameters of the variety (Futsuhara and Kikuchi, 1997). Kikuchi et al. (1985), identified several *sd-1* mutants in rice and these mutants has been used in several breeding programs. It was found that *sd-1* gene was responsible for the production GA20 oxidase-2 enzyme involved in the catalytic steps of gibberellin (GA) bio-synthesis (Spielmeyer et al., 2002; Sasaki et al., 2002;). A defect in the production of GA was one of the key determinants for semi-dwarf plant type in most of the *sd-1* mutants (Sakamoto et al., 2004) caused by low GA production due to varied (either loss or reduced) function of GA20 oxidase-2. In *indica* variety IR8, an allelic form of *sd-1* contains 382 bp deletions from exonic regions of *sd-1* locus of 1 to 2 resulting in formation of stop codon, which ultimately modifies the gene function. Whereas in the cases of some *japonica* varieties namely Jikkoku, Calrose76 and Reimei, single base substitutions lead to a single amino acid change in the *sd-1* gene (Spielmeyer et al., 2002; Ashikari et al., 2002).

Rice, as a facultative short-day plant and early heading or flowering, has a significant impact on the regional adaptability of the rice varieties. Number of QTLs has been found in

crosses among wild strains of rice, *japonica* and *indica* strains. Apart from genetic analysis, induced mutations played a pivotal role for improving rice architecture by developing a large number of variants such as, early, dwarf, high tillering, blast resistance, low amylose and high yielding mutants (Soomro *et al.*, 2006). The basic requirement for direct improvement target agronomic trait, available genetic variability is required to meet the demand of the breeder. Therefore, induced mutations with the discovery of an array of radiation mutagen and improved treatments methods offer the possibility for the induction of desired changes in various attributes, which can be exploited as such or through recombination breeding (Cheema and Atta 2003). Hence, the primary objective of this study is to induce variations in Improved White Ponni (IWP) by using gamma irradiation and to develop desirable semi-dwarf, early high yielding mutants with improved grain quality parameters.

Materials and Methods

Genetic material

Improved White Ponni, an important medium duration (115 days to flowering) and quality rice variety in southern parts of India for its fine slender grains but had a problem of tallness (> 150 cm) which make the crop susceptible to lodging and grain loss. The seeds of Improved White Ponni were subjected to gamma irradiation at different doses (100Gy to 500Gy) by using Gamma Chamber – Model GC 1200 installed at Tamil Nadu Agricultural University, Coimbatore. The experimental plots were raised at Agricultural College and Research Institute (Killikulam) and Agricultural Research Station (Thirupathisaram) during the year 2011 to 2014.

Mutagenic treatment, selection and evaluation

Five hundred well-filled seeds of IWP were treated with gamma rays at various doses from 100 to 500 Gy with an interval of 100Gy. After treatment, M₁ seeds were immediately sown in raised nursery beds along with control seeds. On 25th day after germination, the seedlings were planted in the main field where standard cultural practices were followed and harvested on single plant basis (M₂ seeds). The M₂ generation was raised from individual M₁ plant following plant to progeny method. A total set around 184 M₁ (families) plants seeds were forwarded to M₂ generation which was raised during *rabi* 2012 to summer 2012 (September 2012 to April 2013) without replication. The selection was imposed for dwarf plant type and earliness in flowering along with other desirable characters. A set of 152 mutants were identified in M₂ and forwarded to M₃ generation for their evaluation and validation. These mutants were sown in raised nursery beds and transplanted to the main field 28 days after sowing in three replications during *Kharif* 2013 (May 2013 to November 2013). M₃ generation was evaluated for various

traits associated with the trait of interest, yield component traits and traits controlling quality parameters fetching higher consumer's preference by the methods given in International Rice Research Institute, Standard Evaluation System (SES) 2013. Estimation of amylose content of identified mutants was carried out by calorimetric method ((Juliano, 1979).

Study of GA₃ response in identified mutants

Mutant seeds of IWP along with control were surface sterilized for 30 minutes using with 2 per cent HgCl₂ (Mercuric chloride) and washed with sterile water. After that, seeds were placed over wet filter paper for two days under dark at 30°C for proper germination. Ten uniformly germinated seeds were placed on the plate containing 1 per cent concentration of agar and kept at 25°C for hastening the proper emergence of the second leaf sheath. Then, 1 µl of GA₃ solution containing 10 mg/ml was dropped to the coleoptile region by using micropipette on the rice seedlings. The length of the second-leaf sheath of five seedlings was measured after five days of treatment to calculate GA₃ response (GAR) (Murakami, 1968).

α- amylase activity assay

The embryo-less half seed of the IWP Mutant and control seeds were surface sterilized using 2 per cent HgCl₂ for 15 minutes, washed using sterile distilled water for five minutes and placed perpendicularly on plates containing 2 per cent agar medium containing soluble starch (0.2 per cent), sodium acetate (10 mM) and CaCl₂ (2 mM) with the pH of 5.3. One micromole of GA₃ was added after autoclaving and incubated for 3 days in the dark at 30°C. After incubation, plates were flooded with I₂-KI (0.72 g/l I₂ + 7.2 g/l KI) solution in 0.2 N HCl. Transparent halos around the seed was noticed and it gives an indication of production of α-amylase, which results in the digestion of starch in the plat (Lanahan and Ho, 1988).

Scanning electron microscope (SEM)

Transverse sections of leaf, nodal region and intermodal regions of the unique mutants (Dwarf mutant, narrow rolled leaf mutants and control) were studied for their difference in internal cell arrangement patterns under Scanning Electron Microscope (SEM) (Model: FEI quanta 200 SEM) built in the Department of Nano Science and Technology, Tamil Nadu Agricultural University, Coimbatore.

PCR amplification and Electrophoresis

A set of 55 microsatellite markers associated with semi-dwarfism and earliness were used to characterize mutants in the M₃ generation (**Table 1**). The PCR amplification was carried out using thermal cycler (Applied Biosystems/BioRad) and amplified products were separated by agarose gel matrix (1.5%) containing 1X Tris-Borate EDTA and electrophoresed at 80 volts for 2 hours and visualized with the help of gel documentation system (BioRad).

Statistical analysis

The estimation of mean, analysis of variance and standard error of the traits were worked out by adopting the standard methods (Panse and Sukhatme, 1961). Analysis of phenotypic, genotypic variances and heritability was formulated by Lush (1940), variability parameters like PCV and GCV was determined by the formula given by Burton (1952). Estimation of genetic advance and correlation coefficient among the studied traits was given by Johnson *et al.* (1955). Polymorphism information content (PIC) value of each SSR marker was estimated by using marker scoring data (Anderson *et al.* 1993). The Marker -Trait association analysis between marker data and traits was carried out using Simple linear regression analysis (SLRA).

Results

LD₅₀ Determination

Probit analysis was carried out using seed germination values to determine the Lethal Dose (LD₅₀) of gamma radiation against Improved White Ponni. The expected LD₅₀ value for gamma rays in Improved White Ponni was 354.8 0 Gy. M1 plants were harvested individually in all the treatments and used for next season crop (M₂) where variability was studied for most of the morphological traits.

Variability for quantitative traits in M₂ and M₃ population

M₂ population

In the M₂ generation, the mean days to fifty per cent flowering ranged from 106.23 (100 Gy) to 112.87 (400 Gy) days whereas wildtype recorded 111.50 days (**Table 2a**). The lowest mean value of plant height was recorded by 300 Gy (138.68 cm) and the highest mean value of plant height was recorded by 100 Gy (153.43 cm) and these two values were lesser when compared to the mean value of wildtype (162.39 cm). The maximum panicle length was recorded in 400 Gy (25.08 cm) which was followed by 300 Gy (24.86 cm) and 200 Gy (23.73 cm) and the lowest value was noticed in 100 Gy (23.27 cm). A number of grains per panicle ranged from 179.47 (400 Gy) to 197.43 (100 Gy). Primary culm length was ranged from 113.82 cm (300 Gy) to 130.16 cm (100 Gy) and this was less when compared to wildtype (138.14 cm). The trait secondary culm length ranged from 108.52 cm (300 Gy) to 121.37 (100 Gy). The trait thousand grain weight ranged from 16.52 gram (300 Gy) to 16.64 gram (100 Gy and 400 Gy). Single plant yield ranged from 41.38 gram (300 Gy) to 51.11 gram (200 Gy).

183 **M₃ population**

184 In gamma irradiated population of Improved White Ponni, the mean days to fifty per cent
185 flowering ranged from 104.12 (100 Gy) to 114.26 (200 Gy). The treatment 200 Gy had recorded
186 the highest PCV (9.92), GCV (9.84) and GA as per cent of the mean value of 20.09 with the
187 heritability percentage of 98.32 and this was followed by 100 Gy with heritability and the genetic
188 advance of 99.26 and 17.80 per cent (**Table 2b**). The lowest plant height was recorded by 100
189 Gy (113.65 cm) and the highest mean value of plant height was recorded by 300 Gy (126.94 cm)
190 and these two values were lesser when compared to the mean value of wildtype (142.59 cm). The
191 treatment 100 Gy had recorded the highest PCV (15.60), GCV (15.53), heritability (99.14) and
192 genetic advance (31.86). The highest mean value of panicle length was recorded in 400 Gy
193 (26.17 cm) which was followed by 300 Gy (25.73 cm) and 100 Gy (25.02) and these values were
194 higher than IWP (24.92 cm). PCV (12.25), GCV (11.15), heritability (82.85) and GA% of the
195 mean (20.21) was found to be higher in 200 Gy when compared to all other treatments. The trait
196 number of productive tillers per plant ranged from 10.13 (400 Gy) to 16.04 (100 Gy). The
197 treatment 300 Gy had registered the highest PCV (31.05), GCV (28.36) and GA% of the mean
198 (53.36) whilst heritability (86.52) was higher in 100 Gy.

199 A number of grains per panicle ranged from 169.70 (400 Gy) to 191.15 (300 Gy). The
200 treatment 100 Gy registered higher value of PCV (14.80), GCV (13.64), heritability (84.92) and
201 GA% of the mean (25.90). The trait primary culm length ranged from 88.77 CM (100 Gy) to
202 101.52 cm (300 Gy) and this was less when compared to wild-type (116.54 cm). The phenotypic
203 coefficient of variance (18.59), the genotypic coefficient of variation (18.47), heritability (98.70)
204 and GA% of the mean (37.80) was registered higher in 100 Gy. Secondary culm length ranged
205 from 83.93 cm (100 Gy) to 97.80 (300 Gy). First internodal length ranged from 30.33 cm (200

Gy) to 41.52 (100 Gy) and this was lesser than wildtype (43.56 cm). PCV (23.02), GCV (22.47), heritability (95.28) and GA% of the mean (45.19) were noticed to be higher in 200 Gy. The 2nd internodal length ranged from 20.21 cm (200 Gy) to 26.27 cm (100 Gy). The treatment 100 Gy registered the highest PCV (26.76), GCV (25.83), heritability (93.19) and GA% of the mean (51.36). 3rd internodal length ranged from 11.57 cm (100 Gy) to 16.57 (400 Gy). PCV (32.91), GCV (31.16) and GA% of the mean (60.76) were registered higher in 200 Gy whereas heritability (91.48) was found to be higher in 100 Gy. The 4th internodal length ranged from 5.77 cm (200 Gy) to 8.11 cm (300 Gy). The treatment 200 Gy recorded the highest PCV (35.15), GCV (32.87) and GA% of the mean (63.34) whereas the heritability (88.68) was higher in 100 Gy. The trait thousand grain weight ranged from 16.32 gram (100 Gy) to 16.70 gram (400 Gy). The treatment 300 Gy had recorded the highest PCV (1.53), GCV (1.35), heritability (77.63) and GA% of the mean (2.45). Single plant yield ranged from 33.14 gram (200 Gy) to 44.04 gram (400 Gy). The treatment 200 Gy had recorded the highest PCV value of 30.22 whereas other parameters were higher in 100 Gy.

Identification of unique mutants for agronomic traits

All the mutants were raised under field conditions and screened for novel altered phenotypes in the morphological traits *viz.*, flowering, plant height, tillering habit, narrow rolled leaf, upper albino leaf, grassy and extreme dwarf, lodging resistant, lanky culm, *etc.*, (**Table 3**) (**Fig 1**). Early flowering mutants had registered 79 to 92 days for days to flowering when compared to wild-type which had recorded 112 days to flower. Semi-dwarf mutants identified in this study were early in flowering. High tillering mutants were identified which possess 32 to 44 productive tillers per plant whereas Improved White Ponni had only 18 productive tillers.

Narrowly rolled leaf mutants had less panicle length and number of grains with fine grains when compared to wildtype genotype.

Morphological and quality trait evaluation of desirable semi-dwarf and early mutants in the M₃ generation

The ANOVA was computed for all the component traits studied (**Table 4**) and in case of days to fifty per cent flowering, IWP E-4-3 had recorded lower days of 86.00 and IWP 11-2 recorder higher days of 119.00 days to flower with mean value of 96.93 days. With respect to plant height, the range of 82.21 cm (IWP 59-1) to 142.39 cm (IWP 50-4) was noticed with the mean of 111.18, whereas wildtype had the plant height of 143.26 cm (**Fig 2**). The trait panicle length exhibited the range of 20.46 (IWP 48-2) to 28.69 (IWP 18-1) with the mean value of 24.17 cm. The trait number of productive tillers ranged from 8.50 (IWP D-1) and 25.50 (IWP 1-12) with a mean value of 13.71. Among the 30 mutants studied, the highest primary culm length was found to be low in IWP 59-1 (59.50 cm) and high in IWP 50-4 (117.33 cm) whereas wildtype had registered higher primary culm length of 114.86 cm. The trait secondary culm length had recorder the area of 83.79 cm with the range of 53.54 (IWP 48-4) to 109.74 (IWP 50-4). With respect to 1st internodal length, the mutant IWP 1-9 had recorded a lower value of 16.63 cm and IWP 51-4 had registered higher value 42.08 cm. The trait 2nd internodal length ranged from 11.54 (IWP 48-4) to 40.06 (IWP 1-9) with the mean of 21.17 cm. In case of the trait 3rd internodal length, the minimum and maximum length were found to be reported in IWP 48-3 (4.77 cm) and IWP 1-9 (21.72 cm) whereas wildtype genotype reported 11.50 cm. The trait 4th internodal length had registered 5.78 cm as a mean with the range of 3.33 (IWP E-4-1) to 10.69 (IWP 1-9) (**Fig 3**). A number of grains per panicle ranged from 156.00 (IWP 31-2) to 272.00 (IWP 7-1) with the grand mean of 206.07. The trait thousand grain weight was found to be high

in IWP 1-9 (21.34 gram) and low in IWP 7-1 (13.26 gram) with the mean of 17.04 gram. In case of single plant yield, the mutants IWP 43-1 and IWP 50-4 had recorded higher and lower yield of 46.46 and 25.75 gram, respectively.

Apart from morphological parameters, this study also involved accessing the quality performance of desirable mutants. The trait kernel length before cooking exhibited maximum in IWP E-2 (6.30 mm) and minimum in IWP 11-1 (4.87 mm) whereas wildtype registered the trait value of 5.58 mm. Kernel breadth before cooking had ranged from 1.83 mm (IWP 48-2 and IWP 7-1) to 2.20 mm (IWP 1-9) with the mean of 1.9 mm. With respect to the L/B ratio, the minimum and maximum ratio were found to be observed in IWP 47-2 (2.49), IWP 33-2 (2.49) and IWP 16-6 (3.09). The trait kernel length after cooking revealed that, the mean value of 6.90 with the range of 5.60 mm (IWP D-1) to 7.70 mm (IWP 1-1). In case of kernel breadth after cooking, the minimum and maximum value was found to be noticed in IWP D-1 (2.55 mm) and IWP 1-9 (3.35 mm) with the mean value of 2.93 mm. The trait L/B ratio after cooking showed the mean value of 2.36 with the range of 1.94 (IWP 33-2) to 2.75 (IWP E-3) whereas wildtype genotype recorded 2.37. The maximum and minimum linear elongation ratio was observed in IWP E-3 (1.51) and IWP D-1 (0.99) with the wildtype value of 1.20. The trait BER ranges from 1.28 (IWP 51-4) to 1.69 (IWP 48-2). The maximum and minimum amylose content was found to be noticed in IWP 48-3 (21.26%) and IWP 43-1 (10.40 %) with the mean value of 16.26 %. Gel consistency was ranged from 56.25 mm to 81.25 mm in Improved White Ponni mutants.

Association analysis

Association analysis found out the relationship among the traits studied and reported that plant height had significantly higher positive association with the traits *viz.*, panicle length (0.480), primary culm length (0.970), secondary culm length (0.959), 1st internodal length

274 (0.713), 2nd internodal length (0.691), 3rd internodal length (0.767), 4th internodal length (0.452)
275 and thousand grain weight (0.647) (**Table 5a**). Panicle length had registered high and significant
276 positive correlation with 2nd internodal length (0.477), 3rd internodal length (0.483), 4th
277 internodal length (0.568) and single plant yield (0.505). Primary culm length had a positive
278 relationship with secondary culm length (0.995), 1st internodal length (0.697), 2nd internodal
279 length (0.692), 3rd internodal length (0.790), 4th internodal length (0.383) and thousand grain
280 weight (0.627). Secondary culm length had registered significant positive correlation with 1st
281 internodal length (0.683), 2nd internodal length (0.730), 3rd internodal length (0.793), 4th
282 internodal length (0.366) and thousand grain weight (0.617). The trait 2nd internodal length had
283 recorded the significant positive correlation with 3rd internodal length (0.816), 4th internodal
284 length (0.517) and thousand grain weight (0.588) whereas it has negative association with a
285 number of grains per panicle. The character 3rd internodal length had high and significantly
286 positively correlated with 4th internodal length (0.719) and thousand grain weight (0.508). The
287 trait 4th internodal length was positively with thousand grain weight (0.399) and single plant
288 yield (0.492). Among the quality traits studied, KLBC had high and significant positive
289 correlation with KBBC (0.614), L/B BC (0.757), KLAC (0.633) and L/B AC (0.407). KBBC had
290 registered significant positive and negative correlation with KLAC (0.365) and BER (-0.600),
291 respectively (**Table 5b**). L/B BC ratio was positively correlated with KLAC (0.501), amylose
292 content (0.426) and negatively correlated with single plant yield (-0.441). KLAC had a high and
293 positive association with the traits namely L/B AC (0.820) and LER (0.650), respectively.
294 KBAC had a significant positive correlation with BER (0.617) and GC (0.512). L/B AC had
295 registered positive and negative significant association with LER (0.412) and BER (-0.550). The

character breadth wise expansion ratio exhibited significantly positively correlated with gel consistency (0.442).

Molecular characterization

Polymorphism information content (PIC)

Out of 154 mutants, 30 unique mutants (11 confirmed semi-dwarf and early mutants, 19 early mutants) were selected and subjected to molecular analysis in an M_3 generation. Fifty-six SSR primer pairs were used, which includes three gene-specific markers for the *sd1* locus. Out of these 56 SSR primer pairs, 45 primers exhibited polymorphism between the mutants and found that a total of 96 alleles (**Table 6a**) (**Fig 4**). Alleles per locus ranged from 1 to 3 with a mean of 2.13. It was found that PIC value ranges from 0.037 (RM246) to 0.949 (GA20 Oxi_2) with an average of 0.382. There were several markers which recorded high PIC value and they are highly polymorphic in nature. The markers viz., GA20 Oxi_2 (0.949), RM310 (0.662), RM140 (0.648), RM7365 (0.633), RM3431 (0.616) had recorded PIC value more than 0.6. Several other markers namely GA20 Oxi_3 (0.584), RM25 (0.570), RM3555 (0.537), RM493 (0.531), RM5720 (0.529), RM240 (0.523) and RM3912 (0.513) had registered PIC value more than 0.5.

Marker-trait associations

Single marker analysis was also done and out of 45 polymorphic SSR primer pairs, only 25 primers had a significant correlation with traits studied (**Table 6b**). An amount of phenotypic variation (R^2) accounted by markers were estimated and this clearly explains the existence of phenotypic variation to the total variance. RM 310 showed the maximum phenotypic variance of 65.20 per cent and associated with the trait primary culm length whereas the marker RM7365 explained minimum phenotypic variance and linked with a gel consistency. The marker

associated with the trait days to 50% flowering was RM335 and RM3452 with an R^2 value of 37.30 and 31.30 per cent. The marker RM 252 was highly associated with the traits plant height and 1st internodal length with the R^2 value of 62.40 and 49.20 per cent and several other markers were also associated. The marker RM310 was highly associated with the traits primary culm length, secondary culm length, 4th internodal length, thousand grain weight, KLBC, L/B BC and LER. The marker RM 3912 recorded high association with the traits 2nd and 3rd internodal length with high R^2 value. The marker RM587 was highly associated with the traits number of grains and KLAC with the R^2 value of 36.20 and 37.30 per cent. The marker RM167 exhibited a high association with the trait number of productive tillers with the R^2 value of 21.80 per cent. The marker RM7365 had exhibited a highly significant correlation with the kernel breadth before cooking and explains 21.10 per cent variation to the total variance. The marker RM3431 showed significant association with the traits KBAC and BER with the variability percentage of 22.70 and 30.30 per cent to the total variance. The markers RM551, RM38 and RM246 recorded significantly associated with the traits L/B ratio after cooking, amylose content and gel consistency.

The GA₃ response of dwarf and early mutants using the micro-drop method

Second leaf sheath length was measured in all identified dwarf mutants after the application of GA₃ to the coleoptiles of the seeds after germination (**Table 7**). The GA₃ response was estimated based on the length of second leaf sheath of GA₃ treated and non-treated seedlings as a control. The GA₃ response was estimated on two different durations of 5th day and 15th after treatment. The mean 2nd leaf sheath length of mutants on 5th DAT in non-treated seedlings was 1.60 with the range of 0.92 (IWP E-4) to 1.93 (IWP 48-2) whereas in GA₃ treated seedlings it was ranged from 1.52 (IWP E-4) to 2.98 (IWP 59-1). On 15th DAT, the mean second leaf sheath

length of mutants in non-treated seedling was ranged from 2.30 (IWP 1-12) to 3.03 (IWP 48-2) whereas in GA₃ treated seedlings it was ranged from 2.98 (IWP 7-1) to 3.89 (IWP 43-1, IWP 33-2). The maximum GA₃ response was exhibited by the mutant IWP 18-1 (189.80) and minimum GA₃ response had recorded by the mutant IWP 50-11 (112.26) on 5th DAT whereas in 15th DAT the mutant IWP 18-1 (146.84) recorded the maximum response to GA₃ application and mutant IWP 48-2 (104.29) had registered the minimum response. The mutants IWP 47-2 (120.23), IWP 11-2 (134.04), IWP 1-1 (139.48), IWP 48-2 (122.45), IWP 7-1 (133.39), IWP 1-12 (140.35) and IWP 50-11 (112.26) had recorded less GA₃ response when compared wildtype (149.12) on 5th DAT. On 15th DAT, the mutants *viz.*, IWP 47-2 (107.44), IWP 59-1 (117.82), IWP 1-1 (120.38), IWP 48-2 (104.29), IWP 7-1 (110.78) and IWP 50-11 (117.10) exhibited less GA₃ response when compared to wildtype genotype (159.69). In 5th DAT, the maximum and minimum shoot length of non-treated seedlings was recorded by the mutant IWP 43-1 (10.40 cm) and IWP E-4 (3.60 cm) whereas in treated seedlings the maximum and minimum shoot length was exhibited by the mutants IWP 1-12 (14.50 cm) and NLM (7.67 cm) (**Table 8**) (**Fig 5**). In 15th DAT, shoot length of non-treated seedlings of mutants were ranged from 4.60 cm (IWP 18-1) to 12.00 cm (IWP 43-1) whereas the shoot length of treated seedlings was ranged from 10.73 cm (IWP 50-11) to 21.33 cm (IWP 48-3).

α – amylase activity

The identified dwarf and early mutants of Improved White Ponni were subjected to study α – amylase activity induced by the application of GA₃ induction by GA application. In this study, α – amylase production and secretion were observed as plaques in the wild-type as well as in dwarf and early mutants *viz.*, IWP 43-1, IWP 47-2, IWP 59-1, IWP D-1, IWP 18-1, IWP E-3 and IWP E-4. Based on the staining pattern of the starch plate with iodine solution revealed that

the mutants namely IWP 50-11 and IWP 33-2 showed there is no α – amylase production which indicated that these two mutants were related to the GA pathway (Fig 6). The other mutants viz., IWP 43-1, IWP 47-2 and IWP 18-1 exhibited α – amylase production was significantly decreased when compared to wild type. The mutants IWP 59-1, IWP D-1, IWP E-3 and IWP E-4 were produced more α – amylase than wild-type.

Study on cell arrangement pattern in unique mutants of Improved White Ponni

Internodal regions of dwarf and early mutants and leaf sections of identified (IWP 59-1) and of Improved White Ponni along with wild-type were subjected to observe under scanning electron microscope to study the pattern of internal cell arrangements. Cell patterning differences were observed and revealed that the semi-dwarf mutant of White Ponni exhibited larger cell size and less number of cells arrangement per unit area than the wild type (Fig 7).

Discussion

Micro-mutational events with the least deleterious effects were considered to be the most reliable and effective mutations in providing variability for quantitative traits. The mutagens which offered to induce variability is utmost important for any crop breeding program since events created by mutation may alter some large or small phenotypic expression. Besides the observation of frequency of chlorophyll mutations across the seedlings of M_2 generations, changes in quantitative traits were also observed in both M_2 and M_3 generations. Gaul (1964) classified viable mutations as macro and micro mutations, while Swaminathan (1965) grouped them as macromutations and systematic mutations. Besides the usage of chlorophyll mutations observed in Improved White Ponni to estimate the effectiveness and efficiency of mutagen, viable mutants for plant type *viz.*, early/late flowering, dwarf, high tillering habit, narrow rolled leaf, upper albino, lodging resistant, grassy and extreme dwarf and lanky culms were observed in M_2 population. In case of plant type modifications, semi-dwarf and narrowly rolled leaf mutants appeared in mutated population. The frequency of semi-dwarf mutants was high in lower doses of gamma radiation when compared to higher doses in gamma-irradiated populations. Shadakshari *et al.* (2001) reported the occurrence of higher frequency of dwarf/semi-dwarf non-lodging early flowering and high yielding mutants in five rice varieties treated with gamma rays. Yankulav *et al.* (1980) and Reddi and Rao (1988), who reported that reduction on mutagen's effectiveness were higher with increase in their dosage and our study also reported the same. Apart from estimating the mutagen efficacy and efficiency, chlorophyll and viable mutation frequency estimation become useful for selecting the suitable mutagen in crop breeding (Nilan *et al.*, 1965).

The viable mutations isolated in the study showed changes in major traits which could be utilized in the future breeding programme where reshuffling of traits may be tried by

conventional breeding methods. However, most of the morphological mutants identified in M_2 generation failed to inherit in M_3 generation. According to the statements of Luo *et al.* (2012), these characters may be controlled by recessive genes or are susceptible to the environment. Moreover, whatever the changes occurred in the plants due to mutation is an error according to the plant's geometry. They tend to rectify it in due course through recombinational events. That is why most of the observed mutants were not inherited in future generations. Thus, evolving a new phenotype with consistent expression through mutation is a chance event rather than a choice.

Induced variability on quantitative traits in M_2 and M_3 generations

In the present investigation, comparison on means and variances (phenotypic and genotypic) of various quantitative traits including the quality parameters in M_2 and M_3 generations indicated a considerable shift in mean and variability in the treatments and this could be because of recombination happened in the M_1 plants. Siddiqui and Singh (2010), who found out that the variability may increase significantly with respect to increase or decrease in the mean values of the trait studied. Our results were also accorded with earlier reports and this may be expected due to the reason that, recombinational events due to mutation creates more variations in the living systems (Johnston, 2001).

The rate at which different mutations occurring at a specified stage is mutation rate and this estimate can be made when the targeted mutation is very obvious and detectable. But most of the quantitative traits do not follow this pattern and making the situation of detecting mutations very complex. Under these circumstances, mutational heritability [genetic variance increase by a single generation of mutation (V_m)/the environmental variance of the trait (V_e)] is estimated

(Houle *et al.*, 1996) and one has to measure the amount of new genetic variation arising in each generation due to mutation. Moreover, the variations for quantitative traits observed in the M_1 generation have no significant importance in deciding the number of mutations for the targeted trait. The detection of mutations for a quantitative trait has to be decided based on the progeny values *i.e.* M_2 generation. The mutant population which exhibits high mean coupled with a high variance for a trait is the first choice of selection. Usually, mutations are detected when there is huge phenotypic effect on a trait has been observed in large mutant population of a variety or elite cultivar.

Investigation on M_2 generation had shown higher variation in 100 Gy for the trait days to fifty per cent flowering. The genetic parameters were very low in M_3 generation when compared to M_2 generation treated with gamma rays. Nayudu *et al.* (2007) and Anilkumar (2008) had also reported similar findings of our study. The trait plant height showed wider differences of all genetic parameters and the selection was effective in M_2 and M_3 as reported by Singh *et al.* (2006). The traits namely panicle length, productive tillers and grains per panicle shown moderate to high variability in M_2 and M_3 indicating the scope for selection and improvement (Kumar *et al.*, 2006). These traits expresses moderate to high h^2 and low to high GA in M_2 and M_3 generation and these results are lined with Ahmed *et al.* (2010). Primary culm length and secondary culm lengths had exhibited moderate variability, moderate to high h^2 and low to moderate GA (Bin-mei *et al.*, 2006). A similar trend was also observed in M_3 generation for the traits *viz.*, 1st internodal length, 2nd internodal length, 3rd internodal length and 4th internodal length. Variability study clearly revealed that these traits showed a slightly wider variation in the mutant population (Zou *et al.*, 2005; Bin-mei *et al.*, 2006). Thousand grain weight and single plant yield exhibited minimum variability, low to high h^2 and GA in M_2 and M_3 generations

(Kishor *et al.*, 2008; Yadav *et al.*, 2010). These two parameters (h^2 and GA) was a trustworthy real estimate ifor selection than heritability alone. Conversely, it is not essential criteria that a trait conveying high heritability will also display high genetic advance. This is because; the heritability calculation leads to estimation errors and largely depends on genotype-environment interactions. Estimation of the coefficient of variation specifies only the available variability of the particular trait and it does not provide any information about heritability.

Genetic parameters for different quantitative traits were estimated and most of the traits exhibited varied and significant mean performance with a high level of variation among the mutants. The traits namely, plant height, days to 50 % flowering, panicle length, productive tillers per plant, grains per panicle, thousand grain weight and grain yield per plant exhibited slightly wider variability, higher h^2 and moderate to high GA among the mutants. These genetic parameters could be valuable measure for the effective selection towards yield enhancement is possible (Kishor *et al.*, 2008; Yadav *et al.*, 2010; Akinwale *et al.*, 2011). Among the grain quality traits studied, most of the traits exhibited higher variation among the mutants and some of the putative mutants had on par quality with the wildtype. The results were accorded with the findings of Siddiqui and Sanjeeva (2010) and Subbaiah (2011). In this study, major emphasis and attention were given to the selection of dwarf and early mutants, which determines crops per year and adaptation towards maximizing the yield over seasonal and regional specification. The identified semi-dwarf and early mutants had shown approximately 35 to 45 per cent reduction in plant height and 4 to 33 days earlier in duration than wildtype.

The magnitude of association of component traits assists the plant breeder to improve the yield and other important traits. Among the biometric traits studied, the traits panicle length and 4th internodal length had a significant positive relationship with grain yield as reported as well

(Sankar *et al.*, 2006; Anilkumar, 2008; Immanuel *et al.*, 2011; Bagheri *et al.*, 2011; Akinwale *et al.*, 2011). Saif-ur-Rasheed *et al.* (2002a) reported the association of a number of tillers and productive tillers had an optimistic relationship with grain yield. The trait plant height was positively associated with the traits panicle length, primary culm length, secondary culm length, 1st internodal length, 2nd internodal length, 3rd internodal length, 4th internodal length and thousand grain weight. The selection based on the above traits highly influences the grain yield of the mutants and provides an indirect selection of these traits would ultimately increase the yield quantum (Rashid *et al.*, 2013). Another trait primary culm length exhibited positive and significant correlation with the traits secondary culm length, 1st internodal length, 2nd internodal length, 3rd internodal length, 4th internodal length and thousand grain weight. Another trait secondary culm length exhibited positive and significant correlation with the traits 1st internodal length, 2nd internodal length, 3rd internodal length, 4th internodal length and thousand grain weight as reported by Muhammad *et al.* (1982). The trait 4th internodal length had registered significant positive association with thousand grain weight. Apart from morphological traits, quality parameters of the mutants were also shown on par in most of the traits observed with the positive relationship among them and these mutants would be resulted to improve those traits through effective breeding strategies (Khatun *et al.*, 2003; Veni *et al.*, 2003).

Molecular characterization and Marker-trait Associations

Semi-dwarf and early mutants, narrowly rolled leaf mutants of Improved White Ponni generated through gamma radiation were identified in M₂ were forwarded to M₃ (30 mutants) generations. These mutants were surveyed using 55 microsatellite markers to detect the polymorphism among mutants and it was noticed that 96 alleles were detected. Among the SSR markers, RM246 and GA20 Oxi_2 found to be have lower and higher PIC value of 0.037 and

0.949 with a mean value of 0.382. . The microsatellites *viz.*, GA20 Oxi_2, RM310, RM140, RM7365, RM3431, GA20 Oxi_3, RM25, RM3555, RM493, RM5720, RM240 and RM3912 had recorded PIC value more than 0.5 (Chen *et al.*, 2011; Kumar and Bhagwat, 2013). The PIC value made the reflection of allelic diversity between the mutants and not found to be uniformly high for the SSR loci tested (Wang *et al.*, 2009; Pervaiz *et al.*, 2010). These findings were supportive that, microsatellites are more potent and informative to study the genetic divergence and variability pattern of closely related individuals (Xu *et al.*, 2004).

Marker data were also correlated with traits studied in rice mutants for determining the informative SSR markers associated with these traits. Out of 45 polymorphic SSR primer pairs, only 25 primers had a significant association with different traits studied. Single marker analysis exhibited that, the markers RM3912 and RM430 was highly associated with the trait plant height. The marker RM302 had exhibited a high association with the traits plant height, primary culm length, secondary culm length, 1st internodal length and 4th internodal length. The marker RM302 on chromosome 1 had shown putative linkage with semi-dwarfism (Subashri *et al.*, 2008, Wang *et al.*, 2009). Another gene (*sd-1*) based marker, GA20Oxi_2 was positioned near to the marker was also found linked with semi-dwarf trait. Asano *et al.* (2007) also reported that the dwarfism is due to complete loss of *sd-1* gene function. The marker RM310 showed association with primary culm length, secondary culm length, 1st internodal length, 2nd internodal length, and 3rd internodal length. Several grain-quality traits were observed in mutants selected from M₃ generation of White Ponni and marker-trait association trait study revealed that the marker RM7365 had exhibited a highly significant correlation with the KBBC, the marker RM3431 showed significant association with the traits KBAC and BER. The markers RM551, RM38 and RM246 recorded significantly associated with the traits L/B ratio AC, amylose and GC. Roy *et*

al. (2006) reported that an association study in wheat showed a total of 99 of the 221 polymorphic SSR bands and explained a maximum of R^2 value of 8.12 per cent for tiller numbers to 27.95 per cent for harvest index (Monfared *et al.*, 2008; Kalivas *et al.*, 2011).

The response of mutants to GA₃ application

GA₃ responses of semi-dwarf and early mutants were determined using a micro-drop method by GA₃ application on the coleoptiles region of the seeds after three days of germination. In the present investigation of GA₃ response, all the mutants were recorded moderate to high response to the GA₃ application. Narrowly rolled leaf mutant also recorded low response to the GA₃ application (Ueguchi-Tanaka *et al.*, 2000). The shoot length was also observed in all the mutants five and fifteen days after GA₃ application which revealed that the increase in shoot length of semi-dwarf and early mutants showed variation in their responsiveness. Thus, this study clearly emphasized that all the mutants identified as GA₃ response mutants except IWP 11-2, IWP 48-2, IWP 50-11 and IWP 33-2 which showed a very minimal response to the GA₃ application after fifteen days and these were GA₃ insensitive mutants. Ogi *et al.* (1993) discovered that the *sd-1* gene reduced the internode length by preventing cell division in internodes. Other researchers have proved that the *sd-1* gene was sensitive to GA₃, which influenced the content of GA₃ and the vigour of peroxidase in the plant (Shi and Shen, 1996; Gao *et al.*, 2009 and Asano *et al.*, 2009). Application of GA₃ in *iga-1* semi-dwarf mutant resulted in the partial restoration of wildtype plant height and this was confirmed as insensitive and similar result was also reported by the current study, were both sensitive and insensitive mutants were identified (Wang *et al.*, 2009).

α – amylase activity of semi-dwarf and early mutants

The α -amylase production in assay plates and shoot elongation by the exogenous application of GA₃ are GA- mediated process (Matsukura *et al.*, 1998). The staining pattern of agar plate revealed that the mutant's *viz.*, IWP 50-11, IWP 33-2, IWP 43-1, IWP 47-2 and IWP 18-1 had exhibited low production of α - amylase which ultimately infers that these mutants have some defects in the GA related pathway (Qin *et al.*, 2008). The secretion of α -amylase was found to be noticed as low around the seed placed in the assay plates containing GA₃ in both wildtype and mutants and this was lined with the earlier report in the d62 mutant (Li *et al.*, 2010). On overall conclusions, it was found that the identified semi-dwarf early mutants were either GA₃ responsive or non-responsive in nature (Lanahan and Ho, 1988; Chandler, 1988 Zou *et al.*, 2005).

Cell structural pattern analysis in internodal regions of Semi-dwarf mutants

Intercalary meristem cell division and elongation are the major cause for internodal elongation in rice and flaw in these processes severely affects the plant height. A longitudinal section from the internodal regions of the semi-dwarf mutants and wild-type was examined under a scanning electron microscope (SEM) to analyse the inter-cell arrangement patterns. This study revealed that the semi-dwarf mutant exhibited larger cell size (more length and breadth) and a lesser number of cells arrangement per unit area than the wildtype. These results ultimately highlighted that these mutants are defective in GA mediated pathway and therefore cell division was minimized which would ultimately result in the reduced internodal lengths (Wang *et al.*, 2009).

Conclusion

The overall study concluded that the semi-dwarf mutants identified had a significant reduction in the 2nd, 3rd and 4th internodal regions and thereby reduces the plant height (up to 50 per cent reduction), imparted lodging resistance and provided significant yield and quality improvement over the wild-type. The identified early mutants of White Ponni exhibited 4 to 33 days of earliness than the wildtype and thereby aiding to reduce crop duration. The molecular study on these mutants using markers linked to plant height like RM302, GA20Oxi_1 and GA20Oxi_2 on chromosome 1 had allelic variations between the mutants. GA20Oxi_2 showed null alleles in the dwarf mutants and this clearly emphasized that there are some base deletions occurred in the region of exon 2 of *sd-1* region and it can be further studied for the expression profiling. GA₃ response and α -amylase activity were studied which reported that the identified semi-dwarf mutants were more or less sensitive to GA₃. These desired mutants of White Ponni might have the potential for further improvement of rice production through future breeding programmes.

Acknowledgement

The authors thank the BARC (*Board of Research in Nuclear Sciences*) *BARC, Mumbai* for financial support.

Competing interests

The authors declare that they have no competing interest.

Author Contributions

MAP has designed the research plan. RC, APL and MAP conducted the research. RC, APL, KKV and YJK analyzed and interpreted the data. RC wrote the manuscript. The authors approved the final version of the manuscript.

References:

Ahmed MF, Iqbal M, Masood MS, Rabbani MA, Munir M. Assessment of genetic diversity among Pakistani wheat (*Triticum aestivum* L.) advanced breeding lines using RAPD and SDS-PAGE. Electron J Biotechn. 2010 May; 13(3):1-2.

Cheema AA, Atta BM. Radio sensitivity studies in basmati rice. Pak J Bot. 2003 Jun 1;35(2):197-207.

Akinwale MG, Gregorio G, Nwilene F, Akinyele BO, Ogunbayo SA, Odiyi AC. Heritability and correlation coefficient analysis for yield and its components in rice (*Oryza sativa* L.). African Journal of Plant Science. 2011 Mar 31;5(3):207-12.

Muhammad A, Rehman A, Cheema AA. Correlation between yield and yield attributing characters in some induced dwarf mutants of rice (*Oryza sativa* L.). Pakistan Journal of Agricultural Research. 1982;3(3):141-4.

Anderson JA, Churchill GA, Autrique JE, Tanksley SD, Sorrells ME. Optimizing parental selection for genetic linkage maps. Genome. 1993 Feb 1;36(1):181-6.

Anilkumar CV. Genetic analysis of economic traits in segregating population of Rice (*Oryza sativa* L.) [dissertation]. Coimbatore: Tamil Nadu Agricultural University; 2008.

Asano K, Hirano K, Ueguchi-Tanaka M, Angeles-Shim RB, Komura T, Satoh H, Kitano H, Matsuoka M, Ashikari M. Isolation and characterization of dominant dwarf mutants, Slr1-d, in rice. *Mol Genet Genomics*. 2009 Feb 1;281(2):223-31.

Asano K, Takashi T, Miura K, Qian Q, Kitano H, Matsuoka M, Ashikari M. Genetic and molecular analysis of utility of sd1 alleles in rice breeding. *Breed sci*. 2007;57(1):53-8.

Ashikari M, Sasaki A, Ueguchi-Tanaka M, Itoh H, Nishimura A, Datta S, Ishiyama K, Saito T, Kobayashi M, Khush GS, Kitano H. Loss-of-function of a rice gibberellin biosynthetic gene, GA20 oxidase (GA20ox-2), led to the rice ‘green revolution’. *Breed Sci*. 2002; 52(2): 143-50.

Bagheri N, Babaeian-Jelodar N, Pasha A. Path coefficient analysis for yield and yield components in diverse rice (*Oryza sativa* L.) genotypes. *Biharean biol*. 2011 Jun 1;5(1):32-5.

Bin-mei LI, Can CH, Yue-jin WU, Ji-ping TO, Jin-hua WU, Ying ZH, Qin YU. Effect of dominant semi-dwarf gene on plant height and its related traits and sensitivity to gibberellic acid in rice. *Rice Sci*. 2006;13(3):179-84.

Burton GW. Quantitative inheritance in grasses. *Proceedings of VI International Grassland Congress*; 1952. Chandler PM. Hormonal regulation of gene expression in the “slender” mutant of barley (*Hordeum vulgare* L.). *Planta*. 1988 Jul 1;175(1):115-20.

664 Chen CX, Li SC, Wang SQ, Liu HN, Deng QM, Zheng AP, Zhu J, Wang LX, Li, P. Assessment
665 of the Genetic Diversity and Genetic Structure of Rice Core Parent Guichao 2, its Parents and
666 Derivatives. Journal of Plant Sciences. 2011;6(2):66-76
667

668 Fang Y, Wu W, Zhang X, Jiang H, Lu W, Pan J, Hu J, Guo L, Zeng D, Xue D. Identification of
669 quantitative trait loci associated with tolerance to low potassium and related ions concentrations
670 at seedling stage in rice (*Oryza sativa* L.). Plant Growth Regul. 2015 Nov 1;77(2):157-66.
671

672 He F, Xi Z, Zeng R, Talukdar A, Zhang G. Identification of QTLs for plant height and its
673 components by using single segment substitution lines in rice (*Oryza sativa* L.). Rice Sci.
674 2005;12(3):151-6.
675

676 Fujino K, Sekiguchi H. Mapping of quantitative trait loci controlling heading date among rice
677 cultivars in the northernmost region of Japan. Breed Sci. 2008;58(4):367-73.
678

679 Futsuhara Y, Kikuchi F. Inheritance of morphological characters. 2. Inheritance of dwarfism.
680 Science of the rice plant. Volume 3:Genetics. Tokyo(JP): Food and Agriculture Policy Research
681 Center; 1997.
682

683 Futsuhara Y, Toriyama K, Tsunoda K. Breeding of a new rice variety "Reimei" by gamma-ray
684 irradiation. Japanese Journal of Breeding. 1967 Jun 25;17(2):85-90.
685

686 Gaul H. Mutations in plant breeding. Radiation botany. 1964 Jan 1;4(3):155-232.

Hori K, Kataoka T, Miura K, Yamaguchi M, Saka N, Nakahara T, Sunohara Y, Ebana K, Yano M. Variation in heading date conceals quantitative trait loci for other traits of importance in breeding selection of rice. *Breed sci.* 2012;62(3):223-34.

Houle D, Morikawa B, Lynch M. Comparing mutational variabilities. *Genetics.* 1996 Jul 1;143(3):1467-83.

Immanuel SC, Pothiraj N, Thiagarajan K, Bharathi M, Rabindran R. Genetic parameters of variability, correlation and pathcoefficient studies for grain yield and other yield attributes among rice blast disease resistant genotypes of rice (*Oryza sativa* L.). *Afr J Biotechnol.* 2011;10(17):3322-34.

Johnson HW, Robinson HF, Comstock R. Estimates of Genetic and Environmental Variability in Soybeans. *Agron J.* 1955 Jul 1;47(7):314-8.

Johnston MO. Mutations and new variation: overview. *Encyclopedia of Life Sciences.* 2009.

Juliano BO. A simplified assay for milled rice amylose. *Cereal science today.* 1971;16:334-60.

Kalivas A, Xanthopoulos F, Kehagia O, Tsaftaris AS. Agronomic characterization, genetic diversity and association analysis of cotton cultivars using simple sequence repeat molecular markers. *Genet Mol Res.* 2011 Feb 8;10(1):208-17.

709 Kanbe T, Sasaki H, Aoki N, Yamagishi T, Ebitani T, Yano M, Ohsugi R. Identification of QTLs
710 for improvement of plant type in rice (*Oryza sativa* L.) using Koshihikari/Kasalath chromosome
711 segment substitution lines and backcross progeny F2 population. Plant Prod Sci. 2008 Jan
712 1;11(4):447-56.

713

714 Kebriyaae D, Kordrostami M, Rezadoost MH, Lahiji HS. QTL analysis of agronomic traits in
715 rice using SSR and AFLP markers. Notulae Scientia Biologicae. 2012 May 10;4(2):116-23.

716

717 Khatun MM, Ali MH, Dela Cruz QD. Correlation studies on grain physicochemical
718 characteristics of aromatic rice. Pak J Biol Sci. 2003;6(5):511-3.

719

720 Khin TN. Rice mutation breeding for varietal improvement in Myanmar. Plant Mutation Reports.
721 2006 May;1(1)31-6.

722

723 Kikuchi F, Itakura N, Ikehashi H, Yokoo M, Nakane A, Maruyama K. Genetic analysis of
724 semidwarfism in high-yielding rice varieties in Japan. Japan: Bulletin of the National Institute of
725 Agricultural Sciences, Series D. 1985.

726

727 Kishor C, Prasad Y, Haider ZA, Kumar R, Kumar K. Quantitative analysis of upland rice.
728 ORYZA-An International Journal on Rice. 2008;45(4):268-72.

729

730 Veni BK, Rani NS, Prasad AS, Prasad GS. Character association and path analysis studies for
731 quality traits in aromatic rices. Andhra Agricultural Journal (India). 2003.

732 Kumar S, Gautam AS, Chandel S. Estimates of genetic parameters for quality traits in rice
733 (*Oryza sativa* L.) in mid hills of Himachal Pradesh. *Crop Research-Hisar*-. 2006;32(2):206-8.
734

735 Kumar V, Bhagwat SG. Microsatellite (SSR) based assessment of genetic diversity among the
736 semi-dwarf mutants of elite rice variety ‘WL112’. *International Journal of Plant Breeding and*
737 *Genetics*. 2012 Jan 1;6:195-205.
738

739 Lanahan MB, Ho TH. Slender barley: a constitutive gibberellin-response mutant. *Planta*. 1988
740 Jul 1;175(1):107-14.
741

742 Li W, Wu J, Weng S, Zhang Y, Zhang D, Shi C. Identification and characterization of dwarf 62,
743 a loss-of-function mutation in *DLT/OsGRAS-32* affecting gibberellin metabolism in rice. *Planta*.
744 2010 Nov 1;232(6):1383-96.
745

746 Lin YR, Wu SC, Chen SE, Tseng TH, Chen CS, Kuo SC, Wu HP, Hsing YC. Mapping of
747 quantitative trait loci for plant height and heading date in two inter-subspecific crosses of rice
748 and comparison across *Oryza* genus. *Bot Stud*. 2011 Jan 1;52:1-4.
749

750 Luo WX, Li YS, Wu BM, Tian YE, Zhao B, Zhang L, Yang K, Wan P. Effects of electron beam
751 radiation on trait mutation in azuki bean (*Vigna angularis*). *Afr J Biotechnol*.
752 2012;11(66):12939-50.
753

754 Lush JL. Intra-sire correlation and regression of offspring on dams as a method of estimating
755 heritability of characters. Record of Proceedings. American Society of Animal Production; 1940
756

757 Lincoln M, Kumar VS, Kotasthane AS and Verulkar SB. Mapping Quantitative Trait loci for
758 tillers number, plant height and their correlation in rice [*Oryza sativa* L.]. International Journal of
759 Genetics. 2017;9(10):309-13.
760

761 Marathi B, Guleria S, Mohapatra T, Parsad R, Mariappan N, Kurungara VK, Atwal SS, Prabhu
762 KV, Singh NK, Singh AK. QTL analysis of novel genomic regions associated with yield and
763 yield related traits in new plant type based recombinant inbred lines of rice (*Oryza sativa* L.).
764 BMC plant biology. 2012 Dec;12(1):137.
765

766 Matsukura C, Itoh SI, Nemoto K, Tanimoto E, Yamaguchi J. Promotion of leaf sheath growth by
767 gibberellic acid in a dwarf mutant of rice. Planta. 1998 May 1;205(2):145-52.
768

769 Moncada P, Martinez CP, Borrero J, Chatel M, Gauch Jr H, Guimaraes E, Tohme J, McCouch
770 SR. Quantitative trait loci for yield and yield components in an *Oryza sativa* × *Oryza rufipogon*
771 BC2F2 population evaluated in an upland environment. Theor Appl Genet. 2001 Jan
772 1;102(1):41-52.
773

774 Rashidi Monfared S, Mardi M, Hoseinzadeh AH, Naghavi MR. Association analysis of
775 important agronomic traits to retrotransposon markers SSAPs in durum wheat accessions. J
776 Modern Genet. 2008;3:29-35.

777 Nayudu KS, Vasline YA, Vennila S. Studies on variability, heritability and genetic advance for
778 certain yield components in rice (*Oryza sativa* L.) var. Jeeraga Samba. Crop Improvement -
779 India-. 2007;34(2):142-4
780
781 Nilan RA, Konzak CF, Wagner J, Legault RR. Effectiveness and efficiency of radiations for
782 inducing genetic and cytogenetic changes. Radiation Botany.1965;71-89.
783
784 Ogi Y, Kato H, Maruyama K, Kikuchi F. The effects on the culm length and other agronomic
785 characters caused by semidwarfing genes at the sd-1 locus in rice.Japanese Journal of Breeding.
786 1993 Jun 1;43(2):267-75.
787
788 Panse VG, Sukhatme PV. Statistical methods for agricultural workers. 2nd ed. New Delhi(India):
789 Indian Council of Agricultural Research;1961.
790
791 Pervaiz ZH, Rabbani MA, Khaliq I, Pearce SR, Malik SA. Genetic diversity associated with
792 agronomic traits using microsatellite markers in Pakistani rice landraces. Electron J Biotechnol.
793 2010 May;13(3):4-5.
794
795 Qin R, Qiu Y, Cheng Z, Shan X, Guo X, Zhai H, Wan J. Genetic analysis of a novel dominant
796 rice dwarf mutant 986083D. Euphytica. 2008 Apr 1;160(3):379-87.
797

798 Rabiei B, Kordrostami M, Sabouri A, Sabouri H. Identification of QTLs for Yield Related Traits
799 in Indica Type Rice Using SSR and AFLP Markers. *Agriculturae Conspectus Scientificus*. 2015
800 Dec 1;80(2):91-9.
801
802 Rabiei B. Linkage map of SSR markers and QTLs detection for heading date of Iranian rice
803 cultivars. *J AgricSci Technol*. 2007 Jan 28;9:235-42.
804
805 Saif-ur-Rasheed M, Sadaqat HA, Babar M. Correlation and path analysis for yield and its
806 components in Rice (*Oryza sativa* L.). *Asian J Plant Sci*. 2002;1(3), 241-244.
807
808 Rashid KA, Kahliq I, Farooq MO, Ahsan MZ. Correlation and cluster analysis of some yield and
809 yield related traits in Rice (*Oryza Sativa*). *Int J Agric Sci Res (Chennai)*. 2013;3(4):25-30.
810
811 Reddi TVV, Rao DR. Relative effectiveness and efficiency of single and combination treatments
812 using gamma rays and sodium azide in inducing chlorophyll mutations in rice. *Cytologia*
813 (Tokyo). 1988 Sep 25;53(3):491-8.
814
815 Roy JK, Bandopadhyay R, Rustgi S, Balyan HS, Gupta PK. Association analysis of
816 agronomically important traits using SSR, SAMPL and AFLP markers in bread wheat. *Curr Sci*.
817 2006 Mar 10;90(5):683-9.
818

819 Sakamoto T, Miura K, Itoh H, Tatsumi T, Ueguchi-Tanaka M, Ishiyama K, Kobayashi M,
820 Agrawal GK, Takeda S, Abe K, Miyao A. An overview of gibberellin metabolism enzyme genes
821 and their related mutants in rice. *Plant Physiol.* 2004 Apr 1;134(4):1642-53.
822
823 Sangodele EA, Hanchinal RR, Hanamaratti NG, Shenoy V, Kumar VM. Analysis of drought
824 tolerant QTL linked to physiological and productivity component traits under water-stress and
825 non-stress in rice (*Oryzasativa* L.). *Int J Curr Res Acad Rev.* 2014;2(5):108-13.
826
827 Sankar PD, Sheeba A, Anbumalarmathi J. Variability and character association studies in rice
828 (*Oryza sativa* L.). *Agricultural Science Digest.* 2006;26(3):182-4.
829
830 Sasaki A, Ashikari M, Ueguchi-Tanaka M, Itoh H, Nishimura A, Swapan D, Ishiyama K, Saito
831 T, Kobayashi M, Khush GS, Kitano H, Matsuoka M. Green revolution: a mutant gibberellin-
832 synthesis gene in rice. *Nature.* 2002 Apr;416(6882):701.
833
834 International Rice Research Institute. Standard Evaluation System For Rice-International Rice
835 Testing Program. Manila (Philippines); 2013.
836
837 Shadakshari YG, Chandrappa HM, Kulkarni RS, Shashidhar HE. Induction of beneficial
838 mutants in rice (*Oryza sativa* L.). *Indian J Genet Plant Breed.* 2001;61(3):274-6.
839
840 Shi C, Shen Z. Effects of semidwarf gene *sd1* on agronomic traits in rice (*Oryza sativa* subsp.
841 *indica*). *Zhongguoshuidaokexue.* 1996;10(1):13-8.

842 Siddiqui SA, Singh S. Induced genetic variability for yield and yield traits in basmati rice. World
843 Journal of Agricultural Sciences. 2010;6(3):331-7.
844

845 Singh SP, Singhara GS, Pararay GA, Bhat GN. Genetic variability and character association
846 studies in rice (*Oryzasativa* L.). Agricultural Science Digest. 2006;26(3):212-4.
847

848 Soomro AM, Naqvi MH, Bughio HR, Bughio MS. Sustainable enhancement of rice production
849 through the use of mutation breeding. Plant Mutation Reports. 2006;1:13-7.
850

851 Spielmeyer W, Ellis MH, Chandler PM. Semidwarf (sd-1), “green revolution” rice, contains a
852 defective gibberellin 20-oxidase gene. Proceedings of the National Academy of Sciences. 2002
853 Jun 25;99(13):9043-8.
854

855 Subashri M, Robin S, Vinod KK, Rajeswari S, Mohanasundaram K, Raveendran TS. Trait
856 identification and QTL validation for reproductive stage drought resistance in rice using selective
857 genotyping of near flowering RILs. Euphytica. 2009 Mar 1;166(2):291-305.
858

859 Subbaiah PV, Sekhar MR, Reddy KH, Reddy NPE. Variability and genetic parameters for grain
860 yield and its components and kernel quality attributes in CMS based rice hybrids (*Oryza sativa*
861 L.). Int J Appl Biol Pharm. 2011;2:603-9.
862

863 Subudhi PK, De Leon TB, Tapia R, Chai C, Karan R, Ontoy J, Singh PK. Genetic interaction
864 involving photoperiod-responsive Hd 1 promotes early flowering under long-day conditions in
865 rice. *Sci Rep*. 2018 Feb 1;8(1):2081.
866

867 Susanto U, Aswidinnoor H, Koswara J, Setiawan A, Lopena V, Torizo L, Parminder VS. QTL
868 mapping of yield, yield components, and morphological traits in rice (*Oryza sativa* L.) using SSR
869 marker. *Buletinul Institutului Agronomic Cluj-Napoca. Seriaagricultură..* 2008;36(3).
870

871 Swaminathan MS. Report of meeting of the symposium in the use of induced mutations in plant
872 breeding. *Radiation botany*. 1965 Jan 1;5(1):65-9.
873

874 Takeuchi Y, Hori K, Suzuki K, Nonoue Y, Takemoto-Kuno Y, Maeda H, Sato H, Hirabayashi H,
875 Ohta H, Ishii T, Kato H. Major QTLs for eating quality of an elite Japanese rice cultivar,
876 Koshihikari, on the short arm of chromosome 3. *Breed Sci*. 2008;58(4):437-45.
877

878 Ueguchi-Tanaka M, Fujisawa Y, Kobayashi M, Ashikari M, Iwasaki Y, Kitano H, Matsuoka M.
879 Rice dwarf mutant dl, which is defective in the α subunit of the heterotrimeric G protein, affects
880 gibberellin signal transduction. *Proc Natl Acad Sci U S A*. 2000 Oct 10;97(21):11638-43.
881

882 Wang H, Chen Z, Guo T, Liu Y, Li H. Genetic analysis and gene mapping of dwarf mutant rice
883 CHA-1. In *Induced plant mutations in the genomics era. Proceedings of an International Joint*
884 *FAO/IAEA Symposium,2009 (pp. 431-433). Joint FAO/IAEA Division of Nuclear Techniques*
885 *in Food and Agriculture, International Atomic Energy Agency, Rome.*

Xu Y, Beachell H, McCOUCH SR. A marker-based approach to broadening the genetic base of rice in the USA. Crop Sci. 2004 Nov 1;44(6):1947-59.

Yadav P, Rangare NR, Anurag PJ, Chaurasia AK. Quantitative analysis of rice (*Oryza sativa* L.) in Allahabad agro climate zone. Journal of Rice research. 2010;3(1):16-8.

Yankulov MT, Isasi EM, Abreu F. Some aspects of the sensitivity and mutability of two French bean varieties under the influence of gamma radiation from ⁶⁰Co and ethyl methanesulphonate (EMS).Ciencias de la Agricultura. 1980;(7):59-64.

Gao Z, Liu X, Guo L, Liu J, Dong G, Hu J, Han B, Qian Q. Identification of a novel tillering dwarf mutant and fine mapping of the TDDL (T) gene in rice (*Oryza sativa* L.). Chin Sci Bull. 2009 Jun 1;54(12):2062-8.

Zhao X, Daygon VD, McNally KL, Hamilton RS, Xie F, Reinke RF, Fitzgerald MA. Identification of stable QTLs causing chalk in rice grains in nine environments. Theor appl genet. 2016 Jan 1;129(1):141-53.

Zou J, Chen Z, Zhang S, Zhang W, Jiang G, Zhao X, Zhai W, Pan X, Zhu L. Characterizations and fine mapping of a mutant gene for high tillering and dwarf in rice (*Oryza sativa* L.). Planta. 2005 Nov 1;222(4):604-12.

Legends

Tables

Table 1. List of microsatellite markers used in this study

Table 2a. Mean, variability and heritability estimates of morphological traits in the M2 generation of improved white ponni generated by gamma irradiation

Table 2b. Mean, variability and heritability estimates of morphological traits in the M3 generation of improved white ponni generated by gamma irradiation

Table 3. Mutants exhibiting altered morphological traits observed in M2 and M3 generation (gamma rays) of improved white ponni

Table 4. Mean performance of semi-dwarf and early mutants of improved white ponni in M3 generation

Table 5a. Genotypic correlation coefficients among different quantitative traits of selected mutants (Semi-dwarf and early mutants) in M3 generation (gamma rays) of improved white ponni

Table 5b. Genotypic correlation coefficient among different quality traits of selected mutants (Semi-dwarf and early mutants) in M3 generation (gamma rays) of improved white ponni

934 Table 6a. Allelic distribution and PIC values of SSR markers in this study

935

936 Table 6b. Marker-Trait Associations estimated by simple linear regression analysis (SLRA)

937

938 Table 7. GA3 response of dwarf and early mutants using the microdrop method

939

940 Table 8. Shoot length variation after application of GA3 by the microdrop method

941

942 **Figures**

943 Figure1. Viable morphological mutants observed in the M₂ generation of Improved White Ponni

944 1) Early mutant 2) Tall early mutant 3) Early Semi-dwarf mutant 4) Extremely dwarf mutant 5)

945 High tillering late mutant 6) High tillering mutant 7) Narrow rolled leaf mutant 8) Upper albino

946 mutant

947

948 Figure 2: Variation on Morphological traits in identified mutants in M₃ generation

949 Flowering and plant height variation

950 Variation on culm length - 1st, 2nd, 3rd and 4th Intermodal length 1) Improved White Ponni

951 2) IWP 43-1 3) IWP 59-1 and 4) IWP 50-11

952

953 Figure 3: Variation in internodal lengths of Improved White Ponni mutants compared to

954 wildtype

955

956 Figure 4: Genotypic data of novel Improved White Ponni mutants using microsatellite markers

957 Genotyping result of RM252

Genotyping result of RM310

Figure 5: Variation on Shoot length in early semi-dwarf mutants of Improved White Ponni with and without GA₃ application by micro -drop method

Figure 6: Variation in the alpha-amylase activity of unique semi-dwarf mutant in this study
a) Wildtype- Improved White Ponni b) Identified mutants. -GA₃ – without GA₃ application and +GA₃ – with GA₃ application

Figure 7: Variation in cell structural pattern of the third internodal region in Improved White Ponni and Semi-dwarf Mutant (IWP 59-1). This image explaining the cell size (length and breadth of each cell between wildtype and mutant (green markings)

bioRxiv preprint doi: <https://doi.org/10.1101/500637>; this version posted December 18, 2018. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Fig 2a

Table 1. List of Microsatellite Markers Used in this study

S.No.	Name of the primer	Chrom.	Position (cM)	QTL associated	References
1.	RM580	1	68.2	<i>qPH-1, qLFI-1</i>	Feng-hua <i>et al.</i> , 2005
2.	RM246	1	115.2	<i>qPH1.1, qGP-1</i>	Lincoln <i>et al.</i> , 2017, Rabiei <i>et al.</i> , 2007
3.	RM543	1	145.6	<i>qHd1</i>	Rabiei, 2007
4.	RM302	1	147.8	<i>qTGW-1, qDFF1-1</i>	Sangodele <i>et al.</i> , 2014, Marathi <i>et al.</i> , 2012
5.	GA20 Oxi_1	1	147.8	<i>Sd-1</i>	Spielmeier <i>et al.</i> , 2002
6.	GA20 Oxi_2	1	147.8	<i>Sd-1</i>	Spielmeier <i>et al.</i> , 2002
7.	GA20 Oxi_3	1	147.8	<i>Sd-1</i>	Spielmeier <i>et al.</i> , 2002
8.	RM263	2	127.5	<i>qPh2.2</i>	Lin <i>et al.</i> , 2011
9.	RM250	2	170.1	<i>qPh2.2</i>	Lin <i>et al.</i> , 2011
10.	RM5430	2	111.5	<i>wt100-vb2.1</i>	Susanto <i>et al.</i> , 2008
11.	RM5699	2	42.1	<i>qHPH2d</i>	Zhao <i>et al.</i> , 2016,
12.	RM240	2	158	<i>qHd2</i>	Lin <i>et al.</i> , 2011
13.	RM207	2	191.2	<i>qHd2</i>	Lin <i>et al.</i> , 2011
14.	RM5849	3	18.4	<i>qHA3-3</i>	Takeuchi <i>et al.</i> , 2008
15.	RM7365	3	49.3	<i>qTGW3-1</i>	Hori <i>et al.</i> , 2012
16.	RM545	3	35.3	<i>qHD-3</i>	Liu <i>et al.</i> , 2012, Lin <i>et al.</i> , 2011
17.	RM7	3	64	<i>qDTH3.1</i>	Moncada <i>et al.</i> , 2001
18.	RM3203	3	2.2	<i>qHd3</i>	Lin <i>et al.</i> , 2011
19.	RM252	4	99	<i>qPh4</i>	Lin <i>et al.</i> , 2011
20.	RM567	4	153.6	<i>qPh4</i>	Lin <i>et al.</i> , 2011
21.	RM5709	4	109.9	<i>qPHT4-3</i>	Marathi <i>et al.</i> , 2012
22.	RM551	4	0	<i>qPHT4-1</i>	Marathi <i>et al.</i> , 2012
23.	RM430	5	78.7	<i>qPh5</i>	Lin <i>et al.</i> , 2011
24.	RM480	5	130.6	<i>qPh5</i>	Lin <i>et al.</i> , 2011
25.	RM541	6	75.5	<i>qPh6.1</i>	Lin <i>et al.</i> , 2011
26.	RM30	6	125.4	<i>qPh6.1</i>	Lin <i>et al.</i> , 2011
27.	RM253	6	37.0	<i>qHD6.1</i>	Zhao <i>et al.</i> , 2016
28.	RM7023	6	51.3	<i>qGT6d</i>	Lapitan <i>et al.</i> , 2009
S.No.	Name of the primer	Chrom.	Position (cM)	QTL associated	References