Rapid evolution by spontaneous mutation increases genetic diversity facilitating plant population survival

Henning Nottetbrock¹, Mao-Lun Weng¹, Matthew T. Rutter² & Charles B. Fenster¹

¹Department of Biology and Microbiology, South Dakota State University, Brookings, SD USA

²Department of Biology, College of Charleston, Charleston, SC USA

Abstract

Using a mechanistic eco-evolutionary trait-based neighborhood-model, we quantify the impact of mutations on spatial interactions to better understand the potential effect of niche evolution through mutations on the population dynamics of *Arabidopsis thaliana*. We use 100 twenty-fifth generation mutation accumulation (MA) lines (genotypes) derived from one founder genotype to study mutational effects on neighbor responses in a field experiment. We created individual-based maps (15,000 individuals), including phenotypic variation, to quantify mutational effects within genotypes versus between genotypes on reproduction and survival. At small-scale, survival is enhanced but reproduction is decreased when a genotype is surrounded by different genotypes. At large-scale, seed set is facilitated by different genotypes while the same genotype has either no effect or negative effects. Mutations may provide a mechanism for plants to quickly evolve niches and may drive competition, facilitation and selection with profound consequences for future population and community dynamics.

Introduction

A complete understanding of population and community dynamics requires linking intraspecific genetic diversity with spatial ecological interactions (Bolnick et al. 2011, Genung et al. 2011). Evolutionary mechanisms -- mutation, drift, gene flow and selection -- are responsible for intraspecific genetic variation contributing to both ecological structure and species diversity (Hart et al. 2016). These community properties impact ecological mechanisms such as competition and facilitation (Whitlock 2014). Competition and facilitation in turn affect population and community dynamics that depend on variation in demographic rates (Chase et al 2002, Solivers et al 2015). Demographic rates are influenced by spatial interactions of genetic variation and this variation may contribute to species coexistence: genotypes can hinder or favor the survival of each other by using similar or different resources with negative or positive consequences for coexisting species (Hart et al. 2016, Hausch et al. 2018). Intraspecific genetic variation contributing to individual trait variation and environmental adaptability may promote species coexistence by both increasing habitat heterogeneity and altering competitive hierarchies (Violle et al. 2012, Ehlers et al. 2016). However, intraspecific genetic variation may hinder species coexistence when intraspecific genetic variation is diminished by competition between individuals (Hart et al 2016). To fully understand the causes of coexistence mechanisms, we need to integrate intraspecific evolutionary mechanisms with intraspecific ecological mechanisms underlying spatial species interactions (Bolnick et al. 2011, Ehlers et al. 2016).

The number of studies exploring eco-evolutionary dynamics and their potential feedbacks on population and community dynamics has dramatically increased over the last two decades (Shefferson and Salguero-Gomez 2015). An important current issue in community and ecosystem genetics research is determining the relevance of intraspecific genetic variation and genetic differentiation (divergence) to ecological and evolutionary processes at the community and ecosystem level (Genung et al. 2011, Pujol et al. 2018). Quantifying intraspecific trait variation defining the fundamental niches of species is an important link between ecology and evolution (Violle and Jiang 2009). Spatial structure is another important component of realistic eco-evolutionary dynamics, such that in spatially structured populations, selection is determined by the interplay between demographic and genetic structures (Lion et al. 2011). Demographic structure describes the spatial distribution of individuals through birth, death and migration resulting in spatial patterns, while genetic structure describes the spatial distribution of genotypes. The amount and spatial pattern of genetic variation may constrain evolution of traits influencing competitive ability of individuals (Wilson 2014). Moreover, competitive abilities depend on species' niches. The strength of competition is determined by how much individual niches overlap with each other (Hutchinson 1957, Holt 2009) and how long coexistence has occurred (Connell 1980).

Species niche breadths are reflected by individual trait variability and intraspecific and interspecific genetic and spatial interactions driven by environmental conditions. Thus, intraspecific trait variability shaped by intraspecific genetic variation can influence ecological mechanisms driving variation among individual persistence (Lankau 2009). Only when the

spatial relationship between traits and demographic parameters is explicitly described can robust hypotheses about the effects of individual variation on competitive outcomes be accurately formulated (Hart et al. 2016). Especially in controlled common garden environments, genetic variation in one species can have predictable and heritable effects on associated communities and ecosystems (Carr and Dudash 1995; Whitham et al. 2003, 2006; Johnson & Stinchcombe 2007; Bailey et al. 2009; Johnson, Vellend & Stinchcombe 2009). Hence, coupling evolutionary genetics with community ecology may advance our understanding of species interactions and population and community dynamics (Baron et al. 2016), especially for populations and communities that suffer environmental change (Bellard et al. 2012).

An important gap in our understanding of the link between evolutionary and ecological processes is the nearly complete absence of data quantifying how rapidly genetic variation governing within species competitive hierarchies evolves (Hausch et al. 2018). An important source of novel population genetic variation is mutation. Therefore, we aim to investigate the link between evolutionary genetics and spatial ecological interactions of mutation accumulation lines (MA lines) in the model plant organism *Arabidopsis thaliana* (Brassicaceae) to advance our understanding of the genetic origins of population and community dynamics, contributing to a fuller understanding of the maintenance of biodiversity. We take advantage of 25th generation *A. thaliana* mutation accumulation (MA) lines that were planted under field conditions in years 2004 and 2005 with spatial records of each individual each year (e.g., Rutter et al. 2010; 2012, Rutter et al. 2018) (Fig. 1).

Here we focus on the neighborhood effects of MA lines on focal plants from the same MA line (within genotypes) or different MA lines (between genotypes) to quantify the evolution of intraspecific genetic variation through the accumulation of spontaneous mutations. We implement trait-based neighborhood models (Nottebrock et al. 2017a, Nottebrock et al. 2017b, Lachmuth et al. 2017) to understand ecological mechanisms of within and between MA lines genetic variation to examine eco-evolutionary dynamics (Lion 2018). Specifically, we ask: 1) Do spontaneous mutations rapidly introduce enough genetic variation among MA lines to influence competition and facilitation of *A. thaliana*? 2) Do spontaneous mutations of MA lines contribute to competition, facilitation and selection promoting differential MA line survival and reproduction at different spatial scales? 3) How may competitive and facilitative effects potentially determine population and community dynamics?

Methods

MA lines are generated from a single nearly homozygous individual founder and cultivated via limited effective population number. In the case of *A. thaliana*, this occurs through single seed descent, resulting in $N_e = 1$. Thus typical MA line cultivation results in an unbiased sample of mutation effects ranging from deleterious to advantageous, although lethal mutations are excluded (Lynch and Walsh 1998). Each MA line accumulates independent spontaneous mutations. After the propagation of a set of MA lines through multiple generations, the genetic differences among the MA lines and between those lines and the founder reflect the input of mutation. Significant MA line effects for multiple traits, including performance and trait measures, were found under both field and greenhouse conditions (Rutter et al. 2010; Roles et al. 2016, Rutter et al. 2018). Each of the MA lines in our experiment is fixed for an average of 20 different sequence level mutations, single nucleotide mutations (SNMs) and indels combined (Ossowski et al 2010, Rutter et al. 2012, Weng et al. 2019).

MA lines and field experiments

We used survival and seed set data of *A. thaliana* MA lines and the founder as assessed in field experiments in 2004 and 2005 from Rutter et al. (2010, 2012 and 2018) planted in a randomized design (Fig. 1). Rutter et al. (2010, 2012) planted seedlings of 100 MA lines and the founder at the four-leaf stage, approximately two weeks post germination, into a secondary successional field at Blandy Experimental Farm (BEF) in Virginia (39°N, 78°W). Each of the 100 MA lines was used to found up to five sublines to minimize biases due to maternal effects

introduced by the specific location within the greenhouse. We founded six sublines from each of the six lines representing the premutation founder genotype. In 2003, subline plants were used to generate all seed utilized in all field experiments. In each planting, 7504 individuals were planted, 7000 individuals of 100 MA lines (70 replicates per MA line, 14 replicates per subline) and 504 individuals of the founder (14 replicates per subline). The planting environment corresponds to a spring ephemeral life-history, where plants germinate and complete the life-cycle in the spring. At the time of planting, vegetation was scant but present. By harvest, the *A. thaliana* individuals were dwarfed by naturally occurring vegetation.

The plot was arranged in 14 spatial blocks with each containing 12 subblocks (Fig. 1B) (total plot area approximately 35 x 25 m). Each block included one seedling from each subline and in total 7504 individuals. We used the spatial information of each individual within the described design and created a raster of all plant individuals with R packages (raster, maps, maptools). We used individual-based maps neighborhood matrices with exact spatial and trait information of each genotype. If all five sublines did not produce enough seedlings to distribute in all blocks, seedlings from other sublines of the same MA line were overrepresented in blocks to maintain the same overall number of plants per MA line. Plants dying within the first 3 days of transplant (about 50 plants) were considered to have died from transplant shock and were replaced with another plant from the same MA line. Plants were censused weekly for survival. Plants were harvested by late May, by which time they had senesced. In 2004 a total of 5915 individuals including 394 founders survived. In 2005, a total of 4506 individuals survived

140 including 302 founders. Plants were oven dried and biomass was measured. All fruits produced
 141 by each plant were counted and in combination with seed production used as the
 142 measurements of seed set. For our analysis, we used measurements of two response variables
 143 representing an important part of the life history of *A. thaliana* to calculate direct
 144 neighborhood effects on: a) the survival rate of plant individuals and b) individual seed set from
 145 plants that survived and produced fruits.

146

Statistical Analyses

We use eco-evolutionary trait-based neighborhood models that include intraspecific genetic variation and phenotypic variation expressed by plant trait biomass to analyze competition and facilitation between individuals of *A. thaliana*. We analyzed plant survival rate and focal seed set measurements from years 2004 and 2005 separately. For each year, we considered all individuals as focal plants in the analysis based on individual-based neighborhood matrices to analyze 1) rate of plant survival to reproduction and 2) seed set of those plants that survived to reproduction. We analyzed neighboring plants to focal plants in a radius of 80 cm (small-scale) or 200 cm (large-scale) of a given focal plant as two spatial scales in the neighborhood analyses (Fig. 1) to quantify selection, competition or facilitation between plants depending on their genotypic and phenotypic variation.

We used extensions of linear mixed models (package lme4, Bates et al. 2014) in R ver. 3.3.3 (www.r-project.org) to conduct neighborhood analyses of focal seed set and survival. We assumed binomial errors for the analyses of plant survival and Poisson errors for analyses of seed set. The mixed models described interactions among plants by including neighborhood indices as explanatory variables at two spatial scales in separate models. Neighborhood indices are spatial density effects of surrounding neighborhood plants that affect focal seed set and survival. For each plant, we used the Euclidian distance between the focal plant and the neighboring plants to compute response effects of intra- (same genotype) or inter-genotypic (different genotype) neighbors in a given radius around focal plants (Nottebrock et al. 2017). Moreover, we used a neighborhood index that accounts for the decline of neighbor effects with

distance from the focal plant (Uriarte et al. 2010) and summed the amount of biomass from all individuals in a radius of 80 cm or 200 cm respectively by a Gaussian interaction kernel (Lachmuth et al. 2018, Nottebrock et al. 2017, Damgaard 2004). We used random effects of block and subblock to correct for environmental variation between local heterogeneous conditions. Importantly, we correct for between MA line effects by including MA lines as a random effect. In addition, including a random slope of biomass on each random intercept corrects for the intraspecific phenotypic variation depending on local conditions of plant focal individuals. Moreover, the weighted neighborhood density by plant biomass accounts for environmental variation between neighboring plants. Our model parameters and a detailed model description of subblock and block models can be found in S1 supplementary text (Supplementary Material).

Neighborhood matrices of all individuals (individual-based maps) were used to analyze the effect of intra- and inter-genotypic neighbors on survival and focal seed set with spatial interaction kernels of neighborhood (plant biomass) density. By incorporating different genotypes and phenotypic variation, we can quantify how important genetic variation is for neighborhood models and if the phenotypic variation explains spatial interactions between individuals. We assume the consequences of genetic differences to be larger between MA lines than between any MA line with the founder. This is a valid assumption since each MA line differs from the other by approximately $20 + 20 = 40$ mutations, while any two replicates within a MA line will differ by one generation, ≤ 2 mutations. Thus, we simulated line effects from

parameters derived from MA lines as random effects with the R package ‘merTools’ and the function ‘plotREsim’ in R 2018.

We used the trait values of neighbors to calculate trait-based neighborhood indices including plant biomass as a trait (Goldberg & Fleetwood, 1987; Goldberg & Landa, 1991; Cahill et al., 2005). We fitted eco-evolutionary trait-based neighborhood models at two different spatial scales for response variables (survival and seed set) for each of the two and both years. To address our objectives, we first analyzed models with differential effect in which intra-genotypic neighbors (within all MA lines and founder) had a different effect on survival (A1, A2, Table 1) and seed set (B1, B2, Table 1) than inter-genotypic neighbors at small-scale (s) and large-scale (l). In addition, we analyzed models with neutral effects on survival (A1, A2, Table 1) and seed set (B1, B2, Table 1) that included total neighbor density without the split between intra-genotypic and inter-genotypic neighbors at small-scale (s) and large-scale (l). To this end, all models were fitted with two separate neighborhood indices that were calculated from intra- and inter-genotypic neighbors. To justify the inclusion of individual plant biomass as trait-values for interacting plants in the model, we used AICc to compare the models with and without the trait-proxy (Burnham and Anderson 2002). We found that all models perform better including biomass as a trait-proxy ($\Delta AICc > 2$). All eco-evolutionary trait-based neighborhood models contained random effects of subblock nested in block at block scale and subblock scale on the intercept, MA line identity on the intercept and the focal trait-value (plant biomass) on the slope. Additionally, because direct environmental variables were not measured during the field experiments, we included in each model the individual’s biomass to correct for environmental

212 conditions for spatial autocorrelation. All variables are scaled and centered to assure
 213 comparability between predictor variables. Models of differential and neutral effects for 2004
 214 and 2005 (Table 1, A1-A2, B1-B2) are fitted at small-scale (80cm scale) and at large-scale
 215 (200cm scale). Hereafter, the 80 cm scale models are referred to as “small-scale” models and
 216 the 200 cm scale models are referred to as “large scale” models. Neighborhood indices, intra-
 217 and intergenotypic variation and total variation of biomass density are included as inverse
 218 density variables ($1/1+\text{density}$). We compared models of differential and neutral effects
 219 through likelihood ratio tests (LRTs).

220

Results

Focal plants surrounded by plants of the same genotypes varied from 0 to 6 individuals at small-scale (sub-block) and 5 to 6 individuals at large-scale (block). After determining that evolutionary trait-based models perform better when including plant biomass as a trait proxy to calculate trait-based neighborhood indices we found that all models including plant biomass perform better than models including only neighborhood density ($\Delta AIC > 2$, Supplementary Material S2).

Weighted neighbor effects of biomass density on genotypes for plant survival

At small-scale, survival rate is larger when surrounded by inter-genotypic than intra-genotypic neighbors for both years 2004 and 2005 (model A1 and A2; Table 2 and Fig. 2a). This difference of survival rate between intra- and inter-genotypic plants indicates that inter-genotypic neighbors select for genotypic diversity and show stronger competitive effects on the survival of intra-genotypic neighbors at small-scale. This finding is demonstrated by the superior performance of the differential model relative to the neutral model (year 2004: LRT: $\chi^2_{1DF} = 5.21$, $p < 0.05$; year 2005: LRT: $\chi^2_{1DF} = 14.73$, $p < 0.001$). The differential and neutral models of survival at large-scale have only non-significant effects (Table 2). In addition, comparing the AIC we found that models at large-scale perform worse than at small-scale for both 2004 and 2005 (Table 2). We therefore will only discuss small-scale effects on survival. The distance kernel (alpha) at small-scale shows that neighboring plants of different genotypes reduce plant survival by 50% at 53 cm in 2004 and 48 cm in 2005.

Weighted neighbor effects of biomass density on genotypes for seed set

At both small and large scales, the differential model performs better than the neutral model: at small scales (year 2004: LRT: $\chi^2_{1DF} = 52.48$, $p < 0.001$; year 2005: LRT: $\chi^2_{1DF} = 17.37$, $p < 0.001$; respectively) and at large scale (year 2004: LRT: $\chi^2_{1DF} = 679.44$, $p < 0.001$; year 2005: LRT: $\chi^2_{1DF} = 21.17$, $p < 0.001$; respectively). The small-scale model of seed set that included weighted biomass densities demonstrates the reduction of seed set is larger when surrounded by inter-genotypic than intra-genotypic neighbors for both years 2004 and 2005 (model B1 and B2; Table 2 and Fig. 2a). The difference between intra- and inter-genotypic plants indicates that inter-genotypic neighbors have a stronger competitive effect on seed set than intra-genotypic neighbors at small-scale. In contrast, at large-scale seed set increases when surrounded by inter-genotypic neighbors but decreases when surrounded by intra-genotypic neighbors for both years 2004 and 2005 (model B1 and B2; Table 2 and Fig. 2b). Comparing the AIC between small and large-scales, models at large-scale for 2004 and 2005 perform better (Table 2).

Genotype effects on plant survival and seed set

The distance kernel at small-scale in 2004 indicates a reduction of seed set (competition) whereas at large-scale are consistent with an increase of seed set (facilitation). Neighboring plants from all genotypes of *A. thaliana* at small-scale reduce 50% of focal seed set at 23 cm and at large-scale facilitate 50% of focal seed set at 86 cm. The estimation of genotype effects (MA lines and founder) simulated as conditional means and expressed as odds ratios show hierarchical orders of MA line and founder competitive effects at small-scale (Fig. 3). We found

no significance of simulated competitive effects of all MA lines and founder calculated from survival and seed set models at small-scale (ANOVA, $p > 0.1$, Fig. 4a). In addition, we found no significant correlation between competitive effects of MA lines and founder at large-scale. In addition, we did not find selective effects of MA lines and founder on the survival at small-scale (ANOVA, $p > 0.1$, Fig. 4b). However, we found strong correlation between competitive effects of MA lines and founder at small-scale and large-scale indicating a genetic trade-off (ANOVA, $F=460.07$, $P < 0.0001$, Fig. 4c).

Discussion

We are able to quantify individual spatial interactions between and within genotypes (MA lines and founder) using a mechanistic eco-evolutionary trait-based neighborhood model. We demonstrate intraspecific genetic variation due to spontaneous mutations can shape competitive abilities of genetically different individuals of *A. thaliana*. Notably, these differences arose in just 25 generations of mutation and in the absence of natural selection. Many of the mutations differentiating the MA lines also appear in nature, contributing to *A. thaliana* genetic polymorphism (Weng et al., 2019). Plant survival is higher when surrounded by inter-genotypic neighbors supporting genetic diversity. Effects on focal seed sets were reversed between small and large scales; inter-genotypic neighbors have stronger negative (competitive) or positive (facilitative) effects than intra-genotypic neighbors on focal seed sets at small-scale or large-scale, respectively. Moreover, at small-scale, competitive effects of different MA lines have similar impacts on survival and seed set. At large-scale, the intra-genotypic effect on survival turns into less competitive effects on seed set. In contrast, different genotypes show stronger facilitative effects within the population at large-scale, maybe because plants are more competitive against other species. Below we discuss how these scale effects provide insight on community assembly and potential coexistence.

Selection, Competition and Facilitation

Because of stronger statistical signals of effect sizes between genotypes than within genotypes, niche variation may result in niche expansion or reduction (Ehlers et al. 2016). However, these

competitive effects only occur between plants that survive to reproduction and affect seed-set. The neighbor effect on plant survival suggests kin recognition despite our randomized field experiment that likely reduced the spatial autocorrelation of genotypes (Hamilton 1964, Kubisch et al. 2013). Seed-dispersal is limited in space because the *A. thaliana* fruit do not dehiscently explode as they do in related species (Hofhuis and Hay 2017), although a seed bank ensures some dispersal through time (Falahati-Anbaren et al. 2014). The neighbor effect increasing genetic diversity of *A. thaliana* populations can have a positive effect on the coexistence of competing species (Vellend 2006). If we assume that the increase of genetic diversity due to spontaneous mutations of *A. thaliana* affects the extent of individual plant niches, at large-scale the inclusion of other plant species should result in stronger competitive effects of *A. thaliana* with other plant species (Hausch et al. 2018). We demonstrate that *A. thaliana* individuals increase their competitive abilities with an increase of genetic diversity, perhaps due to the expansion of their individual niches due to more variation in plant traits.

Antagonistic interactions between genotypes on different spatial scales may result in genetic trade-offs when mutations are advantageous or deleterious, which over many generations could provide a strong stabilizing force maintaining both species and genetic diversity in this system and promote coexistence (Lankau 2008). Thus, spontaneous mutation could also provide an additional evolutionarily stabilizing effect on community dynamics. Our study demonstrates the potential for a fundamental evolutionary process, mutation, to have profound consequences for community structure. When *A. thaliana* is rare, selection would favor genotypes that compete well and enhance the population's survival relative to that of its

interspecific competitor (Lankau 2009). In contrast, when *A. thaliana* is common and the interspecific competitor is rare, selection would favor *A. thaliana* genotypes that are good intraspecific competitors. This trade-off may result in a decrease in the interspecific competitive ability of *A. thaliana*, effectively increasing the competitor's fitness relative to that of *A. thaliana* (Lankau 2009, Lankau and Strauss 2007). However, the genetic trade-off between intra-and interspecific competitive abilities due to mutation remains unknown because our dataset only includes performance of *A. thaliana* individuals without quantifying the surrounding intraspecific environment (Chesson 2000, Adler et al. 2007).

Competitive Hierarchies of MA lines and Founder lines

Our result show scale dependent competition and facilitation (Nottebrock et al. 2017b) providing further evidence that in a relative short evolutionary time scale spontaneous mutations may change the competitive hierarchies between founders and specific MA lines due to advantageous or deleterious mutations (Rutter et al. 2010). In contrast to Masclaux et al. 2010, we found differential responses to similar genotypes vs different genotypes in *A. thaliana* depending on the neighbor effect on the plant survival or seed set. However, the interaction between similar genotypes and different genotypes seems to depend on the strength of the competitive abilities of the accessions (this study). Comparisons between MA lines and founder demonstrate that competitive hierarchies follow different orders and competitive effects occur at different life-history stages at different spatial scales. Eco-evolutionary processes might reflect spatial selection for diversified genotypes because of niche evolution and individuals of different genotypes that survived to reproduction have stronger competitive abilities (Ehlers et

al. 2016). Moreover, the hierarchical order of competitive effects between MA lines and founders shows strong variation.

Eco-evolutionary dynamics

Populations can adapt evolutionarily to their environment on a time scale equivalent to that of ecological processes and affect present day species interactions, i.e., coexistence (Fussmann et al. 2007). Yet, sustained and rapid climate change could deplete genetic variance faster than it can be replenished by mutation (Fournier-Level et al. 2016). In our study, we show that only 25 generations are enough to influence present day plant interactions of *A. thaliana* in the presence of other plant species. Possible effects of mutations on trait mechanisms might reflect the increased seed set of *A. thaliana* at large-scale, because of the presence of another plant species allowing the establishment and growth of *A. thaliana* (Ehlers et al. 2016). Many traits can influence competition and facilitation between individuals of different genotypes (e.g. Chapin et al. 1993; Caradus and Woodfield 1998; Hausmann et al. 2005). Intraspecific competition between *A. thaliana* individuals tends to be higher than between interspecific competitors (Lankau and Strauss 2007, Van Dam and Baldwin 1998; Tiffin 2002, Strauss and Irwin 2004). Such spatial mechanisms might be described by the scale difference of competition at small-scale and facilitation at large-scale between different *A. thaliana* genotypes. Competition and facilitation are therefore dependent on the extension or reduction of individual niches, which is based on the ability to advance or distract traits due to spontaneous mutations. However, we are not able to directly link our results to benefactor or antagonistic plants, because we focused in our experiment on *A. thaliana* densities and fitness components.

Nevertheless, genotypic-specific interactions are well studied. A number of studies of *A. thaliana* MA lines demonstrate mutation effects on plant traits such as leaf weight, flowering time, trichome density, number of leaves at bolting, duration of vegetative and flowering period (Camara and Pigliucci 1999, Rutter et al. 2010, Stearns and Fenster 2016). It is conceivable that the increase in variance of these traits through spontaneous mutations could contribute to the evolution of competitive hierarchies.

Consequences for population and community dynamics

Competition between genetically different individuals of *A. thaliana* at small-scale and facilitation at large-scale has important implications for population structure (Cahill et al. 2005). Plants may cooperate by competing less, or act selfishly by competing more (Dudley et al. 2013). However, kin recognition might be the result of kin selection when occurring in a heterogeneous environment with various genotypes (Hamilton 1964). Especially, at large-scale, we show that different genotypes facilitate seed set that stabilize the population dynamics by increasing the performance of neighboring plant individuals (Latzel et al. 2013, Castellanos et al. 2014). Thus, selection for diversified genotypes might reveal a coexistence mechanism, because mutations at a short time scale (25 generations) are able to alter the competitive hierarchies of *A. thaliana* individuals of different genotypes. These genotypes might expand or reduce their niches and can therefore act more competitive against strangers (Ehlers et al 2016). Moreover, variation in competitive ability among genotypes due to mutation can lead to intransitive competitive hierarchies at a small-scale, and allow coexistence of competitors at large-scale when there is no single dominant competitor (Vellend and Geber 2005, Taylor and Aarssen

1990, Laure et al. 2017). This might also reflect the genetic trade-off between competitive abilities of MA lines of small-scale competition and of large-scale facilitation (Timan 2004). Low competition between *A. thaliana* individuals at small-scale transmits into low facilitation between different species at larger scale. In turn, strong competition within *A. thaliana* at small-scale results in weaker competition between different species at large-scale. Thus, the constant input of mutation to genetic variation of competitive hierarchies suggests that genetic variation is likely never a limiting factor in the evolution of those traits that influence plant community dynamics.

Implementation of eco-evolutionary trait-based model for population and community ecology

In ecology, two primary contrasting models are crucial to population and community dynamics for understanding the maintenance of biodiversity: the niche and neutral theories of ecology (Fisher and Mehta 2015). The niche theory claims that niches are determined by the difference of abiotic or biotic ecological processes e.g. competition or facilitation (Hutchinson 1957, Bruno et al. 2003, Colwell and Rangel 2009) whereas the neutral theory predicts that ecological processes are neutral and are solely modified by stochasticity (Hubbel 2005). Although it is still under debate which model predicts the dynamic properties of communities and determines the maintenance of biodiversity (Violle et al. 2017), both models miss the importance of evolutionary mechanisms and individual variation influencing population and community dynamics (Violle et al. 2012). The difference of competition and facilitation between or within genotypes demonstrates that evolutionary mechanisms that shape individual variation could

change hierarchies of intra-specific interactions in natural plant populations over a short evolutionary time scale. Our results are concordant with the niche theory stressing the changing of competitive hierarchies among phenotypes in a given environment (Levine and HilleRisLambers 2009, Nottebrock et al. 2017). Moreover, selection and trait evolution favor population survival of *A. thaliana* in the wild, which stresses the importance of combining spatial ecological and evolutionary mechanisms for our understanding of population and community dynamics.

Conclusion

Rapid evolution of *A. thaliana* due to spontaneous mutation alone has profound consequences for population and community dynamics. Competition of *A. thaliana* individuals is genotype-dependent and *A. thaliana* is not a single weak competitor among co-occurring plant species (Ehlers et al. 2016, Soliveres et al 2017). Moreover, the discovery of the underlying eco-evolutionary nature of competition in *A. thaliana* supports a shift from species-based to individual-based community ecology. This would lead to a more predictive ecological theory (Violle et al 2017). We show that including the intraspecific genetic variation and phenotypic variation as separate aspects to a trait-based neighborhood model increases the predictive power to understand population and community dynamics. In particular, non-neutral intraspecific processes may determine species coexistence, because genetic diversity is promoted by having stronger competitive abilities at small-scale and stabilizing population survival at large-scale (Clark et al. 2010). Additionally, our study demonstrates higher genetic diversity increases population survival due to rapid evolution with implications to forecast the fate of species and functional diversity in response to environmental changes (Violle et al 2012). Depending on the time of environmental changes, species may adapt to environmental change by shifting their fundamental niches (Clark 2010). The combination of intraspecific genotypic variation and spatial interactions might advance our understanding of community dynamics, especially of rapid evolution (Koch et al. 2014, Turcotte and Levine 2016). Incorporating genetic variation and the eco-evolutionary process for determining standing levels of genetic variation will provide a better understanding of species interactions underlying the maintenance of biodiversity.

431

432 Keywords (10 for indexing purpose): trait-based community ecology, spontaneous mutations,
 433 competition, facilitation, niche evolution, intraspecific genetic variation, intraspecific
 434 phenotypic variation, *Arabidopsis thaliana*, mutation accumulation lines, eco-evolutionary
 435 dynamics.

436

	Year 2004	Year 2005	Model
Individual	A1 (s,l)	A2 (s,l)	Differential
survival	A1 (s,l)	A2 (s,l)	Neutral
Seed set	B1 (s,l)	B2 (s,l)	Differential
	B1 (s,l)	B2 (s,l)	Neutral

448

449 Table 1. Table of all different eco-evolutionary trait-based neighborhood models including
450 intraspecific variation to test our three objectives of density effects on survival and seed set of
451 *A. thaliana*. Differential models split the neighbor responses into within genotypes and
452 between genotypes. Neutral models describe the overall neighbor response without
453 differentiating between genotypes. All models include random effects of plant biomass on the
454 slope, subblock or block and MA-line on the intercept (see Fig. 1 for design), and respectively
455 depending on small (s) or large-scale (l) (for more details see method section).

Survival model							
(A1) 2004		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		<i>Intra</i>	<i>Inter</i>		<i>Intra</i>	<i>Inter</i>	
Weighted density							
MA	neutral	-0.02 ^{ns}		7551.7	0.004 ^{ns}		7644.5
Lines	differential	-0.02 ^{ns}	-0.37**	7548.6	0.15 ^{ns}	-0.30 ^{ns}	7644.2
(A2) 2005		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		<i>Intra</i>	<i>Inter</i>		<i>Intra</i>	<i>Inter</i>	
Weighted density							
MA	neutral	0.03 ^{ns}		9637.3	0.03 ^{ns}		9872.9
lines	differential	-0.02 ^{ns}	-0.28***	9626.1	0.05 ^{ns}	-0.10 ^{ns}	9873.3
Seed set model							
(B1) 2004		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		<i>Intra</i>	<i>Inter</i>		<i>Intra</i>	<i>Inter</i>	
Weighted density							
MA	neutral	-0.02***		52982	0.03		52944
lines	differential	-0.02**	-0.07***	52948	0.03***	0.15***	52939

(B2) 2005		(s) Small scale (80cm)		AIC	(l) Larger scale (200cm)		AIC
		<i>Intra</i>	<i>Inter</i>		<i>Intra</i>	<i>Inter</i>	
Weighted density							
MA	neutral	-0.008*		49103	-0.008*		49049
lines	differential	-0.009**	-0.09***	49095	-0.01*	0.16***	49026

456

457 Table 2. The coefficient and AIC from eco-evolutionary trait-based neighborhood models of the
458 *Arabidopsis thaliana* field experiment in the years 2004 and 2005. Models include intraspecific
459 genetic variation defined as intra-genotypic (within genotypes) and inter-genotypic (between
460 genotypes) variation. Models are shown at two different spatial scales: weighted subblock
461 density (80cm radius) and weighted block density (200cm radius). All scales include variables of
462 neutral effects (absence of intra- and inter-genotypic variation (total sum of all neighboring
463 genotypes)) and differential effects (presence of intra- and inter-genotypic variation (split of
464 neighboring genotypes in intra- and inter-genotypic genotypes). Thus, neighbor identity is split
465 into intra- and inter-genotypes as differential effects or are combined as neutral effects, which
466 includes all neighbors. P-value levels <0.05*, <0.005**, <0.001*** and >0.05^{ns}.

Figure 2. The barplots show standardized coefficient effect sizes and associated standard errors of eco-evolutionary trait-based neighborhood models at small-scale (a, c) and large-scale (b, d) for all MA-lines for the survival model (a, b) and the seed set model (c, d). Intra-genotypic effects (yellow) and inter-genotypic effects (grey) are presented for 2004 and 2005. Coefficient effect sizes indicate reduction or an increase of seed set depending on biomass density of intra- or inter-genotypic neighbors. In addition, coefficient effect sizes indicate a reduction of survival of intra or inter-genotypic neighbors.

477 Figure 3. The effect range on seed set (a), (b) and survival (c) computed from eco-evolutionary
 478 trait-based neighborhood models shown as odds ratios for year 2004 (for 2005 see
 479 Supplementary Material). (a) Shows the hierarchical competitive order of MA lines and founder
 480 for the seed set model including range effects of each MA line and founder as random
 481 intercepts and the trait proxy biomass as random slopes at small-scale and (b) at large-scale. (c)
 482 Shows the hierarchical selective order of MA lines and founder for the survival model including
 483 range effects of each MA line and founder as random intercepts at small-scale.
 484

Correlation between MA line effect sizes of significant seed set and survival model

Figure 4. Correlation of the simulated conditional mean represented as odds ratio of same genotypes (MA lines) reducing or increasing the effect sizes of the seed set model and of the survival model at small-scale or large-scale for 2004, respectively. For year 2005 see supplementary material. Each point represents different genotypes of *A. thaliana*. Negative values means lower effect sizes on the survival or seed set and positive values means increasing the effect sizes on the survival or seed set. a) Relationship between survival and seed set model. b) Relationship between survival and seed set model. c) Relationship between small-scale survival and large-scale seed set model. Since we are plotting effect sizes, the positive correlation in panel c reflects a trade-off between competition and facilitation (see results).

References

- Adler P., B., HilleRisLambers Janneke & Levine J. M. (2007). A niche for neutrality. *Ecology Letters*, 10, 95–104.
- Agrawal, A.A., Hastings, A.P., Johnson, M.T.J., Maron, J.L. & Salminen, J.-P. (2012). Insect herbivores drive real-time ecological and evolutionary change in plant populations. *Science*, 338, 113–116.
- Antonovics, J. (1976). The Nature of Limits to Natural Selection. *Annals of the Missouri Botanical Garden*, 63, 224–247.
- Bailey JK et al. 2009 From genes to ecosystems: a synthesis of the effects of plant genetic factors across levels of organization. *Phil. Trans. R. Soc. B* **364**, 1607–1616. (doi:10.1098/rstb.2008.0336)
- Baron E., Richirt J., Villoutreix R., Amsellem L., Roux F. & Bennett A. (2015). The genetics of intra- and interspecific competitive response and effect in a local population of an annual plant species. *Functional Ecology*, 29, 1361–1370.
- Bataillon, T. (2003). Shaking the ‘deleterious mutations’ dogma? *Trends in Ecology & Evolution*, 18, 315–317.
- Bates, D., Mächler, M., Bolker, B. & Walker, S. (2014). Fitting Linear Mixed-Effects Models using lme4. *arXiv:1406.5823 [stat]*.
- Bellard C., Bertelsmeier C., Leadley P., Thuiller W. & Courchamp F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, 15, 365–377.

516 Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., *et al.* (2011). Why
517 intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26,
518 183–192.

519 Bossdorf Ol., Prati D., Auge H. & Schmid B. (2004). Reduced competitive ability in an invasive
520 plant. *Ecology Letters*, 7, 346–353.

521 Burnham and Anderson (2002). Model Selection and Multimodel Inference: A Practical
522 Information-Theoretical Approach. 2d ed. New York: Springer-Verlag

523 Bruno, J.F., Stachowicz, J.J. & Bertness, M.D. (2003). Inclusion of facilitation into ecological
524 theory, 7.

525 Cahill, F. J., Kembel, S. W. and Gustafson, D. J. (2005), Differential genetic influences on
526 competitive effect and response in *Arabidopsis thaliana*. *Journal of Ecology*, 93: 958-967.
527 doi:10.1111/j.1365-2745.2005.01013.x

528 Cahill, J.F., Elle, E., Smith, G.R. & Shore, B.H. (2008). Disruption of a belowground mutualism
529 alters interactions between plants and their floral visitors. *Ecology*, 89, 1791–1801.

530 Camara, M. d. and M. Pigliucci. (1999). Mutational contributions to genetic variance-covariance
531 matrices: An experimental approach using induced mutations in *Arabidopsis thaliana*.
532 *Evolution* 53 (6), 1692-1703

533 Caradus, J. & Woodfield, D. *Plant and Soil* (1998) 200: 63.
534 <https://doi.org/10.1023/A:1004296707631>

535 Carr, D. E. and Dudash, M. R. (1996), Inbreeding depression in two species of *Mimulus*
536 (Scrophulariaceae) with contrasting mating systems. *American Journal of Botany*, 83: 586-
537 593. doi:10.1002/j.1537-2197.1996.tb12743.x

538 Castellanos Maria Clara, Donat-Caerols Santiago, González-Martínez Santiago C., Verdú Miguel &
539 Gómez-Aparicio Lorena. (2014). Can facilitation influence the spatial genetics of the
540 beneficiary plant population? *Journal of Ecology*, 102, 1214–1221.

541 Chapin, F. S. III, Autumn, K., & Pugnaire, F. (1993). Evolution of suites of traits in response to
542 environmental stress. *American Naturalist*, 142, S78–S92.

543 Chase, J. M. et al. 2002. The interaction between predation and competition: a review and
544 synthesis. – *Ecol. Lett.* 5: 302–315.

545 Chave, J., Muller-Landau, H.C. & Levin, S.A. (2002). Comparing Classical Community Models:
546 Theoretical Consequences for Patterns of Diversity. *The American Naturalist*, 159, 1–23.

547 Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology*
548 *and Systematics*, 31, 343–366.

549 Clark, J.S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest Trees.
550 *Science*, 327, 1129–1132.

551 Colwell, R.K. & Rangel, T.F. (2009). Hutchinson’s duality: The once and future niche. *PNAS*, 106,
552 19651–19658.

553 Connell, J. (1980). Diversity and the Coevolution of Competitors, or the Ghost of Competition
554 Past. *Oikos*, 35(2), 131-138. doi:10.2307/3544421

555 Damgaard, C. (2004). Inference from plant competition experiments: the effect of spatial
556 covariance. *Oikos*, 107, 225–230.

557 Dudley, S. A., Murphy, G. P., File, A. L. and Robinson, D. (2013), Kin recognition and competition
558 in plants. *Funct Ecol*, 27: 898-906. doi:[10.1111/1365-2435.12121](https://doi.org/10.1111/1365-2435.12121)

559 Ehlers, B.K., Damgaard, C.F. & Laroche, F. (2016). Intraspecific genetic variation and species
560 coexistence in plant communities. *Biology Letters*, 12, 20150853.

561 Exposito-Alonso M, Becker C, Schuenemann VJ, Reiter E, Setzer C, Slovak R, et al. (2018) The rate
562 and potential relevance of new mutations in a colonizing plant lineage. *PLoS Genet* 14(2):
563 e1007155. <https://doi.org/10.1371/journal.pgen.1007155>

564 Falahati-Anbaran, M., Lundemo, S. and Stenøien, H. K. (2014), Seed dispersal in time can
565 counteract the effect of gene flow between natural populations of *Arabidopsis thaliana*. *New*
566 *Phytol*, 202: 1043-1054. doi:[10.1111/nph.12702](https://doi.org/10.1111/nph.12702)

567 Fisher C.K. and Mehta P.(2014). The transition between the niche and neutral regimes in ecology.
568 *Proc Natl Acad Sci USA* 111:13111–13116

569 Fournier-Level A, et al.(2016) Predicting the evolutionary dynamics of seasonal adaptation to
570 novel climates in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 113(20):E2812–E2821.

571 Fridley, J.D. & Grime, J.P. (2010). Community and ecosystem effects of intraspecific genetic
572 diversity in grassland microcosms of varying species diversity. *Ecology*, 91, 2272–2283.

573 Fussmann G. F., Loreau M. & Abrams P. A. (2007). Eco-evolutionary dynamics of communities and
574 ecosystems. *Functional Ecology*, 21, 465–477.

575 Genung, M. A., Schweitzer, J. A., Úbeda, F. , Fitzpatrick, B. M., Pregitzer, C. C., Felker-Quinn, E.
576 and Bailey, J. K. (2011), Genetic variation and community change – selection, evolution, and
577 feedbacks. *Functional Ecology*, 25: 408-419. doi:10.1111/j.1365-2435.2010.01797.x

578 Hamilton, W. D. 1964. The genetical evolution of social behaviour. *I & II. J. Theor. Biol.* 7:1–52.

579 Hart, S. P., Schreiber, S. J., Levine, J. M. and Coulson, T. (2016), How variation between individuals
580 affects species coexistence. *Ecol Lett*, 19: 825-838. doi:[10.1111/ele.12618](https://doi.org/10.1111/ele.12618)

581 Hart, S. P., Freckleton, R. P., Levine, J.M. & Kroon, H. (2018). How to quantify competitive ability.
582 *Journal of Ecology*, 0.

583 Hausmann N. J., Juenger T. E., Sen S., Stowe K. A., Dawson T. E., et al., 2005. Quantitative trait loci
584 affecting δC^{13} and response to differential water availability in *Arabidopsis thaliana*. *Evolution*
585 59: 81–96

586 Hofhuis, H. and Hay, A. (2017), Explosive seed dispersal. *New Phytol*, 216: 339-342.
587 doi:10.1111/nph.14541

588 Holt, R.D. (2009). Bringing the Hutchinsonian niche into the 21st century: Ecological and
589 evolutionary perspectives. *PNAS*, 106, 19659–19665.

590 Hubbell S.P. (2005). Neutral theory in community ecology and the hypothesis of functional
591 equivalence. *Functional Ecology*, 19, 166–172.

592 Hutchinson G.E. (1957). *Concluding remarks. Cold Spring Harbor Symp Quant Biol* 22:415–427.

593 Johnson, M.T.J. Stinchcombe, J.R. An emerging synthesis between community ecology and
594 evolutionary biology *Trends Ecol. Evol.*, 22 (2007), pp. 250-257

595 Johnson, M.T.J., Vellend, M. & Stinchcombe, J.R. (2009). Evolution in plant populations as a driver
596 of ecological changes in arthropod communities. *Philosophical Transactions of the Royal
597 Society of London B* 364:1593-1605.

598 Kerwin et al. (2015) Natural genetic variation in *Arabidopsis thaliana* defense metabolism genes
599 modulates field fitness | eLife Lens. <https://lens.elifesciences.org/05604/>. Last accessed 4
600 May 2018.

601 Koch, H., Frickel, J., Valiadi, M. & Becks, L. (2014). Why rapid, adaptive evolution matters for
602 community dynamics. *Front. Ecol. Evol.*, 2.

603 Kubisch A. *et al.* (2013). Kin competition as a major driving force for invasions. *Am. Nat.* 181: 700–
604 706

605 Lachmuth S., Henrichmann C., Horn J., Pagel J., Schurr F.M. & Rafferty N. (2018). Neighbourhood
606 effects on plant reproduction: An experimental–analytical framework and its application to
607 the invasive *Senecio inaequidens*. *Journal of Ecology*, 106, 761–773.

608 Lankau, R.A. (2011). Rapid Evolutionary Change and the Coexistence of Species. *Annual Review of
609 Ecology, Evolution, and Systematics*, 42, 335–354.

610 Lankau, R.A. & Strauss, S.Y. (2007). Mutual feedbacks maintain both genetic and species diversity
611 in a plant community. *Science*, 317, 1561–1563.

612 Lankau R. A., Wheeler E., Bennett A. E. & Strauss S. Y. (2010). Plant–soil feedbacks contribute to
613 an intransitive competitive network that promotes both genetic and species diversity. *Journal*
614 *of Ecology*, 99, 176–185.

615 Latzel, V., Allan, E., Bortolini Silveira, A., Colot, V., Fischer, M. & Bossdorf, O. (2013). Epigenetic
616 diversity increases the productivity and stability of plant populations. *Nat Commun*, 4, 2875.

617 Laure, G. Zimmermann N. E., Levine J. M., Adler P. r B. & Wootton T (2017). The effects of
618 intransitive competition on coexistence. *Ecology Letters*, 20, 791–800.

619 Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance of
620 species diversity. *Nature*, 461, 254–257.

621 Lion, S. (2018) Theoretical approaches in evolutionary ecology: Environmental feedback as a
622 unifying perspective. *Am. Nat.*191, 21–44.

623 Lynch, M., and B. Walsh. (1998). Genetics and analysis of quantitative traits. Sinauer Associates,
624 Inc., Sunderland, MA.

625 Masclaux, F., Hammond, R. L., Meunier, J. , Gouhier-Darimont, C. , Keller, L. and Reymond, P.
626 (2010), Competitive ability not kinship affects growth of *Arabidopsis thaliana* accessions. *New*
627 *Phytologist*, 185: 322-331. doi:10.1111/j.1469-8137.2009.03057.x

628 Mukai, T. (1964). The Genetic Structure of Natural Populations of *Drosophila Melanogaster*. I.
629 Spontaneous Mutation Rate of Polygenes Controlling Viability. *Genetics*, 50, 1–19.

630 Nei, M. (2013). Mutation-driven evolution. Oxford University Press

631 Nottebrock H., Schmid B., Mayer K., Devaux C., Esler K. J., Böhning-Gaese K., *et al.* (2017a). Sugar
632 landscapes and pollinator-mediated interactions in plant communities. *Ecography*, 40, 1129–
633 1138.

634 Nottebrock H., Schmid B., Treurnicht M., Pagel J., Esler K. J., Böhning-Gaese K., *et al.* (2017b).
635 Coexistence of plant species in a biodiversity hotspot is stabilized by competition but not by
636 seed predation. *Oikos*, 126.

637 Ossowski, S., K. Schneeberger, J. I. Lucas-Lledó, N. Warthmann, R. M. Clark, R. G. Shaw, D. Weigel,
638 and M. Lynch. (2010). The rate and molecular spectrum of spontaneous mutations in
639 *Arabidopsis thaliana*. *Science* 327:92–94.

640 Roles, A. J., and J. K. Conner. (2008). Fitness effects of mutation accumulation in a natural
641 outbred population of wild radish (*Raphanus raphanistrum*): comparison of field and
642 greenhouse environments. *Evolution*, 62: 1066-1075. doi:[10.1111/j.1558-5646.2008.00354.x](https://doi.org/10.1111/j.1558-5646.2008.00354.x)

643 Roles, A. J., M. T. Rutter, C. B. Fenster and J. K. Conner. (2016). Field measurement of genotype by
644 environment interaction for fitness caused by spontaneous mutations in *Arabidopsis thaliana*.
645 *Evolution*, 70: 1039-1050.

646 R Development Core Team (2011). R: A language and environment for statistical computing.
647 Vienna, Austria: R Foundation for Statistical Computing.

648 Rutter, M. T., F. H. Shaw and C. B. Fenster. (2010). Spontaneous mutation parameters for
649 *Arabidopsis thaliana* measured in the wild. *Evolution*, 64: 1825-1835.

650 Rutter, M., A. Roles, J. Conner, R. Shaw, F. Shaw, K. Schneeberger, S. Ossowski, D. Weigel and C.
651 B. Fenster. (2012). Brief Communication: Fitness of *Arabidopsis thaliana* mutation
652 accumulation lines whose spontaneous mutations are known. *Evolution*, 66: 2335-2339.

653 Rutter, M. T., A. J. Roles, and C. B. Fenster. (2018). Quantifying natural seasonal variation in
654 mutation parameters with mutation accumulation lines. *Ecology and Evolution*,
655 <https://doi.org/10.1002/ece3.4085>

656 Schaack, S., Allen, D.E., Latta, L.C., Morgan, K.K. & Lynch, M. (2013). The effect of spontaneous
657 mutations on competitive ability. *Journal of Evolutionary Biology*, 26, 451–456.

658 Shaw, R.G., Byers, D.L. & Darms, E. (2000). Spontaneous Mutational Effects on Reproductive
659 Traits of *Arabidopsis thaliana*. *Genetics* 155:369–378.

660 Shaw, F. H., C. J. Geyer, and R. G. Shaw. (2002). A comprehensive model of mutations affecting
661 fitness and inferences for *Arabidopsis thaliana*. *Evolution* 56:453–463.

662 Shefferson Richard P., Salguero-Gómez Roberto & Rees Mark. (2015). Eco-evolutionary dynamics
663 in plants: interactive processes at overlapping time-scale and their implications. *Journal of*
664 *Ecology*, 103, 789–797.

665 Soliveres, S., Smit, C. & Maestre, F.T. (2015). Moving forward on facilitation research: response to
666 changing environments and effects on the diversity, functioning and evolution of plant
667 communities. *Biol Rev Camb Philos Soc*, 90, 297–313.

668 Soliveres Set al. (2015). Intransitive competition is widespread in plant communities and
669 maintains their species richness. *Ecol. Lett.* 18, 790–798. doi:10.1111/ele.12456

670 Stearns, F. W. and C. B. Fenster. (2016). The effect of induced mutations on quantitative traits in
671 *Arabidopsis thaliana*: Natural versus artificial conditions. *Ecology and Evolution*, doi:
672 10.1002/ece3.2558

673 Strauss, S.Y. & Irwin, R.E. (2004). Ecological and evolutionary consequences of multispecies
674 plant-animal interactions. *Annu. Rev. Ecol. Evol. Syst.*, 35, 435–466.

675 Taylor, D.R., Aarssen, L.W. & Loehle, C. (1990). On the Relationship between r/K Selection and
676 Environmental Carrying Capacity: A New Habitat Templet for Plant Life History Strategies.
677 *Oikos*, 58, 239–250.

678 Tiffin, P. (2002). Competition and time of damage affect the pattern of selection acting on plant
679 defense against herbivores. *Ecology*, 83: 1981-1990. doi:10.1890/0012-
680 9658(2002)083[1981:CATODA]2.0.CO;2

681 Tilman D. (2004). Niche tradeoffs, neutrality, and community structure: a stochastic theory of
682 resource competition, invasion, and community assembly. *Proc. Natl Acad. Sci. USA* 101, 10
683 854–10 861. doi:10.1073/Pnas.0403458101

684 Turcotte, M.M. & Levine, J.M. (2016). Phenotypic Plasticity and Species Coexistence. *Trends Ecol.*
685 *Evol. (Amst.)*, 31, 803–813.

686 Uriarte, M., Swenson, N.G., Chazdon, R.L., Comita, L.S., John Kress, W., Erickson, D., *et al.* (2010).
687 Trait similarity, shared ancestry and the structure of neighbourhood interactions in a
688 subtropical wet forest: implications for community assembly. *Ecol. Lett.*, 13, 1503–1514.

689 Van Dam, N. M. and Baldwin, I. T. (1998), Costs of jasmonate-induced responses in plants
690 competing for limited resources. *Ecology Letters*, 1: 30-33. doi:10.1046/j.1461-
691 0248.1998.00010.x

692 Vellend Mark & Geber Monica A. (2005). Connections between species diversity and genetic
693 diversity. *Ecology Letters*, 8, 767–781.

694 Violle, C., Enquist, B.J., McGill, B.J., Jiang, L., Albert, C.H., Hulshof, C., *et al.* (2012). The return of
695 the variance: intraspecific variability in community ecology. *Trends in Ecology & Evolution*, 27,
696 244–252.

697 Violle, C. & Jiang, L. (2009). Towards a trait-based quantification of species niche. *J Plant Ecol*, 2,
698 87–93.

699 Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N.J.B., Cadotte, M.W., *et al.* (2017).
700 Functional Rarity: The Ecology of Outliers. *Trends in Ecology & Evolution*, 32, 356–367.

701 Weng, M.-L., Becker, C., Hildebrandt, J., Rutter, M.T., Shaw, R.G., Weigel, D. and C. B. Fenster.
702 Fine-Grained Analysis of Spontaneous Mutation Spectrum and Frequency in *Arabidopsis*
703 *thaliana* Genetics. Early online (2018); <https://doi.org/10.1534/genetics.118.301721>

704 Whitham, T. G., Young, W. P., Martinsen, G. D., Gehring, C. A., Schweitzer, J. A., Shuster, S. M.,
705 Wimp, G. M., Fischer, D. G., Bailey, J. K., Lindroth, R. L., Woolbright, S. and Kuske, C. R. (2003),
706 Community and ecosystem genetics: A consequence of extended phenotype. *Ecology*, 84:
707 559-573. doi:10.1890/0012-9658(2003)084[0559:CAEGAC]2.0.CO;2

708 Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy, C.J. *et al.* (2006).

709 A framework for community and ecosystem genetics: from genes to ecosystems. *Nat. Rev.*

710 *Genet.*, 7, 510–523.

711 Whitlock R. & Lortie C. (2014). Relationships between adaptive and neutral genetic diversity and

712 ecological structure and functioning: a meta-analysis. *Journal of Ecology*, 102, 857–872.

713

714

715

716