An interaction map of transcription factors controlling gynoecium development in Arabidopsis

Humberto Herrera-Ubaldo¹, Sergio E. Campos¹, Valentín Luna-García¹, Víctor M. Zúñiga-Mayo^{1#}, Gerardo Armas-Caballero¹, Alexander DeLuna¹, Nayelli Marsch-Martínez², Stefan de Folter^{1*}

1 Unidad de Genómica Avanzada (UGA-LANGEBIO), Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional (CINVESTAV-IPN), Irapuato 36824, Guanajuato, México.

2 Departamento de Biotecnología y Bioquímica, Unidad Irapuato, CINVESTAV-IPN, Irapuato 36824, Guanajuato, México.

Current address: Instituto de Fitosanidad, Colegio de Postgraduados, Campus Montecillo, Texcoco, México.

* Corresponding author: stefan.defolter@cinvestav.mx; phone +52 462 1663011

Running title: Interactome for gynoecium development

Key words: Dynamic interactions, protein interactions, Y2H, interactome, network, transcription factors, gynoecium, fruit

ABSTRACT

Flowers are composed of different organs, whose identity is defined at the molecular by the combinatorial activity of transcription factors (TFs). MADS-box TFs interact forming complexes that have been schematized in the quartet model. The gynoecium is the female reproductive part in the flower, crucial for plant reproduction, and fruit and seed production. Once carpel identity is established, a gynoecium containing many tissues arises. Several TFs have been identified as regulators of gynoecium development, and some of these TFs form complexes. However, broad knowledge about the interactions among these TFs is still scarce. In this work, we used a systems biology approach to understand the formation of a complex reproductive unit as the gynoecium by mapping binary interactions between well-characterized TFs. We analyzed over 3500 combinations and detected more than 200 protein-protein interactions (PPIs), resulting in a process specific interaction map. Topological analyses suggest hidden functions and novel roles for many TFs. Furthermore, a relationship between TFs involved in auxin and cytokinin signaling pathways and other TFs was observed. We analyzed the network by combining PPI data, expression and genetic data, allowing us to dissect it into several dynamic spatio-temporal sub-networks related to gynoecium development subprocesses.

INTRODUCTION

Fruits and seeds are the basis of the food supply. Fruit and flower development in *Arabidopsis* has been studied for more than 20 years (Roeder and Yanofsky, 2006; Alvarez-Buylla et al., 2010). The accepted model for floral organ formation relies on the combinatorial action of mostly MADS-box proteins, resulting in the specification of the four whorls of the flower: sepals, petals, stamens and carpels (Coen and Meyerowitz, 1991). At the molecular level, MADS-box transcription factors (TFs) form complexes in order to regulate gene expression. For the female reproductive organs, the C-class and E-class proteins physically interact to give the identity to the gynoecium (Honma and Goto, 2001; Theissen and Saedler, 2001).

Gynoecium formation starts at stage 6 of flower development, when its identity is established (Smyth et al., 1990). After that, several TFs participate in the subsequent formation of the tissues. These TFs belong to a significant number of TF families such as MADS-box, HOMEODOMAIN, bHLH, Zinc Finger, bZIP, factors mediating hormone transcriptional responses (ARF, ARR, TCP), and others (Roeder and Yanofsky, 2006; Reyes-Olalde et al., 2013).

Current regulatory networks underlying gynoecium patterning include many protein-encoding genes, mostly TFs, acting in a coordinated fashion for the formation of all the tissues (Chavez-Montes et al., 2015; Marsch-Martinez and de Folter, 2016; Zúñiga-Mayo et al., 2019). The involvement of several TFs in the same process generates robustness. At the very early stages, the establishment of two domains, the lateral and the medial, is essential for further gynoecium development. Besides TFs, the hormones auxin and cytokinin are involved (Bowman et al., 1999; Sehra and Franks., 2015; Marsch-Martinez and de Folter, 2016; Muller et al., 2017; Victor-Zúñiga et al., 2019).

Subsequently, the establishment of the carpel margin meristem or CMM is critical (Wynn et al., 2011; Reyes-Olalde et al., 2013). All the medial tissues will be formed from the CMM, including the placenta and ovules, septum, transmitting tract, replum, style and stigma. During these early developmental processes, important TFs are SHOOT MERISTEMLESS (STM), CUP-SHAPED COTYLEDON (CUCs), SPATULA (SPT), ETTIN (ETT), and some B-type ARABIDOPSIS RESPONSE REGULATORS (ARRs) (Sessions et al., 1997; Heisler et al., 2001; Nahar et al., 2012; Kamiuchi et al., 2014; Scofield et al., 2007; Reyes-Olalde et al., 2017).

After that, in the formation of the transmitting tract another set of TFs are involved such as NO TRANSMITTING TRACT (NTT) (Crawford et al., 2007), the HECATE proteins (HEC1, HEC2 and HEC3), with SPT (Gremski et al., 2007), HALF FILLED (HAF), BRASSINOSTEROID ENHANCED EXPRESSION 1 and 3 (BEE1, BEE3), and the AUXIN RESPONSE FACTORS 6 and 8 (ARF6, ARF8) (Crawford and Yanofsky, 2011).

In the medial-lateral patterning of the gynoecium, this is, the formation of valves, valve margins, and replum, a repression-action between lateral and medial factors is necessary, ASYMMETRIC LEAVES 1 and 2 (AS1, AS2), JAGGED (JAG), FILAMENTOUS FLOWER (FIL), YABBY 3 (YAB3), and FRUITFULL (FUL) in the lateral domain; on the other hand, NTT (Marsch-Martinez et al., 2014), BREVIPEDICELLUS (BP), SHOOT MERISTEMLESS (STM) and REPLUMLESS (RPL) (Roeder et al., 2003), promote replum formation in the medial domain (Alonso-Cantabrana et al., 2007; Gonzalez-Reig et al., 2012); while, in between those tissues, SHATTERPROOF 1 and 2 (SHP1, SHP2), ALCATRAZ (ALC), INDEHISCENT (IND), and SPT are involved in the formation of the valve margins at later stages of gynoecium formation (Roeder and Yanofsky, 2006; Girin et al., 2011).

Although there is information about how the above-mentioned TFs perform their activities, most of that information has come from studies done at the individual level. However, virtually all TFs form protein complexes (Vidal et al., 2011; Herrera-Ubaldo et al., 2014; Bemer et al., 2017). For all the TF families listed above, some essential feature of the protein functioning is achieved via PPIs, although, probably we know only the tip of the iceberg. Therefore, generating more PPI data, i.e. direct binary interaction mapping in a high-throughput, standardized manner, would boost our understanding on TF functioning during plant development.

As expected, also for gynoecium and fruit development, information on PPIs is rather scarce (Herrera-Ubaldo et al., 2014). Studies have been reported on the generation of PPI data for some TF families such as MADS-box (de Folter et al., 2005; Immink et al., 2009), Homeobox (Hackbusch et al., 2005), TCP (Danisman et al., 2013), and proteins involved in hormone response like ARF-Aux/IAA (Vernoux et al., 2011) or ARRr (Dortay et al., 2006). In other cases, information comes from the study of a specific tissue or a specific developmental process, e.g. transmitting tract (Gremski et al., 2007) or fruit elongation (Ripoll et al., 2015). Some putative TF complexes have been predicted, such as SEU-LUG-FIL-ANT for the control of the adaxial fate in the gynoecium (Azhakanandam et al., 2008), and ETT-IND-BP-RPL (Simonini et al., 2018) for style development. In summary, current

information about physical interactions important for gynoecium and fruit development is the result of very focused studies.

In this work, we made a systematic effort to boost the information on PPI data that can give us insights into gynoecium and fruit development from the perspective of PPI networks of TFs. The formation of the gynoecium is a complex process, where many actors participate, so the study of this complex process as a system is necessary. We generated a PPI map of TFs that are expressed in the gynoecium and/or fruit and tissues with meristematic activity, and combined this with functional information and the sequence of developmental events that occur to obtain a proper gynoecium and fruit. This is an important step towards a comprehensive gynoecium gene regulatory network, and, furthermore, will be a useful resource for understanding how TFs can function. A deep understanding of molecular mechanisms and the regulatory networks controlling fruit development could be a tool for facing the future problems in crops and food production related to climate change (Zhao et al., 2017).

RESULTS

Here we present a protein-protein interaction network (PPI network) with many of the well-characterized transcription factors (TFs) involved in gynoecium development in *Arabidopsis thaliana* (hereafter *Arabidopsis*). The combination of functional information and the physical interaction data allowed us to extract networks most likely reflecting protein complexes directing the formation of tissues, and with the addition of manually curated expression data we studied the dynamics of interactions related to temporal cues. With these, we dissected the interaction map into 13 sub-networks, each network representing a region and a progressive time during gynoecium development. These sub-networks are likely biologically relevant.

Transcription factors involved in gynoecium development physically interact

The first question we wanted to address was, how the TFs involved in gynoecium development physically interact between them. We selected well-studied TFs (and some co-factors and other TFs) related to this process (most of them are listed in Reyes-Olalde et al., 2013), in total, our study included 72 proteins (gyn set; listed in [Supplemental Table 1](#)). We performed a matrix-based yeast two-hybrid (Y2H) assay with Arabidopsis protein-encoding ORF clones. To date, this is still the technique that has generated most PPI data for many model organisms, in a high-throughput, standardized and reproducible manner (Walhout and Vidal, 2001; Vidal et al., 2011; Braun et al., 2013). Most of the possible combinations were tested, but some of the BD clones were not used due to high autoactivation (results of the autoactivation tests are shown in [Supplemental Table 2](#)). In total, 3648 combinations were tested ([Supplemental Table 2](#)). For the Y2H assay, three interaction reporters were used; two of them allowing yeast growth (-HIS and -ADE), and colonies were confirmed with a third colorimetric reporter (LacZ). The interaction dependent growth was monitored with four yeast crosses (biological replicates), with one technical replicate, giving a total of 8 colonies per marker; one replicate from each marker was used for the LacZ assay, adding four more score points.

In summary, each combination was tested 24 times. Plates were incubated during six days at 22°C. Colony growth was scored on the 6th day after inoculation. In total, yeast growth was detected in 294 combinations. For the construction of the network and further analysis we used the top scored interactions, these are, interactions detected with at least 2 out of the 3 reporters, with at least half of the colonies; this list includes 239 of high confident interactions (all the interactions are listed in [Supplemental Table 3](#)), which were used to generate the network. Almost one-third of the interactions had the highest score (24 points). In some cases, due to autoactivation of the BD clone, just two reporters could be used; in these cases, the highest score was 12 points (70 interactions). After removing duplicated edges, the network has 239 interactions (edges) between 58 proteins (nodes) ([Figure 1](#) and [Supplemental Figure 1](#)).

We detected one major network containing 56 nodes, in addition to the interaction PAN-ARR15 that is not connected to the main network. A basic topological analysis was performed with the major network. Nodes with the highest number of connections (degree) and the value of other topological indexes are listed in [Supplemental Tables 4 and 5](#). Interestingly, PPIs are not restricted to members of the same family of TFs ([Figure 1A](#)), and members of the HOMEBOX and Zinc-finger family could interact with members of all the TF families we tested.

Proteins involved in the same developmental processes are related at the physical interaction level

The PPI map we generated shows a single major component with practically all the interactions, so we wondered if TFs grouped according to their functions (i.e., the formation of a tissue). Since most of the TFs selected for this study have been previously characterized, we could add a tag related to their function or based on the phenotypes observed in the mutants, i.e., determinacy, style formation, etc. ([Supplemental Table 1](#)). With these, we generated networks highlighting proteins participating in 10 different developmental processes ([Figure 2, Supplemental Figure 2](#)). In general, at the level of physical interactions, proteins involved in the same processes do not cluster together in the network ([Figure 2A-C, Supplemental Figure 2](#)), but when those nodes are extracted from the network, we can see sub-networks or so-called modules, probably representing protein complexes ([Figure 2D-F, Supplemental Figure 2](#)).

With the tested interactions, we observe that sub-networks related to carpel number establishment or determinacy are not as complex (number of nodes and interactions) as those related to ovule or abaxial-adaxial patterning. On the other hand, the networks corresponding to valve margins and replum are very similar.

Dynamic interactions during gynoecium development

The sub-networks generated with the separation of interactions according to a functional process is an indicator of the complexity of the processes. It is likely that all the interactions do not occur at the same time. To address the temporal issue, and for a better visualization of gynoecium development through time, we generated a graphic outline of this process ([Figure 3](#)). Its construction is based on the progressive formation of tissues over time (Larsson et al., 2014; Reyes-Olalde et al., 2013), each region delimits a tissue or group of cells that can be defined as different ([Figure 3A](#)). This scheme uses lines as a representation of the different zones within the gynoecium; at the very beginning, the gynoecium primordium is represented as a single line, shortly after is divided in 2 (lateral and medial domains, visible at floral stage 7), and after that many other lines arise (floral stages 8 to 11), as a representation of all the tissues composing a mature gynoecium observed at floral stage 12. With this, we defined more than 10 spatio-temporal contexts (or zones) in which different protein interactions could be taking place during gynoecium development ([Figure 3B](#)).

This scheme helped us to address questions such as where and when the identified interactions are likely to occur during gynoecium development. We took advantage of all the previous studies performed for each gene; most of them at the individual level, a large proportion of this data was previously collected (Reyes-Olalde et al., 2013). In this study we updated the information, added data for other genes, manually curated gene expression data and represented it according to our division of tissues (listed in [Supplemental Table 1](#)).

This graphical representation is useful for displaying individual gene expression patterns during gynoecium development ([Figure 3C](#)). The overlapping of expression patterns with proteins that physically interact is an approach to study interaction profiles of two proteins, revealing where and when that dimer may be performing its functions ([Figure 3D](#)). In some cases, near perfect matches in expression pattern coincidence are observed, such as AG-SEP3 and SEU-LUG; in other cases, as WUS-ETT or CRC-NGA3 partial matches in

expression patterns suggest that those interactions could occur in specific regions and stages of the gynoecium, and interactions change if different contexts.

The use of expression data allowed to dissect the PPI network in a spatio-temporal manner; nodes representing TFs were placed in each of the contexts we defined, so we can have a PPI network for each context (Figure 4, Supplemental Figure 3), these networks represent the subset of interactions that could be occurring in those tissues at specific times. The networks are different in node and edge number; the differences in complexity between them can be easily observed (Table 1). A large fraction of the complete network is observed at the beginning of gynoecium formation. After that, the lateral domain recruits a small part and most of the nodes and interactions are restricted to the medial domain. This network remains very similar for the carpel margin meristem (CMM), ovules and septum; and near floral stage 10 is separated into 4 networks, representing style, stigma, transmitting tract and septum. The networks for valves and replum are similar, they start as small networks, and some nodes and interactions appear over time.

TF interactions and hormonal pathways

Transcriptional responses to hormones (i.e., auxin and cytokinin) are mediated by a set of response regulators: Auxin Response Factors (ARFs) and Arabidopsis Response Regulators (ARR, A and B-types) (Schaller et al., 2015). Some of these proteins were included in our interaction test. We detected interactions between the hormone response regulators and TFs from various families (Figure 5A). Interestingly, five TFs, WUS, HEC2, CRC, AS1, and BP can physically interact with regulators of both hormonal signaling pathways, suggesting that these nodes may serve as links to connect hormonal pathways.

Another interesting phenomenon in the context of protein interactions and hormones is a recent report showing that the interaction-dependent yeast growth could be modified due to auxin addition to the culture medium (Simonini et al., 2016). In order to test if this phenomenon is occurring also with our set of interactions, we used the colonies recovered (representing a dimer or an interaction) from the -ADE experiments (171 interactions, listed

in [Supplemental Table 3](#)). We used the -ADE set because the entire set of interactions could be tested using the same medium (test with -HIS would involve the addition of different concentrations of 3-AT). First, we tested the effect of two hormones on the wild type PJ69 yeast strain. The addition of cytokinin (BAP: 6-Benzylaminopurine) did not affect the growth of the yeast, even at the maximum tested concentration (250 μ M) ([Figure 5B](#)). On the other hand, the effect of the addition of auxin (IAA: Indole-3-Acetic Acid) on the yeast growth is observed in the dilution series; the highest concentration of inoculum seems not to change yeast growth at 20, 50 or 100 μ M IAA, but is affected when 250 μ M IAA was tested ([Supplemental Figure 5](#)). For the Y2H test, we inoculated the colonies in hormone-containing media at different concentrations, and incubated them at 22°C during 6 days. In our conditions, no significant changes in yeast growth were detected due to cytokinin addition ([Figure 5C](#)), or auxin addition ([Supplemental Figure 5](#)).

The extended PPI network and novel actors of gynoecium development

The PPI network we generated includes well-characterized TFs controlling gynoecium development (72 protein-encoding genes we selected for this study, gyn set, [Figure 6A](#)), so it can be considered as a process specific interaction map. Our map contains 239 interactions between 58 proteins ([Figure 6B](#), listed in [Supplemental Table 6](#)). For some proteins we did not detect interactions in our system, which can be due to various reasons (e.g., missing TFs in our gyn set, weak interaction strength, formation of higher-order complex needed, etc.)

So, to partially overcome this and identify novel TF interactors, we generated an extended PPI network ([Figure 6C](#)). The extended PPI network includes reported interactions (IntAct database; Orchard et al., 2014) between the gyn set of proteins (this study) and other TFs, most of the times uncharacterized proteins and obtained in studies not focused on gynoecium development. In this extended network, we included only interactions detected by Y2H assays and furthermore, this list was filtered to include just TF-encoding genes expressed (at least) in carpels ([Supplemental Table 6](#); [purple colored selection](#), [Supplemental Figure 6](#)). This extended network can be considered as a comprehensive

interaction network of TFs involved in gynoecium development (724 interactions, [Figure 6C](#)). All the interactions in the extended network are listed in the [Supplemental Table 6](#). Because our gyn set consists out of well-studied TFs, in the extended network, for every dimer in the list at least one partner has been characterized, meaning we can predict a role for the other TF. In conclusion, this paves the way for focused functional studies of over 200 TFs present in the extended network as candidates for having a role in the regulation of gynoecium development. We present a list of 26 TFs (of the extended PPI network) that have the highest number of interactions (> 4) with the TFs of the gyn set (see [Supplemental Table 6](#)). In general, these TFs share various interactors, and, notably, in the top 10 of TFs with most interactions, there are 9 TFs of the TCP family.

An Aux/IAA TF affects reproductive development

One of the proteins found to interact with known gynoecium TFs in the extended PPI network is IAA27, an Aux/IAA protein not reported as regulator of Arabidopsis gynoecium development, but with potential roles in this process based on the interactions found ([Figure 6C](#), [Supplemental Table 5](#)). We obtained the AD and BD clones and included it in the Y2H assay, where reported interactions were confirmed, and novel interactions were detected ([Figure 1](#) and [Supplemental Figure 5](#)). Subsequently, to study the potential function of IAA27, wild type Arabidopsis plants were transformed with an antisense construct to knock-down the expression of this gene (*anti-IAA27*). As presumed, different phenotypes were observed during the reproductive phase of *anti-IAA27* plants ([Supplemental Figure 5](#)). Expression analysis showed that the level of *IAA27* silencing was correlated with the severity of the observed phenotypes. Independent lines with a severe phenotype were characterized in more detail. The inflorescences of *anti-IAA27* plants produced fewer floral buds compared with wild type inflorescences. Furthermore, the floral organs showed a yellowish aspect, the anthers were shorter with a reduced number of mature pollen grains, and the short fruits produced on average only two seeds.

Different analyses were made to determine why *anti-IAA27* plants developed short fruits. Most of the ovules analyzed showed defects in their morphology, mainly an arrest in

embryo sac development. Transverse sections showed that anthers of *anti-IAA27* plants produced fewer pollen grains compared with wild type anthers. Finally, the *LAT52::GUS* reporter line was used to visualize pollen tube growth. Gynoecia from wild type and *anti-IAA27* plants were hand-pollinated with *LAT52::GUS* pollen. Pollen tubes grew efficiently through wild type gynoecia; however, pollen tubes grew only in the style but not in the ovary of *anti-IAA27* gynoecia ([Supplemental Figure 5](#)). These results indicate that both male and female reproductive development is affected in *anti-IAA27* plants. Consistent with this, reproductive development is also affected when the function of its interactor ARF8 is abolished (Nagpal et al., 2005; Wu et al., 2006). Furthermore, it was recently shown that silencing of the homologue of *IAA27* (*SlIAA27*) in tomato also affects male and female reproduction (Bassa et al., 2012).

DISCUSSION

In the past years, gynoecium development has been studied in a gene-by-gene fashion. Current knowledge for gynoecium development includes dozens of genes and their genetic interactions, though information on protein-protein interactions (PPI) is available but still rather scarce. With the PPI map generated in this study, we want to add a solid basis for the molecular understanding and the generation of a more complete gynoecium GRN. Recently, we reported some steps towards a comprehensive and dynamic GRN for gynoecium formation (Chavez Montes et al., 2015). Furthermore, all the data presented here will be a useful resource for people studying reproductive development.

A network for gynoecium development

With the interactions among TFs reported here, we have generated the most complete PPI map for gynoecium development in *Arabidopsis* to date. Sub-networks or modules were considered as basic units, or fundamental units guiding morphogenesis, as seen in biological networks (Alon, 2003; Zhu et al., 2007). We identified 239 interactions using the gyn set ([Figure 1](#)), placing our PPI network at the position of various other PPI networks in

plants (Braun et al., 2013). In addition, as discussed below, we generated an extended PPI network for gynoecium development containing 724 interactions ([Figure 6](#)).

The yeast 2-hybrid (Y2H) system is one of the most used techniques for the identification of binary protein interactions, with the advantage of being high-throughput, highly standardized, and reproducible, especially when using ORF collections (Walhout and Vidal, 2001; Vidal et al., 2011; Braun et al., 2013). The most complete interaction network in Arabidopsis to date, has been built using the Y2H system (AI-2, Arabidopsis Interactome Map Consortium, 2012), with 5664 interactions between 2661 proteins. Several other networks for Arabidopsis have been constructed with this type of data e.g., ARF-Aux/IAA (Vernoux et al., 2011), TCS (Dortay et al., 2006 & 2008), TOPLESS proteins (Causier et al., 2012), protein-G complex (Klopfleisch et al., 2011), ESCRT proteins (Richardson et al., 2011), cell cycle protein (Boruc et al., 2010), TCP family (Danisman et al., 2013), and MADS-box proteins (de Folter et al., 2005), among others.

The construction of a PPI map guiding gynoecium and fruit development and the GRNs underlying the process is a starting point for the study of the formation of the tissues, comparison of fruit development between species, the study of network rewiring allowing the formation of novel structures or processes in a global way. It can be used for the study of the evolutionary history of this important reproductive unit.

Emerging properties for well-studied transcription factors

The network we generated, has two components, with the exception of the dimer PAN-ARR15, all the nodes form a major network component ([Figure 1](#)). This is an indicator of the close relationship between all the proteins and the processes involved in gynoecium development. Topological analysis is useful to get insights in network architecture and performance (Zhu et al, 2007). Analysis of the main component indicates that it is a compact network (network diameter =5; radius =1; characteristic path length =2.26) ([Supplemental Table 5](#)). We compared our network to other plant PPI networks, and its density is similar to these other networks (density =0.13) ([Supplemental Table 5](#)).

Furthermore, most of the other analyzed topological characteristics are also in the range of the reported biological plant PPI networks, supporting our findings. One interesting aspect is that the nodes are very different in connections (network heterogeneity =0.91), in the case of TFs involved in gynoecium development this heterogeneity is observed in the effect of mutations on the genes, since not all cause the same defect.

Some topological indexes (see [Supplemental Table 4](#)), revealed hidden features for well-known TFs exposing interesting properties/activities for them, which will be an exciting topic for *in planta* and functional analysis. For instance, the high value in node degree (i.e. many interactions) for WUS, BP, STM, YAB1, HEC2 and HEC3, many involved in meristematic activity and all of them involved in early events of gynoecium development (Reyes-Olalde et al., 2013), and the network generated for this early stage ([Figure 4, Table 1](#)), is a sign of how complex this early event of gynoecium formation is.

Furthermore, the node TCP15 has also a high degree; this property is maybe the cause of the severe defects that occur in gynoecium development when *TCP15* is ectopically expressed (Lucero et al., 2015; Kieffer et al., 2011). Another similar example is ARR14 (B-type ARR) (Kieber and Schaller, 2014), a node with many interactions, suggesting essential roles in female reproductive development, though functional and expression data are still scarce, but the PPI data suggests a central role in gynoecium development.

The nodes WUS, TCP15, BP, and ARR14 have high values for stress and betweenness centrality ([Supplemental Table 4](#)), indicating a possible role as a mediator of interactions and maybe for the formation of high-order complexes, interestingly, SEPALLATA3 (SEP3) has similar values. This crucial protein for flower development has been described as a bridge for MADS-box protein interactions (Immink et al., 2009).

The values for clustering coefficient highlight proteins such as SPT, BEE1, and HEC1, as members of a highly connected module, probably reflecting their involvement in transmitting tract formation as a protein complex (Heisler et al., 2001; Gremski et al., 2007;

Crawford et al., 2011). Other proteins with high values of clustering coefficient are AS2 and NGA3, suggesting that probably they can form higher-order complexes.

Neighborhood connectivity is an attractive property of nodes, meaning having few interactors but those interactors function as a hub (having many interactors); some of these proteins are ETT, AINTEGUMENTA (ANT), STYLISH 2 (STY2), and ARR4. In the case of gynoecium development, this differential interaction (heterogeneity) is maybe the cause for the formation of one tissue. The critical interaction could be the difference in the case of very similar networks, for instance, those from the medial region and the tissues it produces.

Biologically relevant protein interactions

The occurrence of PPIs among TFs was tested in a heterologous expression system, namely yeast. As mentioned before, the Y2H system has many advantages and is an attractive technique for binary interaction mapping, which will continue (Vidal and Fields, 2014). As with these studies, next steps will be necessary to explore the functional relevance of detected interactions. The identification of physical interactions between proteins is just a starting point for functional studies and for understanding important developmental and cellular processes. Of course, starting with more focused functional analysis, various complementary techniques are available for PPI confirmation and to obtain further insight in e.g. protein complex formation, which are future perspectives of this work (Xing et al., 2016; Bontinck et al., 2018; Lampugnani et al., 2018).

Nevertheless, based on coexpression of the protein-encoding genes, we consider that most of the interactions could be biologically relevant because for 87% of the dimers, the genes are coexpressed (190 out of 219; based on the sum of the sub-networks in [Supplemental Figure 3](#)). This number is still an underestimation when we would consider e.g. more specific cell-type expression data, protein stability, and protein movement. Furthermore, another kind of evidence supporting the physical interactions detected here, is the interaction of proteins participating in the same biological process.

Extended Network, a directed search of novel players in gynoecium development

The PPI network we generated in this work includes most of the well-characterized TFs involved in gynoecium development, though, it is highly likely that more TFs are involved. A rough estimation sets the expression of more than 500 TFs during carpel formation (de Folter et al., 2004; Chavez Montes et al., 2015). So, it is a challenge to characterize all of them. We made a first step to identify more TFs by exploring PPIs reported for our gyn set of TFs (IntAct database; Orchard et al., 2014). Many non-characterized proteins interact with the TFs of our network, however, we included only those TFs expressed in flower and fruit, which resulted in the extended PPI network with 724 interactions between 284 nodes (Figure 6). This approach proved to be valid since we validated at least one of them, the Aux/IAA IAA27 protein (Supplemental Figure 5). We found similar functions as one of its interactors ARF8, which is involved in reproductive development (Nagpal et al., 2005; Wu et al., 2006). Furthermore, an example of another biologically relevant PPI present in the extended network is between NO TRANSMITTING TRACT (NTT) and SEEDSTICK (STK), which we recently reported as important for medial domain development and plant reproductive competence (Herrera-Ubaldo et al., 2018).

A combinatorial model for gynoecium development?

The quartet model is an elegant representation of the molecular mechanism underlying floral organ specification (Theissen and Saedler, 2001). One of our goals here was the proposal of a combinatorial model for gynoecium formation, an extension of the ABC model. In this sense, testing binary interactions between TF guiding gynoecium development is the first step towards a model explaining gynoecium patterning.

However, there are some crucial differences between the specification of the different floral organs in a flower versus the specification and formation of the different tissues in the gynoecium (i.e. after carpel specification), which, in general, is more complex: 1) The implication of several TF families (flower: MADS and AP2; and more than 16 TF families for gynoecium); 2) The number of TFs involved in the process (Flower <10; gynoecium

>70), this is related to the redundancy observed between many of them; 3) The effect of mutations of the genes, for the flower, mutations in the homeotic genes produce dramatic changes in the architecture of the flower, while in gynoecium formation (i.e. after carpel specification), the mutations (often) generate less dramatic changes. 4) The time of the developmental processes, floral organ specification takes around four days; on the other hand, most of the tissues of the gynoecium (floral stages 6 to 12) are formed during more than ten days.

Considering those four main differences, a possible `model` would involve a succession of networks, controlled in time. For flower formation, spatial organization is a key feature, observed with the spatially restricted interactions, while for gynoecium formation, both time and spatial organization are important during development.

Inserting hormones into the networks

Hormones play a significant role during gynoecium development, particularly auxin and cytokinin (Marsch-Martinez and de Folter, 2016; Zuñiga-Mayo et al., 2019). Given the importance of these two hormones, we included various proteins involved in the transcriptional response into the network (ARFs and ARR). There are other TFs that do not participate directly in the primary response to hormones, but their relationship with hormones has been demonstrated, e.g. SPT (Schuster et al., 2015; Reyes-Olalde et al., 2017), TCP15 (Lucero et al., 2015), HECs (Gaillochet et al., 2018), FUL (Ripoll et al., 2015), STM (Jasinski et al., 2005; Yanai et al., 2005), CRC (Yamaguchi et al., 2017), and AG (Ó'Maoiléidigh et al., 2018), among others.

Here, we found interactions between ARFs, ARRs, and TFs of different families. Especially, we detected five TFs (WUS, HEC2, CRC, AS1, and BP) that can interact with both auxin and cytokinin related TFs. These TFs are expressed in meristems and boundaries, so they may have a role as decision-makers, in the control of proliferation and differentiation. Other studies have observed similar cases (e.g. Oh et al., 2014; Yazaki et al., 2016; Gaillochet et al., 2018). Our detected PPIs give an opportunity to explain molecularly

how these TFs can do so. It would be interesting to detect the dynamics of protein complex formation during different stages of gynoecium development, to know whether the identified interactions occur simultaneously or change in time.

Another interesting phenomenon involving PPIs and hormones is the ability of a hormone to modify the physical interaction of proteins, which has been observed recently for auxin with ETT (an ARF member) (Simonini et al., 2016), as well as affecting the transcriptional output (Simonini et al., 2017). Since we recovered all the colonies that grew in our Y2H experiments, we had the entire dimer collection in yeast colonies. So, we tested whether the gynoecium development PPI network could be modulated by the addition of auxin or cytokinin (IAA or BAP, respectively), however, in our conditions we did not detect alterations in PPI formation based on changes in yeast growth. Still this is an interesting topic to further explore, as well as directly in plants.

The fruit is in the net

In this work, we started the study of gynoecium development from the systems biology perspective, the generation of a PPI map with many of the major players involved in this process, which gave us a vision of how complex the interplay could be between the TFs involved in the formation of tissues.

The next step, besides continuing with functional studies *in planta*, is to study the dynamical aspects (i.e. in spatio-temporal manner) of the interactions during gynoecium development. Moreover, TFs that could function as hubs in the network also deserve attention and should be further studied in this regard. Intriguing are the observed PPIs of TFs that interact with auxin and cytokinin related signaling proteins, and it will be very interesting to unravel the molecular meaning of this. Furthermore, we started studying the effects on PPIs by temporal cues and hormones. On the other hand, the protein complexes we proposed here, illustrated as sub-networks or modules need to be studied with complementary techniques such as bimolecular fluorescence complementation (BiFC) in different tissues *in planta* (Smaczniak et al., 2012; Long et al., 2017), co-

immunoprecipitation (co-IP), protein arrays (Yazaki et al., 2016), and/or immuno or affinity purification followed by mass spectrometry analysis (IP-MS/AF-MS) (Smaczniak et al., 2012; Gadeyne et al., 2014; Huang et al., 2016).

We hope the information we presented here will serve as a resource and will be useful for other people studying plant reproductive development. Furthermore, since many TFs also participate in other developmental programs, some of these interactions may be conserved in other contexts, where they can be further explored.

MATERIALS AND METHODS

Clones

Information about the collection and generation of the *Arabidopsis* 72 ORF clones (the gyn set) is provided in [Supplemental Table 2](#); many come from the REGIA collection (Paz-Ares and The REGIA Consortium, 2002). The gyn set is collection of mainly TFs that are expressed in the gynoecium and/or fruit and tissues with meristematic activity. Many of the well-characterized TFs involved in gynoecium development in *Arabidopsis thaliana* are present. Furthermore, a few cofactors are present as well.

Yeast 2-hybrid assay

A matrix-based interaction assay was conducted as previously reported (de Folter and Immink, 2011). The GAL4 system (Invitrogen) was used; the coding sequences of the selected genes in entry clones were recombined with pDEST22 for GAL4-AD fusions or with pDEST32 for BD-fusions. Yeast transformation was performed with the PEG/LiAc method using the PJ69 yeast strain (James et al., 1996). All the tested combinations were generated by mating (see [Supplemental Table 2](#)), four AD colonies per construct were arranged in a single-well omnitray plate in a 384-colonies array. BD clones were grown individually in single-well plates. Mating was performed using a floating pin replicator (V&P scientific) in plates with YPAD medium. For diploid selection, colonies were transferred to plates with SD-TRP-LEU medium, the latter step was done twice. For the

protein interaction test, diploid colonies were transferred to SD-LEU-TRP-ADE and to SD-LEU-TRP-HIS with different concentrations of 3-Amino triazole (see [Supplemental Table 2](#)), each selection marker was tested in duplicate. Plates were incubated at 22°C. Yeast growth was scored 6 days after inoculation. For the LacZ assay the RO-BLUE medium was used (de Folter and Immink, 2011) for these tests, one plate per protein was tested. Yeast colonies from both markers were recovered and cryo-conserved for future analysis (i.e., hormone dependent interactions).

Hormone-dependent interaction assays

PJ69 yeast strain (James et al., 1996) and the colonies identified and conserved from the Y2H assay (-ADE set, [Supplemental Table 3](#)) were recovered and inoculated in medium SD-GLUC-TRP-LEU-ADE with increasing concentrations of cytokinin (6-Benzylaminopurine, BAP) or auxin (Indole-3-Acetic Acid, IAA). NaOH was used to help to dissolve the hormones to prepare stock solutions; 100 µM NaOH was tested as negative control for both, auxin and cytokinin experiments.

Expression data

Data about gene expression for the selected protein-encoding genes were collected from literature; most of this information is an updated version of a previous report (Reyes-Olalde et al., 2013). For floral stage 6, RNA-seq data was collected from a previous report (Jiao and Meyerowitz, 2010), an expression value of >1.0 RPKM was considered. Separation in different regions during development was done according to previous works (Reyes-Olalde et al., 2013; Larsson et al., 2014). Functional processes were assigned according to the literature. All the information about expression and phenotypes is available in [Supplemental Table 1](#).

Interaction data collection and extended network construction

Protein interaction data was downloaded from IntAct (last accessed on 31.10.2018) (Orchard et al., 2014), the locus ID list of the 72 genes in the gyn set was used as query ([Supplemental Table 1](#)). Three filters were applied to the interaction list, 1) Interactions detected in Y2H assays (MI:2277, Cr-two hybrid; MI: 0726, Reverse two hybrid; MI:0018,

two hybrid; MI:0397, two hybrid array; MI: 1112, two hybrid pooling approach; MI: 1356, validated two hybrid) were considered. 2) Interactions between transcription factors; a list was downloaded from the Plant Transcription Factors Database V4.0 (Jin et al., 2017) (last accessed 08.11.2018). 3) Expression in carpel; RNA-seq expression data was collected from ARAPORT11 (Krishnakumar et al., 2015) (last accessed 08.11.2018), genes with values of >0.5 TPM in carpel or stage 12 inflorescence samples were considered. Information about expression, GO classification and protein domains in the 221 TFs in the extended network ([Supplemental Table 6](#) and [Supplemental Figure 6](#)) was collected from ARAPORT11 (Krishnakumar et al., 2015). The networks shown in Figure 6 are available in a Cytoscape network file (see [Supplemental Figure 7](#)).

Network construction and topological analysis

Networks were generated using Cytoscape V3.6 (Shannon et al., 2003). Network analyses were performed with the Cytoscape built-in application NetworkAnalyzer (release 2.7; Assenov et al., 2008).

Generation of IAA27 knock-down lines and phenotypical characterization

A 293 bp fragment upstream of the stop codon of the *IAA27* gene (At4g29080) was PCR-amplified from cDNA using the primers 1KDL-*IAA27* and 2KDL-*IAA27* (see [Supplemental Table 7](#)), and subcloned into pENTR/D-TOPO (Invitrogen, Carlsbad, CA, USA). The pENTR/D-TOPO clone was sequence verified and recombined with the binary vector pHELLSGATE8 (Helliwell et al., 2002), resulting in the antisense construct *anti-IAA27*. The *anti-IAA27* lines were classified according to severity of fruit size phenotypes: strong if the fruit reached up to 50% of its normal size, medium if the fruit reached between 50 and 75% of its normal size, and weak if the fruit growth between 76-90% of its normal size.

For RT-PCR, plant tissue was collected for total RNA isolation according to the method described by Verwoerd et al. (1989). Total RNA was treated with DNaseI (Invitrogen) and reverse-transcribed using M-MLV (Invitrogen). Semiquantitative expression analysis for *IAA27* was performed using the primers 1RT-*IAA27* and 2RT-*IAA27* (see [Supplemental Table 7](#)). For pollination analysis, *LAT52::GUS* was used as pollen tube marker (Johnson et

al., 2004). Gynoecia from *anti-IAA27* plants were emasculated 24 h before anthesis. The next day, anti-IAA27 gynoecia were hand-pollinated with pollen from *LAT52::GUS* marker line. 24 h after pollination, the gynoecia were collected and incubated 7 h at 37°C with a 5-bromo-4-chloro-3-indolyl-b-glucuronic acid solution (Gold Biotechnology, St Louis, MO, USA; Jefferson et al., 1987), placed in 90% ethanol for 1 h, followed by 70% ethanol for 1 h, and overnight in Hoyer solution (Anderson, 1954). Cross-section and GUS images were taken using a LEICA CTR6000 (Wetzlar, Germany) microscope in Differential Interference Contrast (DIC) mode.

AUTHOR CONTRIBUTIONS

HHU and SdF conceived and designed the study; HHU, SEC, VLG, VMZM, and GAC performed the research; HHU and SdF analyzed the data; HHU performed visualization and prepared the figures; HHU, AD, NMM, and SdF discussed the research; HHU and SdF wrote the paper.

ACKNOWLEDGMENTS

We would like to thank Jeanneth Pablo-Villa for initial assistance in the yeast 2-hybrid experiments in the lab. Furthermore, we thank the following people for clones: Cristina Ferrándiz, Alexander Heyl, Froukje van der Wal and Richard Immink, Eva Zanchetti and Lucia Colombo, Mitsuhiro Aida, Juan Carlos Ochoa-Sánchez, Irepan Reyes-Olalde, and the Arabidopsis Biological Resource Center (ABRC). We thank Karla González-Aguilera for technical assistance in the lab. HHU and VLG were supported by the Mexican National Council of Science and Technology (CONACyT) with a PhD fellowship (243380 and 487657, respectively). Work in the SdF laboratory was financed by the CONACyT grants CB-2012-177739 and FC-2015-2/1061, and NMM by the CONACyT grant CB-2015-255069. SdF acknowledges support of the Marcos Moshinsky Foundation and the European Union H2020-MSCA-RISE-2015 project ExpoSEED (grant no. 691109). No conflict of interest is declared.

REFERENCES

- Alon, U. (2003). Biological networks: the tinkerer as an engineer. *Science* 301:1866-1867.
- Alonso-Cantabrana, H., Ripoll, J.J., Ochando, I., Vera, A., Ferrandiz, C., and Martinez-Laborda, A. (2007). Common regulatory networks in leaf and fruit patterning revealed by mutations in the Arabidopsis ASYMMETRIC LEAVES1 gene. *Development* 134:2663-2671.
- Alvarez-Buylla, E.R., Benitez, M., Corvera-Poire, A., Chaos Cador, A., de Folter, S., Gamboa de Buen, A., Garay-Arroyo, A., Garcia-Ponce, B., Jaimes-Miranda, F., Perez-Ruiz, R.V., et al. (2010). Flower development. *The arabidopsis book* 8:e0127.
- Anderson, L.E. (1954). Hoyer's Solution as a Rapid Permanent Mounting Medium for Bryophytes. *The Bryologist* 57:242-244.
- Arabidopsis Interactome Mapping Consortium. (2011). Evidence for Network Evolution in an Arabidopsis Interactome Map. *Science* 333:601.
- Assenov, Y., Ramirez, F., Schelhorn, S.E., Lengauer, T., and Albrecht, M. (2008). Computing topological parameters of biological networks. *Bioinformatics* 24:282-284.
- Azhakanandam, S., Nole-Wilson, S., Bao, F., and Franks, R.G. (2008). SEUSS and AINTEGUMENTA mediate patterning and ovule initiation during gynoecium medial domain development. *Plant physiology* 146:1165-1181.
- Bassa, C., Mila, I., Bouzayen, M., and Audran-Delalande, C. (2012). Phenotypes associated with down-regulation of SI-IAA27 support functional diversity among Aux/IAA family members in tomato. *Plant & cell physiology* 53:1583-1595.
- Bemer, M., van Dijk, A.D.J., Immink, R.G.H., and Angenent, G.C. (2017). Cross-Family Transcription Factor Interactions: An Additional Layer of Gene Regulation. *Trends in plant science* 22:66-80.
- Bontinck, M., Van Leene, J., Gadeyne, A., De Rybel, B., Eeckhout, D., Nelissen, H., and De Jaeger, G. (2018). Recent Trends in Plant Protein Complex Analysis in a Developmental Context. *Frontiers in plant science* 9:640.
- Boruc, J., Van den Daele, H., Hollunder, J., Rombauts, S., Mylle, E., Hilson, P., Inze, D., De Veylder, L., and Russinova, E. (2010). Functional modules in the Arabidopsis core cell cycle binary protein-protein interaction network. *The Plant cell* 22:1264-1280.
- Bowman, J.L., Baum, S.F., Eshed, Y., Putterill, J., and Alvarez, J. (1999). Molecular genetics of gynoecium development in Arabidopsis. *Current topics in developmental biology* 45:155-205.
- Braun, P., Aubourg, S., Van Leene, J., De Jaeger, G., and Lurin, C. (2013). Plant protein interactomes. In: *Annual Review of Plant Biology*. 161-187.
- Causier, B., Ashworth, M., Guo, W., and Davies, B. (2012). The TOPLESS interactome: a framework for gene repression in Arabidopsis. *Plant physiology* 158:423-438.
- Chávez Montes, R.A., Herrera-Ubaldo, H., Serwatowska, J., and de Folter, S. (2015). Towards a comprehensive and dynamic gynoecium gene regulatory network. *Current Plant Biology* 3-4:3-12.

- Coen, E.S., and Meyerowitz, E.M. (1991). The war of the whorls: genetic interactions controlling flower development. *Nature* 353:31-37.
- Crawford, B.C.W., Ditta, G., and Yanofsky, M.F. (2007). The NTT Gene Is Required for Transmitting-Tract Development in Carpels of *Arabidopsis thaliana*. *Current Biology* 17:1101-1108.
- Crawford, B.C.W., and Yanofsky, M.F. (2011). HALF FILLED promotes reproductive tract development and fertilization efficiency in *Arabidopsis thaliana*. *Development (Cambridge, England)* 138:2999-3009.
- Danisman, S., van Dijk, A.D., Bimbo, A., van der Wal, F., Hennig, L., de Folter, S., Angenent, G.C., and Immink, R.G. (2013). Analysis of functional redundancies within the *Arabidopsis* TCP transcription factor family. *Journal of experimental botany* 64:5673-5685.
- de Folter, S., Busscher, J., Colombo, L., Losa, A., and Angenent, G.C. (2004). Transcript profiling of transcription factor genes during silique development in *Arabidopsis*. *Plant molecular biology* 56:351-366.
- de Folter, S., and Immink, R.G. (2011). Yeast protein-protein interaction assays and screens. *Methods Mol Biol* 754:145-165.
- de Folter, S., Immink, R.G., Kieffer, M., Parenicova, L., Henz, S.R., Weigel, D., Busscher, M., Kooiker, M., Colombo, L., Kater, M.M., et al. (2005). Comprehensive interaction map of the *Arabidopsis* MADS Box transcription factors. *The Plant cell* 17:1424-1433.
- Dortay, H., Gruhn, N., Pfeifer, A., Schwerdtner, M., Schmülling, T., and Heyl, A. (2008). Toward an interaction map of the two-component signaling pathway of *Arabidopsis thaliana*. *Journal of Proteome Research* 7:3649-3660.
- Dortay, H., Mehnert, N., Burkle, L., Schmulling, T., and Heyl, A. (2006). Analysis of protein interactions within the cytokinin-signaling pathway of *Arabidopsis thaliana*. *The FEBS journal* 273:4631-4644.
- Gadeyne, A., Sanchez-Rodriguez, C., Vanneste, S., Di Rubbo, S., Zauber, H., Vanneste, K., Van Leene, J., De Winne, N., Eeckhout, D., Persiau, G., et al. (2014). The TPLATE adaptor complex drives clathrin-mediated endocytosis in plants. *Cell* 156:691-704.
- Gaillochet, C., Jamge, S., van der Wal, F., Angenent, G., Immink, R., and Lohmann, J.U. (2018). A molecular network for functional versatility of HECATE transcription factors. *The Plant journal : for cell and molecular biology*.
- Girin, T., Paicu, T., Stephenson, P., Fuentes, S., Korner, E., O'Brien, M., Sorefan, K., Wood, T.A., Balanza, V., Ferrandiz, C., et al. (2011). INDEHISCENT and SPATULA interact to specify carpel and valve margin tissue and thus promote seed dispersal in *Arabidopsis*. *The Plant cell* 23:3641-3653.
- Gonzalez-Reig, S., Ripoll, J.J., Vera, A., Yanofsky, M.F., and Martinez-Laborda, A. (2012). Antagonistic gene activities determine the formation of pattern elements along the mediolateral axis of the *Arabidopsis* fruit. *PLoS genetics* 8:e1003020.
- Gremski, K., Ditta, G., and Yanofsky, M.F. (2007). The HECATE genes regulate female reproductive tract development in *Arabidopsis thaliana*. *Development* 134:3593-3601.
- Hackbusch, J., Richter, K., Muller, J., Salamini, F., and Uhrig, J.F. (2005). A central role of *Arabidopsis thaliana* ovate family proteins in networking and subcellular localization of 3-aa loop extension

- homeodomain proteins. *Proceedings of the National Academy of Sciences of the United States of America* 102:4908-4912.
- Heisler, M.G., Atkinson, A., Bylstra, Y.H., Walsh, R., and Smyth, D.R. (2001). SPATULA, a gene that controls development of carpel margin tissues in Arabidopsis, encodes a bHLH protein. *Development* 128:1089-1098.
- Helliwell, C.A., Wesley, S.V., Wielopolska, A.J., and Waterhouse, P.M. (2002). High-throughput vectors for efficient gene silencing in plants. *Functional Plant Biology* 29:1217-1225.
- Herrera-Ubaldo, H., Lozano-Sotomayor, P., Ezquer, I., Di Marzo, M., Chavez Montes, R.A., Gómez-Felipe, A., Pablo-Villa, J., Diaz-Ramirez, D., Ballester, P., Ferrandiz, C., et al. (2018) New roles of NO TRANSMITTING TRACT and SEEDSTICK during medial domain development in Arabidopsis fruits. *Development*: doi: 10.1242/dev.172395.
- Herrera-Ubaldo, H., Zanchetti, E., Colombo, L., and de Folter, S. (2014). Protein interactions guiding carpel and fruit development in Arabidopsis. *Plant Biosystems* 148:169-175.
- Honma, T., and Goto, K. (2001). Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* 409:525-529.
- Huang, H., Alvarez, S., Bindbeutel, R., Shen, Z., Naldrett, M.J., Evans, B.S., Briggs, S.P., Hicks, L.M., Kay, S.A., and Nusinow, D.A. (2016). Identification of Evening Complex Associated Proteins in Arabidopsis by Affinity Purification and Mass Spectrometry. *Molecular & cellular proteomics : MCP* 15:201-217.
- Immink, R.G., Tonaco, I.A., de Folter, S., Shchennikova, A., van Dijk, A.D., Busscher-Lange, J., Borst, J.W., and Angenent, G.C. (2009). SEPALLATA3: the 'glue' for MADS box transcription factor complex formation. *Genome biology* 10:R24.
- James, P., Halladay, J., and Craig, E.A. (1996). Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast. *Genetics* 144:1425-1436.
- Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., Phillips, A., Hedden, P., and Tsiantis, M. (2005). KNOX action in Arabidopsis is mediated by coordinate regulation of cytokinin and gibberellin activities. *Current biology : CB* 15:1560-1565.
- Jefferson, R.A., Kavanagh, T.A., and Bevan, M.W. (1987). GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *The EMBO journal* 6:3901-3907.
- Jiao, Y., and Meyerowitz, E.M. (2010). Cell-type specific analysis of translating RNAs in developing flowers reveals new levels of control. *Molecular systems biology* 6:419.
- Jin, J., Tian, F., Yang, D.C., Meng, Y.Q., Kong, L., Luo, J., and Gao, G. (2017). PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic acids research* 45:D1040-D1045.
- Johnson, M.A., von Besser, K., Zhou, Q., Smith, E., Aux, G., Patton, D., Levin, J.Z., and Preuss, D. (2004). Arabidopsis hapless mutations define essential gametophytic functions. *Genetics* 168:971-982.

- Kamiuchi, Y., Yamamoto, K., Furutani, M., Tasaka, M., and Aida, M. (2014). The CUC1 and CUC2 genes promote carpel margin meristem formation during Arabidopsis gynoecium development. *Frontiers in plant science* 5:165.
- Kieber, J.J., and Schaller, G.E. (2014). Cytokinins. *The Arabidopsis book* / American Society of Plant Biologists 12:e0168.
- Kieffer, M., Master, V., Waites, R., and Davies, B. (2011). TCP14 and TCP15 affect internode length and leaf shape in Arabidopsis. *The Plant journal : for cell and molecular biology* 68:147-158.
- Klopfleisch, K., Phan, N., Augustin, K., Bayne, R.S., Booker, K.S., Botella, J.R., Carpita, N.C., Carr, T., Chen, J.G., Cooke, T.R., et al. (2011). Arabidopsis G-protein interactome reveals connections to cell wall carbohydrates and morphogenesis. *Molecular systems biology* 7:532.
- Krishnakumar, V., Hanlon, M.R., Contrino, S., Ferlanti, E.S., Karamycheva, S., Kim, M., Rosen, B.D., Cheng, C.Y., Moreira, W., Mock, S.A., et al. (2015). Araport: the Arabidopsis information portal. *Nucleic acids research* 43:D1003-1009.
- Lampugnani, E.R., Wink, R.H., Persson, S., and Somssich, M. (2018). The Toolbox to Study Protein-Protein Interactions in Plants. *Critical Reviews in Plant Sciences*:1-27.
- Larsson, E., Roberts, C.J., Claes, A.R., Franks, R.G., and Sundberg, E. (2014). Polar auxin transport is essential for medial versus lateral tissue specification and vascular-mediated valve outgrowth in Arabidopsis gynoecia. *Plant physiology* 166:1998-2012.
- Long, Y., Stahl, Y., Weidtkamp-Peters, S., Postma, M., Zhou, W., Goedhart, J., Sanchez-Perez, M.I., Gadella, T.W.J., Simon, R., Scheres, B., et al. (2017). In vivo FRET-FLIM reveals cell-type-specific protein interactions in Arabidopsis roots. *Nature* 548:97-102.
- Lucero, L.E., Uberti-Manassero, N.G., Arce, A.L., Colombatti, F., Alemano, S.G., and Gonzalez, D.H. (2015). TCP15 modulates cytokinin and auxin responses during gynoecium development in Arabidopsis. *The Plant Journal* 84:267-282.
- Marsch-Martinez, N., and de Folter, S. (2016). Hormonal control of the development of the gynoecium. *Current opinion in plant biology* 29:104-114.
- Marsch-Martinez, N., Zuniga-Mayo, V.M., Herrera-Ubaldo, H., Ouwerkerk, P.B.F., Pablo-Villa, J., Lozano-Sotomayor, P., Greco, R., Ballester, P., Balanza, V., Kuijt, S.J.H., et al. (2014). The NTT transcription factor promotes replum development in Arabidopsis fruits. *The Plant Journal* 80:69-81.
- Muller, C.J., Larsson, E., Spichal, L., and Sundberg, E. (2017). Cytokinin-Auxin Crosstalk in the Gynoecial Primordium Ensures Correct Domain Patterning. *Plant physiology* 175:1144-1157.
- Nagpal, P., Ellis, C.M., Weber, H., Ploense, S.E., Barkawi, L.S., Guilfoyle, T.J., Hagen, G., Alonso, J.M., Cohen, J.D., Farmer, E.E., et al. (2005). Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. *Development* 132:4107-4118.
- Nahar, M.A., Ishida, T., Smyth, D.R., Tasaka, M., and Aida, M. (2012). Interactions of CUP-SHAPED COTYLEDON and SPATULA genes control carpel margin development in Arabidopsis thaliana. *Plant & cell physiology* 53:1134-1143.

- O'Maoileidigh, D.S., Stewart, D., Zheng, B., Coupland, G., and Wellmer, F. (2018). Floral homeotic proteins modulate the genetic program for leaf development to suppress trichome formation in flowers. *Development* 145.
- Oh, E., Zhu, J.Y., Bai, M.Y., Arenhart, R.A., Sun, Y., and Wang, Z.Y. (2014). Cell elongation is regulated through a central circuit of interacting transcription factors in the Arabidopsis hypocotyl. *eLife* 3.
- Orchard, S., Ammari, M., Aranda, B., Breuza, L., Briganti, L., Broackes-Carter, F., Campbell, N.H., Chavali, G., Chen, C., Del-Toro, N., et al. (2014). The MIntAct project - IntAct as a common curation platform for 11 molecular interaction databases. *Nucleic Acids Research* 42:D358-D363.
- Paz-Ares, J., and The REGIA Consortium. (2002). REGIA, an EU project on functional genomics of transcription factors from Arabidopsis thaliana. *Comparative and functional genomics* 3:102-108.
- Reyes-Olalde, J.I., Zuniga-Mayo, V.M., Chavez Montes, R.A., Marsch-Martinez, N., and de Folter, S. (2013). Inside the gynoecium: at the carpel margin. *Trends in plant science* 18:644-655.
- Reyes-Olalde, J.I., Zuniga-Mayo, V.M., Serwatowska, J., Chavez Montes, R.A., Lozano-Sotomayor, P., Herrera-Ubaldo, H., Gonzalez-Aguilera, K.L., Ballester, P., Ripoll, J.J., Ezquer, I., et al. (2017). The bHLH transcription factor SPATULA enables cytokinin signaling, and both activate auxin biosynthesis and transport genes at the medial domain of the gynoecium. *PLoS genetics* 13:e1006726.
- Richardson, L.G., Howard, A.S., Khuu, N., Gidda, S.K., McCartney, A., Morphy, B.J., and Mullen, R.T. (2011). Protein-Protein Interaction Network and Subcellular Localization of the Arabidopsis Thaliana ESCRT Machinery. *Frontiers in plant science* 2:20.
- Ripoll, J.J., Bailey, L.J., Mai, Q.A., Wu, S.L., Hon, C.T., Chapman, E.J., Ditta, G.S., Estelle, M., and Yanofsky, M.F. (2015). microRNA regulation of fruit growth. *Nature plants* 1:15036.
- Roeder, A.H.K., Ferrandiz, C., and Yanofsky, M.F. (2003). The role of the REPLUMLESS homeodomain protein in patterning the Arabidopsis fruit. *Current biology* 13:1630-1635.
- Roeder, A.H.K., and Yanofsky, M.F. (2006). Fruit development in Arabidopsis. In: *The arabidopsis book--* Somerville, C.R., and Meyerowitz, E.M., eds. Rockville, MD: American Society of Plant Biologists. 1-50.
- Schaller, G.E., Bishopp, A., and Kieber, J.J. (2015). The yin-yang of hormones: cytokinin and auxin interactions in plant development. *The Plant cell* 27:44-63.
- Schuster, C., Gaillochet, C., and Lohmann, J.U. (2015). Arabidopsis HECATE genes function in phytohormone control during gynoecium development. *Development* 142:3343-3350.
- Scofield, S., Dewitte, W., and Murray, J.A. (2007). The KNOX gene SHOOT MERISTEMLESS is required for the development of reproductive meristematic tissues in Arabidopsis. *The Plant journal : for cell and molecular biology* 50:767-781.
- Sehra, B., and Franks, R.G. (2015). Auxin and cytokinin act during gynoecial patterning and the development of ovules from the meristematic medial domain. *Wiley interdisciplinary reviews. Developmental biology* 4:555-571.

- Sessions, A., Nemhauser, J.L., McColl, A., Roe, J.L., Feldmann, K.A., and Zambryski, P.C. (1997). ETTIN patterns the Arabidopsis floral meristem and reproductive organs. *Development* 124:4481-4491.
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* 13:2498-2504.
- Simonini, S., Bencivenga, S., Trick, M., and Ostergaard, L. (2017). Auxin-Induced Modulation of ETTIN Activity Orchestrates Gene Expression in Arabidopsis. *The Plant cell* 29:1864-1882.
- Simonini, S., Deb, J., Moubayidin, L., Stephenson, P., Valluru, M., Freire-Rios, A., Sorefan, K., Weijers, D., Friml, J., and Ostergaard, L. (2016). A noncanonical auxin-sensing mechanism is required for organ morphogenesis in Arabidopsis. *Genes & development* 30:2286-2296.
- Simonini, S., Stephenson, P., and Ostergaard, L. (2018). A molecular framework controlling style morphology in Brassicaceae. *Development* 145.
- Smaczniak, C., Immink, R.G., Muino, J.M., Blanvillain, R., Busscher, M., Busscher-Lange, J., Dinh, Q.D., Liu, S., Westphal, A.H., Boeren, S., et al. (2012). Characterization of MADS-domain transcription factor complexes in Arabidopsis flower development. *Proceedings of the National Academy of Sciences of the United States of America* 109:1560-1565.
- Smyth, D.R., Bowman, J.L., and Meyerowitz, E.M. (1990). Early flower development in Arabidopsis. *The Plant cell* 2:755-767.
- Theissen, G., and Saedler, H. (2001). Plant biology. Floral quartets. *Nature* 409:469-471.
- Vernoux, T., Brunoud, G., Farcot, E., Morin, V., Van den Daele, H., Legrand, J., Oliva, M., Das, P., Larrieu, A., Wells, D., et al. (2011). The auxin signalling network translates dynamic input into robust patterning at the shoot apex. *Molecular systems biology* 7:508.
- Verwoerd, T.C., Dekker, B.M., and Hoekema, A. (1989). A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic acids research* 17:2362.
- Vidal, M., Cusick, M.E., and Barabasi, A.L. (2011). Interactome networks and human disease. *Cell* 144:986-998.
- Vidal, M., and Fields, S. (2014). The yeast two-hybrid assay: still finding connections after 25 years. *Nature methods* 11:1203-1206.
- Walhout, A.J.M., and Vidal, M. (2001). High-Throughput Yeast Two-Hybrid Assays for Large-Scale Protein Interaction Mapping. *Methods* 24:297-306.
- Wu, M.F., Tian, Q., and Reed, J.W. (2006). Arabidopsis microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development* 133:4211-4218.
- Wynn, A.N., Rueschhoff, E.E., and Franks, R.G. (2011). Transcriptomic characterization of a synergistic genetic interaction during carpel margin meristem development in Arabidopsis thaliana. *PloS one* 6:e26231.
- Xing, S., Wallmeroth, N., Berendzen, K.W., and Grefen, C. (2016). Techniques for the Analysis of Protein-Protein Interactions in Vivo. *Plant physiology* 171:727-758.

- Yamaguchi, N., Huang, J., Xu, Y., Tanoi, K., and Ito, T. (2017). Fine-tuning of auxin homeostasis governs the transition from floral stem cell maintenance to gynoecium formation. *Nature communications* 8:1125.
- Yanai, O., Shani, E., Dolezal, K., Tarkowski, P., Sablowski, R., Sandberg, G., Samach, A., and Ori, N. (2005). Arabidopsis KNOXI proteins activate cytokinin biosynthesis. *Current biology : CB* 15:1566-1571.
- Yazaki, J., Galli, M., Kim, A.Y., Nito, K., Aleman, F., Chang, K.N., Carvunis, A.R., Quan, R., Nguyen, H., Song, L., et al. (2016). Mapping transcription factor interactome networks using HaloTag protein arrays. *Proceedings of the National Academy of Sciences of the United States of America* 113:E4238-4247.
- Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D.B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., et al. (2017). Temperature increase reduces global yields of major crops in four independent estimates. *Proceedings of the National Academy of Sciences of the United States of America* 114:9326-9331.
- Zhu, X., Gerstein, M., and Snyder, M. (2007). Getting connected: analysis and principles of biological networks. *Genes & development* 21:1010-1024.
- Zúñiga-Mayo, V.M., Gómez-Felipe, A., Herrera-Ubaldo, H., and de Folter, S. (2019). Gynoecium development: networks in Arabidopsis and beyond. *Journal of experimental botany*:accepted.

FIGURE LEGENDS

Figure 1. Transcription factors (TFs) involved in gynoecium development interact physically. (A) Matrix representing the interactions detected in the Y2H assay, proteins are grouped in families. (B) Protein-protein (PPI) network constructed with the information in (A).

Figure 2. Proteins involved in the same process interact physically. (A-C) Representation of processes comprising gynoecium development within the interaction network, the proteins involved are colored in yellow (D-F).

Figure 3. Gynoecium development over time. (A) Stages of gynoecium development, false colors represent different tissues within the gynoecium. (B) Schematic representation of gynoecium development, lines represent regions depicted in (A); scale bar represents 1 day. (C) Expression patterns of *FRUITFULL* and *SPATULA*, lines represent those regions indicated in (B). (D) Interaction patterns of the dimers AG-SEP3, SEU-LUG, CRC-NGA3 and WUS-ETT, colored lines represent the expression pattern of the protein-encoding gene in the same color.

Figure 4. Succession of protein interaction networks during gynoecium development. The combination of gene expression data and physical protein-protein interaction (PPI) information allowed the dissection of the network into 13 sub-networks underlying the formation of tissues or developmental sub-processes.

Figure 5. The effect of cytokinin on protein interactions. (A) Sub-network including transcription factors involved in auxin and cytokinin signaling pathways and their interactions with other proteins. (B) The addition of BAP to the culture medium does not affect the growth of the PJ69 yeast strain, neither when diluted yeast concentrations are inoculated (indicated with the triangle; dilution series 1x, 10x, 100x, 1000x, and 10000x). (C) The addition of BAP to the culture medium does not affect the interaction-dependent

growth of yeast; some of the combinations shown in (A) are presented on medium containing 0 μ M, 50 μ M, or 100 μ M BAP.

Figure 6. The extended network of transcription factors controlling gynoecium development in Arabidopsis. (A) Number of proteins used in this study, the gyn set. (B) PPI data identified for the gyn set (this work). (C) The extended network, the PPI data of this work combined with PPI data from the IntAct database (based on Y2H for our gyn set nodes and only of protein-encoding genes with expression in carpels).

Table 01. Topological comparison of the 13 sub-networks

SUPPLEMENTAL INFORMATION

Supplemental Table 1. List of genes selected for this study. Gene expression information and phenotype information. Related to Figures 1-4.

Supplemental Table 2. Yeast 2-hybrid experiment. List of clones, results of the autoactivation test, and tested combinations. Related to Figure 1.

Supplemental Table 3. Results of the Yeast 2-hybrid experiment. List of all the interactions detected, list of top-scored interactions used for the network construction, and list of interactions tested in the hormone assay. Related to Figure 1 and 5.

Supplemental Table 4. Topological analysis of the interaction map. Related to Figure 1.

Supplemental Table 5. Topological comparison of plant interactomes. Related to Figure 1 and Table 1.

Supplemental Table 6. Extended network. List of interactions in the extended network, list of proteins with >4 interactions, list of the 221 TF candidates for gynoecium development, and its classification in GO and protein domains. Related to Figure 6.

Supplemental Table 7. Primer list.

Supplemental Figure 1. High-resolution protein interaction network of transcription factors controlling gynoecium development in Arabidopsis. Related to Figure 1.

Supplemental Figure 2. Networks related to phenotypes. Related to Figure 2. A) Nodes representing genes associated with a phenotype are selected in each network. B) Extraction of the selected nodes.

Supplemental Figure 3. Networks for the 13 contexts (or zones) during gynoecium development. Related to Figures 3-4.

Supplemental Figure 4. The effect of auxin on protein interactions. Related to Figure 5. A) The addition of IAA to the culture medium does not affect the growth of the PJ69 yeast strain when in high concentrations inoculated. However, a growth effect is observed when dilutions of yeast are inoculated (indicated with the triangle; dilution series 1x, 10x, 100x, 1000x, and 10000x). B) The addition of IAA to the culture medium does not affect the interaction-dependent growth of yeast (high yeast concentration used); some of the tested combinations are presented on medium containing 50 μ M and 100 μ M IAA.

Supplemental Figure 5. The gene *IAA27* is involved in reproductive development. Related to Figure 6. (A) Overview of the Y2H assay, the plate for BD-*IAA27* is shown. (B) Interactions detected for the BD-*IAA27* clone based on the 3 interaction markers. (C) Expression of *IAA27* during reproductive development in Arabidopsis. (D) Expression of *IAA27* is reduced at different levels in independent *anti-IAA27* lines. The lines were classified according to the severity of affected fruit size (strong phenotype if the fruit reached up to 35% of its normal size; medium if the fruit reached between 35 and 75% of

its normal size, and weak if the fruit reached between 76-90% of its normal size). Flower (E, F) and fruit (G-H) phenotypes of wildtype (WT) plants (E,G,I) and *anti-IAA27* lines (F,H,J). Pollen tube growth visualization in WT (K) and *anti-IAA27* plants (L) after pollination with *LAT52::GUS* pollen. Transverse section of WT (M) and *anti-IAA27* (N) anthers showing that *anti-IAA27* anthers produced fewer pollen grains (pollen degeneration observed) compared with WT anthers. WT ovules (O) compared with *anti-IAA27* (P) ovules showing morphology defects in *anti-IAA27* ovules (affected embryo sac development).

Supplemental Figure 6. Expression data for the 221 TF candidates of the extended PPI network. Related to Figure 6.

Supplemental Figure 7. Cytoscape data file for the extended network. Related to Figure 6.

Supplemental References.

Table 01. Topological comparison of the 13 sub-networks

Zone		Nodes	Edges	Clustering coefficient	Diameter	Radius	Network centralization	Shortest paths	Char. path length	Avg. number of neighbors	Density	Heterogeneity	Isolated nodes	Self-loops
1	Gynoecium primordium	46	162	0.34	5	1	0.40	1894	2.226	6.61	0.147	0.867	0	10
2	Lateral domain	10	4	0.00	2	1	0.19	8	1.25	0.60	0.067	1.106	5	1
3	Medial domain	30	43	0.21	7	4	0.24	552	2.953	2.47	0.085	0.986	6	6
4	Valves	14	17	0.09	3	2	0.35	90	1.844	2.14	0.165	0.786	4	2
5	Young replum	6	6	0.44	3	2	0.30	20	1.5	2.00	0.4	0.577	1	0
6	Carpel margin meristem	33	58	0.19	7	4	0.23	756	2.86	3.09	0.097	0.947	5	7
7	Replum	11	22	0.50	3	2	0.39	72	1.444	3.82	0.382	0.589	2	1
8	Medial region	25	48	0.35	5	3	0.48	462	2.247	3.36	0.14	0.959	3	6
9	Ovules	32	72	0.18	5	3	0.38	812	2.296	4.06	0.131	0.911	3	7
10	Stigma	8	10	0.44	2	1	0.43	20	1.3	1.75	0.25	0.892	3	3
11	Style	12	12	0.18	3	2	0.36	72	2.028	1.67	0.152	0.894	3	2
12	Septum	13	20	0.39	3	2	0.36	56	1.571	2.31	0.192	0.952	5	5
13	Transmitting tract	6	10	0.67	1	1	0.30	12	1	2.00	0.4	0.707	2	4

Figure 1. Transcription factors (TFs) involved in gynoecium development interact physically. (A) Matrix representing the interactions detected in the Y2H assay, proteins are grouped in families. **(B)** Protein-protein (PPI) network constructed with the information in (A).

Figure 2. Proteins involved in the same process interact physically. (A-C) Representation of processes comprising gynoecium development within the interaction network, the proteins involved are colored in yellow (D-F).

Figure 3. Gynoecium development over time. (A) Stages of gynoecium development, false colors represent different tissues within the gynoecium. (B) Schematic representation of gynoecium development, lines represent regions depicted in (A); scale bar represents 1 day. (C) Expression patterns of *FRUITFULL* and *SPATULA*, lines represent those regions indicated in (B). (D) Interaction patterns of the dimers AG-SEP3, SEU-LUG, CRC-NGA3 and WUS-ETT, colored lines represent the expression pattern of the protein-encoding gene in the same color.

Figure 4. Succession of protein interaction networks during gynoecium development. The combination of gene expression data and physical protein-protein interaction (PPI) information allowed the dissection of the network into 13 sub-networks underlying the formation of tissues or developmental sub-processes.

Figure 5. The effect of cytokinin on protein interactions. (A) Sub-network including transcription factors involved in auxin and cytokinin signaling pathways and their interactions with other proteins. (B) The addition of BAP to the culture medium does not affect the growth of the PJ69 yeast strain, neither when diluted yeast concentrations are inoculated (indicated with the triangle; dilution series 1x, 10x, 100x, 1000x, and 10000x). (C) The addition of BAP to the culture medium does not affect the interaction-dependent growth of yeast; some of the combinations shown in (A) are presented on medium containing 0 μ M, 50 μ M, or 100 μ M BAP.

Figure 6. The extended network of transcription factors controlling gynoecium development in Arabidopsis. (A) Number of proteins used in this study, the gyn set. (B) PPI data identified for the gyn set (this work). (C) The extended network, the PPI data of this work combined with PPI data from the IntAct database (based on Y2H for our gyn set nodes and only of protein-encoding genes with expression in carpels).

Supplemental Figure 1. High-resolution protein interaction network of transcription factors controlling gynoecium development in Arabidopsis. Related to Figure 1.

Supplemental Figure 2. Networks related to phenotypes. Related to Figure 5.

A) Nodes representing genes associated with a phenotype are selected in each network. B) Extraction of the selected nodes.

A

B

Zone 03: Medial domain

Zone 04: Valves

Zone 05: Young replum

Zone 06: Carpel Margin Meristem (CMM)

Zone 07: Replum

Zone 08: Medial region

Zone 09: Ovules

Zone 10: Stigma

Zone 11: Style

Zone 12: Septum

Zone 13: Transmitting tract

Supplemental Figure 4. The effect of auxin on protein interactions. Related to Figure 5. A) The addition of IAA to the culture medium does not affect the growth of the PJ69 yeast strain when in high concentrations inoculated. However, a growth effect is observed when dilutions of yeast are inoculated (indicated with the triangle; dilution series 1x, 10x, 100x, 1000x, and 10000x). B) The addition of IAA to the culture medium does not affect the interaction-dependent growth of the yeast (high yeast concentration used); some of the tested combinations are presented on medium containing 50 μ M and 100 μ M IAA.

Supplemental Figure 5. *IAA27* is involved in reproductive development. Related to Figure 6. (A) Overview of the Y2H assay, the plate for BD-IAA27 is shown. (B) Interactions detected for the BD-IAA27 clone based on the 3 interaction markers. (C) Expression of *IAA27* during reproductive development in Arabidopsis. (D) Expression of *IAA27* is reduced at different levels in independent *anti*-*IAA27* lines. The lines were classified according to the severity of affected fruit size (strong phenotype if the fruit reached up to 35% of its normal size; medium if the fruit reached between 35 and 75% of its normal size, and weak if the fruit reached between 76-90% of its normal size). Flower (E, F) and fruit (G-H) phenotypes of wildtype (WT) plants (E,G,I) and *anti*-*IAA27* lines (F,H,J). Pollen tube growth visualization in WT (K) and *anti*-*IAA27* plants (L) after pollination with *LAT52::GUS* pollen. Transverse section of WT (M) and *anti*-*IAA27* (N) anthers showing that *anti*-*IAA27* anthers produced fewer pollen grains (pollen degeneration observed) compared with WT anthers. WT ovules (O) compared with *anti*-*IAA27* (P) ovules showing morphology defects in *anti*-*IAA27* ovules (affected embryo sac development).

Supplemental Figure 6. Expression data for the 221 TF candidates of the extended PPI network. Related to Figure 6.

Supplemental Table 5. Topological comparison between plant protein interactomes

	A1 (a)	AUX signalling (b)	Cell cycle (c)	MADS (d)	Membrane (e)	Pathogens (f)	G-complex (g)	Gynoecium dev (h)
Clustering coefficient	0.0478	0.5574	0.1161	0.2228	0.1009	0.0239	0.0590	0.3122
Connected components	124	1	2	1	2	25	1	2
Network diameter	16	4	5	6	10	11	6	5
Network radius	1	2	1	3	1	1	4	1
Network centralization	0.0820	0.3682	0.2511	0.3604	0.1140	0.1084	0.3268	0.4016
Shortest paths	5767578	2256	1266	5550	35158	752650	187056	2972
Characteristic path length	4.749	1.726	2.657	2.637	4.226	4.786	3.326	2.259
Avg. number of neighbors	4.156	17.417	3.949	7.040	3.684	2.914	2.480	7.298
Number of nodes	2661	48	39	75	190	926	433	57
Density	0.0016	0.3706	0.1039	0.0951	0.0195	0.0031	0.0057	0.1303
Heterogeneity	2.336	0.578	0.683	0.867	1.170	2.307	3.676	0.906
Isolated nodes	27	0	0	0	0	0	0	0
Self loops	135	15	0	5	9	9	5	11
Multi edge node pairs	0	0	0	3	3	0	136	20

REFERENCES

- (a) Arabidopsis Interactome Mapping Consortium. (2011). Evidence for Network Evolution in an Arabidopsis Interactome Map. *Science* 333:601.
- (b) Vernoux, T., Brunoud, G., Farcot, E., Morin, V., Van den Daele, H., Legrand, J., Oliva, M., Das, P., Larrieu, A., Wells, D., et al. (2011). The auxin signalling network translates dynamic input into robust patterning at the shoot apex. *Mol Syst Biol* 7:508.
- (c) Boruc, J., Van den Daele, H., Hollunder, J., Rombauts, S., Mylle, E., Hilson, P., Inze, D., De Veylder, L., and Russinova, E. (2010). Functional modules in the Arabidopsis core cell cycle binary protein-protein interaction network. *The Plant cell* 22:1264-1280.
- (d) de Folter, S., Immink, R.G., Kieffer, M., Parenicova, L., Henz, S.R., Weigel, D., Busscher, M., Kooiker, M., Colombo, L., Kater, M.M., et al. (2005). Comprehensive interaction map of the Arabidopsis MADS Box transcription factors. *The Plant cell* 17:1424-1433.
- (e) Lalonde, S., Sero, A., Pratelli, R., Pilot, G., Chen, J., Sardi, M.I., Parsa, S.A., Kim, D.Y., Acharya, B.R., Stein, E.V., et al. (2010). A membrane protein/signaling protein interaction network for Arabidopsis version AMPv2. *Frontiers in physiology* 1:24.
- (f) Mukhtar, M.S., Carvunis, A.R., Dreze, M., Eppele, P., Steinbrenner, J., Moore, J., Tasan, M., Galli, M., Hao, T., Nishimura, M.T., et al. (2011). Independently evolved virulence effectors converge onto hubs in a plant immune system network. *Science* 333:596-601.
- (g) Klopffleisch, K., Phan, N., Augustin, K., Bayne, R.S., Booker, K.S., Botella, J.R., Carpita, N.C., Carr, T., Chen, J.G., Cooke, T.R., et al. (2011). Arabidopsis G-protein interactome reveals connections to cell wall carbohydrates and morphogenesis. *Mol Syst Biol* 7:532.
- (h) This work

Supplemental Table 7. Primer list

Used for	Primer name	Sequence	Gene
CDS amplification, entry clone construction	SDF 369	CACCATGCCGATCAACGATCAGTTTC	AT2G01760 (ARR14)
	SDF 370	CTATCTTTGTCTTGAAGATCTTTCC	
	SDF 575	CACCATGGCGAGTGTTGAAGGTGATG	AT3G61830 (ARF18)
	SDF 576	TCACCCCCTACTACGATTTTCGAA	
	SDF 642	CACCATGGCAAATTTTCGAGAATCTTT	AT1G18400 (BEE1)
	SDF 643	TTAAAGGGACCATGTTGATAAATGGA	
	SDF 596	CACCATGGCTGGTCTCGATCTAGGCAC	AT2G35270 (GIK)
	SDF 597	TTAAAACGGAGCCCTACCGGCGCC	
	SDF 644	CACCATGAATAATTATAATATGAACC	AT5G09750 (HEC3)
	SDF 645	TTAGATTAATTCTCCTACTCCTCTTC	
	SDF 592	CACCATGGAGCTGTTTCCTGCTCAGC	AT1G32240 (KAN2)
	SDF 593	TTAGTGAGATCGACCCAGAGTAAACTCA	
	SDF 850	CACCATGAACACAAGAGGAAATTAATC	AT5G60140 (REM11)
	SDF 851	TCATCCGCTGATAATCTTGACCTT	
	SDF 852	CACCATGGAAGAGAACTCAAGTAAAAA	AT3G46770 (REM13)
	SDF 853	TCAGGCCTTGATAATGTGGACCTT	
	IC75	ATGGCCAGAGACGGTGGT	AT1G10470 (ARR4)
	IC76	CTAATCTAATCCGGGACTCCTC	
	IC73	ATGAACAGTTCAGGAGGTTT	AT2G40670 (ARR16)
	IC74	TTAGCTTCTGCAGTTCATGA	
	IC63	ATGGAAAAAGCCTTGAGAAA	AT1G12980 (DRN)
	IC64	CTATCCCCACGATCTTCG	
Knock down line	1KDL-IAA27	CACCGCTACCTTGAGCTCTCTTCTG	AT4G29080 (IAA27)
	2KDL-IAA27	GTTCTGCTTCTGCACTTCTC	
RT-PCR	1RT-IAA27	GCTACCTTGAGCTCTCTTCTG	AT4G29080 (IAA27)
	2RT-IAA27	GTTCTGCTTCTGCACTTCTC	

SUPPLEMENTAL REFERENCES

- Alonso-Cantabrana, H., Ripoll, J. J., Ochando, I., Vera, A., Ferrándiz, C., & Martínez-Laborda, A. (2007). Common regulatory networks in leaf and fruit patterning revealed by mutations in the *Arabidopsis* *ASYMMETRIC LEAVES1* gene. *Development*, 134(14), 2663–2671. <https://doi.org/10.1242/dev.02864>
- Alvarez, J., & Smyth, D. R. (1999). *CRABS CLAW* and *SPATULA*, two *Arabidopsis* genes that control carpel development in parallel with *AGAMOUS*. *Development*, 126(11), 2377–2386.
- Alvarez, J., & Smyth, D. R. (2002). *CRABS CLAW* and *SPATULA* Genes Regulate Growth and Pattern Formation during Gynoecium Development in *Arabidopsis thaliana*. *International Journal of Plant Sciences*, 163(1), 17–41. <https://doi.org/10.1086/324178>
- Alvarez, J. P., Goldshmidt, A., Efroni, I., Bowman, J. L., & Eshed, Y. (2009). The *NGATHA* Distal Organ Development Genes Are Essential for Style Specification in *Arabidopsis*. *The Plant Cell Online*, 21(5), 1373–1393. <https://doi.org/10.1105/tpc.109.065482>
- Argyros, R. D., Mathews, D. E., Chiang, Y.-H., Palmer, C. M., Thibault, D. M., Etheridge, N., Argyros, D. A., Mason, M. G., Kieber, J. J., & Schaller, G. E. (2008). Type B Response Regulators of *Arabidopsis* Play Key Roles in Cytokinin Signaling and Plant Development. *The Plant Cell*, 20(8), 2102–2116. <https://doi.org/10.1105/tpc.108.059584>
- Arnaud, N., & Pautot, V. (2014). Ring the BELL and tie the KNOX: roles for TALEs in gynoecium development. *Frontiers in Plant Science*, 5, 1–7. <https://doi.org/10.3389/fpls.2014.00093>
- Azhakanandam, S., Nole-Wilson, S., Bao, F., & Franks, R. G. (2008). *SEUSS* and *AINTEGUMENTA* Mediate Patterning and Ovule Initiation during Gynoecium Medial Domain Development. *Plant Physiology*, 146, 1165–1181. <https://doi.org/10.1104/pp.107.114751>
- Bao, F., Azhakanandam, S., & Franks, R. G. (2010). *SEUSS* and *SEUSS-LIKE* Transcriptional Adaptors Regulate Floral and Embryonic Development in *Arabidopsis*. *Plant Physiology*, 152, 821–836. <https://doi.org/10.1104/pp.109.146183>
- Bowman, J. L., & Smyth, D. R. (1999). *CRABS CLAW*, a gene that regulates carpel and nectary development in *Arabidopsis*, encodes a novel protein with zinc finger and helix-loop-helix domains. *Development*, 126(11), 2387–2396. <https://doi.org/10.1016/j.palaeo.2004.10.017>
- Brambilla, V., Battaglia, R., Colombo, M., Masiero, S., Bencivenga, S., Kater, M. M., & Colombo, L. (2007). Genetic and Molecular Interactions between *BELL1* and *MADS* Box Factors Support Ovule Development in *Arabidopsis*. *The Plant Cell*, 19(8), 2544–2556. <https://doi.org/10.1105/tpc.107.051797>
- Chandler, J. W. (2018). Class VIIIb APETALA2 Ethylene Response Factors in Plant Development. *Trends in Plant Science*, 23(2), 151–162. <https://doi.org/10.1016/j.tplants.2017.09.016>
- Chandler, J. W., Jacobs, B., Cole, M., Comelli, P., & Werr, W. (2011). *DORNROSCHE*-LIKE expression marks *Arabidopsis* floral organ founder cells

- and precedes auxin response maxima. *Plant Molecular Biology*, 76, 171–185. <https://doi.org/10.1007/s11103-011-9779-8>
- Chandler, J. W., & Werr, W. (2017). DORNROSCHEN, DORNROSCHEN-LIKE, and PUCHI redundantly control floral meristem identity and organ initiation in *Arabidopsis*. *Journal of Experimental Botany*, 68(13), 3457–3472. <https://doi.org/10.1093/jxb/erx208>
 - Chuang, C. F., Running, M. P., Williams, R. W., & Meyerowitz, E. M. (1999). The *PERIANTHIA* gene encodes a bZIP protein involved in the determination of floral organ number in *Arabidopsis thaliana*. *Genes and Development*, 13(3), 334–344. <https://doi.org/10.1101/gad.13.3.334>
 - Colombo, M., Brambilla, V., Marcheselli, R., Caporali, E., Kater, M. M., & Colombo, L. (2010). A new role for the *SHATTERPROOF* genes during *Arabidopsis* gynoecium development. *Developmental Biology*, 337(2), 294–302. <https://doi.org/10.1016/j.ydbio.2009.10.043>
 - Conner, J., & Liu, Z. (2000). *LEUNIG*, a putative transcriptional corepressor that regulates *AGAMOUS* expression during flower development. *PNAS*, 97(23), 12902–12907. <https://doi.org/10.1073/pnas.230352397>
 - Crawford, B. C. W., & Yanofsky, M. F. (2011). *HALFFILLED* promotes reproductive tract development and fertilization efficiency in *Arabidopsis thaliana*. *Development*, 138(14), 2999–3009. <https://doi.org/10.1242/dev.067793>
 - Dinnyen, J. R., Weigel, D., & Yanofsky, M. F. (2006). *NUBBIN* and *JAGGED* define stamen and carpel shape in *Arabidopsis*. *Development*, 133(9), 1645–1655. <https://doi.org/10.1242/dev.02335>
 - Dinnyen, J. R., Weigel, D., & Yanofsky, M. F. (2005). A genetic framework for fruit patterning in *Arabidopsis thaliana*. *Development*, 132(21), 4687–4696. <https://doi.org/10.1242/dev.02062>
 - Dinnyen, J. R., Yadegari, R., Fischer, R. L., Yanofsky, M. F., & Weigel, D. (2004). The role of *JAGGED* in shaping lateral organs. *Development*, 131(5), 1101–1110. <https://doi.org/10.1242/dev.00949>
 - Ditta, G., Pinyopich, A., Robles, P., Pelaz, S., & Yanofsky, M. F. (2004). The *SEP4* Gene of *Arabidopsis thaliana* Functions in Floral Organ and Meristem Identity. *Current Biology*, 14, 1935–1940. <https://doi.org/10.1016/j>
 - Durbak, A. R., & Tax, F. E. (2011). *CLAVATA* Signaling Pathway Receptors of *Arabidopsis* Regulate Cell Proliferation in Fruit Organ Formation as well as Meristems. *Genetics*, 189, 177–194. <https://doi.org/10.1534/genetics.111.130930>
 - Eklund, D. M., Cierlik, I., Staldal, V., Claes, A. R., Vestman, D., Chandler, J., & Sundberg, E. (2011). Expression of *Arabidopsis* *SHORT INTERNODES/STYLISH* Family Genes in Auxin Biosynthesis Zones of Aerial Organs Is Dependent on a GCC Box-Like Regulatory Element. *Plant Physiology*, 157(4), 2069–2080. <https://doi.org/10.1104/pp.111.182253>
 - Elliott, R. C., Betzner, A. S., Huttner, E., Oakes, M. P., Tucker, W. Q. J., Gerentes, D., Perez, P., & Smyth, D. R. (1996). *AINTEGUMENTA*, an *APETALA2*-like Gene of *Arabidopsis* with Pleiotropic Roles in Ovule Development and Floral Organ Growth. *The Plant Cell*, 8(2), 155–168. <https://doi.org/10.1105/tpc.8.2.155>
 - Erdmann, R., Gramzow, L., Melzer, R., Theißen, G., & Becker, A. (2010). *GORDITA* (*AGL63*) is a young paralog of the *Arabidopsis thaliana* B_{sister} MADS

- box gene *ABS* (*TT16*) that has undergone neofunctionalization. *The Plant Journal*, 63, 914–924. <https://doi.org/10.1111/j.1365-313X.2010.04290.x>
- Eshed, Y., Baum, S. F., Perea, J. V., & Bowman J. L. (2001). Establishment of polarity in lateral organs of plants. *Current Biology*, 11, 1251–1260.
 - Favaro, R., Pinyopich, A., Battaglia, R., Kooiker, M., Borghi, L., Ditta, G., Yanofsky M. F., Kater M. M., & Colombo, L. (2003). MADS-Box Protein Complexes Control Carpel and Ovule Development in Arabidopsis. *The Plant Cell*, 15, 2603–2611. <https://doi.org/10.1105/tpc.015123.2>
 - Flanagan, C. A., Hu, Y., & Ma, H. (1996). Specific expression of the *AGL1* MADS-box gene suggests regulatory functions in *Arabidopsis* gynoecium and ovule development. *The Plant Journal*, 10(2), 343–353.
 - Girin, T., Paicu, T., Stephenson, P., Fuentes, S., Korner, E., O'Brien, M., Sorefan, K., Wood, T. A., Balanza, V., Ferrándiz, C., Smyth, D. R., & Ostergaard, L. (2011). INDEHISCENT and SPATULA Interact to Specify Carpel and Valve Margin Tissue and Thus Promote Seed Dispersal in *Arabidopsis*. *The Plant Cell*, 23(10), 3641–3653. <https://doi.org/10.1105/tpc.111.090944>
 - Gonçalves, B., Hasson, A., Belcram, K., Cortizo, M., Morin, H., Nikovics, K., Viallette-Guiraud, A., Takeda, S., Aida, M., Laufs, P., & Arnaud, N. (2015). A conserved role for *CUP-SHAPED COTYLEDON* genes during ovule development. *The Plant Journal*, 83(4), 732–742. <https://doi.org/10.1111/tpj.12923>
 - Gremski, K., Ditta, G., & Yanofsky, M. F. (2007). The *HECATE* genes regulate female reproductive tract development in *Arabidopsis thaliana*. *Development*, 134(20), 3593–3601. <https://doi.org/10.1242/dev.011510>
 - Groszmann, M., Paicu, T., Alvarez, J. P., Swain, S. M., & Smyth, D. R. (2011). *SPATULA* and *ALCATRAZ*, are partially redundant, functionally diverging bHLH genes required for Arabidopsis gynoecium and fruit development. *The Plant Journal*, 68, 816–829. <https://doi.org/10.1111/j.1365-313X.2011.04732.x>
 - Großhardt, R., Lenhard, M., & Laux, T. (2002). *WUSCHEL* signaling functions in interregional communication during *Arabidopsis* ovule development. *Genes & Development*, 16, 1129–1138. <https://doi.org/10.1101/gad.225202>
 - Heisler, M. G. B., Atkinson, A., Bylstra, Y. H., Walsh, R., & Smyth, D. R. (2001). *SPATULA*, a gene that controls development of carpel margin tissues in *Arabidopsis*, encodes a bHLH protein. *Development*, 128(7), 1089–1098. <https://doi.org/10.1105/tpc.9.10.1859>
 - Heyl, A., & Schmülling, T. (2003). Cytokinin signal perception and transduction. *Current Opinion in Plant Biology*, 6(5), 480–488. [https://doi.org/10.1016/S1369-5266\(03\)00087-6](https://doi.org/10.1016/S1369-5266(03)00087-6)
 - Hibara, K. I., Karim, M. R., Takada, S., Taoka, K. I., Furutani, M., Aida, M., & Tasaka, M. (2006). *Arabidopsis CUP-SHAPED COTYLEDON3* Regulates Postembryonic Shoot Meristem and Organ Boundary Formation. *The Plant Cell*, 18(11), 2946–2957. <https://doi.org/10.1105/tpc.106.045716>
 - Hirota, A., Kato, T., Fukaki, H., Aida, M., & Tasaka, M. (2007). The Auxin-Regulated AP2/EREBP Gene *PUCHI* Is Required for Morphogenesis in the Early Lateral Root Primordium of Arabidopsis. *The Plant Cell*, 19(7), 2156–2168. <https://doi.org/10.1105/tpc.107.050674>

- Ishida, T., Aida, M., Takada, S., & Tasaka, M. (2000). Involvement of *CUP-SHAPED COTYLEDON* genes in gynoecium and ovule development in *Arabidopsisthaliana*. *Plant Cell Physiology*, 41(1), 60–67. <https://doi.org/10.1093/pcp/41.1.60>
- Ishida, K., Yamashino, T., Yokoyama, A., & Mizuno, T. (2008). Three Type-B Response Regulators, ARR1, ARR10 and ARR12, Play Essential but Redundant Roles in Cytokinin Signal Transduction Throughout the Life Cycle of *Arabidopsisthaliana*. *Plant Cell Physiology*, 49(1), 47–57. <https://doi.org/10.1093/pcp/pcm165>
- Jiao, Y., & Meyerowitz, E. M. (2010). Cell-type specific analysis of translating RNAs in developing flowers reveals new levels of control. *Molecular Systems Biology*, 6(419), 1–14. <https://doi.org/10.1038/msb.2010.76>
- Kamiuchi, Y., Yamamoto, K., Furutani, M., Tasaka, M., & Aida, M. (2014). The *CUC1* and *CUC2* genes promote carpel margin meristem formation during *Arabidopsis* gynoecium development. *Frontiers in Plant Science*, 5, 1–9. <https://doi.org/10.3389/fpls.2014.00165>
- Kanrar, S., Onguka, O., & Smith, H. M. S. (2006). *Arabidopsis* inflorescence architecture requires the activities of KNOX-BELL homeodomain heterodimers. *Planta*, 224(5), 1163–1173. <https://doi.org/10.1007/s00425-006-0298-9>
- Karim, M. R., Hirota, A., Kwiatkowska, D., Tasaka, M., & Aida, M. (2009). A Role for *Arabidopsis* *PUCHI* in Floral Meristem Identity and Bract Suppression. *The Plant Cell*, 21(5), 1360–1372. <https://doi.org/10.1105/tpc.109.067025>
- Kerstetter, R. A., Bollman, K., Taylor, R. A., Bomblies, K., & Poethig, R. S. (2001). *KANADI* regulates organ polarity in *Arabidopsis*. *Nature*, 411, 706–709.
- Kieffer, M., Master, V., Waites, R., & Davies, B. (2011). TCP14 and TCP15 affect internode length and leaf shape in *Arabidopsis*. *The Plant Journal*, 68, 147–158. <https://doi.org/10.1111/j.1365-313X.2011.04674.x>
- Kirch, T., Simon, R., Grünewald, M., & Werr, W. (2003). *DORNROSCHE/ENHANCER OF SHOOT REGENERATION1* Gene of *Arabidopsis* Acts in the Control of Meristem Cell Fate and Lateral Organ Development. *The Plant Cell*, 15, 694–705. <https://doi.org/10.1105/tpc.009480>
- Koizumi, K., & Gallagher, K. L. (2013). Identification of SHRUBBY, a SHORT-ROOT and SCARECROW interacting protein that controls root growth and radial patterning. *Development*, 140, 1292–1300. <https://doi.org/10.1242/dev.090761>
- Kuusk, S., Sohlberg, J. J., Long, J. A., Fridborg, I., & Sundberg, E. (2002). *STY1* and *STY2* promote the formation of apical tissues during *Arabidopsis* gynoecium development. *Development*, 129, 4707–4717.
- Larue, C. T., Wen, J., & Walker, J. C. (2009). A microRNA-transcription factor module regulates lateral organ size and patterning in *Arabidopsis*. *The Plant Journal*, 58(3), 450–463. <https://doi.org/10.1111/j.1365-313X.2009.03796.x>
- Li, J., Dai, X., & Zhao, Y. (2006). A Role for Auxin Response Factor 19 in Auxin and Ethylene Signaling in *Arabidopsis*. *Plant Physiology*, 140(3), 899–908. <https://doi.org/10.1104/pp.105.070987>
- Liljegren, S. J., Ditta, G. S., Eshed, Y., Savidge, B., Bowman, J. L., Yanofsky, M. F. (2000). *SHATTERPROOF* MADS-box genes control seed dispersal in *Arabidopsis*. *Nature*, 404, 766–770.

- Liljegren, S. J., Roeder, A. H. K., Kempin, S. A., Gremski, K., Østergaard, L., Guimil, S., Reyes, D. K., & Yanofsky, M. F. (2004). Control of Fruit Patterning in *Arabidopsis* by INDEHISCENT. *Cell*, 116(6), 843–853. [https://doi.org/10.1016/S0092-8674\(04\)00217-X](https://doi.org/10.1016/S0092-8674(04)00217-X)
- Liu, Z., Franks, R. G., & Klink, V. P. (2000). Regulation of Gynoecium Marginal Tissue Formation by *LEUNIG* and *AINTEGUMENTA*. *The Plant Cell*, 12, 1879–1891.
- Lozano-Sotomayor, P., Chávez Montes, R. A., Silvestre-Vañó, M., Herrera-Ubaldo, H., Greco, R., Pablo-Villa, J., Galliani, B. M. Diaz-Ramirez, D., Weemen Mieke, Boutilier, K., Pereira, A., Colombo, L., Madueño, F., Marsch-Martínez, N., & de Folter, S. (2016). Altered expression of the bZIP transcription factor DRINK ME affects growth and reproductive development in *Arabidopsis thaliana*. *The Plant Journal*, 88(3), 437–451. <https://doi.org/10.1111/tpj.13264>
- Lucero, L. E., Uberti-manassero, N. G., & Gonzalez, D. H. (2015). TCP15 modulates cytokinin and auxin responses during gynoecium development in *Arabidopsis*. *The Plant Journal*, 84, 267–282. <https://doi.org/10.1111/tpj.12992>
- Mallory, A. C., Bartel, D. P., & Bartel, B. (2005). MicroRNA-Directed Regulation of *Arabidopsis AUXIN RESPONSE FACTOR17* Is Essential for Proper Development and Modulates Expression of Early Auxin Response Genes. *The Plant Cell*, 17, 1360–1375.
- Mandel, M. A., & Yanofsky, M. F. (1995). The *Arabidopsis AGL8* MADS Box Gene is Expressed in Inflorescence Meristems and is Negatively Regulated by *APETALA1*. *Development*, 121, 1763–1771.
- Mantegazza, O., Gregis, V., Mendes, M. A., Morandini, P., Alves-Ferreira, M., Patreze, C. M., Nardeli, S. M., Kater, M. M., & Colombo, L. (2014). Analysis of the *Arabidopsis REM* gene family predicts functions during flower development. *Annals of Botany*, 114, 1507–1515. <https://doi.org/10.1093/aob/mcu124>
- Martínez-Fernández, I., Sanchís, S., Marini, N., Balanzá, V., Ballester, P., Navarrete-Gómez, M., Oliveira, A. C., Colombo, L., & Ferrándiz, C. (2014). The effect of NGATHA altered activity on auxin signaling pathways within the *Arabidopsis* gynoecium. *Frontiers in Plant Science*, 5, 1–11. <https://doi.org/10.3389/fpls.2014.00210>
- Mason, M. G., Li, J., Mathews, D. E., Kieber, J. J., & Schaller, G. E. (2004). Type-B Response Regulators Display Overlapping Expression Patterns in *Arabidopsis*. *Plant Physiology*, 135(2), 927–937. <https://doi.org/10.1104/pp.103.038109>
- Mizzotti, C., Ezquer, I., Paolo, D., Rueda-Romero, P., Fiorella, R., Battaglia, R., Rogachev, I., Aharoni, A., Kater, M. M., Caporali, E., & Colombo, L. (2014). SEEDSTICK is a Master Regulator of Development and Metabolism in the *Arabidopsis* Seed Coat. *PLoS Genetics*, 10(12), 1–15. <https://doi.org/10.1371/journal.pgen.1004856>
- Mizzotti, C., Mendes, M. A., Caporali, E., Schnittger, A., Kater, M. M., Battaglia, R., & Colombo, L. (2012). The MADS box genes *SEEDSTICK* and *ARABIDOPSIS B_{sister}* play a maternal role in fertilization and seed development. *The Plant Journal*, 70, 409–420. <https://doi.org/10.1111/j.1365-3113.2011.04878.x>
- Nagpal, P., Ellis, C. M., Weber, H., Ploense, S. E., Barkawi, L. S., Guilfoyle, T. J., Hagen, G., Alonso, J. M., Cohen, J. D., Farmer, E. E., Ecker, J. R., & Reed, J. W.

- (2005). Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. *Development*, 132(18), 4107–4118. <https://doi.org/10.1242/dev.01955>
- Ng, K. H., Yu, H., & Ito, T. (2009). AGAMOUS Controls *GIANT KILLER*, a Multifunctional Chromatin Modifier in Reproductive Organ Patterning and Differentiation. *PLoS Biology*, 7(11), 1-17. <https://doi.org/10.1371/journal.pbio.1000251>
 - Nahar, M. A. U., Ishida, T., Smyth, D. R., Tasaka, M., & Aida, M. (2012). Interactions of *CUP-SHAPED COTYLEDON* and *SPATULA* genes control carpel margin development in *Arabidopsis thaliana*. *Plant and Cell Physiology*, 53(6), 1134–1143. <https://doi.org/10.1093/pcp/pcs057>
 - Nemhauser, J. L., Feldman, L. J., & Zambryski, P. C. (2000). Auxin and *ETTIN* in *Arabidopsis* gynoecium morphogenesis. *Development*, 127(18), 3877–3888. <http://dev.biologists.org/content/127/18/3877.short>
 - Nole-Wilson, S., Azhakanandam, S., & Franks, R. G. (2010). Polar auxin transport together with AINTEGUMENTA and REVOLUTA coordinate early Arabidopsis gynoecium development. *Developmental Biology*, 346, 181–195. <https://doi.org/10.1016/j.ydbio.2010.07.016>
 - Ohno, C. K., Reddy, G. V., Heisler M. G. B., & Meyerowitz, E. M. (2004). The *Arabidopsis* *JAGGED* gene encodes a zinc finger protein that promotes leaf tissue development. *Development*, 131(5), 1111–1122. <https://doi.org/10.1242/dev.00991>
 - Okushima, Y., Overvoorde, P. J., Arima, K., Alonso, J. M., Chan, A., Chang, C., Ecker, J. R., Hughes, B., Lui, A., Nguyen, D., Onodera, C., Quach, H., Smith, A., Yu, G., & Theologis, A. (2005). Functional Genomic Analysis of the *AUXIN RESPONSE FACTOR* Gene Family Members in *Arabidopsis thaliana*: Unique and Overlapping Functions of *ARR7* and *ARF19*. *The Plant Cell*, 17, 444-463.
 - Pautot, V., Dockx, J., Hamant, O., Kronenberger, J., Grandjean, O., Jublot, D., & Traas, J. (2001). *KNAT2*: Evidence for a Link between Knotted-like Genes and Carpel Development. *The Plant Cell Online*, 13(8), 1719–1734. <https://doi.org/10.1105/tpc.13.8.1719>
 - Pelaz, S., Ditta, G. S., Baumann E., Wisman Ellen & Yanofsky, M. F. (2000). B and C floral organ identity functions require SEPALLATA MADS-box genes. *Nature*, 405, 200–203.
 - Pekker, I., Alvarez, J. P., & Eshed, Y. (2005). Auxin Response Factors Mediate *Arabidopsis* Organ Asymmetry via Modulation of KANADI Activity. *The Plant Cell Online*, 17(11), 2899–2910. <https://doi.org/10.1105/tpc.105.034876>
 - Pinyopich, A., Ditta, G. S., Savidge, B., Liljegren, S. J., Baumann, E., Wisman, E., & Yanofsky, M. F. (2003). Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature*, 424, 85–88.
 - Poulis, S., & Vlachonasios, K. E. (2018). Synergistic action of *GCN5* and *CLAVATA1* in the regulation of gynoecium development in *Arabidopsis thaliana*. *New Phytologist*, 220, 593–608. <https://doi.org/10.1111/nph.15303>
 - Prasad, K., Zhang, X., Tobón, E., & Ambrose, B. A. (2010). The Arabidopsis B-sister MADS-box protein, GORDITA, represses fruit growth and contributes to integument development. *The Plant Journal*, 62(2), 203–214. <https://doi.org/10.1111/j.1365-313X.2010.04139.x>

- Ragni, L., Belles-Boix, E., Günl, M., & Pautot, V. (2008). Interaction of *KNAT6* and *KNAT2* with *BREVIPEDICELLUS* and *PENNYWISE* in *Arabidopsis* Inflorescences. *The Plant Cell*, 20(4), 888–900. <https://doi.org/10.1105/tpc.108.058230>
- Rajani, S., & Sundaresan, V. (2001). The *Arabidopsis* myc/bHLH gene *ALCATRAZ* enables cell separation in fruit dehiscence. *Current Biology*, 11(24), 1914–1922. [https://doi.org/10.1016/S0960-9822\(01\)00593-0](https://doi.org/10.1016/S0960-9822(01)00593-0)
- Reiser, L., Modrusan, Z., Margossian, L., Samach, A., Ohad, N., Haughn, G. W., & Fischer, R. L. (1995). The *BELL1* Gene Encodes a Homeodomain Protein Involved in Pattern Formation. *Cell*, 83, 735–742.
- Reyes-Olalde, J. I., Zúñiga-Mayo, V. M., Serwatowska, J., Chávez Montes, R. A., Lozano-Sotomayor, P., Herrera-Ubaldo, H., Gonzalez-Aguilera, K. L. Ballester, P., Ripoll, J. J., Ezquer, I., Paolo, D., Heyl, A., Colombo, L., Yanofsky, M. F., Ferrandiz, C., Marsch-Martínez, N., & de Folter, S. (2017). The bHLH transcription factor SPATULA enables cytokinin signaling, and both activate auxin biosynthesis and transport genes at the medial domain of the gynoecium. *PLoS Genetics*, 13(4), 1–31. <https://doi.org/10.1371/journal.pgen.1006726>
- Ripoll, J. J., Roeder, A. H. K., Ditta, G. S., & Yanofsky, M. F. (2011). A novel role for the floral homeotic gene *APETALA2* during *Arabidopsis* fruit development. *Development*, 138(23), 5167–5176. <https://doi.org/10.1242/dev.073031>
- Roeder, A. H. K., Ferrándiz, C., & Yanofsky, M. F. (2003). The Role of the REPLUMLESS Homeodomain Protein in Patterning the *Arabidopsis* Fruit. *Current Biology*, 13, 1630–1635.
- Romera-Branchat, M., Ripoll, J. J., Yanofsky, M. F., & Pelaz, S. (2013). The *WOX13* homeobox gene promotes replum formation in the *Arabidopsis thaliana* fruit. *The Plant Journal*, 73(1), 37–49. <https://doi.org/10.1111/tpj.12010>
- Running, M. P., & Meyerowitz, E. M. (1996). Mutations in the *PERANTHIA* gene of *Arabidopsis* specifically alter floral organ number and initiation pattern. *Development*, 122(4), 1261–1269. <http://www.ncbi.nlm.nih.gov/pubmed/8620853>
- Schaller, G. E., Bishopp, A., & Kieber, J. J. (2015). The Yin-Yang of Hormones: Cytokinin and Auxin Interactions in Plant Development. *The Plant Cell*, 27(1), 44–63. <https://doi.org/10.1105/tpc.114.133595>
- Scofield, S., Dewitte, W., & Murray, J. A. H. (2007). The *KNOX* gene *SHOOT MERISTEMLESS* is required for the development of reproductive meristematic tissues in *Arabidopsis*. *The Plant Journal*, 50, 767–781. <https://doi.org/10.1111/j.1365-313X.2007.03095.x>
- Sessions, A., Nemhauser, J. L., McCall, A., Roe, J. L., Feldmann, K. A., & Zambryski, P. C. (1997). *ETTIN* patterns the *Arabidopsis* floral meristem and reproductive organs. *Development*, 124(22), 4481–4491. <https://doi.org/10.1385/0896034437>
- Sessions, R. A., & Zambryski, P. C. (1995). *Arabidopsis* gynoecium structure in the wild type and in *ettin* mutants. *Development*, 121, 1519–1532.
- Shi, C. L., Stenvik, G. E., Vie, A. K., Bones, A. M., Pautot, V., Proveniers, M., Aalen, R. B., & Butenko, M. A. (2011). *Arabidopsis* Class I KNOTTED-Like Homeobox Proteins Act Downstream in the IDA-HAE/HSL2 Floral Abscission

- Signaling Pathway. *The Plant Cell*, 23(7), 2553–2567. <https://doi.org/10.1105/tpc.111.084608>
- Sieber, P., Gheyselinck, J., Gross-Hardt, R., Laux, T., Grossniklaus, U., & Schneitz, K. (2004). Pattern formation during early ovule development in *Arabidopsis thaliana*. *Developmental Biology*, 273, 321–334. <https://doi.org/10.1016/j.ydbio.2004.05.037>
 - Siegfried, K. R., Eshed, Y., Baum, S. F., Otsuga, D., Drews, G. N., & Bowman, J. L. (1999). Members of the *YABBY* gene family specify abaxial cell fate in *Arabidopsis*. *Development*, 126, 4117–4128.
 - Smith, H. M. S., Campbell, B. C., & Hake, S. (2004). Competence to Respond to Floral Inductive Signals Requires the Homeobox Genes *PENNYWISE* and *POUND-FOOLISH*. *Current Biology*, 14, 812–817. <https://doi.org/10.1016/j.cub.2004.04.032>
 - Sorefan, K., Girin, T., Liljegren, S. J., Ljung, K., Robles, P., Galván-Ampudia, C. S., Friml, J., Yanofsky, M. F., & Østergaard, L. (2009). A regulated auxin minimum is required for seed dispersal in *Arabidopsis*. *Nature*, 459(7246), 583–586. <https://doi.org/10.1038/nature07875>
 - Trigueros, M., Navarrete-Gomez, M., Sato, S., Christensen, S. K., Pelaz, S., Weigel, D., Yanofsky M. F., & Ferrándiz, C. (2009). The NGATHA Genes Direct Style Development in the *Arabidopsis* Gynoecium. *The Plant Cell Online*, 21, 1394–1409. <https://doi.org/10.1105/tpc.109.065508>
 - Uberti-Manassero, N. G., Lucero, L. E., Viola, I. L., Vegetti, A. C., & Gonzalez, D. H. (2011). The class I protein AtTCP15 modulates plant development through a pathway that overlaps with the one affected by CIN-like TCP proteins. *Journal of Experimental Botany*, 63(2), 809–823. <https://doi.org/10.1093/jxb/err305>
 - Urbanus, S. L., de Folter, S., Shchennikova, A. V., Kaufmann, K., Immink, R. G. H., & Angenent, G. C. (2009). In planta localisation patterns of MADS domain proteins during floral development in *Arabidopsis thaliana*. *BMC Plant Biology*, 9(5), 1–16. <https://doi.org/10.1186/1471-2229-9-5>
 - Ung, N., Lal, S., & Smith, H. M. S. (2011). The Role of *PENNYWISE* and *POUND-FOOLISH* in the Maintenance of the Shoot Apical Meristem in *Arabidopsis*. *Plant Physiology*, 156(2), 605–614. <https://doi.org/10.1104/pp.110.171462>
 - Villarino, G. H., Flores-Vergara, M., Hu, Q., Sehra, B., Robles, L., Brumos, J., Stepanova, A., Manrique, S., Colombo L., Sundberg, E., Heber, S., & Franks, R. G. (2016). Temporal and spatial domain-specific transcriptomic analysis of a vital reproductive meristem in *Arabidopsis thaliana*. *Plant Physiology*, 171(1), 42–61. <https://doi.org/10.1104/pp.15.01845>
 - Vroemen, C. W., Mordhorst, A. P., Albrecht, C., Kwaaitaal, M. A. C. J., & de Vries, S. C. (2003). The *CUP-SHAPER COTYLEDON 3* Gene is Required for Boundary and Shoot Meristem Formation in *Arabidopsis*. *The Plant Cell*, 15, 1563–1577. <https://doi.org/10.1105/tpc.012203>
 - Wang, B., Xue, J. S., Yu, Y. H., Liu, S. Q., Zhang, J. X., Yao, X. Z., Liu, Z. X., Xu, X. F., Yang, Z. N. (2017). Fine regulation of ARF17 for anther development and pollen formation. *BMC Plant Biology*, 17(1), 1–12. <https://doi.org/10.1186/s12870-017-1185-1>

- Wollmann, H., Mica, E., Todesco, M., Long, J. A., & Weigel, D. (2010). On reconciling the interactions between *APETALA2*, miR172 and *AGAMOUS* with the ABC model of flower development. *Development*, 137(21), 3633–3642. <https://doi.org/10.1242/dev.036673>
- Wu, M. F., Tian, Q., & Reed, J. W. (2006). *Arabidopsis* microRNA167 controls patterns of *ARF6* and *ARF8* expression, and regulates both female and male reproduction. *Development*, 133(21), 4211–4218. <https://doi.org/10.1242/dev.02602>
- Xu, B., Li, Z., Zhu, Y., Wang, H., Ma, H., Dong, A., & Huang, H. (2008). Arabidopsis Genes *ASI*, *AS2*, and *JAG* Negatively Regulate Boundary-Specifying Genes to Promote Sepal and Petal Development. *Plant Physiology*, 146(2), 566–575. <https://doi.org/10.1104/pp.107.113787>
- Yang, J., Tian, L., Sun, M. X., Huang, X. Y., Zhu, J., Guan, Y. F., Jia, Q. S. Yang, Z. N. (2013). AUXIN RESPONSE FACTOR17 Is Essential for Pollen Wall Pattern Formation in Arabidopsis. *Plant Physiology*, 163, 720-731.
- Yanofsky, M. F., Ma, H., Bowman, J. L., Drews, G. N., Feldmann, K. A., & Meyerowitz, E. M. (1990). The protein encoded by the *Arabidopsis* homeotic gene *agamous* resembles transcription factors. *Nature*, 346, 35–39. <https://doi.org/10.1038/346035a0>
- Zuñiga-Mayo, V. M., Marsch-Martínez, N., & de Folter, S. (2012). JAIBA, a class-II HD-ZIP transcription factor involved in the regulation of meristematic activity, and important for correct gynoecium and fruit development in Arabidopsis. *The Plant Journal*, 71, 314–326. <https://doi.org/10.1111/j.1365-313X.2012.04990.x>