Opsin gene evolution in amphibious and terrestrial combtooth blennies (Blenniidae)

Running title: Vision from water to land in blennies

Fabio Cortesi¹, Karen L. Cheney^{1,2}, Georgina M. Cooke³, Terry J. Ord³

¹Queensland Brain Institute and ²School of Biological Sciences, The University of Queensland, Brisbane, QLD 4072, Australia

³School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney 2052, Australia

Correspondence: F Cortesi, Fax: +61 (0)7 33654522, E-mail: fabio.cortesi@uqconnect.edu.au

Abstract

Evolutionary adaptations to life on land include changes to the physiology, morphology and behaviour of an animal in response to physical differences between water and air. The visual systems of amphibious species show pronounced morphological adaptations; yet, whether molecular changes also occur remains largely unknown. Here, we investigated the molecular evolution of visual pigment genes (opsins) in amphibious and terrestrial fishes belonging to the Salaria division of blennies (Blenniidae). We hypothesized that when conquering land, blenny opsins adapt – in terms of sequence variation and/or gene expression – to match both higher light intensities as well as the broader light spectrum. Using retinal transcriptomes in six species ranging from fully aquatic to fully terrestrial, we found very little variation in opsin gene sequences or gene expression between species. All blennies expressed a single rod opsin gene as well as two cone opsin genes sensitive to longer-wavelengths of light: *RH2A-1* (green-sensitive) and *LWS* (red-sensitive). They also expressed one or two short-wavelength-sensitive cone opsin genes (*SWS2Aα*, *SWS2Aβ*; blue-sensitive) in a phylogenetically inert manner. However, based on amino acid predictions, both SWS2A proteins confer similar peak spectral sensitivities and differential expression is therefore unlikely to be ecologically significant. Red-sensitivity is likely beneficial for feeding on algae and detritus, the main food source of Salaria blennies, and could be co-adapted to perceive visual displays in terrestrial species, which often use red dorsal fins to signal during aggressive disputes and courtship. Our data suggests that on the molecular level, the visual systems that evolved in aquatic blennies have been retained in species that have transitioned onto land.

Keywords fish, vision, transcriptomics, gene duplication, visual pigment

Introduction

The evolution of tetrapods provides a textbook example for the transition from water-to-land and occurred in the Devonian around 380 million years ago (MacIver et al. 2017). Interestingly, terrestrial or semi-terrestrial (amphibious) lifestyles, have evolved independently in at least thirty-three osteichthyan families (Ord and Cooke 2016, Wright and Turko 2016). Selective forces that drive fishes to leave the water may include access to food, new or better resources for reproduction (Shimizu et al. 2006), an escape from aquatic predation (Ord et al. 2017) or reduced competition (reviewed in Sayer 2005, Wright and Turko 2016). However, plastic and/or evolutionary adaptations are often needed in response to the extreme environmental changes between water and air. Such adaptations to life on land include: aerial respiration (Mandic et al. 2009, Regan et al. 2011), mucus secretion to prevent dehydration (Smith 1930, Sturla et al. 2002), and shifts in ammonia excretion to avoid NH_3 intoxication (Davenport and Sayer 1986, Chew and Ip 2014). Skeletal modifications are also common and may be triggered due to changes in locomotion (Kawano and Blob 2013, Brunt et al. 2016), an increase in body weight due to increase in apparent gravity (Turko et al. 2017), and by elevated oxygen content (Rossi et al. 2018). Moreover, behavioural modifications may occur, such as shade seeking to avoid desiccation during low tide (Colombini et al. 1995, Ord and Tonia Hsieh 2011).

The sensory systems of animals may also change as they move on to land. Since most fishes lack eyelids or similar structures, many amphibious fish species have modifications such as pigmented corneas and lenses, horizontal pupils and retractable eyes, to cope with increased light intensity on land (Sayer 2005). The refractive index of air versus water is also drastically different requiring modifications of the cornea, the lens or even the retina itself to maintain acuity when switching between environments (Sayer 2005). An extreme example of this can be found in the four-eyed fish, *Anableps anableps*, which has horizontally divided eyes: the top half has a flatter cornea compared to the bottom half to allow simultaneous vision above and below water, respectively (Swamynathan et al. 2003). However, while morphological adaptations have been studied in some detail, very little is known about possible molecular changes to the visual systems of amphibious and terrestrial fishes.

At the core of animal vision lie the opsin proteins that, together with a vitamin A-derived chromophore, form a light-sensitive photopigment located in the photoreceptors of the retina (Wald 1968, Hunt et al. 2014). Ancestral opsin gene duplications and subsequent changes to their amino acid compositions have led to five classes of vertebrate visual opsins that can be defined by their photoreceptor specificities and the different wavelengths of light

they are maximally sensitive to ($\lambda_{\max}$) (Hunt et al. 2014). RH1, the rod opsin is expressed in rod cells and used for dim-light vision, while the four other opsin classes are expressed in cone photoreceptors and used for daylight colour vision. These are: two short-wavelength, ultraviolet (UV) and blue-sensitive opsins (SWS1 and SWS2); one medium-wavelength-sensitive green opsin (RH2); and a long-wavelength-sensitive red opsin (LWS) (Yokoyama 2002). In teleost fishes and amphibians with different cone morphologies, SWS opsins are further expressed in single cones, while RH2 and LWS opsins are found in double cones (two single cones that are fused together) (Hunt et al. 2014).

The evolution of visual opsins in teleost fishes is especially dynamic involving additional gene duplications, deletions, pseudogenizations and gene conversions, which has led to fishes having between one and 40 opsin genes in their genomes (Cortesi et al. 2015, Lin et al. 2017, Musilova et al. 2018 preprint). This diversity is primarily thought to be due to the different light environments fishes inhabit. For example, light at either end of the visible spectrum gets absorbed with increasing depth and consequently deeper living species gradually lose their *SWS1* and *LWS* genes (Musilova et al. 2018 preprint). Species living in murky red-light dominated habitats, on the other hand, often have red-shifted visual systems (e.g., Hofmann et al. 2009, Escobar-Camacho et al. 2016); however, this could also be due to sexual selection (see Sandkam et al. 2018 for a recent review on guppy *LWS* evolution). Mutations in the coding sequence, differential expression of opsin genes and switches between A₁-based and A₂-based chromophores also enable adaptations to more subtle differences in photic environments, such as between seasons (Shimmura et al., 2017), microhabitats (Fuller et al. 2010, Luehrmann et al. in review), feeding habits (Stieb et al. 2017), predation pressure (Sandkam et al., 2018), or in response to sexual selection (Seehausen et al. 2008).

In this study, we focused on the evolution of opsin genes in amphibious and terrestrial combtooth blennies (Blenniidae). Combtooth blennies are small (< 10 cm) scaleless fish that are commonly found in many shallow tropical and warm water marine habitats, including coral reefs, estuaries, mangroves, tide pools, and sometimes on land (Hundt et al. 2014a, Hundt and Simons 2018). They comprise one of the most diverse percomorph families consisting of 400 described species (58 genera; fishbase.org) that fall into 13 phylogenetic clades (Hundt et al. 2014a, Hundt and Simons 2018). Within the Salariini division of blennies, amphibious behaviour is common, and more than twenty species in at least three genera exhibit a highly terrestrial lifestyle (Ord and Cooke 2016). In these species, post-settlement larvae (~30 days from hatching; Platt and Ord 2015) are believed to transition to a

terrestrial lifestyle within the supralittoral splash zone and do not voluntarily return to the aquatic environment (Ord et al. 2017). Terrestrial blennies move freely about the rocks using a tail-twisting behaviour that allows them to efficiently shuffle along the rocks and jump distances of several body lengths (Hsieh 2010). Like most intertidal blennies, amphibious and terrestrial species feed primarily on detritus and algae (Hundt et al. 2014a, Hundt et al. 2014b, Hundt and Simons 2018) and possess a cryptic body colouration and patterning that reduces predation (Morgans and Ord 2013, Ord et al. 2017). However, unlike aquatic and most amphibious blennies, adults of terrestrial species in both sexes display a brightly coloured red-orange dorsal fin during aggressive disputes and courtship (Bhikajee and Green 2002, Shimizu et al. 2006, Ord and Tonia Hsieh 2011). These conspicuous fins contrast against the rocky backgrounds on which the blennies are active (Morgans and Ord 2013, Morgans et al. 2014), and are further accentuated in courting males by a darkening of the body to a largely uniform charcoal black (Ord and Tonia Hsieh 2011).

Morphological adaptation for aerial vision amongst the Salariini has been reported for the amphibious (possibly exclusively terrestrial) Kirk's blenny, *Alticus kirkii*, which, by separating the cornea conjunctiva and cornea propria, has formed an additional eye chamber that is adjustable to accommodate changes in refracting indices of media (Zander 1974). Also, the retina of the mildly amphibious rippled rockskipper, *Istiblennius edentulus*, (see Ord and Cooke 2016) shows prominent swellings and folds, and a central depression into which the lens can be retracted. This allows light to be focused onto the back of the eye in both water and air (Zander 1974).

In terms of the molecular basis for vision, little is known about the evolution of opsin genes in Salariini blennies. However, a close relative to the Salariini, the rock-pool blenny, *Parablennius parvicornis*, was recently found to possess six cone opsins and one rod opsin in its genome (Musilova et al. 2018 preprint). Also, transcriptome sequencing in another aquatic blenny, the bluestriped fangblenny, *Plagiotremus rhinorhynchus*, revealed that this species expresses five orthologous cone opsins in its eyes (Musilova et al. 2018 preprint). Given that amphibious species show morphological adaptations for vision in air, and that blenny visual opsins are extremely diverse, we hypothesized that aerial vision and exposure to full sunlight caused adaptations in the Salariini opsin gene repertoires. More specifically, due to the tissue damage UV-radiation inflicts, we expected to see a lack of *SWS1* expression in the eyes of these fishes. Instead, feeding on detritus and algae (Hundt et al. 2014b), which generally show strong reflection in the red due to their chlorophyll component (Stieb et al. 2017), as well as the use of red dorsal fins for aggressive and courtship displays (see Ord and Tonia

Hsieh 2011, Morgans and Ord 2013), might have caused amphibious and terrestrial blennies to have a more long-wavelength, *LWS*, dominated visual system.

Material and Methods

Study species

Adult blennies belonging to the Salariaiini clade (Hundt et al. 2014a) were collected on snorkel or on foot with hand nets from several sites around the Indo-Pacific. Given that these blennies actively seek shade to avoid desiccation (Ord and Tonia Hsieh 2011), sampling either occurred in the early hours of the morning or late in the afternoon. Immediately post capture, their eyes were enucleated and stored on RNAlater (Life Technologies) for subsequent molecular analysis. From French Polynesia (FP), we collected Lined Rockskipper, *Istiblennius lineatus* (Mo'orea: 17°29'52.05"S, 149°45'17.24"W; November 8-10, 2013; N = 3), Blackmargin Rockskipper, *Praealticus caesius* (Mo'orea: 17°29'52.05"S, 149°45'17.24"W; November 8-10, 2013; N = 3); and Marquesan Rockskipper, *Alticus simplicirrus* (Tahiti: 17°30'57.67"S, 149°24'25.14"W; November 14-15, 2013 N = 3). From the Seychelles (S), we collected *I. lineatus* (4°48'9"S, 53°30'60"E and 4°33'56"S, 55°27'10"E; April 19 and 26, 2014; N = 3); Reef Margin Blenny, *Entomacrodus striatus* (4°48'9"S, 53°30'60"E; April 19 and 22, 2014; N = 3); and the Seychelles Rockskipper, *Alticus anjouanae* (4°33'56"S, 55°27'10"E; April 20, 2014; N = 3). A Jewelled Blenny, *Salarias fasciatus* (N = 1), was collected in May 2017 from Heron Island (23°44'S, 151°91'E), Great Barrier Reef, Australia.

All experimental procedures were approved by Animal Ethics Committees from The University of New South Wales (11/36B and 13/21) and The University of Queensland (QBI/304/16). Fish were collected under permits issued by Protocole D'Accueil (10/10/2013) French Polynesia, Seychelles Bureau of Standards (#A0157), the Great Barrier Reef Marine Park Authority (G17/38160.1) and Queensland Fisheries (#180731).

Transcriptome sequencing, quality filtering and de-novo assembly

Retinae were dissected out of the eyecup and total RNA was extracted using an RNAeasy Mini Kit (Qiagen) including DNase treatment following the manufacturer's instructions. RNA integrity was checked using an Eukaryotic Total RNA Nano chip on a Bioanalyzer 2100 (Agilent Technologies). All retinal transcriptomes, except for *S. fasciatus*, were sequenced in-house at the Queensland Brain Institute's sequencing facility, Brisbane, Australia. Sequencing libraries were prepared from 100-1,000 ng of total RNA using the

TruSeq total stranded mRNA Library Prep Kit protocol (Illumina, San Diego), and library concentrations were measured using a Qubit dsDNA BR Assay Kit (Thermo Fisher). Individual libraries were barcoded and up to 12 libraries/lane were pooled at equimolar ratios. Libraries were sequenced at PE125 on a HiSeq 2000 using Illumina's SBS chemistry version 4. The *S. fasciatus* library preparation (strand-specific, 250~300 bp insert) and sequencing (RNAseq HiSeq PE150) was outsourced to Novogene (<https://en.novogene.com/>).

Filtering and *de novo* assembly of retinal transcriptomes followed the protocol described in (de Busserolles et al. 2017). In short, raw-read transcriptomes were uploaded to the Genomics Virtual Laboratory (GVL 4.0.0) (Afgan et al. 2015) on the Galaxy Australia server (<https://galaxy-qld.genome.edu.au/galaxy/>) and quality filtered using Trimmomatic (Galaxy v.0.32.2) (Bolger et al. 2014) before being *de novo* assembled using Trinity (Galaxy v.0.0.2) (Haas et al. 2013). Two transcriptomes per species were assembled, with the exception of *S. fasciatus* where only one individual was sequenced. Raw-read libraries and assemblies are available on the Sequence Read Archive (<https://www.ncbi.nlm.nih.gov/sra>) and the Transcriptome Shotgun Assembly Database (<https://www.ncbi.nlm.nih.gov/genbank/tsa/>), respectively (Table S1).

Opsin gene mining, phylogenetic reconstructions and opsin gene expression

Opsin gene mining and expression analyses followed the protocol described in de Busserolles et al. 2017. In short, putative Salariini opsin genes were searched for by mapping the assembled transcripts to the opsin coding sequences of the dusky dottyback, *Pseudochromis fuscus* (Cortesi et al. 2016) in Geneious v.11.0.2 (www.geneious.com). *Ps. fuscus* was chosen because it belongs to the closely related Pseudochromidae (Alfaro et al. 2018) and possesses an opsin gene repertoire containing representatives from all ancestral vertebrate opsin gene classes (Cortesi et al. 2016). Mapped contigs were extracted and compared to publicly available opsin gene sequences using BLASTN (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Moreover, because *de novo* assembly based on short-read libraries is prone to create misassemblies (chimeric sequences), and/or to overlook closely related or lowly expressed gene copies, we used a second approach to account for all expressed Salariini visual opsins. This approach entailed the mapping of unassembled reads to the reference *Ps. fuscus* genes, followed by a manual extraction of gene copies using paired-end information to move from single polynucleotide polymorphism (SNP) to SNP along the gene. Extracted reads were *de novo* assembled and if necessary, their consensus was used as a species-specific template

against which unassembled reads were repeatedly re-mapped until the whole coding region could be extracted.

Salariini visual opsins were confirmed and assigned to specific opsin gene classes based on their phylogenetic relationships to a reference dataset obtained from GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and Ensembl (www.ensembl.org/) (Fig. 1, Fig. S1). Gene coding regions were aligned using the L-INS-I settings as part of the Geneious MAFFT plugin v.1.3.7 (Kato and Standley 2013), and jModeltest v.2.1.10 (Darriba et al. 2012) was subsequently used to select the most appropriate model of sequence evolution based on the Akaike information criterion. MrBayes v.3.2.6 (Ronquist et al. 2012) run on the CIPRES platform (Miller et al. 2010) was then used to infer the phylogenetic relationship between opsin genes using the following parameter settings: GTR+I+G model; two independent MCMC searches with four chains each; 10 million generations per run; 1000 generations sample frequency; and, 25% burn-in. Phylogenies were also reconstructed using GTR+G models, to account for the possibility of variable sites, however, no substantial differences in tree structure or node support could be found. Raw trees from either approach and corresponding data alignments have been deposited in Dryad (<https://datadryad.org/>), and GenBank accession numbers for the relevant genes are given either in Fig. S1, or in Table S1.

Quantitative opsin gene expression was measured by mapping the unassembled reads against the extracted opsin coding regions for each species as per de Busserolles et al. 2017. We then compared the rod opsin expression to the combined cone opsin expression, the proportional expression of each cone opsin gene to the combined cone opsin expression and finally, the proportional expressions of single (*SWS2A α* , *SWS2A β*) and double cone (*RH2A-1*, *LWS*) genes amongst themselves (Table 2). Single and double cone opsin gene expression for each species was then plotted onto the Salariini phylogeny taken from Ord and Cooke, 2016 (Fig. 2). Moreover, we also quantified the expression of *cytochrome P450 family 27 subfamily c member 1* (*cyp27c1*) by comparing its gene expression to the total opsin expression of fishes. The zebrafish *cyp27c1* ortholog converts vitamin A₁-based chromophore to the longer shifted A₂-chromophore (Enright et al. 2015) and A₂ has previously been reported in the peacock blenny, *Salarias pavo*, based on microspectrophotometry results (White et al. 2004).

Opsin gene sequence analysis and spectral sensitivity predictions

Opsin gene coding sequences were aligned using the L-INS-I settings in MAFFT (Geneious plugin v.1.3.7) (Kato and Standley 2013) and gene specific opsin trees were reconstructed

using RAxML v.8.2.11 (Geneious plugin v.3.0) (Stamatakis 2014), a GTR+G model and 1000 bootstraps to generate the support values for majority-rule consensus trees (Fig. S2). The alignments and corresponding trees were then used to test for site-specific positive selection using codeml in PAML (Yang 2007) as described in detail by (Hofmann et al. 2012). Briefly, codeml was run on the graphical user interface pamlX v.1.3.1. (Xu and Yang 2013) using likelihood ratio tests (LRT) to compare M1a vs. M2 and M8 vs. M8a. Bayesian Empirical Bayes (BEB) criteria (Yang et al. 2005) were subsequently applied to identify specific amino acid sites under positive selection in case of significant LRTs (Table S2). Next, we imported the alignments into MEGA7 (Kumar et al. 2016), which was used to calculate synonymous (ds) and nonsynonymous substitution rates (dn) using the Nei-Gojobori method (Jukes-Cantor distances)(Table 1).

Salariini opsin amino acid sequences were then aligned to bovine rhodopsin (Palczewski et al. 2000) to assess their variability within the transmembrane and the retinal binding pocket sites as well as at known opsin spectral tuning sites (Fasick and Robinson 1998, Hunt et al. 2001, Yokoyama 2008, Dungan et al. 2015). Since the dn/ds ratio indicated low variability between Salariini opsins, we extended this analysis to also include the opsins of the rock-pool blenny, *Parablennius parvicornis* (Musilova et al. 2018 preprint), as a more distantly related obligate aquatic species from the sister clade of the Parablenniini (Hundt et al. 2014a). Amino acid substitutions were assessed for each opsin gene amongst Salariini species, and between the Salariini and *Pa. parvicornis* taking specific note of sites that differ in polarity between species (as per Hofmann et al. 2012) (Table 1 and Table S3). Finally, we inferred the spectral sensitivities of the Salariini opsins based on changes at known spectral tuning sites compared to the well documented opsin sequences of *Ps. fuscus* (Cortesi et al. 2016) (Table S3). Individual sites are referred to according to their location relative to bovine rhodopsin.

Results

Opsin gene phylogeny and gene expression

Retinal transcriptomes from our studied species contained six cone and one rod opsin (*RH1*) gene. Phylogenetic reconstruction revealed two blue-sensitive *SWS2A* copies (*SWS2Aα*, *SWS2Aβ*), three green-sensitive *RH2* copies (*RH2A-1*, *RH2A-2* and *RH2B*), and one red-sensitive *LWS* gene (Fig. 1, Fig. S1, Table S1).

Rod opsin expression was generally much higher than cone opsin expression (> 73% of total opsin expression), except in *E. striatus* and *I. lineatus* (S) where rod and cone opsin

genes were more equally expressed (~ 50% of total opsin expression each). In terms of cone opsin expression, single cone genes (*SWS2Aα* + *SWS2Aβ*) accounted for ~ 10 – 15% of total cone opsin expression, except for one *Pr. caesius* individual which had 24.4 % single cone gene expression (Table 2). Comparing single and double cone expression separately, the single cone specific *SWS2A* paralogs were expressed at similar ratios (~ 40 – 60 % each) within the subclade containing the two terrestrial *Alticus* spp. as well as the immediate sister group to this genus, the mildly-amphibious *Pr. caesius*. In the remaining mildly-amphibious, amphibious and aquatic species, *SWS2Aβ* was primarily expressed (> 83%) (Fig. 2, Table 2). Neither the UV-sensitive *SWS1* nor the violet-sensitive *SWS2B* single cone genes were found to be expressed in the retinae of adult Salariini. Considering double cone gene expression, all blennies were found to express high amounts of *RH2A-1* and *LWS* (~ 40 – 60% each)(Fig. 2, Table 2). *RH2A-2*, on the other hand, was found to be lowly expressed (< 0.1% of total cone opsin expression) only allowing full coding sequence reconstruction for *A. simplicirrus* and partial sequence reconstructions for *Pr. caesius* and *I. lineatus* (Fig. 1). *S. fasciatus* was the only Salariini species with low levels of *RH2B* expression (< 0.1% of total cone opsin expression). Therefore, gene expression for *RH2A-2* and *RH2B* are unlikely to be relevant for adult Salariini vision and have been excluded from all further analyses (data not shown). *cyp27c1* was found to be lowly expressed in all blennies at < 0.5% compared to total opsin expression (Table 2).

Opsin gene diversity, selection analysis and predicted spectral sensitivities

The Salariini opsin genes were highly similar both when compared between species within the clade, and when compared to the more distant *Pa. parvicornis* sister species (Table 1). Overall, *SWS2Aα* (13/22 amino acid substitutions within Salariini/against *Pa. parvicornis*) had the highest and *RH2A-1* (3/11) the lowest number of amino acid substitutions within transmembrane- and retinal binding pocket sites. Only a few of these amino acid substitutions were found to occur in potential spectral tuning sites, namely those sites that have previously been identified to change the spectral sensitivity of opsin pigments and/or substitutions that incorporated changes in polarity (Tables 1 and S3). Reflective of the low variability between the Salariini genes, using PAML revealed that none of the Salariini genes are under positive selection (Table S2).

Most Salariini opsin genes showed no changes in key tuning sites when compared to their *Ps. fuscus* orthologs (Cortesi et al. 2016; Table S3). Consequently, assuming a purely A₁ chromophore based visual system, the peak spectral sensitivity ($\lambda_{\max}$) for the Salariini

SWS2A α was estimated to be at 448 nm. For SWS2A β we found one substitution at site T269A. The reverse substitution i.e. A269T, causes a 6 nm red-shift in $\lambda_{\max}$ (Yokoyama 2008) and we therefore assumed a similar shift but in the opposite direction in our case. Thus, the Salariaiini SWS2A β was estimated to have a 451 nm $\lambda_{\max}$. The Salariaiini RH2A-1 was found to be identical in key spectral tuning sites to the *Ps. fuscus* RH2A α and its estimated $\lambda_{\max}$ therefore to be 524 nm. The Salariaiini LWS (561 nm $\lambda_{\max}$) was also identical in tuning sites to its *Ps. fuscus* ortholog, except for *P. ceaseus* (554 nm $\lambda_{\max}$) where the substitution S164A is likely to have blue shifted the spectral sensitivity by 7 nm (Yokoyama 2008). Finally, the Salariaiini RH1 (500 nm $\lambda_{\max}$) showed one substitution at N83D likely to cause a 2 nm red-shift (Yokoyama 2008) and in the case of the *S. fasciatus* RH1 (501 – 503 nm $\lambda_{\max}$), showed a second substitution at A124G further red-shifting this visual pigment by 1 – 3 nm (Hunt et al. 2001).

Discussion

In this study, we investigated the molecular evolution of vision in fishes from the Salariaiini division of blennies, which have transitioned from water to land. First, we sequenced the retinal transcriptomes of six species classified as fully aquatic (*S. fasciatus*), mildly amphibious (*I. lineatus* and *Pr. caesius*), amphibious (*E. striatus*), and fully terrestrial (*A. simplicirrus* and *A. anjouanae*) (Ord and Cooke 2016; Fig. 1). We found that within their eyes, blennies express one rod opsin (*RH1*) and between three and four cone opsins (Fig. 2) independent of habitat or geographic region (i.e. South Pacific vs. Indian Ocean). *RH1* expression in fishes shows a strong diurnal pattern, starting with high expression in the morning and gradually decreasing as the day goes on (e.g., Korenbrot and Fernald 1989, Stieb et al. 2016). Hence, differences in sampling time i.e. early morning versus late afternoon, most likely explains the discrepancy of rod to cone opsin expression found in this study (Table 2).

Regarding the cone opsins, all Salariaiini species expressed the double cone genes sensitive to the red (*LWS*) and the green (*RH2A-1*) part of the light spectrum. We also found a second *RH2A* paralog in the genera *Alticus*, *Praealticus* and *Istiblennius* (Fig. 1). The *RH2A-2* gene was not expressed in the closely related *S. fasciatus* nor is it present in the genome of the more distantly related *Pa. parvicornis* (Musilova et al. 2018 preprint). One could therefore assume that this *RH2A* duplication is specific to the clade containing the amphibious and terrestrial Salariaiini species (Ord and Cooke 2016). However, based on our phylogenetic reconstruction, *RH2A-2* is basal to a greater *RH2A* clade which includes gene

orthologs from dottybacks and cichlids (Fig. S1). Both cichlids (Escobar-Camacho et al. 2016) and dottybacks (Cortesi et al. 2016) have two *RH2A* duplicates which have undergone widespread gene conversion and consequently cluster closely together within species/families. The blenny *RH2A* copies, on the other hand, appear unaffected by gene conversion. Rather than being Salariini specific then, it is likely that *RH2A-2* is the blenny ortholog of an ancestral *RH2A* duplication and that *Pa. parvicornis* has lost the copy secondarily. As more and more fish genomes become available (e.g. Musilova et al. 2018 preprint), the patterns surrounding *RH2* evolution are likely to become clear in the near future.

Finally, there was a strong phylogenetic signal in the expression of the blue-sensitive *SWS2A* duplicates: only the clade containing the two *Alticus* species and *Pr. caesi* expressed both of the copies at a similar ratio, while the remaining Salariini species mainly expressed the slightly longer tuned *SWS2A β* copy (Fig. 2). Whether this difference in expression is ecologically significant or whether it is the result of a phylogenetic inertia remains to be investigated. Given that the predicted $\lambda_{\max}$ for the two *SWS2A* paralogs is only 3 nm apart, we currently favour the latter scenario. As such, based on cone opsin expression and chromophore A₁-derived spectral sensitivity predictions, Salariini blennies appear to have a well-developed, potentially trichromatic, colour vision sense which reaches across the visible light spectrum ranging from 448 - 561 nm in $\lambda_{\max}$.

Our study species live in shallow reef environments, are amphibious or have left the aquatic realm altogether. The light environment these species experience differs depending on their habitat with both an increase in light intensity and in the proportion of UV and red wavelengths when moving into shallower water and finally onto land (see e.g., Marshall et al. 2003). Opsin gene expression in fishes has been shown to be influenced by changes in light environment with water depth (e.g., Stieb et al. 2016), and degree of suspended organic material (e.g., Fuller et al. 2005). However, our data shows that there are no large differences in opsin gene expression between Salariini species from different habitats (Fig. 2; Table 2). All of the species appear to have red-shifted visual systems hinting towards the importance of detecting longer-wavelengths of light for survival. Mounting evidence suggests that long-wavelength reception in coral reef fishes is especially beneficial when feeding on algae or similar chlorophyll containing organic matter, which strongly reflect the red part of the light spectrum (for a recent review on the topic see Marshall et al. 2018). It is also possible that the red-orange dorsal fin that appears unique to the terrestrial species (TJ Ord, unpublished data) has evolved to exploit the pre-existing (ancestral) visual state of seeing red as a means of

maximizing the efficiency of territorial and courtship signalling on land (Bhikajee and Green 2002, Shimizu et al. 2006, Ord and Tonia Hsieh 2011). The red-orange coloration of the dorsal fin of terrestrial blennies has been shown to be highly chromatically contrasting against the typical environmental background found on land (Morgans and Ord 2013). Similarly, sexually selected red-orange colour signals evolving in response to an innate red sensory bias in conspecific receivers has also been suggested for a number of other fish species (e.g., see Rodd et al. 2002, Smith et al. 2004, Seehausen et al. 2008).

It is notable that none of the species was found to express the UV-sensitive *SWS1* gene or the violet-sensitive *SWS2B* gene. While it is possible that *SWS2B* was lost in the ancestor of all blennies (Cortesi et al. 2015), *SWS1* is present, at least on the genomic level, in *Pa. parvicornis* (Musilova et al. 2018 preprint). Salariini blennies could have lost *SWS1* independently or alternatively, *SWS1* might simply not be expressed in adult blennies. Instead, *SWS1* and also the *RH2A-2* and *RH2B* paralogs, might be used at different developmental stages, a common feature of opsin gene expression in fishes (e.g., Spady et al. 2006, Carleton et al. 2008, Cortesi et al. 2016, Savelli et al. 2018). Supporting the potential use of *SWS1* at earlier life stages, Siebeck and Marshall (2007) found that adult Salariini blennies have UV blocking eyes while larval stages have UV to violet transmitting eyes. In an interesting parallel, mudskippers (Gobiidae) (You et al. 2014) and the Asian swamp eel (*Monopterus albus*) (Musilova et al. 2018 preprint), both of which are amphibious, have lost their *SWS1* gene with the swamp eel further missing its *SWS2B* copy (Cortesi et al. 2015). It is possible that the tissue damage UV-radiation induces has led to the convergent loss and/or the inexpression of the shorter-shifted *SWS* genes in amphibious species more generally.

Overall, opsin orthologs were highly conserved between Salariini species, and only very few changes in key tuning sites potentially shifting spectral sensitivities could be found (Table 1 and Table S3). Therefore, neither at the opsin-sequence nor at the expression level were there any apparent adaptations to the transition from water to land. This is surprising since morphological adaptations of the visual system when moving out-of-water are common (Sayer 2005). It remains possible that instead of relying on their opsins, Salariini blennies change the chromophore part of the photopigment. For example, most amphibians use red-shifted chromophore A₂ based visual systems during their aquatic life stages, but change to a blue-shifted A₁ based visual system after metamorphosis and moving onto land (Wilt 1959, Liebman and Entine 1968, Bridges 1972). Most Salariini species were found to lowly express *Cyp27c1*, the enzyme responsible for the A₁ to A₂ switch (Enright et al. 2015) (Table 2). It remains to be seen if the use of these chromophores differs between habitats.

In summary, we found no apparent molecular adaptations in terms of opsin gene sequence variability and expression in terrestrial blennies when compared to their amphibious and aquatic sister species. On the contrary, our data suggests that Salariini blennies evolved visual systems early on that were ideal to conquer shallow reef and intertidal habitats, and no further molecular adaptations have been made for life on land.

Acknowledgements

We would like to thank Sara Stieb and Fanny de Busserolles for assistance with RNA extractions and specimen collection, and Janette Edson from the Queensland Brain Institute's sequencing facility for library preparation and RNA sequencing.

Competing Interests

The authors declare no competing interests.

Funding

F.C. was supported by a Swiss National Science Foundation Early Postdoc Mobility Fellowship (165364) and a UQ Development Fellowship. This study was also funded by a Discovery Project grant from the Australian Research Council to T.J.O. (DP120100356).

Author Contributions

T.J.O. conceived the study and designed the experiments together with F.C. and K.L.C. T.J.O., G.M.C. and F.C. collected the specimens. F.C. performed the experiments, analysed the data, and wrote the initial manuscript. All authors reviewed and approved the final version of the manuscript.

Data Accessibility

Raw-read transcriptomes (SRA tba) and single gene sequences (#tba) are available through GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>). Gene alignments and single gene phylogenies can be accessed through Dryad (#tba). All other data is given either in the main manuscript or the supplementary material.

References

- Afgan, E., C. Sloggett, N. Goonasekera, I. Makunin, D. Benson, M. Crowe, S. Gladman, Y. Kowsar, M. Pheasant, and R. Horst. 2015. Genomics virtual laboratory: a practical bioinformatics workbench for the cloud. *PLoS ONE* **10**:e0140829.
- Alfaro, M. E., B. C. Faircloth, R. C. Harrington, L. Sorenson, M. Friedman, C. E. Thacker, C. H. Oliveros, D. Černý, and T. J. Near. 2018. Explosive diversification of marine fishes at the Cretaceous–Palaeogene boundary. *Nature Ecology & Evolution* **2**:688-696.
- Bhikajee, M., and J. Green. 2002. Behaviour and habitat of the Indian Ocean amphibious blenny, *Alticus monochrus*. *African Zoology* **37**:221-230.
- Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**:2114-2120.
- Bridges, C. 1972. The rhodopsin-porphyrin visual system. Pages 417-480 *Photochemistry of vision*. Springer.
- Brunt, E. M., A. J. Turko, G. R. Scott, and P. A. Wright. 2016. Amphibious fish jump better on land after acclimation to a terrestrial environment. *Journal of Experimental Biology* **219**:3204.
- Carleton, K. L., T. C. Spady, J. T. Streelman, M. R. Kidd, W. N. McFarland, and E. R. Loew. 2008. Visual sensitivities tuned by heterochronic shifts in opsin gene expression. *BMC Biology* **6**:22.
- Chew, S. F., and Y. K. Ip. 2014. Excretory nitrogen metabolism and defence against ammonia toxicity in air-breathing fishes. *Journal of Fish Biology* **84**:603-638.
- Colombini, I., R. Berti, A. Ercolini, A. Nocita, and L. Chelazzi. 1995. Environmental factors influencing the zonation and activity patterns of a population of *Periophthalmus sobrinus* Eggert in a Kenyan mangrove. *Journal of Experimental Marine Biology and Ecology* **190**:135-149.
- Cortesi, F., Z. Musilová, S. M. Stieb, N. S. Hart, U. E. Siebeck, K. L. Cheney, W. Salzburger, and N. J. Marshall. 2016. From crypsis to mimicry: changes in colour and the configuration of the visual system during ontogenetic habitat transitions in a coral reef fish. *Journal of Experimental Biology* **219**:2545-2558.
- Cortesi, F., Z. Musilová, S. M. Stieb, N. S. Hart, U. E. Siebeck, M. Malmstrøm, O. K. Tørresen, S. Jentoft, K. L. Cheney, N. J. Marshall, K. L. Carleton, and W. Salzburger. 2015. Ancestral duplications and highly dynamic opsin gene evolution in percomorph fishes. *Proceedings of the National Academy of Sciences* **112**:1493-1498.
- Darriba, D., G. L. Taboada, R. Doallo, and D. Posada. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature methods* **9**:772-772.
- Davenport, J., and M. Sayer. 1986. Ammonia and urea excretion in the amphibious teleost *Blennius pholis* (L.) in sea-water and in air. *Comparative Biochemistry and Physiology Part A: Physiology* **84**:189-194.
- de Busserolles, F., F. Cortesi, J. V. Helvik, W. I. Davies, R. M. Templin, R. K. Sullivan, C. T. Michell, J. K. Mountford, S. P. Collin, and X. Irigoien. 2017. Pushing the limits of photoreception in twilight conditions: the rod-like cone retina of the deep-sea pearlsides. *Science Advances* **3**:eaao4709.
- Dungan, S. Z., A. Kosyakov, and B. S. Chang. 2015. Spectral tuning of killer whale (*Orcinus orca*) rhodopsin: evidence for positive selection and functional adaptation in a cetacean visual pigment. *Molecular Biology and Evolution* **33**:323-336.
- Enright, J. M., M. B. Toomey, S.-y. Sato, S. E. Temple, J. R. Allen, R. Fujiwara, V. M. Kramlinger, L. D. Nagy, K. M. Johnson, Y. Xiao, M. J. How, S. L. Johnson, N. W. Roberts, V. J. Kefalov, P. F. Guengerich, and J. C. Corbo. 2015. Cyp27c1 red-shifts

- the spectral sensitivity of photoreceptors by converting vitamin A 1 into A 2. *Current Biology* **25**:3048-3057.
- Escobar-Camacho, D., E. Ramos, C. Martins, and K. L. Carleton. 2017. The Opsin Genes of Amazonian Cichlids. *Molecular Ecology* **26**:1343-4356.
- Fasick, J. I., and P. R. Robinson. 1998. Mechanism of spectral tuning in the dolphin visual pigments. *Biochemistry* **37**:433-438.
- Fuller, R. C., K. L. Carleton, J. M. Fadool, T. C. Spady, and J. Travis. 2005. Genetic and environmental variation in the visual properties of bluefin killifish, *Lucania goodei*. *Journal of Evolutionary Biology* **18**:516-523.
- Fuller, R. C., L. A. Noa, and R. S. Strellner. 2010. Teasing apart the many effects of lighting environment on opsin expression and foraging preference in bluefin killifish. *The American Naturalist* **176**:1-13.
- Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B. Couger, D. Eccles, B. Li, and M. Lieber. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* **8**:1494.
- Hofmann, C. M., N. J. Marshall, K. Abdilleh, Z. Patel, U. E. Siebeck, and K. L. Carleton. 2012. Opsin evolution in damselfish: convergence, reversal, and parallel evolution across tuning sites. *Journal of Molecular Evolution* **75**:79-91.
- Hofmann, C. M., K. E. O'Quin, N. J. Marshall, T. W. Cronin, O. Seehausen, and K. L. Carleton. 2009. The eyes have it: regulatory and structural changes both underlie cichlid visual pigment diversity. *PLoS biology* **7**:e1000266.
- Hsieh, S.-T. T. 2010. A locomotor innovation enables water-land transition in a marine fish. *PLoS ONE* **5**:e11197.
- Hundt, P. J., S. P. Iglésias, A. S. Hoey, and A. M. Simons. 2014a. A multilocus molecular phylogeny of combtooth blennies (Percomorpha: Blennioidei: Blenniidae): multiple invasions of intertidal habitats. *Molecular Phylogenetics and Evolution* **70**:47-56.
- Hundt, P. J., Y. Nakamura, and K. Yamaoka. 2014b. Diet of combtooth blennies (Blenniidae) in Kochi and Okinawa, Japan. *Ichthyological Research* **61**:76-82.
- Hundt, P. J., and A. M. Simons. 2018. Extreme dentition does not prevent diet and tooth diversification within combtooth blennies (Ovalentaria: Blenniidae). *Evolution* **72**:930-943.
- Hunt, D. M., K. S. Dulai, J. C. Partridge, P. Cottrill, and J. K. Bowmaker. 2001. The molecular basis for spectral tuning of rod visual pigments in deep-sea fish. *Journal of Experimental Biology* **204**:3333-3344.
- Hunt, D. M., M. W. Hankins, S. P. Collin, and N. J. Marshall. 2014. *Evolution of visual and non-visual pigments*. Springer.
- Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**:772-780.
- Kawano, S. M., and R. W. Blob. 2013. Propulsive forces of mudskipper fins and salamander limbs during terrestrial locomotion: implications for the invasion of land. *Integrative and Comparative Biology* **53**:283-294.
- Korenbrod, J. I., and R. D. Fernald. 1989. Circadian rhythm and light regulate opsin mRNA in rod photoreceptors. *Nature* **337**:454.
- Kumar, S., G. Stecher, and K. Tamura. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**:1870-1874.
- Liebman, P., and G. Entine. 1968. Visual pigments of frog and tadpole (*Rana pipiens*). *Vision Research* **8**:761-775.

- Lin, J.-J., F.-Y. Wang, W.-H. Li, and T.-Y. Wang. 2017. The rises and falls of opsin genes in 59 ray-finned fish genomes and their implications for environmental adaptation. *Scientific Reports* **7**:15568.
- Luehrmann, M., F. Cortesi, K. L. Cheney, F. De Busserolles, and N. J. Marshall. in review. Microhabitat partitioning correlates with eye morphology and opsin gene expression in coral reef cardinalfishes (Apogonidae)
- MacIver, M. A., L. Schmitz, U. Mugan, T. D. Murphey, and C. D. Mobley. 2017. Massive increase in visual range preceded the origin of terrestrial vertebrates. *Proceedings of the National Academy of Sciences* **114**:2374-2384.
- Mandic, M., K. A. Sloman, and J. G. Richards. 2009. Escaping to the surface: a phylogenetically independent analysis of hypoxia-induced respiratory behaviors in sculpins. *Physiological and Biochemical Zoology* **82**:730-738.
- Marshall, N. J., F. Cortesi, F. de Busserolles, U. E. Siebeck, and K. L. Cheney. 2018. Colours and colour vision in reef fishes: past, present and future research directions. *Journal of Fish Biology*:doi:<https://doi.org/10.1111/jfb.13849>
- Marshall, N. J., K. Jennings, W. N. McFarland, E. R. Loew, G. S. Losey, and W. L. Montgomery. 2003. Visual biology of hawaiian coral reef fishes. III. Environmental light and an integrated approach to the ecology of reef fish vision. *Copeia* **2003**:467-480.
- Miller, M. A., W. Pfeiffer, and T. Schwartz. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Pages 1-8 *in* Gateway Computing Environments Workshop (GCE), 2010. IEEE.
- Morgans, C. L., G. M. Cooke, and T. J. Ord. 2014. How populations differentiate despite gene flow: sexual and natural selection drive phenotypic divergence within a land fish, the Pacific leaping blenny. *BMC Evolutionary Biology* **14**:97.
- Morgans, C. L., and T. J. Ord. 2013. Natural selection in novel environments: predation selects for background matching in the body colour of a land fish. *Animal Behaviour* **86**:1241-1249.
- Musilova, Z., F. Cortesi, M. Matschiner, W. I. L. Davies, S. M. Stieb, F. de Busserolles, M. Malmstroem, O. K. Toerresen, J. K. Mountford, R. Hanel, K. S. S. Jakobsen, K. L. Carleton, S. Jentoft, J. N. Marshall, and W. Salzburger. 2018. Vision using multiple distinct rod opsins in deep-sea fishes. *bioRxiv*, 424895:doi:<https://doi.org/10.1101/424895>
- Ord, T. J., and G. M. Cooke. 2016. Repeated evolution of amphibious behavior in fish and its implications for the colonization of novel environments. *Evolution* **70**:1747-1759.
- Ord, T. J., T. C. Summers, M. M. Noble, and C. J. Fulton. 2017. Ecological release from aquatic predation is associated with the emergence of marine blenny fishes onto land. *The American Naturalist* **189**:570-579.
- Ord, T. J., and S. Tonia Hsieh. 2011. A highly social, land-dwelling fish defends territories in a constantly fluctuating environment. *Ethology* **117**:918-927.
- Palczewski, K., T. Kumasaka, T. Hori, C. A. Behnke, H. Motoshima, B. A. Fox, I. L. Trong, D. C. Teller, T. Okada, R. E. Stenkamp, M. Yamamoto, and M. Miyano. 2000. Crystal structure of rhodopsin: a G protein-coupled receptor. *Science* **289**:739-745.
- Platt, E. R., and T. J. Ord. 2015. Population Variation in the Life History of a Land Fish, *Alticus arnoldorum*, and the Effects of Predation and Density. *PLoS ONE* **10**:e0137244.
- Regan, K. S., M. G. Jonz, and P. A. Wright. 2011. Neuroepithelial cells and the hypoxia emersion response in the amphibious fish *Kryptolebias marmoratus*. *Journal of Experimental Biology* **214**:2560-2568.

- Rodd, F. H., A. Hughes Kimberly, F. Grether Gregory, and T. Baril Colette. 2002. A possible non-sexual origin of mate preference: are male guppies mimicking fruit? *Proceedings of the Royal Society of London. Series B: Biological Sciences* **269**:475-481.
- Ronquist, F., M. Teslenko, P. van der Mark, D. L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M. A. Suchard, and J. P. Huelsenbeck. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**:539-542.
- Rossi, G. S., A. J. Turko, and P. A. Wright. 2018. Oxygen drives skeletal muscle remodeling in an amphibious fish out of water. *Journal of Experimental Biology* **221**:jeb180257.
- Sandkam, B., B. Dalton, F. Breden, K. Carleton, and H. e. B. Fuller. 2018. Reviewing guppy color vision: integrating the molecular and physiological variation in visual tuning of a classic system for sensory drive. *Current Zoology* **64**:535-545.
- Savelli, I., I. N. Flamarique, T. Iwanicki, and J. S. Taylor. 2018. Parallel opsin switches in multiple cone types of the starry flounder retina: tuning visual pigment composition for a demersal life style. *Scientific Reports* **8**:4763.
- Sayer, M. D. 2005. Adaptations of amphibious fish for surviving life out of water. *Fish and Fisheries* **6**:186-211.
- Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. Mrosso, R. Miyagi, I. van der Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai, and N. Okada. 2008. Speciation through sensory drive in cichlid fish. *Nature* **455**:620-626.
- Shimizu, N., Y. Sakai, H. Hashimoto, and K. Gushima. 2006. Terrestrial reproduction by the air-breathing fish *Andamia tetradactyla* (Pisces; Blenniidae) on supralittoral reefs. *Journal of Zoology* **269**:357-364.
- Siebeck, U. E., and N. J. Marshall. 2007. Potential ultraviolet vision in pre-settlement larvae and settled reef fish - A comparison across 23 families. *Vision Research* **47**:2337-2352.
- Smith, C., I. Barber, J. Wootton Robert, and L. Chittka. 2004. A receiver bias in the origin of three-spined stickleback mate choice. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**:949-955.
- Smith, H. W. 1930. Metabolism of the lung-fish, *Protopterus aethiopicus*. *Journal of Biological Chemistry* **88**:97-130.
- Spady, T. C., J. W. Parry, P. R. Robinson, D. M. Hunt, J. K. Bowmaker, and K. L. Carleton. 2006. Evolution of the cichlid visual palette through ontogenetic subfunctionalization of the opsin gene arrays. *Molecular Biology and Evolution* **23**:1538-1547.
- Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**:1312-1313.
- Stieb, S. M., K. L. Carleton, F. Cortesi, N. J. Marshall, and W. Salzburger. 2016. Depth dependent plasticity in opsin gene expression varies between damselfish (Pomacentridae) species. *Molecular Ecology* **25**:3645-3661.
- Stieb, S. M., F. Cortesi, L. Sueess, K. L. Carleton, W. Salzburger, and N. J. Marshall. 2017. Why UV-and red-vision are important for damselfish (Pomacentridae): structural and expression variation in opsin genes. *Molecular Ecology* **26**:1323-1342.
- Sturla, M., P. Paola, G. Carlo, M. M. Angela, and U. B. Maria. 2002. Effects of induced aestivation in *Protopterus annectens*: a histomorphological study. *Journal of Experimental Zoology* **292**:26-31.
- Swamynathan, S. K., M. Crawford, W. G. Robison JR, J. Kanungo, and J. Piatigorsky. 2003. Adaptive differences in the structure and macromolecular compositions of the air and water corneas of the “four-eyed” fish (*Anableps anableps*). *The FASEB journal* **17**:1996-2005.

- Turko, A. J., D. Kultz, D. Fudge, R. P. Croll, F. M. Smith, M. R. Stoyek, and P. A. Wright. 2017. Skeletal stiffening in an amphibious fish out of water is a response to increased body weight. *Journal of Experimental Biology* **220**:3621-3631.
- Wald, G. 1968. The molecular basis of visual excitation. *Nature* **219**:800.
- White, E., D. Goncalves, J. C. Partridge, and R. F. Oliveira. 2004. Vision and visual variation in the peacock blenny. *Journal of Fish Biology* **65**:227-250.
- Wilt, F. H. 1959. The differentiation of visual pigments in metamorphosing larvae of *Rana catesbeiana*. *Developmental Biology* **1**:199-233.
- Wright, P. A., and A. J. Turko. 2016. Amphibious fishes: evolution and phenotypic plasticity. *Journal of Experimental Biology* **219**:2245-2259.
- Xu, B., and Z. Yang. 2013. PAMLX: a graphical user interface for PAML. *Molecular Biology and Evolution* **30**:2723-2724.
- Yang, Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution* **24**:1586-1591.
- Yang, Z., W. S. Wong, and R. Nielsen. 2005. Bayes empirical Bayes inference of amino acid sites under positive selection. *Molecular Biology and Evolution* **22**:1107-1118.
- Yokoyama, S. 2002. Molecular evolution of color vision in vertebrates. *Gene* **300**:69-78.
- Yokoyama, S. 2008. Evolution of dim-light and color vision pigments. *Annu. Rev. Genomics Hum. Genet.* **9**:259-282.
- You, X., C. Bian, Q. Zan, X. Xu, X. Liu, J. Chen, J. Wang, Y. Qiu, W. Li, and X. Zhang. 2014. Mudskipper genomes provide insights into the terrestrial adaptation of amphibious fishes. *Nature Communications* **5**:5594.
- Zander, C. 1974. Beziehungen zwischen Körperbau und Lebensweise bei Blenniidae (Pisces) aus dem Roten Meer. III. Morphologie des Auges. *Marine Biology* **28**:61-71.

Tables

Table 1. Summary of Salariini opsin gene variation

	<i>SWS2Aα</i>	<i>SWS2Aβ</i>	<i>RH2A-1</i>	<i>LWS</i>	<i>RH1</i>
Total number of nucleotides	1053	1056	1059	1074	1065
Total number of amino acids	350	351	353	357	354
Variable nucleotide sites	140	105	64	71	85
Indels	0	0	0	0	0
<u>Between Salariini species (amino acids)</u>					
Variable transmembrane sites	13(6)	10(2)	3(2)	10(3)	11(3)
Variable retinal binding pocket sites	1(0)	1(0)	0	1(1)	2(0)
N° of substitutions at known opsin tuning sites	0	4(0*)	0	4(1*)	1(1*)
Nonsynonymous substitutions (<i>dn</i>)	0.014	0.009	0.003	0.006	0.008
Synonymous substitution (<i>ds</i>)	0.205	0.147	0.093	0.107	0.110
<i>dn/ds</i>	0.068	0.061	0.032	0.056	0.073
<u>vs. <i>Parablennius parvicornis</i> (amino acids)</u>					
Variable transmembrane sites	22(10)	18(4)	10(2)	12(3)	14(4)
Variable retinal binding pocket sites	2(1)	1(0)	1(0)	1(1)	3(1)
Number of substitutions at known tuning sites	1(0*)	5(0*)	0	5(1*)	2(2*)

() substitution with changes in polarity; * substitutions at known tuning sites of gene in question.

668
669
670

Table 2. Summary of transcriptomes, opsin mapping, and proportional opsin gene expression. # raw reads refers to the total number of paired-end fragments. *RH1* = rod opsin, *SWS2* = short-wavelength sensitive 2, *RH2* = rhodopsin like 2, *LWS* = long-wavelength sensitive

RNA sequencing			Mapping # reads					Proportional opsin and <i>cyp27c1</i> expression %										
Transcriptome			Rod	Single cones (SC)		Double cones (DC)		Rod vs Cone		Cone opsin vs total cone expression				SC		DC		<i>Cyp27c1</i>
Species	ID	# raw reads	<i>RH1</i>	<i>SWS2Aα</i>	<i>SWS2Aβ</i>	<i>RH2A-1</i>	<i>LWS</i>	R	C	<i>SWS2A</i> <i>α</i>	<i>R</i> <i>H</i> 2	<i>L</i> <i>W</i> <i>S</i>		<i>SWS2A</i> <i>α</i>		<i>R</i> <i>H</i> 2	<i>L</i> <i>W</i> <i>S</i>	
<i>Alticus simplicirrus</i> French Polynesia Terrestrial	B7	20,291,698	199,410	2,338	2,144	19,738	8,713	85.7	14.3	7.2	6.5	60.1	26.2	52.2	47.8	69.7	30.3	0.15
	B8	15,763,272	147,622	1,433	1,358	13,712	9,508	84.9	15.1	5.6	5.3	53.0	36.2	51.4	48.6	59.4	40.6	0.13
	B9	22,851,082	211,302	2,597	1,721	22,119	16,189	83.1	16.9	6.2	4.1	52.1	37.6	60.2	39.8	58.1	41.9	0.14
	Mean	19,635,351	186,111	2,123	1,741	18,523	11,470	84.6	15.4	6.3	5.3	55.1	33.3	54.6	45.4	62.4	37.6	0.14
	Se	2,072,226	19,548	353	227	2,502	2371	0.8		0.5	0.7	2.5	3.6	2.8		3.7		0.01
<i>Alticus anjouanae</i> Seychelles Terrestrial	B10	22,500,526	289,057	1,785	1,391	13,544	6,791	92.4	7.6	7.7	6.0	57.8	28.6	56.3	43.7	66.9	33.1	0.06
	B11	37,810,316	450,726	2,341	2,417	20,911	14,142	91.8	8.2	5.9	6.1	52.8	35.2	49.3	50.7	60.0	40.0	0.11
	B12	6,118,452	80,619	483	613	4,297	2,834	90.7	9.3	5.9	7.5	52.5	34.1	44.1	55.9	60.6	39.4	0.18
	Mean	22,143,098	273,467	1,536	1,474	12,917	7,922	91.6	8.4	6.5	6.5	54.3	32.6	49.9	50.1	62.5	37.5	0.12
	Se	9,150,398	107,125	551	522	4,806	3,313	0.5		0.6	0.5	1.7	2.0	3.5		2.2		0.03
<i>Praealticus caesi</i> French Polynesia Mildly Amphibious	B4	19,353,526	226,133	1,262	1,882	7,221	2,620	94.5	5.5	9.8	14.6	55.7	19.9	40.2	59.8	73.7	26.3	0.12
	B5	19,211,400	155,934	875	1,466	8,382	7,093	89.7	10.3	5.0	8.3	47.3	39.5	37.4	62.6	54.5	45.5	0.15
	B6	27,900,670	259,074	4,217	8,959	36,687	39,094	74.3	25.7	4.8	10.2	41.5	43.6	32.1	67.9	48.8	51.2	0.03
	Mean	22,155,199	213,714	2,118	4,102	17,430	16,269	86.1	13.9	6.5	11.0	48.2	34.3	36.6	63.4	59.0	41.0	0.10
	Se	2,873,029	30,415	1,055	2,431	9,634	11,485	6.1		1.6	1.9	4.1	7.3	2.4		7.5		0.04

<i>Entomacrodus striatus</i> Seychelles Amphibious	B13	27,001,052	220,729	4,115	21,024	93,036	104,454	49.6	50.4	1.9	9.5	42.0	46.6	16.4	83.6	47.5	52.5	0.02
	B14	30,325,336	425,996	4,523	42,449	168,400	196,605	50.6	49.4	1.1	10.4	41.1	47.4	9.7	90.3	46.5	53.5	0.01
	B15	20,757,818	223,445	2,576	28,409	111,942	112,151	46.5	53.5	1.0	11.2	44.1	43.6	8.3	91.7	50.3	49.7	0.02
	Mean	26,028,069	290,057	3,738	30,627	124,459	137,737	48.9	51.1	1.3	10.4	42.4	45.8	11.5	88.5	48.1	51.9	0.02
	Se	2,804,423	67,974	593	6,284	22,638	29,518	1.2	0.3	0.5	0.9	1.1	2.5	1.1	0.00			
<i>Istiblennius lineatus</i> French Polynesia Mildly Amphibious	B1	11,510,186	127,409	401	5,351	18,821	13,210	77.0	23.0	1.1	14.3	50.0	34.6	7.0	93.0	59.1	40.9	0.19
	B2	6,765,526	56,929	273	2,728	11,077	8,243	71.6	28.4	1.2	12.3	49.9	36.6	9.1	90.9	57.7	42.3	0.19
	B3	3,343,260	28,355	154	1,310	5,102	3,810	73.0	27.0	1.5	12.7	49.4	36.4	10.5	89.5	57.6	42.4	0.52
	Mean	7,206,324	70,898	276	3,130	11,667	8,421	73.9	26.1	1.3	13.1	49.8	35.9	8.9	91.1	58.1	41.9	0.30
	Se	2,367,868	29,435	71	1,184	3,971	2,715	1.6	0.1	0.6	0.2	0.6	1.0	0.5	0.11			
<i>Istiblennius lineatus</i> Seychelles Mildly Amphibious	B16	30,354,518	367,265	1,628	33,763	125,652	101,135	58.1	41.9	0.6	13.0	48.2	38.2	4.6	95.4	55.8	44.2	0.03
	B17	56,930,548	607,878	2,689	60,877	213,029	178,091	57.0	43.0	0.6	13.5	47.1	38.8	4.2	95.8	54.8	45.2	0.04
	B18	7,165,396	89,373	438	9,546	33,658	27,844	55.3	44.7	0.6	13.5	47.3	38.6	4.4	95.6	55.1	44.9	0.02
	Mean	31,483,487	354,839	1,585	34,729	124,113	102,357	56.8	43.2	0.6	13.3	47.5	38.5	4.4	95.6	55.2	44.8	0.03
	Se	14,377,048	149,808	650	14,826	51,786	43,377	0.8	0.0	0.2	0.3	0.2	0.1	0.3	0.01			
<i>Salarias fasciatus</i> Australia Aquatic		44,650,228	795,112	4,592	25,534	83,252	84,588	79.9	20.1	2.3	13.0	42.3	42.4	15.3	84.7	50.0	50.0	0.09

672 Figures

Fig. 1. Bayesian consensus phylogeny for vertebrate opsin genes. The Salariini retinal transcriptomes contained seven opsin genes. One rod opsin (*RH1*) and six cone opsin genes belonging to three different cone opsin classes: short-wavelength-sensitive 2 (*SWS2Aα*, *SWS2Aβ*), rhodopsin-like 2 (*RH2A-1*, *RH2A-2*, *RH2B*), and long-wavelength-sensitive (*LWS*). Note that *RH2A-2* was found at very low expression levels in three out of the six Salariini species, while *RH2B* was lowly expressed in *Salarias fasciatus* alone (also see Fig. 2, Table 2). Dark shading indicates genes from various Salariini species, light shading genes from sister species (Blenniidae). Black and grey spheres indicate Bayesian posterior probabilities > 0.9 and 0.7, respectively.

Fig. 2. Salariini species phylogeny and associated opsin gene expression. a) Salariini species (n = 3 per species, except for *S. fasciatus* where n = 1) expressed two paralogs of the short-wavelength-sensitive 2A opsin gene (*SWS2Aα*, *SWS2Aβ*), and b) one rhodopsin-like 2 (*RH2A-1*) and a long-wavelength-sensitive (*LWS*) opsin gene. The mean proportional gene expression was similar between species independent of habitat or sampling location, except for the *SWS2A* paralogs in the clade containing *Pr. caesioides* and the two *Alticus* spp. Displayed are the mean expression values separated by cone specificity (Hunt et al., 2014). For details on individual expression values and gene expression in relation to total cone and total opsin expression see Table 2. A red circle indicates terrestrial *Alticus* spp., which use red dorsal fins for aggressive and courtship displays (Bhikajee and Green 2002, Shimizu et al. 2006, Ord and Tonia Hsieh 2011). AQ = aquatic, AP = amphibious, M-AP = mildly-amphibious, T = Terrestrial, definitions and phylogeny as per (Ord & Cooke, 2016). † Seychelles, ‡ French Polynesia, § Heron Island (Australia).