Co-allocation Action of Azospirillum brasilense Inoculation and Seed Priming for Improving Germination and Seedling Growth in Aged Grass Seeds

Short title: *Azospirillum brasilense* Inoculation and Aged Grass Seeds

Xv Liu¹, Zhao Chen¹, Yani Gao¹, Qian Liu¹, Wennan Zhou¹, Tian Zhao¹, Wenbo Jiang¹, Xuewen Cui¹, Jian Cui² and Quanzhen Wang^{1*}

¹Department of Grassland Science, College of Animal Sci. and Techn., Northwest A&F University, Yangling, Shaanxi Province, China.

²Department of Plant Science, College of Life Science, Northwest A&F University, Yangling, Shaanxi Province, China.

* Corresponding author:

Prof. Dr. Wang Quanzhen

E-mail: wangquanzhen191@163.com

Abbreviations: DAS, day after sowing; Gi, germination index; GP, germination percentage; RL, root length; SL, shoot length; SVi, the seed vigor index.

No. of figures: **7** (Colours were not in published paper vision but electric on line);

No. of tables: **4**;

No. of supplementary tables: **3**;

1 Abstract

2 Germination of seeds during the transportation or after prolonged storage naturally and inevitably
3 decreases because of ageing, but germination potential can be partially restored with seed priming
4 treatments. A novel attempt was made to investigate the effects of combined treatments and to
5 optimize the conditions for naturally aged seeds of tall fescue (*Festuca arundinacea* Schreb.),
6 orchardgrass (*Dactylis glomerata* L.) and Russian wild rye (*Psathyrostachys juncea* (Fisch.)
7 Nevski) using an orthogonal activity level experimental design [factor A: *Azospirillum brasilense*
8 concentration, factor B: three seed priming treatments (H₂O, MgSO₄ and H₂O₂) and factor C:
9 different priming times]. Multivariate regression model analysis was applied to determine the
10 interactive effects of pairwise factors (A and C) and to optimize experimental conditions. The
11 results showed that the mixed treatments positively affected seed germination and seedling growth.
12 The three seed priming treatments were the dominant factors for germination promotion, whereas
13 the bacterial concentration had the largest effect on seedling growth, especially root elongation.
14 The combined results of all determined attributes showed that *A. brasilense* bio-priming with
15 H₂O₂ priming constituted the optimal combination. The optimal bacterial concentration of *A.*
16 *brasilense* and the time of seed priming were 52.3×10^6 colony forming units (CFU) mL⁻¹ and
17 17.0 h, respectively. We also discussed the physiological mechanisms that could repair aged seeds:
18 the malondialdehyde content and antioxidant enzyme (superoxide dismutase, peroxidase, catalase
19 and ascorbate peroxidase) activities were affected.

20 Introduction

21 The successful completion of seed germination and the establishment of a normal seedling
 22 determine the survival and propagation of plant species ^[1]. However, seeds inevitably and
 23 irreversibly deteriorate and age during storage. This deterioration and aging lead to substantial
 24 economic losses and decreased genetic diversity ^[1]. Seed aging can result in reduced germination,
 25 seedling abnormalities and even total loss of seed viability, which are correlated with disturbances
 26 in various cellular metabolic and biochemical processes, including cell membrane disintegration,
 27 mitochondrial dysfunction, enzyme inactivation, Amadori and Maillard reactions, telomere
 28 shortening and damage to biological key players such as carbohydrates, DNA and RNA ^[2].
 29 Currently, the excessive production of reactive oxygen species (ROS; H₂O₂, O₂⁻, etc.) and their
 30 attack on lipids and proteins is thought to be the main cause of seed aging ^[2]. Plants are well
 31 equipped to minimize the damaging effects of seed aging via their antioxidant defense system,
 32 which consists of superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT), etc.
 33 However, it is not always possible for the antioxidant defense system to scavenge excessive ROS
 34 by themselves. Seeds always have greater lipid contents than other plant tissues, particularly
 35 forage seeds such as those of tall fescue (*Festuca arundinacea* Schreb.) and Russian wild rye
 36 (*Psathyrostachys juncea* (Fisch.) Nevski) ^[3, 4]. Besides, it is difficult to control the ambient
 37 temperature, humidity and seed moisture content during storage. These factors can hasten the
 38 aging process of seeds ^[5, 6]. Previous reports have shown that under ambient storage conditions in
 39 eastern Serbia in 2009, the reduction in germination of Italian ryegrass (*Lolium multiflorum* L.)
 40 and timothy (*Phleum pratense* L.) seeds occurred from 180th and 330th days after harvest,
 41 respectively; the germination rates of *Jatropha curcas* L. and lettuce (*Lactuca sativa*) seeds stored

for one year decreased by 89.29 and 47.12%, respectively ^[7-9]. Therefore, exploring effective and pragmatic techniques for seed recovery from aging is urgently warranted for plant germplasm conservation and better agriculture development.

Recent attempts to repair aged seeds through conventional strategies have been met with limited advancement due to the complex nature of seed aging. Behtari and Tilaki reported that the germination percentage of osmoprimed tall fescue seeds stored at 25°C for 1 year decreased significantly, most reducing by 60.1% ^[10]. In this scenario, the possibility of the application of plant growth-promoting rhizobacteria (PGPR) has emerged as a hot research topic. Various studies have reported that PGPR are capable of alleviating abiotic stress and enhancing growth in plants ^[11, 12]. *Azospirillum* is versatile and one of the most widely studied genus of PGPR (e.g. *Azotobacter* sp., *Bacillus* sp., *Pseudomonas* sp. and *Herbaspirillum* sp.). It can exert a positive influence on plant growth via complex regulatory networks involved in biological nitrogen fixation, enzyme activation, the production of growth regulatory substances (auxins, gibberellins, polyamines, etc.) and the regulation of the expression of genes in roots ^[13, 14].

Seed bio-priming with PGPR can allow enough numbers of bacteria to infect seeds, which is the most promising method for the application of PGPR ^[15]. Numerous studies have been conducted using *A. brasilense* bio-priming or a consortium of PGPR strains on plants against diverse abiotic stresses, including salinity, osmotic stress and suboptimal temperature ^[16, 17]. Currently, the fact that the combination of bio-priming with PGPR and another type of seed priming approach (e.g., hydropriming, osmopriming, or redox priming) results in better plant performance is attracting increasing amounts of attention ^[18, 19]. These combinations are more practical and simpler than

co-inoculation of PGPR strains, for which direct antagonism, competition and other factors must be considered. However, the effect of bio-priming via *Azospirillum* on aged seeds has been rarely evaluated, and no reports are available on the repair mechanisms of this genus or optimal combinations with other priming treatments on aged seeds.

Tall fescue, orchardgrass (*Dactylis glomerata* L.) and Russian wildrye are considered to be of great importance among perennial forages due to their high productivity, nutritive value, durability and tolerance to unfavorable conditions. However, poor storability constrains their widespread application [20]. Therefore, the goals of this study were to investigate the effects of *A. brasilense* and various seed priming approaches, including hydropriming, osmopriming and redox priming on the germination and seedling growth of naturally aged forage seeds of tall fescue, orchardgrass and Russian wildrye stored for different years at room temperature and to identify the optimal combination using an orthogonal matrix design and model analysis. Aging-related physiological changes were also examined to better understand the mechanisms of co-treatments on the aged seeds. This study could potentially provide new insights into identifying novel PGPR as elicitors of aging-related repair.

Materials and Methods

Bacterial Culture, Seed Materials and Experimental Design

Azospirillum brasilense Yu62 was kindly provided by the College of Life Science, Northwest A&F University. The Yu62 culture was prepared on agar-Congo red medium at 37°C for 3 days [21], a single colony of Yu62 was picked and transferred to nitrogen-free semisolid malate (NFb)

liquid medium containing 0.1% NH₄Cl, and incubated for 48 h at 30°C under vigorous agitation (120 rpm). The cells were harvested by centrifugation (8142 × g, 10 min) and were resuspended in autoclaved 0.12 M phosphate-buffered saline (pH 7.2). The prepared bacterial suspensions were adjusted to 10⁸ colony forming units (CFU) mL⁻¹, 10⁷ CFU mL⁻¹ and 10⁶ CFU mL⁻¹ by adjusting OD values and serially diluting.

Seeds of tall fescue, orchardgrass and Russian wild rye were acquired from the China Agriculture University Grassland Research Station located at the Hexi Corridor in Jiuquan, Gansu Province, China. All seeds were stored for 1 to 10 years in paper bags at room temperature in the same laboratory. Six groups of the three grass species, namely, tall fescue 2015 the year harvested, tall fescue 2014, tall fescue 2006, Russian wild rye 2009, orchardgrass 2013 and orchardgrass 2006 that had initial seed moisture contents of 9.3%, 10.1%, 8.2%, 7.9%, 7.7% and 7.9%, respectively, composed the nested experiments. Each group was fixed in an [L₉ (3⁴)] orthogonal matrix design for seed priming treatments.

Three factors were selected: the bacterial concentration of *A. brasilense* (factor A); the seed priming approaches viz., hydropriming (autoclaved distilled water), osmopriming (0.37 mol/L MgSO₄), and redox priming (H₂O₂-priming: 0.1% H₂O₂) were selected based on numerous previous studies and pre-experiments (factor B)^[4, 8]; and priming time (factor C). Each factor was assigned three levels, and the nine treatment combinations of different parameters were laid out in an [L₉ (3⁴)] orthogonal matrix design replicated thrice (Table 1). Additionally, a control was included without any priming treatment (Table 1).

103

104 **Table 1. The L_9 (3^4) orthogonal matrix in the experimental design.**

Treatments	A: the concentration of <i>A. brasilense</i> (CFU mL ⁻¹)		B: seed priming treatment	C : priming time (h)
	matrix	Code (X_1)	matrix	matrix
T1	1 (10^6)	1	1 (hydropriming)	3 (24)
T2	2 (10^7)	10	1	1 (12)
T3	3 (10^8)	100	1	2 (18)
T4	1	1	2 (MgSO ₄ -priming)	2
T5	2	10	2	3
T6	3	100	2	1
T7	1	1	3 (H ₂ O ₂ -priming)	1
T8	2	10	3	2
T9	3	100	3	3

105

106 Seed Treatments and Germination Tests

107 Seeds were surface-sterilized with 2% NaOCl solution for 5 min followed by 70% ethanol for 10
 108 min, after which the seeds were rinsed five times with sterile distilled water. According to the
 109 orthogonal design [L_9 (3^4)], each group of seeds that were coated with gauze were primed at
 110 ambient temperature. Inoculation was performed by soaking the sterilized seeds in prepared
 111 bacterial suspensions or only sterile phosphate-buffered saline for control seeds, for 2 h [15]. After

hydropriming, osmopriming, or redox priming followed by *A. brasilense* inoculation, the primed seeds were air-dried to reduce their moisture content. Three replicates were used for each treatment and 50 healthy seeds from each replication were evenly spaced on two layers of moistened Whatman No. 1 filter paper in each 150-mm Petri dish. Equal volumes of distilled water were added daily. Petri dishes were placed in a climate incubator (RQH-250, Jing Hong Laboratory Instrument Co., Ltd, Shanghai, China) that maintained an alternating diurnal regimen of 16 h of light at $25 \pm 0.5^\circ\text{C}$ and 8 h of dark at $20 \pm 0.5^\circ\text{C}$ at 85% relative humidity for 28 days. Seed germination was recorded daily, and the seed was considered germinated when its radicle extended at least 0.5 mm. The shoot and root lengths of 10 randomly selected seedlings from each replication were measured in accordance with the rules recommended by the International Seed Testing Association (ISTA) at 28th day after sowing (DAS). When the total number of seedlings was fewer than 10, all seedlings were measured. To more precisely reflect the vigor of seeds, the germination index (Gi) was estimated as follows:

$$\text{Germination index (Gi)} = \sum_{i=1}^t Gt / Dt \quad (i = 1, 2, \dots, t)$$

where Gt is the number of germinated seeds within a day, Dt is the corresponding number of germination days and t is the number of total germination period (28 days).

The seed vigour index (SVi) was determined using the following formula:

$$\text{SVi} = \text{Gi} \cdot R$$

where R is length of root.

Determination of Physiological Parameters in Aged Grass

Seeds

Lipid peroxidation was assessed in terms of the malondialdehyde (MDA) content in fresh seedlings according to the thiobarbituric acid protocol [22]. Each group of sample containing 0.5 g crumbled seedlings was mixed with 5 mL trichloroacetic acid (TCA; 0.5%) and was centrifuged at $10,000 \times g$ for 25 min. For every 1 mL of the aliquot of the supernatant, 4 mL of 20% TCA containing 0.5% thiobarbituric acid was added. The mixture was heated at 95 °C for 30 min and then the tubes were cooled immediately in an ice-bath. After the tube was centrifuged at $10\,000 \times g$ for 10 min, the absorbance of supernatant was measured at 450, 532, and 600 nm. The MDA content was expressed as $\mu\text{mol} \cdot \text{g}^{-1} \text{FW}$.

The activities of the antioxidant enzymes SOD, POD, CAT and ascorbate peroxidase (APX) in fresh seedlings were determined. The extraction of antioxidant enzymes was carried out on seedling samples in accordance with the procedures described by Zhang and Nan, with minor modifications [23]. Each group of seedlings (0.5 g) was homogenized in 5 mL of 50 mM phosphate buffer (pH 7.8 for SOD and POD containing 1% (w/v) polyvinylpyrrolidone, pH 7 for CAT and APX containing 0.1 mM Na_2EDTA). The homogenate was centrifuged at $10000 \times g$ for 15 min at 4°C, after which the supernatant was collected for the determination of enzyme activities. The activities of SOD, POD, CAT and APX were analyzed as reported by Zhang and Nan [23].

Statistical Analysis

The data were statistically analyzed by means of variance analysis (ANOVA) using the software

SPSS version 20.0 and were compared using the Duncan test ($P < 0.05$).

The below-mentioned analyses and graphical procedures were performed using SAS version

8.2. The germination percentage (GP) was simulated using the following logistic model [24, 25]:

$$Y = \frac{c}{1 + e^{a - b \cdot X}}$$

where a , b , and c are constants and X represents the germination days. The

models were significant at $P < 0.05$.

For the generic results, the bacterial concentration of *A. brasilense* and seed priming time

(factors A and C) were denoted by X_1 and X_3 . The dependent variables were denoted by Y_i , which

included GP, G_i , SV_i , root length, shoot length, seedling length, the shoot/root ratio (the index of

seed vigor during aging), MDA content, and the activity of SOD, POD, CAT and APX. These

dependent variables were approached and analyzed via two-variable (X_1 and X_3) quadratic

regression models described as follows:

$$Y = \sum_{i=1}^2 (\beta_{i \times j + 1} X_i^j) + u \quad (i = 1, 2; j = 1, 2)$$

where β is a constant. The critical values of X_1 and X_3 were obtained from the quadratic models

that were significant at $P < 0.05$.

Results

The effect of co-treatment on the vigor of aged seeds

None of the seeds of tall fescue 2006 germinated in the whole nested experiment, only 2-4% of

seeds germinated in treatments 1-4 (T1-T4) and T7 for orchardgrass 2006 (data not shown). The

models regarding germination dynamics (Fig 1) showed that the germination of T1-T5, and T8 in

tall fescue 2015 occurred earlier than it did for the control (CK), and the greatest germination percentage (28 DAS) occurred in T1 (98.7%). The germination from all treatments occurred earlier than it did from the controls for both orchardgrass 2013 and Russian wildrye 2009. The other 8 treatments increased germination of orchardgrass 2013 except for T3, the germination percentages from T2 and T6 (92.0%) were the highest. The germination percentages of all treatments were higher than that of the CK, the highest germination percentage was in T1 (62.7%) among them in Russian wildrye 2009. Whereas, the co-treatments had a negative effect on germination of tall fescue 2014 except for T7 (72.0%) compared with the CK. But the germination started earlier in T2 and T7. The lowest germination percentages of tall fescue 2015, tall fescue 2014 and orchardgrass 2013 occurred in T9, T3 and T3, respectively. While the lowest germination percentage of Russian wildrye 2009 occurred in T9, which increased by 2.7 times compared with that of the CK.

Fig 1. The curves of logistic models for germination percentage under the combined treatments for tall fescue 2015 (A), 2014 (B), orchardgrass 2013 (C) and Russian wild rye 2009 (D). From T1 to T9 are the treatments, and T10 is the control.

The range analyses revealed that factor C (seed priming time) exhibited the largest effect on germination percentage among three factors, with the exception that factor B (seed priming approaches) had the highest range value (R value; 4.0) for tall fescue 2015 (Table 2). And the

most effective factors on the Gi and SVi of tall fescue 2015, orchardgrass 2013 and Russian wildrye 2009 were factors B, A (the bacterial concentration) and C, individually, which were the same for every species. The most important factors on the Gi and SVi of tall fescue 2014 were factors C and B, which were different. The analysis of variance indicated that the germination percentage was significantly ($P < 0.05$) affected by both grass species and factor C, and their interactive effects were also significant ($P < 0.05$) both in pairwise factors and among three factors, with the exceptions of the grass species $\times$ factor A and grass species $\times$ factor B interaction (Table 3). Both factors B and C significantly ($P < 0.01$) affected the Gi and SVi in different-aged seeds, also their interactive effect was significant ($P < 0.05$).

Table 2. Average responses of each level and the range analyses of germination percentage (GP), germination index (Gi), seed vigor index (SVi), root (RL) and shoot (RL) length in tall fescue 2015, tall fescue 2014, orchardgrass 2013 and Russian wild rye 2009.

Treatment	Tall fescue 2015			Tall fescue 2014			Orchardgrass 2013			Russian wildrye 2009		
	A(CFU/ml)	B	C(h)	A(CFU/ml)	B	C(h)	A(CFU/ml)	B	C(h)	A(CFU/ml)	B	C(h)
GP (%)												
Level I ^a	93.78	95.56	88.44	88.00	89.99	37.33	41.75	30.88	94.00	62.89	61.56	64.82
Level II ^a	92.89	94.67	87.78	87.99	86.67	41.17	36.67	28.47	92.67	59.11	55.48	57.11
Level III ^a	95.11	91.56	87.99	88.22	87.56	28.64	28.71	47.78	95.11	57.60	62.56	57.67
R ^m	2.22	4.00	0.67	0.24	3.32	12.52	13.04	19.30	2.44	5.29	7.08	7.71
Order ^a	3	1	2	3	1	3	2	1	2	3	2	1
Gi												
Level I ^a	68.30	84.67	76.90	72.26	73.10	29.00	30.50	20.86	74.08	53.37	53.95	59.30
Level II ^a	72.16	72.85	67.01	70.56	68.88	30.21	28.39	20.44	68.29	48.91	42.80	43.53
Level III ^a	75.84	58.79	67.66	68.75	69.59	19.42	19.74	37.33	73.94	47.93	53.47	47.39
R ^m	7.54	25.89	9.89	3.51	4.22	10.79	10.75	16.89	5.79	5.43	11.15	15.77
Order ^a	2	1	1	3	2	2	3	1	3	3	2	1
SVi												
Level I ^a	446.87	558.92	433.14	432.53	432.59	205.08	207.51	158.01	490.92	420.57	433.19	481.89
Level II ^a	482.40	499.92	376.99	405.79	388.55	205.92	195.94	142.77	440.31	392.52	334.05	362.45
Level III ^a	501.74	372.18	413.19	385.00	402.18	141.18	148.72	251.40	499.79	416.44	462.29	385.20
R ^m	54.87	186.74	56.15	47.52	44.03	64.74	58.79	108.63	59.48	28.05	128.24	119.44
Order ^a	3	1	1	2	3	2	3	1	2	3	1	2
RL												
Level I ^a	4.85	4.74	2.70	4.18	4.03	2.14	2.10	2.31	6.24	5.17	5.84	5.28
Level II ^a	5.49	4.69	4.90	3.73	4.24	2.46	2.01	2.27	4.45	7.11	5.72	6.00

Level III ^a	5.39	6.30	3.80	3.49	3.13	2.20	2.69	2.23	5.04	5.17	5.89	6.17
R ^m	0.64	1.61	2.20	0.68	1.11	0.32	0.68	0.08	1.79	1.94	0.17	0.89
Order ^a	3	2	1	3	2	2	1	3	1	1	3	2
SL												
Level I ^a	6.53	6.61	5.63	5.99	5.92	7.18	6.74	7.61	6.63	7.74	8.03	8.17
Level II ^a	6.67	6.81	5.62	5.76	5.64	6.98	7.09	6.91	6.42	7.99	7.60	8.15
Level III ^a	6.55	6.33	6.10	5.59	5.78	7.26	7.59	6.90	6.70	8.71	8.81	8.11
R ^m	0.14	0.48	0.48	0.39	0.28	0.27	0.85	0.71	0.28	0.98	1.20	0.06
Order ^a	3	1	1	2	3	3	1	2	2	2	1	3
SOD												
Level I ^a	18.86	14.90	2.82	4.16	2.72	20.42	30.24	25.38	22.18	16.07	10.64	16.18
Level II ^a	13.83	20.62	5.13	2.28	12.81	25.01	30.33	33.60	17.65	14.37	12.84	5.35
Level III ^a	21.92	19.21	16.36	12.04	7.77	35.67	21.62	22.34	15.71	10.02	16.85	18.14
R ^m	8.09	5.72	13.54	9.77	10.09	15.26	8.70	11.26	6.46	6.05	6.21	12.79
Order ^a	1	3	1	3	2	1	3	2	2	3	2	1
CAT												
Level I ^a	56.67	164.93	329.03	421.44	370.68	423.31	224.05	465.16	179.71	158.47	59.18	174.32
Level II ^a	181.86	168.77	318.00	241.62	310.00	133.30	208.48	125.16	138.60	141.53	174.17	102.34
Level III ^a	171.23	76.06	280.55	221.45	246.90	489.45	613.52	455.72	91.44	120.14	186.79	143.48
R ^m	125.19	92.71	48.48	200.00	123.78	356.15	405.04	340.00	88.27	38.32	127.61	71.97
Order ^a	1	2	3	1	2	2	1	3	3	3	1	2

204 † A, B and C represent the concentration of *Azospirillum brasilense*, the seed priming treatment and priming time.

205 ‡ⁿ Average response of each level regarding experimental indicators.

206 §^m R means the average range between three average responses of each level regarding experimental indicators.

207 ¶* The ordinal numeral for the range sequence of the three factors in decreasing order.

208

209 Seedling growth

210 Compared with the CK, T7 resulted in the biggest increase in root length, and the shoot length
 211 increased in T5 and T6 in tall fescue 2015 group (Fig 2). The root length in T8 was the longest
 212 significantly, and the shoot lengths in both T8 and T9 were markedly improved in tall fescue
 213 2014. In group of orchardgrass 2013, the root and shoot lengths were increased by T3 and T6,
 214 respectively. While in Russian wildrye 2009, the changes of root elongation was not significant.
 215 The root length was the largest in T8, the shoot length in T6 was significantly higher than that of
 216 the CK.

217

218 **Figure 2. Log ratio of root length (A) and shoot length (B).**

219 * indicates the difference was significant between the treatment and the control at $P < 0.05$.

220

221 Factor A markedly affected the root length, and the effect of factor A was the largest for tall

222 fescue 2014 and orchardgrass 2013 (Tables 2 and 3). The interactive effects were also significant
 223 ($P < 0.01$) on root length, except for the factors $A \times B$. Although factors A, B and C did not
 224 significantly influence the shoot length, their pairwise interactive effects were significant ($P <$
 225 0.05; Table 3).

226 **Table 3. Variance analyses for the model, the effects of each experimental factors and their**
 227 **interaction effects on germination percentage, gemination index, seed vigor index, root, and**
 228 **shoot length. Columns A, X_1 , X_2 and X_3 represent grass species, bacteria concentration,**
 229 **seed priming and priming duration, respectively.**

Source		Gemaiaon percentage		Gemaiaon index		Seed vigor index		Root length		Shoot length	
(factors)	DF	F Value	Pr>F	F Value	Pr>F	F Value	Pr>F	F Value	Pr>F	F Value	Pr>F
GS	3	327.95	<.001	184.96	<.001	110.32	<.001	40.25	<.001	53.18	<.001
X_1	2	1.73	0.184	2.12	0.126	1.44	0.243	12.99	<.001	2.95	0.058
X_2	2	2.31	0.105	12.92	<.001	12.30	<.001	2.51	0.088	1.42	0.248
X_3	2	4.62	0.013	7.18	0.001	5.41	0.006	0.85	0.433	1.63	0.203
GS* X_1	6	1.63	0.151	2.85	0.014	3.23	0.007	2.40	0.035	1.02	0.420
GS* X_2	6	1.97	0.080	5.52	<.001	3.91	0.002	2.01	0.074	3.16	0.008
GS* X_3	6	4.33	0.001	4.73	<.001	1.79	0.112	3.33	0.006	0.63	0.706
X_1 * X_2	4	4.70	0.002	7.91	<.001	9.80	<.001	2.04	0.096	2.60	0.042
X_1 * X_3	4	3.55	0.010	10.78	<.001	13.24	<.001	2.87	0.028	2.49	0.049
X_2 * X_3	4	3.26	0.016	5.38	0.001	7.81	<.001	8.11	<.001	3.26	0.016
GS* X_1 * X_2	12	3.51	<.001	2.93	0.002	1.35	0.206	1.96	0.036	1.41	0.177
GS* X_1 * X_3	12	2.32	0.013	3.32	0.001	2.42	0.010	2.646	0.005	2.68	0.004

GS*X ₂ *X ₃	12	2.15	0.022	1.99	0.036	2.08	0.028	2.18	0.020	1.61	0.106
Model	39	29.63	<.001	19.73	<.001	13.54	<.001	7.75	<.001	6.29	<.001
R ²		0.94		0.91		0.87		0.79		0.75	

230

231 **Characterization of physiological responses**

232 The MDA content of CK in Russian wildrye 2009 was significantly ($P < 0.05$) higher than those
233 of CKs in tall fescue 2015, tall fescue 2014 and orchardgrass 2013. And the SOD activity of CK in
234 Russian wildrye 2009 was noticeably lower than those in the three other controls. These results
235 indicated that the seed of Russian wildrye 2009 was the most severely damaged by aging (Fig 3A;
236 S1 Table). In all treatments, the MDA content sharply decreased relative to that of the CKs in
237 Russian wildrye 2009 and tall fescue 2015. The lowest MDA content in tall fescue 2014 and
238 orchardgrass 2013 occurred in T6 and T7 (Fig 3A). The changing trends regarding the MDA
239 content and SOD activity were opposite in Russian wildrye 2009 and tall fescue 2014 (T1, T4, T5
240 and CK). The highest activities of SOD and CAT were observed in T6 for tall fescue 2015 and in
241 T7 for tall fescue 2014. While the highest SOD and CAT activities were in T3 and T2 for
242 orchardgrass 2013, and were in T8 and T9 for Russian wildrye 2009, respectively (Fig 3A, B and
243 D). The highest POD values for tall fescue 2015 and Russian wildrye 2009 were found in T3 and
244 T2, separately (Fig 3C).

245

246 **Fig 3. MDA content (A), activity of SOD (B), POD (C) and CAT (D) of tall fescue (2015 and 2014),**

orchardgrass (2013) and Russian wildrye (2009). Data are presented as mean $\pm$ SD. The different lowercase letters indicate significant differences at $P < 0.05$ within the years among the treatments.

Factor A and the interactive effects significantly ($P < 0.05$) influenced SOD activity, and factor A was the most important factor for those of tall fescue 2015, orchardgrass 2013 and Russian wildrye 2009 (Tables 2 and 4). The experimental factors did not significantly affect the MDA content, except seed species (Table 4). The experimental factors markedly affected the POD and CAT activities with the exceptions of factor A, $A \times B$, $B \times C$ and grass species $\times A \times B$ on the POD activity as well as factors A and B on the CAT activity.

Table 4. Variance analyses for the model, the effects of each experimental factors and their interaction effects on MDA, POD, SOD and CAT.

Source		MDA		POD		SOD		CAT	
(factors)	DF	F Value	Pr>F	F Value	Pr>F	F Value	Pr>F	F Value	Pr>F

GS	3	72.064	<.001	36.214	<.001	36.349	<.001	16.816	<.001
X ₁	2	0.947	0.392	2.871	0.063	14.803	<.001	2.410	0.096
X ₂	2	0.277	0.759	9.357	<.001	2.649	0.077	1.588	0.211
X ₃	2	0.144	0.866	3.445	0.037	1.474	0.235	4.796	0.011
GS*X ₁	6	1.259	0.286	3.294	0.006	6.447	<.001	4.941	<.001
GS*X ₂	6	1.302	0.266	3.333	0.006	3.063	0.009	7.143	<.001
GS*X ₃	6	1.330	0.254	2.299	0.042	8.441	<.001	3.490	0.004
X ₁ *X ₂	4	0.926	0.453	2.482	0.0502	6.727	<.001	7.216	<.001
X ₁ *X ₃	4	0.993	0.416	5.438	0.001	7.314	<.001	5.612	<.001
X ₂ *X ₃	4	1.328	0.267	2.195	0.077	13.392	<.001	6.023	<.001
GS*X ₁ *X ₂	12	1.103	0.369	1.771	0.067	8.083	<.001	4.525	<.001
GS*X ₁ *X ₃	12	1.089	0.381	2.288	0.015	5.394	<.001	6.351	<.001
GS*X ₂ *X ₃	12	1.067	0.398	2.268	0.016	7.086	<.001	5.250	<.001
Model	39	8.009	<.0001	6.827	<.001	8.838	<.001	5.591	<.001
R ²		0.796		0.769		0.812		0.732	

263

264 Model analysis for interactive effects and optimized 265 conditions

266 Given the interactive effects of factors A and C and exploring optimized conditions, regression
267 models were constructed. As shown in Figure 4, the root and seedling length in the four groups of
268 aged seeds showed via three-dimensional parabolic trends that there were maximum values with
269 increase in factor A. The root and seedling length were sharply increased when the bacterial

concentration was in the range of $30.0\text{--}75.0 \times 10^6$ CFU mL⁻¹. The SVi also had maximum values with a bacterial concentration of $30.0\text{--}75.0 \times 10^6$ CFU mL⁻¹ when priming duration was less than 24.0 h, although the SVi of orchardgrass 2013 exhibited a completely opposite trend (Fig 4E-H). The Gi of Tf 2015 continuously increased with the increase in factor A under short priming time (less than 8.0 h), and the Gi of three other seed groups showed similar trends as those of the SVi of orchardgrass2013(Fig 4A-D).

Fig 4. Response surface plots showing the interactions between the bacterial concentration and priming duration on the germination index (A, B, C and D), seed vigor index (E, F, G and H), root length (I, J, K and L) and seedling length (M, N, O and P).

The MDA content in Russian wildrye 2009 and the APX activity in tall fescue 2015 had minimum values at 52.8×10^6 CFU mL⁻¹, 19.2 h and 52.1×10^6 CFU mL⁻¹, 15.6 h, respectively (Figs 5B, 6F and S2 Table). The MDA content in tall fescue 2015 decreased with the increase in factor A from 60.0×10^6 CFU mL⁻¹ and showed opposite changing pattern as that of SOD activity. When factor A was less than 60.0×10^6 CFU mL⁻¹, the MDA content in tall fescue 2014 was sharply reduced with the increase in factor A (Figs 5A, C and 6D). The POD activity in Russian wildrye 2009 exhibited a parabolic trend that peaked at 61.1×10^6 CFU mL⁻¹ at 3.9 h; the CAT activity in Russian wildrye 2009 was positively correlated with bacterial concentration when factor A was larger than 30.0×10^6 CFU mL⁻¹ and when the priming time was longer than 16.0 h

(Figs 5E, 6B and S2 Table). The CAT activity in tall fescue 2015 was the highest within the factor A range of $45.0\text{--}90.0 \times 10^6 \text{ CFU mL}^{-1}$ and when the priming time was less than 4.0 h (Fig 5D). The APX and CAT activities showed analogous trends, and the APX and SOD activities were nearly complementary in orchardgrass 2013 (Figs 5F, 6C and 6E).

Fig 5. Response surface plots showing the interactions between the bacterial concentration and priming duration on the MDA content (A, B and C) and CAT activity (D, E and F).

Fig 6. Response surface plots showing the interaction between the bacterial concentration and priming duration on the activity of POD (A, B), SOD (C, D) and APX (E, F).

A total of 48 pairs of critical values composed of factors A and C were obtained from the model analysis, as shown in Figure 7. The mode of critical values was considered the largest value and was used as the final optimal value [4, 25]. The final optimal conditions for factors A and C were the bacterial concentration of *A. brasilense* of $52.3 \times 10^6 \text{ CFU mL}^{-1}$ and the seed priming time of 17.0

h.

Fig 7. Scatter plot showing the stationary points of optimal models related to the bacterial concentration and priming duration.

Discussion

Seed aging has become an intractable problem in agricultural science, which negatively affects the external morphology and internal physiology of seeds and seedlings [9]. Seed pre-treating with PGPR is increasingly favored by scholars because of its eco-friendly and high efficiency characteristic. The present study showed that the combined treatments of *A. brasilense* inoculation and the three seed priming treatments counteracted the adverse effects from natural aging on the four groups of seeds. The results revealed that the three seed priming treatments played the dominant role in improving the seed germination and vigor. And the pairwise interactions among three factors were also significant, especially between bacterial concentration of *A. brasilense* and three seed priming treatments for the germination percentage and germination index. However, the sole function of bacterial concentration was not significant. In this regard, Carrozzi et al. noted that hydration treatment might affect only the early stages of imbibitions, while the function of *A. brasilense* is additive to the hydration process, which could result in repair actions for the promotion of aged seed germination [8]. It was also found that the germination of Russian wildrye 2009 during the early phases in all treatments lagged far behind that of the other seeds. This

phenomenon indicates that the germination percentage might have been intimately linked to the reparation of seed deterioration during the lag phase of germination [8]. Accordingly, for the older seed, a longer lag phase was required to reverse deterioration by seed priming. Previous studies have reported that hydropriming, osmopriming and redox priming can invigorate aged seed associated with the initiation of repair- and germination-related processes, such as increased α -amylase activity, the activation and resynthesis of some ROS scavengers, the repair of chromosomal damage, protein, DNA and RNA [4, 26]. On the other hand, *A. brasilense* may contribute to phytohormone communication between PGPR and aged seeds, mainly GA and IAA produced by PGPR. IAA might assist GA biosynthesis which could activate some genes responsible for α -amylase mRNA transcription in aged seeds during germination [27, 28]. Therefore, we speculated here that *A. brasilense* could synergize the reinvigorating action of the three seed priming treatments for seed germination and vigor modulation in aged seeds; the latter treatments would play the main role in the early stages.

The co-treatments expressed remarkable root elongation-promoting activity, particularly under the bacterial concentration of 10^7 CFU mL⁻¹ and H₂O₂-priming for 18.0 h (T8) for all three aged species. The study revealed that the effect of bacterial concentration of *A. brasilense* was the greatest among the factors on root length. The root elongation response might be mainly due to the positive feedback loop between root and bacterial IAA. As reported previously, hormones produced by *A. brasilense* could affect root cell wall dynamics [29]; *A. brasilense* inoculation can stimulate root carboxylate exudation acting as precursors of phytohormones, which indirectly and positively affected maize root length and area [30]. In addition, under the interactive effects of

factors A and C, the root and seedling lengths first increased but then decreased with the increase in factor A, there were somehow critical thresholds between the factors for the co-treatments. A moderate bacterial concentration ($50.0\text{--}65.0 \times 10^6$ CFU mL⁻¹) and priming duration (5.8–26.3 h) could effectively facilitate the root system development in aged seeds. Similarly, *A. brasilense* Sp 7-S and Sp 245 at 10^7 CFU mL⁻¹ stimulated root and seedling growth in cucumber and lettuce ^[31].

To further elucidate the underlying physiological mechanisms of the combined treatments, both antioxidant enzymatic activities and lipid peroxidation were investigated. Seed aging inevitably generates the overproduction of ROS, which accelerate the disintegration of membranes. In the present study, the corresponding treatments effectively increased the activities of SOD, POD and CAT and lowered the MDA contents in the four groups of aged seeds. Especially, in severely aged seed of Russian wildrye 2009, the maximum POD activity occurred at 61.1×10^6 CFU mL⁻¹ and 3.9 h, and the minimum value of the MDA content occurred at 52.8×10^6 CFU mL⁻¹ and 19.2 h, under the strong interactive effect. It was found that the germination percentage and root length of aged seed were negatively correlated with MDA content, and were positively correlated with the SOD activity (S3 Table), which is in line with the finding of Pereira et al. ^[32]. Wojtyla et al. pointed out that in aged seeds, the deleterious effects derived from excessive ROS can hinder cell wall loosening to allow cell elongation in root apical meristems ^[33]. Coincidentally, SOD can directly modulate the amount of ROS ^[4]. The increased activities of antioxidant enzymes confer the ability of plants to scavenge excessive ROS ^[34]. Therefore, reduction of MDA content and improvement of antioxidant enzyme activities in aged seeds after the co-treatment can be related to better germination and seedling growth in treated seeds compared with the untreated ones. It

was clarified here that *A. brasilense* inoculation was the major modulator to improve the SOD activity in aged seeds. And this mode of action is consistent with the promoting effect for root length mentioned above. Recent studies have shown that PGPR regulate the expression of the antioxidant-responsive genes *MnSOD* and *APX* [12]. Wang et al. also indicated that a consortium of PGPR that induce drought tolerance involves SOD and *cAPX*, not POD and CAT, in cucumber [35]. In this regard, the present results indicated that the three seed priming approaches were the most effective factors for enhancing POD and CAT activities. It was observed that the activities of APX and SOD were complementary in orchardgrass 2013 and that the APX activity in tall fescue 2015 had a minimum value under the interactive effects of factors A and C, which contradicts the results of some studies. However, this finding suggested that an operation mode of co-allocation might exist among antioxidant enzymes during the repair process in aged seeds. It was speculated that APX might complement the activity of SOD when the amount of ROS over-accumulated in aged seeds and that it might not play a relevant role in antioxidant enzyme system in relatively new seeds. This assumption is also supported by the study from Bosco de Oliveira et al. [36]. Additionally, the reduction in MDA content after the combined treatment suggested the capability for attenuating the extent of membrane lipid peroxidation under aging stress. Although the experimental factors did not significantly affect the MDA content, the changing trends were complementary between the MDA content and SOD activity in Russian wildrye and tall fescue. Any sort of treatments reduced the MDA levels in the seedlings, which could be explained by the enhancement of antioxidant enzyme activities [32].

The combined treatments were more effective for boosting the growth of relatively aged seed

(Russian wildrye 2009) compared with that of new seed. This result could confirm the repair effect of the co-treatments on aged seeds. The repair effect of co-treatments on the SOD activities, mitochondria, associated enzymes or proteins might be less/negative in new seeds with respect to old seeds, which resulted from different physiological states such as seed moisture levels, water transport and redox status ^[4]. And seeds are expected to exhibit differences in experimental factors, which are manifested by the different genes in the different aged seeds. For tall fescue 2014, the co-treatment had a small positive impact, even negative effect on germination maybe because of its own characteristic. But it is worth noting that the germination in T2 and T7 of tall fescue 2014 occurred earlier, and T1, T2, T5, T6 and T8 significantly increased the root length, which means the co-treatment had potentially positive effect on seeds of tall fescue 2014. Therefore, the co-treatments have to be considered carefully in this species and to require further study. In addition, inter- and intra-species differences existed for the effect of factor B. The choice of seed priming treatment largely depends on plant species, variety and storage duration. The greatest average root length for tall fescue 2015, the highest average GP, Gi, and SVi for orchardgrass 2013, and those of GP, SVi, root length, shoot length, the activity of SOD and CAT for Russian wildrye 2009 occurred in the H₂O₂-priming. H₂O₂ of an appropriate level could function as a common stress and developmental signaling molecule that regulates a common set of transcription factors related to the antioxidant defense system and can eventually restore redox homeostasis and oxidative membrane damage ^[37]. The effects of H₂O₂ were also correspondingly manifested by T8 (H₂O₂-priming) stimulating root elongation for all three aged species mentioned above. However, the germination percentage was the lowest in T9 (H₂O₂ priming), the germination percentage in

Russian wildrye 2009 of severely aged seeds was enhanced by all nine treatments. This result implied that the level of ROS in T9 in Russian wildrye 2009 that already have an abundance of ROS, including H₂O₂, was not within the optimum range of concentrations, but the range was still within the “oxidative window” [38]. Therefore, the H₂O₂ concentration lower than 10⁸ CFU mL⁻¹ or priming duration shorter than 24h may be required for severely aged seeds. It is worth noting that the germination percentage of Russian wildrye 2009 in T8 was relatively higher than the remaining six treatments, and the seeds of orchardgrass 2006 germinated in T7. Taking into account the overall analysis of results of all experimental indicators, H₂O₂ was still the most effective condition for aged forage seeds. Similarly, H₂O₂ priming has improved the germination and seedling growth of pea, rice as well as aged maize, squash and tomato seeds [39-41]; *A. brasilense* inoculation with H₂O₂-priming also enhanced the grain yield of wheat under dryland conditions [19]. Therefore, *A. brasilense* bio-priming combined with H₂O₂-priming constituted the optimal combination for repairing aged seeds.

Overall, a physiological mechanistic model of the combined treatments for aged grass seeds was proposed based on the current results. The PGPR bio-priming and the three seed priming treatments are coordinated but have distributed responsibilities regarding the reinvigoration of aged grass seeds during the different developmental periods. During the initiation of germination, the three seed priming treatments were the dominant factors and *A. brasilense* strains aided these activities that could lead to repair processes. For the enhancement of seedling growth (especially root length) and SOD activity, *A. brasilense* was the main driving force.

Conclusion

Azospirillum brasilense inoculation and the three seed priming treatments repaired aged grass seeds by the specific operation mode of co-allocation during the different developmental stages. The three seed priming methods were the most effective factors for stimulating germination, whereas the bacterial concentration had the largest effect on root elongation. The combined treatments that elicited aging-related repair reduced the MDA content and improved antioxidant enzyme activities. Overall, *A. brasilense* bio-priming with H₂O₂-priming was the optimal combination for reinvigorating aged seeds. The final optimal conditions consisted of the bacterial concentration of *A. brasilense* of 52.3×10^6 CFU mL⁻¹ and a seed priming time of 17.0 h, given the interactive effects. However, the interactional mechanisms of combined treatments are just beginning to be unraveled. Whether the co-treatments affect aging-related proteins or genes as well as antioxidant enzyme genes still requires further exploration.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgements

National Key R&D Program of China (2017YFE0111000) and The National Natural Science Foundation of China (31472138) funded this work.

Author Contributions

Q.W. and X.L. carried out the conception and designed the experiments. X.L., Z.C., W.C., Y.G., Q.L., W.Z., T.Z. and W.J. performed the experiments. Q.W., X.L., Z.C. and J.C. analyzed the data and co-wrote the manuscript. All authors reviewed the manuscript.

References

1. Chen L, Chen QZ, Kong LQ, Xia FS, Yan HF, Zhu YQ, Mao PS. Proteomic and physiological analysis of the response of oat (*Avena sativa*) seeds to heat stress under different moisture conditions. *Front. Plant Sci.* 2016; 7: 896. doi: 10.3389/fpls.2016.00896
2. Parkhey S, Naithani SC, Keshavkant S. Protein metabolism during natural ageing in desiccating recalcitrant seeds of *Shorea robusta*. *Acta Physiol Plant.* 2014; 36 (7): 1649-1659. doi: 10.1007/s11738-014-1540-x
3. Yao Z, Liu L, Gao F, Rampitsch C, Reinecke DM, Ozga JA, Ayele BT. Developmental and seed aging mediated regulation of antioxidative genes and differential expression of proteins during pre- and post-germinative phases in pea. *J Plant Physiol.* 2012; 169 (15): 1477-1488. doi: 10.1016/j.jplph.2012.06.001
4. Liu J, Wang QZ, Karagić Đ, Liu X, Cui J, Gui J, Gu MY, Gao W. Effects of ultrasonication on increased germination and improved seedling growth of aged grass seeds of tall fescue and Russian wildrye. *Sci Rep.* 2016; 6: 22403. doi: 10.1038/srep22403
5. Ellis RH, Roberts EH. An investigation into the possible effects of ripeness and repeated threshing on barley seed longevity under six different storage environments. *Ann Bot.* 1981; 48 (1):93-96.
6. Ellis RH, Hong TD, Roberts EH. The low-moisture-content limit to the negative logarithmic relation between seed longevity and moisture content in three subspecies of rice. *Ann Bot.* 1992; 69(1):53-58.
7. Stanisavljevic R, Djokic D, Milenkovic J, Đukanovic L, Stevovic V, Simic A, Dodi D. Seed germination and seedling vigour of italian ryegrass, cocksfoot and timothy following harvest and storage. *Ciência E Agrotecnologia.* 2011; 35: 1141-1148.
8. Carrozzi LE, Creus CM, Barassi CA, Monterubbianesi G, Di-Benedetto A. Reparation of aged lettuce (*Lactuca sativa*) seeds by osmotic priming and *Azospirillum brasilense* inoculation. *Botany.* 2012; 90(11):

1093-1102.

9. Moncaleano-Escandon J, Silva BCF, Silva SRS, Granja JAA., Alves MCJL, Pompelli MF. Germination responses of *Jatropha curcas* L. seeds to storage and aging. *Ind Crops Prod.* 2013; 44(1): 684-690. doi: 10.1016/j.indcrop.2012.08.035

10. Behtari B, Tilaki GAD. Changes in germination characteristics and seedling growth between storage and non-storage of primed tall fescue seed. *Afr J Biotechnol.* 2012; 11: 4317-4322.

11. Garcíaicristobal J, Garcíaavillaraco A, Ramos B, Gutierrez-Mañero J, Lucas JA. Priming of pathogenesis related-proteins and enzymes related to oxidative stress by plant growth promoting rhizobacteria on rice plants upon abiotic and biotic stress challenge. *J Plant Physiol.* 2015; 188: 72-79. doi: 10.1016/j.jplph.2015.09.011

12. Bharti N, Pandey SS, Barnawal D, Patel VK, Kalra A. Plant growth promoting rhizobacteria *Dietzia natronolimnaea* modulates the expression of stress responsive genes providing protection of wheat from salinity stress. *Sci Rep.* 2016; 6: 34768.

13. Zawoznik MS, Ameneiros M, Benavides MP, Vazquez S, Groppa MD. Response to saline stress and aquaporin expression in *Azospirillum*-inoculated barley seedlings. *Appl Microbiol Biotechnol.* 2011; 90(4): 1389-1397.

14. Rozier C, Hamzaoui J, Lemoine D, Czarnes S, Legendre L. Field-based assessment of the mechanism of maize yield enhancement by *Azospirillum lipoferum* CRT1. *Sci Rep.* 2017; 7(1): 7416. doi: 10.1038/s41598-017-07929-8

15. Mahmood A, Turgay OC, Farooq M, Hayat R. Seed biopriming with plant growth promoting rhizobacteria: a review. *FEMS Microbiol Ecol.* 2016; 92(8): 112. doi: 10.1093/femsec/fiw112

16. Fibach-Paldi S, Burdman S, Okon Y. Key physiological properties contributing to rhizosphere adaptation and plant growth promotion abilities of *Azospirillum brasilense*. *FEMS Microbiol Lett.* 2012; 326: 99-108. doi: 10.1111/j.1574-6968.2011.02407.x

17. Curá J, Franz D, Filosofía J, Balestrasse K, Burgueño L. Inoculation with *Azospirillum* sp. and *Herbaspirillum* sp. bacteria increases the tolerance of maize to drought stress. *Microorganisms.* 2017; 5(3): 41. doi: 10.3390/microorganisms5030041

18. Zulueta-Rodríguez R, Hernández-Montiel LG, Murillo-Amador B, Rueda-Puente EO, Capistrán LL, Troyo-Diéguez E, Córdoba-Matson MV. Effect of hydropriming and biopriming on seed germination and growth of two Mexican fir tree species in danger of extinction. *Forests.* 2015; 6(12): 3109-3122. doi: 10.3390/f6093109

- 510 19. Jafariyan T, Zarea MJ. Hydrogen peroxide affects plant growth promoting effects of Azospirillum. J Crop
511 Sci Biotech. 2016; 19(2): 167-175. doi: 10.1007/s12892-015-0127-4
- 512 20. Stajković-Srbinić O, Delić D, Kuzmanović D, Sikirić B, Rasulić N, Nikolić B, Knežević-Vukčević J.
513 Growth and nutrient uptake of orchardgrass (*Dactylis glomerata* L.) and meadow fescue (*Festuca pratensis*
514 Huds.) as affected by rhizobacteria. Not Bot Horti Agrobo Cluj-Napoca. 2016; 44(1): 296-301. doi:
515 10.15835/nbha44110252
- 516 21. Caceres EAR. Improved medium for isolation of Azospirillum spp. Appl Environ Microb. 1982; 44:
517 990-991.
- 518 22. Bao AK, Wang SM, Wu GQ, Xi JJ, Zhang JL, Wang CM. Overexpression of the Arabidopsis H⁺-PPase
519 enhanced resistance to salt and drought stress in transgenic alfalfa (*Medicago sativa* L.). Plant Sci. 2009;
520 176: 232-240.
- 521 23. Zhang YP, Nan ZB. Growth and anti-oxidative systems changes in *Elymus dahuricus* is affected by
522 Neotyphodium endophyte under contrasting water availability. J Agron Crop Sci. 2007; 193: 377-386. doi:
523 10.1111/j.1439-037X.2007.00279.x
- 524 24. Riva A. Statistical methods and the geographer by S. Gregory. 1978; Longman.
- 525 25. Lattin JM, Carroll JD, Green PE. 2003. Analyzing Multivariate Data. Pacific Grove, CA: Thomson
526 Brooks/Cole; 2003.
- 527 26. Ghassemi-Golezani K, Hosseinzadeh-Mahootchy A, Zehtab-Salmasi S, Turchi M. Improving field
528 performance of aged chickpea seeds by hydro-priming under water stress. Int J Plant Animal Environ Sci.
529 2012; 2: 168-176.
- 530 27. Siadat SA, Moosavi SA, Zadeh MS, Fotouhi F, Zirezadeh M. Effects of halo and phytohormone seed
531 priming on germination and seedling growth of maize under different duration of accelerated ageing
532 treatment. Afr J Agr Res. 2011; 6(31): 6453-6462.
- 533 28. Mangmang JS, Deaker R, Rogers G. Germination characteristics of cucumber influenced by plant growth-
534 promoting rhizobacteria. Int J Vegetable Sci. 2015a; 22: 66-75. doi: 10.1080/19315260.2014.938850
- 535 29. Richards SL, Wilkins KA, Swarbreck SM, Anderson AA, Habib N, Smith AG, McAinsh M, Davies JM. The
536 hydroxyl radical in plants: from seed to seed. J Exp Bot. 2015; 66: 37-46.
- 537 30. D'Angioli AM, Viani RAG, Lambers H, Sawaya ACHF, Oliveira RS. Inoculation with Azospirillum
538 brasilense (Ab-V4, Ab-V5) increases Zea mays root carboxylate-exudation rates, dependent on soil
539 phosphorus supply. Plant Soil. 2016; 410: 499-507. doi: 10.1007/s11104-016-3044-5
- 540 31. Mangmang JS, Deaker R, Rogers G. Optimal plant growth-promoting concentration of Azospirillum

541 brasilense inoculated to cucumber, lettuce and tomato seeds varies between bacterial strains. Israel J Plant
542 Sci. 2015b; 62: 145-152.

543 32. Pereira SIA, Moreira H, Argyras K, Castro PML, Marques APGC. Promotion of sunflower growth under
544 saline water irrigation by the inoculation of beneficial microorganisms. Appl Soil Ecol. 2016; 105: 36-47.
545 doi: 10.1016/j.apsoil.2016.03.015

546 33. Wojtyla L, Lechowska K, Kubala S, Garneczarska M. Molecular processes induced in primed
547 seeds-increasing the potential to stabilize crop yields under drought conditions. J Plant Physiol. 2016; 203:
548 116-126. doi: 10.1016/j.jplph.2016.04.008

549 34. Ibrahim EA. Seed priming to alleviate salinity stress in germinating seeds. J Plant Physiol. 2016; 192: 38-46.

550 35. Wang CJ, Yang W, Wang C, Gu C, Niu DD, Liu HX, Wang YP, Guo JH. Induction of drought tolerance in
551 cucumber plants by a consortium of three plant growth-promoting rhizobacterium strains. Plos One. 2012;
552 7(12): e52565.

553 36. Bosco de Oliveira A, Gomes-Filho E, Enéas-Filho J, Prisco JT, Alencar NLM. Seed priming effects on
554 growth, lipid peroxidation, and activity of ROS scavenging enzymes in NaCl-stressed sorghum seedlings
555 from aged seeds. J Plant Interactions. 2012; 7(2): 151-159. doi: 10.1080/17429145.2011.582590

556 37. Bhattacharjee S. An inductive pulse of hydrogen peroxide pretreatment restores redox-homeostasis and
557 oxidative membrane damage under extremes of temperature in two rice cultivars. Plant Growth Regul. 2012;
558 68(3):395-410. doi: 10.1007/s10725-012-9728-9

559 38. Bailly C, El-Maarouf-Bouteau H, Corbineau F. From intracellular signaling networks to cell death: the dual
560 role of reactive oxygen species in seed physiology. C R Biol. 2008; 331 (10): 806-814. doi:
561 10.1016/j.crv.2008.07.022

562 39. Barba-Espin G, Diaz-Vivancos P, Job D, Belghazi M, Job C, Hernández JA. Understanding the role of H₂O₂
563 during pea seed germination: a combined proteomic and hormone profiling approach. Plant Cell Environ.
564 2011; 34: 1907-1919. doi: 10.1111/j.1365-3040.2011.02386.x

565 40. Liu G, Porterfield DM, Li YC, Klassen K. Increased oxygen bioavailability improved vigor and germination
566 of aged vegetable seeds. Hortscience. 2012; 47(12): 1714-1721.

567 41. Hussain S, Khan F, Hussain HA, Nie L. Physiological and biochemical mechanisms of seed priming-induced
568 chilling tolerance in rice cultivars. Front Plant Sci. 2016; 7: 116. doi: 10.3389/fpls.2016.00116

Supporting information

S1 Table. The MDA content and the activity of SOD and POD in tall fescue 2015, 2014, orchardgrass 2013 and Russian wild rye 2009.

S2 Table. Canonical response surface analysis for germination percentage, germination index, seed vigor index, root, shoot and seedling length, ratio of S/R, MDA content, the activity of SOD, POD, CAT and APX of the seedlings based on coded data.

S3 Table. Pearson correlation coefficients for activity of SOD, APX, POD, CAT, and MDA content.

Figure 1

Figure 4

Figure 5

Figure 6

Figure 7

Figure 2

Figure 3