Genome sequence of *Pseudomonas aeruginosa* PAO1161, a PAO1 derivative with the ICEPae1161 integrative and conjugative element

Adam Kawalek^{1*}, Karolina Kotecka¹, Magdalena Modrzejewska¹, Grazyna Jagura-Burdzy¹, Aneta Agnieszka Bartosik^{1*}

¹ Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Department of Microbial Biochemistry, Warsaw, Poland

* To whom correspondence should be addressed: anetab2@ibb.waw.pl or a.kawalek@ibb.waw.pl.

ABSTRACT

Pseudomonas aeruginosa is a cause of nosocomial infections, especially in patients with cystic fibrosis and burn wounds. PAO1 strain and its derivatives are widely used to study the biology of this bacterium, however recent studies demonstrated differences in the genomes and phenotypes of derivatives from different laboratories.

Here we report the genome of *P. aeruginosa* PAO1161 laboratory strain, a *leu*⁻, Rif^R, restriction-modification defective PAO1 derivative, described as the host of IncP-8 plasmid FP2, conferring the resistance to mercury. Comparison of PAO1161 genome with PAO1-UW sequence revealed an inversion of a large genome segment between rRNA operons and 100 nucleotide polymorphisms, short insertions and deletions. These included a change in *leuA*, resulting in E108K substitution, which caused leucine auxotrophy and a mutation in *rpoB*, enhancing rifampicin resistance. Nonsense mutations were detected in *PA2735* and *PA1939* encoding a DNA methyltransferase and a putative OLD family endonuclease, respectively. Moreover, a 12 kb RPG42 prophage and a novel 108 kb PAPI-1 like integrative conjugative element (ICE) encompassing a mercury resistance operon were identified. The ICE_{Pae1161} was transferred to *Pseudomonas putida* cells, where it integrated in the genome and conferred the mercury resistance. This indicates that the FP2 plasmid is in fact an ICE.

INTRODUCTION

Pseudomonas aeruginosa is a Gram-negative gammaproteobacterium commonly found in various ecological niches and characterized by the ability to survive in unfavourable, frequently changing environmental conditions. This opportunistic human pathogen is often a source of nosocomial infections in immuno-compromised patients. In cystic fibrosis patients *P. aeruginosa* chronically colonizes the lungs and is a major factor causing mortality^{1,2}.

Majority of research on this metabolically versatile bacterium uses sublines or derivatives of *P. aeruginosa* PAO1 strain, originally isolated from a wound of a patient in Holloway's laboratory, Melbourne, Australia³. Over the years, the strain was shipped to laboratories worldwide and its different attenuated derivatives, including auxotrophic strains and strains with mobile genetic elements were obtained⁴. In 1999 the genome of *P. aeruginosa* PAO1 stored at the University of Washington (PAO1-UW) was sequenced⁵, providing a reference for studies on *P. aeruginosa* genomes. Up to December 2018, the *Pseudomonas* Genome Database, a database devoted to the information on *Pseudomonas* species⁶, contained 2226 sequenced *P. aeruginosa* genomes, including 11 PAO1 derivatives (sublines). Remarkably, sequencing of the PAO1 subline (MPAO1) as well as PAO1-DSM strain stored at the German Collection for Microorganisms and Cell Cultures revealed presence of multiple nucleotide polymorphisms and short insertions-deletions relative to the reference PAO1-UW⁷. A major feature differing genomes of PAO1 derivatives is the occurrence of a large inversion resulting from the homologous recombination between two rRNA operons *rrnA* and *rrnB*⁵, which is present in the reference PAO1-UW genome and not in e.g., MPAO1 and PAO1-DSM⁷. Despite an asymmetrical positioning of the *dif* region in PAO1-UW, this inversion does not seem to affect chromosome segregation and such large rearrangements might be common among bacteria⁸. The presence of multiple short sequence modifications could lead to the variations in e.g. virulence and fitness between strains used in different laboratories⁷. This data indicate an ongoing micro- and macro- evolution of bacterial genomes and suggests that sequence diversification in laboratory strains should be taken into consideration in the analysis of phenotypic data^{9,10}.

In this work we focus on the genome of *P. aeruginosa* PAO1161 strain, a PAO1 derivative requiring leucine for growth on minimal media and selected as lacking the restriction-modification system (*rmo*-10 mutation)¹¹. This strain is described as the host for plasmid FP2 of IncP-8, the only known member of this incompatibility group, conferring resistance to mercury^{12,13}. The FP2 factor demonstrated the chromosome-mobilizing ability (Cma) and was extensively used in interrupted mating technique for preparation of the genetic map of *P. aeruginosa* chromosome^{4,14}. The PAO1161 derives from the PAO38 *leu*-38 mutant obtained by treatment of PAO1 with manganese chloride and search for leucine auxotrophs³ (Figure 1A). PAO38 acquired the FP2 element from PAT (*P. aeruginosa* strain 2)¹⁵ to yield strain PAO170¹⁶. Following mutagenesis of PAO170 with N-

methyl-N'-nitro-N-nitrosoguanidine PAO1161 was selected as defective in restriction and modification systems (*r^m*) on the basis of the altered susceptibility to phage infection^{11,17}. To facilitate the use of PAO1161 in conjugation experiments spontaneous rifampicin resistant clone was obtained¹⁸. The PAO1161 strain was used in studies on chromosome segregation and gene expression in *P. aeruginosa* using genome wide approaches^{19–21} as well as in other physiological and genetic studies^{22–28}.

Here we report the genome sequence of *P. aeruginosa* PAO1161 strain. Comparison with PAO1-UW reference sequence revealed the presence of a large number of single-nucleotide polymorphisms (SNPs), insertions-deletions (indels), as well as inversion of large genome segment between rRNA genes. Moreover a functional PAPI-1 like integrative conjugative element (ICE), containing a mercury resistance operon was identified in PAO1161 genome, indicating that the FP2 is not a plasmid but an ICE (designated ICE*Pae*1161).

RESULTS AND DISCUSSION

Comparison of *P. aeruginosa* PAO1161 genome with PAO1 reference assembly

P. aeruginosa PAO1161 genome was sequenced and assembled, resulting in a single circular chromosome of 6383803 bp. A comparison with the reference PAO1-UW genome (NC_002516) revealed three major structural differences (Figure 1B). The PAO1161 genome lacks the large inversion between ribosomal RNA operons *rrnA* and *rrnB* similarly to other PAO1 derivatives like MPAO1 and PAO1-DSM⁷, but observed in PAO1-UW⁵. Remarkably, PAO1161 possesses two large insertions. The 107796 bp insertion in *tRNA_{Lys}* gene between *lepA* (*PA4541*) and *clpB* (*PA4542*), flanked by 48 bp repeated sequences, displays a significant similarity to PAPI-1 like integrative conjugative elements (see below)^{29,30}. The second 11981 bp insertion between *tRNA_{Met}* (*PA4673.1*) and *higA* (*PA4674*), flanked by 82 bp repeats, is identical to the prophage-like RGP42 element identified also in MPAO1 and PAO1-DSM⁷. Additionally, PAO1161 lacks a 280 bp fragment containing *PA1796.3* and *PA1796.4* tRNA genes and has an 107 bp insertion downstream of *PA2327*.

A comparison of PAO1161 and PAO1-UW genome sequences using NucleDiff³¹ revealed 251 single nucleotide polymorphisms (SNPs), multiple nucleotide polymorphisms (MNPs) and short insertions / deletions (indels). Since a direct sequence comparison does not provide information about the quality of predicted variants, the outcome can be greatly affected by errors during genome sequence consensus calling, caused for instance by mapping of reads derived from highly similar sequences to another, very similar, parts of the genome. To identify high quality variants we aligned the reads used in genome assembly to the consensus sequence and checked the frequency of the variant at a given genome position (see Materials and Methods). Overall 100 SNPs/MNPs/indels

were called with a high confidence between the PAO1161 and the PAO1-UW reference. The remaining (heterozygous) variants were mostly SNPs (149/151) located in coding sequences of genes belonging to Pseudomonas Orthologous Groups (POG) possessing multiple members in the same strain (multiple orthologs within PAO1 genome)⁶. This group included 17 SNPs in *PA0723-PA0726* genes from the Pf1 cryptic prophage region syntenic to the genes within RGP42 element also present in PAO1161 genome⁷. Eight of these variants were also identified previously in MPAO1 and PAO1-DSM⁷, nevertheless these did not meet our quality criteria.

Effect of SNPs, MNPs and indels

The high-confidence variants encompassed 52 SNPs, 6 MNPs, 15 deletions and 27 insertions. Of these 44 were mapping to the intergenic regions in PAO1-UW genome and 9 were synonymous (silent) mutations (Supplementary Table S1). Three mutations resulted in an introduction of premature stop codons (Table 1). Interestingly, these encompass *PA2735* gene, recently shown to encode a N6-adenosine DNA methyltransferase acting on a conserved sequence³², and *PA1939*, encoding a putative overcoming lysogenization defect (OLD) family nuclease containing an N-terminal ATPase domain and a C-terminal TOPRIM domain³³. Since *P. aeruginosa* PAO1161 strain was selected as defective in its restriction system^{11,17}, it is tempting to speculate that PA1939 could be the protein involved in degradation of the foreign DNA and together with PA2735 they could constitute the chromosomal restriction-modification system of *P. aeruginosa*.

Four of the identified sequence variants resulted in frame shifts, leading to expression of truncated proteins (Table 1). In comparison to PAO1, PAO1161 genome contains a frame shift in the *PA0683* (*hxcY*) gene encoding a component of the Hxc system, a type II secretion system active under phosphate-limiting growth conditions and dedicated to the secretion of low-molecular-weight alkaline phosphatases LapA and LapB³⁴⁻³⁶. The frameshift in PAO1161 genome results in expression of a protein with only 72 aa identical to PA0683 from PAO1-UW.

In contrast, indels identified in *PA1327*, *PA1685* and *PA2141* mapped to the 3' part of the genes, resulting only in short sequence alteration of C-terminal regions of corresponding proteins, unlikely to affect their function (Table 1). This is highly plausible as the analysis of the protein sequence of PA1327, PA1685 or PA2141 orthologs (Pseudomonas Ortholog Groups: POG001263, POG003705, POG003333, respectively), showed that the PAO1161-like C-termini are typical for most orthologs from the corresponding groups.

The effect of 14 indels and 1 SNP is predicted as a shift in start or stop codon of the corresponding gene(s) leading to extension of the protein product in PAO1161 relative to PAO1-UW (Table 2). These also include changes in intergenic regions, like 413850 T→C mutation upstream of *PA0369*, modifying an in frame stop-codon and possibly leading to a translation of a longer protein. Similar, start codon changing role can be attributed to nucleotide changes within

PA2046-PA2047 and PA4874-PA4875 intergenic regions, as well as to 5 other indels (Table 2). The effect of three PAO1161/PAO1-UW sequence differences is fusion of proteins encoded by adjacent genes in PAO1-UW genome into one protein in PAO1161. Additionally, the frame shift observed in PAO1-UW in *PA0748* gene, encoding an AraC type transcriptional regulator, is not observed in PAO1161. Notably, analysis of the sequences composing Pseudomonas Orthologs Group of corresponding PAO1 genes showed that in all cases the PAO1161-like protein products constitute a vast majority of all deposited sequences. One exception is the PAO1161 *relA* gene encoding a GTP pyrophosphokinase, in which a start codon shift was identified, compared to PAO1-UW (Table 2). RelA, together with SpoT, controls the level of guanosine 5'-triphosphate-3'-diphosphate (ppGpp) alarmone involved in bacterial adaptive responses and virulence³⁷. Importantly, previous mass spectrometry analysis and iTRAQ quantification of PAO1161 proteome showed that RelA is produced in PAO1161 cells³⁸.

Except nucleotide changes with a major effect of the corresponding protein products, numerous SNPs and indels resulting in amino acid substitutions or deletions relative to corresponding PAO1-UW proteins were identified (Table 3). One special case is PA2492 (*mexT*), where both a deletion and a SNP were observed in PAO1161 relative to PAO1-UW sequence (Table 1 and Table 3). MexT is a LysR type transcriptional regulator activating expression of the MexEF-OprN multidrug efflux system, extensively studied in the context of quorum sensing signalling and resistance to antimicrobial agents^{39–42}. PAO1161 does not possess the 8-nt insertion within *mexT* (Table 1), observed in some PAO1 sublines⁴³. Concomitantly, PAO1161 contains a SNP in the *mexT* gene, leading to F172I change which is also found in most of the strains with the active *mexT*⁴³.

A sequence variation was also observed in *mexS*, encoding a negative regulator of MexT^{40,44,45}, resulting in D249N amino acid change, which was shown to not affect the protein function⁴⁰.

PAO1161 strain derives from the strain PAO38 which was selected as the leucine auxotroph⁴⁶ (Figure 1A). Genome sequencing of PAO1161 revealed that this strain possesses a mutation in *leuA*, encoding a putative 2-isopropylmalate synthase, resulting in E108K substitution. Analysis of Pseudomonas Ortholog Group of the *leuA* (POG001874) showed that, the only *P. aeruginosa* strains carrying this mutation are PAO579^{46,47} and PAO581⁴⁸, two PAO38 derivatives (Figure 1A). To validate that this substitution causes the leucine auxotrophy, we exchanged PAO1161 *leuA* to the PAO1 allele. The allele replacement fully restored the ability of PAO1161 strain to grow on minimal medium without leucine (Figure 2), confirming that the E108K substitution in LeuA caused leucine auxotrophy.

P. aeruginosa PAO1161 used in this study was a rifampicin resistant clone¹⁸. Rifampicin binds to a conserved pocket on the β -subunit of RNA polymerase therefore blocking RNA transcript elongation⁴⁹. Resistance to this drug results from mutations on the *rpoB* gene that change the

structure of the pocket⁴⁹⁻⁵¹. Our analysis revealed presence of a SNP in *rpoB*, encoding a DNA-directed RNA polymerase subunit beta, resulting in H531L substitution (Table 3). This amino acid change was frequently observed in spontaneous *P. aeruginosa* Rif^R mutants⁵², strongly indicating that this SNP confers PAO1161 strain with rifampicin resistance.

Interestingly, for 8 proteins the same changes were found in PAO1161 strain and in MPAO1 and / or PAO1-DSM⁷ relative to corresponding PAO1-UW proteins (Table 3). Fifteen changes seem to be PAO1161 strain specific (Table 3, bolded) as revealed by comparison of the sequences with other members from the corresponding Pseudomonas Ortholog Groups. These include codon deletions in orthologs of *pauA*, encoding a pimeloyl-CoA synthetase involved in biotin synthesis⁵³ and *PA3067*, a MarR family transcriptional regulator. Amino acids deletion was also found in KinB, forming a two-component system with the transcriptional regulator AlgB involved in control of alginate and virulence factors production in *P. aeruginosa*⁵⁴⁻⁵⁷. The point mutation changing amino acid sequence R221H was also detected in PAO1161 *algB* (Table 3). It is tempting to speculate that the identified changes may attenuate the virulence properties of PAO1161 cells.

Other strain specific substitutions include T103I change in the transcriptional regulator GcdR regulating glutarate utilization and lysine catabolism⁵⁸, A232T in the carbamate kinase ArcC involved in fermentative arginine degradation⁵⁹, V126M in the ribonuclease E (*rne*) involved in mRNA turnover⁶⁰. The G108D change was also identified in Tsi4 of PAO1161 relative to corresponding protein of PAO1-UW. This protein is a part of antibacterial effector-immunity pair together with type VI secretion exported toxin Tse4⁶¹. Overall, the identified substitutions could influence the structure and/or function of the corresponding proteins and should be considered in analyses involving these genes using PAO1161 strain.

PAO1161 genome contains an ICE conferring resistance to mercury

PAO1161 was described as a strain containing the FP2 plasmid of IncP-8 incompatibility group, which confers the cells with mercury resistance¹⁵. Indeed, the strain used in our lab was exceptionally resistant to mercury, growing in LB medium supplemented with up to 200 μ M HgCl₂ (data not shown). Surprisingly, during the genome assembly no extra-chromosomal elements could be identified. Instead, an almost 108 kbp insertion in the chromosome, with a putative mercury resistance operon, was found (Figure 1B, Figure 3A). The insertion shows similarities (in sequence and organization/composition of operons flanking the putative integration site) to the PAPI-1 family of integrative conjugative elements (ICEs)^{29,30,62} abundant in *Pseudomonas* genomes. ICEs are mobile genetic elements, with a modular structure, encoding complete conjugation machinery (usually a type IV secretion system) allowing transfer of their genome to another host. They are reversibly integrated into a host genome and can be passively propagated during bacterial chromosome segregation and cell division⁶²⁻⁶⁴. PAPI-1 (108 kb, 115 orfs, integrated in *tRNA*_{Lys}) was

first described in the genome of highly virulent *P. aeruginosa* PA14 strain⁶⁵.

The element identified within PAO1161 genome, which we named ICEPae1161, has an integration site within *tRNA_{Lys}* and PAPI-1 like organization of boundary operons: an operon starting with a gene encoding a putative ParA protein, at one end and an operon encoding a putative relaxase (TraI) and site-specific recombinase (Int) at the other (Figure 3A). Analysis of gene content (Supplementary Table S2), revealed that out of 120 predicted genes within ICEPae1161, 102 genes are found in at least one other PAPI-1 like element, whereas orthologs of 41 genes are found in all ICEs analysed (Supplementary Table S2).

Integration of ICE into the chromosome as well as its excision is mediated by an ICE encoded site directed recombinase / integrase⁶⁶. Recombination between an attachment site in the chromosome (*attB*) and the corresponding site on a circular ICE (*attP*) leads to integration of the element into the genome, now flanked by identical *attL* and *attR* sequences (Figure 3B). Excision of the ICEPae1161 and the presence of a circular form was analysed using PCR with primers flanking the *att* sequences (Figure 3BC). The analysis confirmed occurrence of the circular ICE in PAO1161 cells (Figure 3C), indicating that the element can exist in two forms.

To verify the transferability of ICEPae1161, we tagged it with a streptomycin resistance cassette (*aadA*). Subsequently, PAO1161 ICE-Sm^R strain was used as a donor in mating in static liquid cultures with *Pseudomonas putida* KT2440 as a recipient. The conjugants were selected on M9 plates supplemented with streptomycin, but lacking leucine to block the growth of donor cells. Streptomycin resistant *P. putida* clones were obtained with a low efficiency of 2×10^{-7} transconjugants / per donor cell.

To confirm that the *P. putida* conjugants show also enhanced resistance to mercury, attributed to the presence of *mer* operon within the ICEPae1161, we analysed the growth of transconjugants in medium containing HgCl₂. The recipient *P. putida* KT2440 cells were unable to grow at a HgCl₂ concentration higher than 2 μ M (data not shown). In contrast, the growth of streptomycin resistant *P. putida* transconjugants was not inhibited by the presence of 40 μ M HgCl₂ in the medium, confirming acquisition of mercury resistance (Figure 3D).

Finally, to confirm that the ICEPae1161 integrated into *P. putida* KT2440 chromosome, we searched for putative *attB* sites in its genome using the 48 bp TGGTGGGTCGTGTAGGATTCTGAACCTACGACCAATTGGTTAAAAGCCA sequence flanking ICEPae1161 in PAO1161 chromosome. *P. putida* KT2440 genome contains three potential attachment sites designated *attB1-3*, two of them the *attB1* and *attB2* adjacent to each other (Figure 3E). A PCR analysis, using primer pairs specific to the ICEPae1161/KT2440 chromosome junctions revealed predictably oriented ICEPae1161 in the *P. putida* genome (Figure 3F). Among six individual transconjugants, the specific PCR products were observed preferentially for ICE

integrated in one of adjacent sites, *attB1* and *attB2* with one clone also positive for ICE integration in *attB3* (Figure 3F). Overall, these data confirm the ability of the ICE*Pae*1161 identified in the PAO1161 genome to transfer to another host, integration into chromosome at a specific site and conferring mercury resistance.

Conclusions

In this work we show that *P. aeruginosa* PAO1161 strain carries a PAPI-1 family integrative conjugative element capable of excision, transfer and integration in the genome of the related *Pseudomonas* strain. The ICE*Pae*1161 contains loci conferring mercury resistance, in the past attributed to the FP2 plasmid of IncP-8 incompatibility group.

The genome sequence of *P. aeruginosa* PAO1161 strain, a derivative of the reference PAO1 strain, was compared with the reference PAO1 sequence. The data indicate a number of sequence variants, likely affecting different aspects of cell physiology. Thus genome sequencing of laboratory strains is highly recommended to provide full insight of the genotype and help in the interpretation of phenotypes observed under the conditions studied.

MATERIALS AND METHODS

Strains and growth conditions

Bacterial strains, plasmids and oligonucleotides used in this work are listed in Supplementary Table S3. *P. aeruginosa* PAO1161 (*leu⁻*, *r⁻*, *m⁻*) was provided by B. M. Holloway (Monash University, Clayton, Victoria, Australia). *Escherichia coli* strain DH5 α was used for plasmid manipulations and S17-1 was used to mate pAKE600⁶⁷ derivatives into *P. aeruginosa*. Standard DNA manipulations were performed as described⁶⁸. Templates for PCRs were prepared by boiling the cells pelleted from overnight cultures and resuspended in water.

Bacteria were grown in L broth⁶⁹ at 37°C or on L agar (L broth with 1.5% w/v agar) supplemented with appropriate antibiotics. For selection of *E. coli* strains 150 μ g ml⁻¹ benzylpenicillin sodium salt in liquid medium, 300 μ g ml⁻¹ for solid media or 30 μ g ml⁻¹ streptomycin was used. *P. aeruginosa* and *P. putida* strains were selected by addition of 300 μ g ml⁻¹ carbenicillin, 300 μ g ml⁻¹ rifampicin or 150 μ g ml⁻¹ streptomycin to the medium. Growth analysis was performed in M9 minimal medium⁶⁸ with 0.25% citrate or 0.1% glucose supplemented with 10 μ g ml⁻¹ leucine or 40 μ M HgCl₂ as indicated. Bacterial growth in 96-well plates was monitored by measurements of optical density at 600 nm (OD₆₀₀) using a Varioskan Lux Multimode Microplate Reader and SkanIt RE 5.0 software (Thermo Fisher Scientific).

Plasmids and strains construction

To construct prototrophic *P. aeruginosa* PAO1161 strain, which expresses LeuA without the E108K substitution, the suicide plasmid pKAB607 was constructed. Sequences flanking the

region in PAO1161 chromosome were amplified using primer pairs 1#, 2# and 3#, 4#, respectively (Supplementary Table S3). Primers 2# and 3# contained sequence lacking the mutation observed in PAO1161 genome relative to PAO1-UW and introduced an AflII site, allowing selection of the allele. The obtained PCR fragments were digested with EcoRI, AflII and AflII, BamHI, respectively, and both fragments were ligated with EcoRI, BamHI digested pAKE600⁶⁷ to yield pKAB607. *E. coli* S17-1 strain, carrying pKAB607 was used in the allele exchange procedure performed as described before¹⁸. Presence of modified allele was verified by AflII digestion of PCR amplified chromosomal *leuA*.

P. aeruginosa PAO1161 ICE-Sm^R strain was constructed by insertion of a streptomycin resistance cassette (*aadA*) between orfs D3C65_24365 and D3C65_24370 within ICE*Pae*1161 (Supplementary Table 2). To this end, first the fragments flanking the region of future insertion were amplified from PAO1161 genomic DNA using primer pairs 5#, 6# and 7#, 8#, respectively (Supplementary Table S3). The obtained PCR fragments were digested using MunI, HindIII and HindIII, BamHI, respectively, and the mixture was ligated with EcoRI, BamHI digested pAKE600 to yield pKAB608. The Sm^R cassette, excised with HindIII from pHP45Ω⁷⁰, was ligated with HindIII digested pKAB608 to yield pKAB609. *E. coli* S17-1 pKAB609 strain was used as a donor in the allele exchange procedure with strain PAO1161 Rif^{R18}. Integration of Sm^R cassette was confirmed by PCR.

Genome sequencing

For genome assembly, reads obtained previously from sequencing of the input sample of *P. aeruginosa* PAO1161 strain from ChIP-seq experiments were used¹⁹. The sequencing yielded 23380926 reads (4676185200 nt of raw data). Obtained reads were quality-filtered using FastX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/) and residual Illumina adapters were removed using Cutadapt (<https://github.com/marcelm/cutadapt>)⁷¹. A subsample (7000 000 reads) was used in draft genome assembly using Spades v3.11.1 (<http://cab.spbu.ru/software/spades/>) to estimate the size of PAO1161 genome.

Concomitantly, long reads were generated using MinION (Oxford Nanopore Technologies, Oxford, UK). Genomic DNA was sheared into 20 kb fragments using Covaris gTube (Covaris, Ltd., Brighton, United Kingdom) and the library was prepared using an ONT 1D ligation sequencing kit (SQK-LSK108) with the native barcoding expansion kit (EXP-NBD103). Nanopore sequencing was performed using the NC_48□h Sequencing_Run_FLO-MIN106_SQK-LSK108 protocol, R9.4.1 MinION flowcell and a MinION Mk1B instrument. Raw nanopore data were basecalled using Albacore v2.3.1 (Oxford Nanopore Technologies, Oxford, UK). Reads were quality-filtered and Porechop (<https://github.com/rrwick/Porechop>) and NanoFilt⁷² were used for adapter removal. Overall, MinION sequencing yielded 161877 reads (2160720766 nt), with a median read length of

12489 nucleotides. Long nanopore reads were assembled in a hybrid mode with the Illumina data using Unicycler v.0.4.6⁷³. Genome assembly resulted in 1 circular replicon of 6383803 bp. Errors identified in the assembly were verified by a PCR amplification of DNA fragments, followed by Sanger sequencing on an ABI3730xl Genetic Analyzer (Life Technologies, USA) using BigDye Terminator Mix v. 3.1 chemistry (Life Technologies, USA) followed by correction of the genome sequence using Seqman software (DNA Star, USA).

The assembled genome was annotated using the NCBI Prokaryotic Genome Annotation Pipeline⁷⁴. The nucleotide sequence has been deposited in NCBI Nucleotide database (accession number CP032126).

Genome analysis

Genome synteny between PAO1161 and PAO1-UW was visualized using EasyFig⁷⁵. Structural variations, SNPs, insertions and deletion between the PAO1161 and PAO1-UW (NC_002516) sequences were identified using Nucdiff³¹. The effect of SNP/MNP and indels was predicted using snpEff⁷⁶. Reads were mapped to the PAO1161 genome using Bowtie v2.3.4.2⁷⁷ followed by a verification of the quality of the assembly in regions differentiating PAO1161 and PAO1-UW genomes. Percentage of a given base, relative to bases from all reads, at positions in PAO1161 genome corresponding to the identified SNPs, MNPs and short insertions was analysed using bam-readcount (<https://github.com/genome/bam-readcount>). Regions with short deletions were inspected using Integrative Genomics Viewer v2.4.9⁷⁸. The variants present in more than 80% of the reads with an average mapping quality > 20, were considered as homozygous. In case of 86/100 variants the base frequency was higher than 99%.

ICE analysis and conjugal transfer

Comparative analysis of ICEs and gene ortholog prediction was performed using Mauve v2015-02-13⁷⁹. Genoplots were used to visualize gene distribution⁸⁰. ICE*Pae*1161 transfer from *P. aeruginosa* PAO1161 ICE-Sm^R to *P. putida* KT2440 was performed by growing the strains in L broth overnight at 37°C, harvesting the cells, resuspending in the same amount of medium and mixing in 1:2 (donor: recipient) ratio in plastic 1.5 ml tubes. The mixtures were incubated for 2 h at 37°C with shaking (300 rpm) and then 3 h without shaking, followed by centrifugation and resuspension of the cells in the same volume of 0.9% NaCl. To estimate the efficiency of ICE*Pae*1161 transfer a suspension was serially diluted in 0.9% NaCl and aliquots were spotted onto M9 minimal medium agar plates with 0.1% glucose and 150 µg ml⁻¹ streptomycin without leucine. Lack of leucine allowed growth of *P. putida* KT2440 transconjugants and counter selected PAO1161 *leu*⁻ donor cells. Donor or recipient strain cultures treated in the same way were used to establish the titre of donor and recipient cells in the conjugation mixture. The transfer frequency was calculated as the number of transconjugants per donor cell.

SUPPLEMENTARY MATERIALS

Supplementary Table S1. Synonymous and intergenic SNPs/MNPs and indels identified in PAO1161 / PAO1-UW genome comparison.

Supplementary Table S2. Gene content of ICE*Pae*1161. Orthologs from other ICE were predicted using Mauve⁷⁹ assuming 70% of identity and at least 70% of the sequence coverage.

Supplementary Table S3. Strains, plasmids and primers used in this study.

ACKNOWLEDGEMENTS

This work was funded by National Science Centre, Poland [2015/2015/18/E/NZ2/00675 granted to A.A.B. and 2013/11/B/NZ2/02555 granted to G.J.B.]. We thank Jan Gawor and Robert Gromadka (DNA Sequencing and Oligonucleotides Synthesis Laboratory, IBB PAS, Warsaw, Poland) for performing the DNA sequencing.

AUTHOR CONTRIBUTIONS

A.A.B., A.K. and G.J.B conceived and designed the experiments. A.K., K.K., M.M and A.A.B. conducted the experiments. A.K., K.K., M.M., G.J.B and A.A.B. analyzed and interpreted the results. A.K. and A.A.B. performed bioinformatics analysis. A.K., G.J.B. and A.A.B. wrote the manuscript. All authors reviewed and approved the manuscript.

COMPETING INTERESTS

The author(s) declare no competing interests.

bioRxiv preprint doi: <https://doi.org/10.1101/494302>; this version posted January 2, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure 2. LeuA E108K substitution causes leucine auxotrophy in *P. aeruginosa*. PAO1161 *leuA* allele, carrying the mutation, was replaced with the PAO1 allele to yield strain PAO1161 rev (*leu+*). Growth of PAO1161 (*leu-*) and PAO1161 rev (*leu+*) strains on solid (A) and liquid (B) minimal medium containing 0.25% citrate with or without 10 $\mu\text{g ml}^{-1}$ leucine. Data represent mean OD_{600nm}±SD for 6 biological replicates.

Figure 3. ICEPae1161 identified in *P. aeruginosa* PAO1161 genome is a functional PAPI-1 family integrative and conjugative element conferring resistance to mercury. (A) Comparative genomics of ICEPae1161 and selected PAPI-1 family ICEs. Mauve alignment of ICEPae1161 and PAPI-1⁶⁵, PAGI-5⁸¹, PFGI-1⁸², pKLC102⁸³ and ICEPaePACS2-1 (position 896693:1002644 of NZ_AAQW01000001)²⁹ is presented. Three blocks of sequences which are free of genome rearrangements, such as inversions and duplications, are marked with rectangles connected with

lines. Green segments indicate sequences conserved in all ICEs (backbone). Regions conserved among subsets of analysed ICEs are color coded. White region is specific only to one analysed ICE. Arrows indicate the location and orientation of coding sequences. Boundary genes and the mercury resistance operon are shown on top. **(B)** Schematic model of linear and excised (circular) ICE*Pae*1161. Primer binding sites are indicated in red. **(C)** PCR analysis of ICE*Pae*1161 excision. PCR was performed with indicated primer pairs flanking *att* sequences using PAO1161 genomic DNA as a template and products were separated on a 1.2% agarose gel. **(D)** Growth of *P. putida* KT2440 strain and transconjugants *P. putida* KT2440 ICE*Pae*1161-Sm^R. Strains were grown in L broth with or without 40 μ M HgCl₂. Data represent mean OD_{600nm}±SD for 6 independent transconjugants analysed in 3 biological replicates. **(E)** Genomic context of three potential ICE*Pae*1161 integration sites (*attB*) in *P. putida* KT2440 genome. The sites were identified based on the presence of 48 bp sequence flanking ICE*Pae*1161 in PAO1161 genome. Blue arrows indicate the orientation of the sequences. Schematic model of ICE*Pae*1161 integration at each *attB* site in orientation corresponding to the one observed in PAO1161 genome. **(F)** PCR analysis of ICE*Pae*1161 presence in *P. putida* KT2440. Genomic DNA of six independent transconjugants was used as a template in PCR using the indicated primers and products were separated on 1% agarose gels.

Table 1 SNPs and indels identified in *P. aeruginosa* PAO1161 genome, resulting in expression of truncated proteins. The effect of a mutation is predicted using the PAO1-UW genome as a reference.

Mutation effect	PAO1-UW position	Nucleotide change	AA change	length PAO1/PAO1161	PAO1 gene	PAO1161 ID	Description
stop gained	2121203	C → T	W340*	665 / 339	PA1939	D3C65_15950	putative ATP-dependent endonuclease of the OLD family
	2356682	CC → C	L173	182 / 172	PA2141	D3C65_14865	CinA family protein
	3097884	G → A	Q209*	792 / 208	PA2735	D3C65_11725	SAM-dependent DNA methyltransferase
frame-shift	740419	G → GC	V73A?	381 / 124	PA0683 (<i>hxcY</i>)	D3C65_22610	putative type II secretion system protein
	1440623	AA → A	K640	656 / 642	PA1327	D3C65_19175	putative protease
	1835045	G → GC [†]	S218S?	249 / 226	PA1685 (<i>masA</i>)	D3C65_17305	enolase-phosphatase E-1
	2807706	CAGCCGGCC → C	aa1-78/35aa	347 / 304	PA2492 (<i>mexT</i>)	D3C65_13040	transcriptional regulator

[†] - SNP at this position in PAO1DSM / MPAO-1⁷ but a nucleotide insertion in our study

Table 2 SNPs and indels identified in *P. aeruginosa* PAO1161 genome influencing the reading frames of genes. The effect of a mutation is predicted using the PAO1-UW genome as a reference. The sequences of PAO1161 proteins were compared with sequences from corresponding Pseudomonas Ortholog Group (POG) for PAO1 proteins.

Position in PAO1-UW	Nucleotide change	PAO1 loci	PAO1161 loci	Effect of mutation
816531	C → CC	PA0748	D3C65_22250	Corrects the reading frame of PA0748 (still frameshift orf).
1215658	G → GG	PA1122	D3C65_20255	Reading frame change after 125 aa. Fusion with 54 aa instead of 22 aa (PAO1). Similar to 498/504 ortologs from POG001082.
2355771	A → AG	PA2139	D3C65_14875	Reading frame change after 29 aa. Fusion with 42 aa instead of 11 aa (PAO1). Similar to 6/12 ortologs from POG008397.
2753522	G → GC	PA2451	D3C65_13240	Fusion of PA2452 and PA2451. No fusion only in PAO1-UW (POG003088).
3083196	A → AG	PA2727	D3C65_11765	Fusion of PA2727 with PA2728. Similar fusion observed for 366/372 proteins from POG002843.
4539468	G → GC	PA4058	D3C65_04690	Fusion of PA4059 with PA4060. No fusion only in PAO1-UW (POG001625).
413850	T → C	IG: PA0369- PA0370	D3C65_01955	Removed TGA codon -12 to -10 from <i>PA0369</i> start codon. Predicted product has 30aa fused to N-terminus of PA0369. Similar to 297/303 ortologs from POG000360.
2239555	A → AG	IG: PA2046- PA2047	D3C65_15395	Change in sequence preceding <i>PA2046</i> . Predicted product has 316 aa fused to N-terminus of PA2046. Similar to 289/298 orthologs from POG003399.
5472415	C → CG	IG: PA4874- PA4875	D3C65_26560	Change in sequence preceding <i>PA4875</i> . Predicted product has 81 aa fused to N-terminus of PA4875. Similar to 292/299 orthologs from POG004503.
1116213	G → GC	PA1029	D3C65_20745	Removes start codon of <i>PA1029</i> . Predicted product has additional 24 aa fused to N- terminus of PA1029. Similar to 310/316 orthologs from POG000991.
4888194	A → AG	PA4360	D3C65_23120	Removes start codon of <i>PA4360</i> . Proposed product has additional 122 aa fused to N- terminus of PA4360. This extension is observed for most members of POG003953.
1275766	GA → G	PA1174	D3C65_19970	First 11 aa of PA1174 replaced by 16aa. Extended sequences in 343/350 ortologs from POG001132.
3016844	G → GC	PA2668	D3C65_12075	Reading frame and start codon changed. Predicted product has last 56 aa identical to C-terminus of PA2668. Extended sequences in 295/301 orthologs from POG008397.
3919508	G → GC	PA3503	D3C65_07595	Predicted product has last 210 aa identical to C-terminus of PA3503. Extended sequences in 69/73 orthologs from POG005142.
1023047	GGCAAGAT → G	PA0934 (<i>relA</i>)	D3C65_21265	Start codon shift and M1VVKGR change. Only in PAO1161.

Table 3 SNPs and indels identified in *P. aeruginosa* PAO1161 genome resulting in amino acid changes relative to corresponding PAO1-UW proteins. PAO1161 strain specific changes, identified by comparison of PAO1161 proteins with sequences in Pseudomonas Ortholog Groups encompassing corresponding PAO1 proteins, are indicated by bolded gene names.

Effect	PAO1-UW position	Nucleotide change	Amino change	acid PAO1 gene	Description
codon change and codon deletion	1102459	CGAGGCGGTGCTGGGCAGTGCCGA A → C	VLGSAEEAV513 V	PA1017 (<i>pauA</i>)	pimeloyl-CoA synthetase
codon deletion	3436070	CGTGCATGGCACG → C	RAMH117	PA3067	transcriptional regulator
	6176156	AGCTCGATCGCCTGCA→A	LDRLQ489	PA5484 (<i>kinB</i>)	histidine kinase
codon insertion	5655229	G → GCGG	W226GW	PA5024	conserved hypothetical protein
missense	183697	T → G*	C310W	PA0159	probable transcriptional regulator
	504428	C → T	T103I	PA0448 (<i>gcdR</i>)	transcriptional regulator
	1589438	G → C [#]	G34A	PA1459	probable methyltransferase
	2669175	G → C [#]	P819A	PA2400 (<i>pvdJ</i>)	pyoverdine side chain peptide synthetase
	2806625	C → T	D249N	PA2491 (<i>mexS</i>)	probable oxidoreductase
	2807982	T → A*	F172I	PA2492 (<i>mexT</i>)	transcriptional regulator
	3132936	C → T	G108D	PA2775 (<i>tsi4</i>)	immunity protein
	3133912	C → T	S68F	PA2776 (<i>pauB3</i>)	FAD-dependent oxidoreductase
	3333256	G → A	V126M	PA2976 (<i>rne</i>)	ribonuclease E
	4212201	A → G*	H636R	PA3760	N-Acetyl-D-Glucosamine phosphotransferase transporter
	4251149	G → A	E108K	PA3792 (<i>leuA</i>)	2-isopropylmalate synthase
	4779026	T → A	H531L	PA4270 (<i>rpoB</i>)	DNA-directed RNA polymerase
	486985	T → G	E158D	PA4341	probable

5				transcriptional regulator
492455 2	CG → GC*	PV177PL	PA4394	mechanosensitive ion channel protein
533648 7	C → G	M405I	PA4751 (<i>ftsH</i>)	cell division protein
574346 1	CG → GC*	T431S	PA5100 (<i>hutU</i>)	urocanase
582548 0	G → A	A232T	PA5173 (<i>arcC</i>)	carbamate kinase
6115455	T → G [#]	K286N	PA5434 (<i>mtr</i>)	tryptophan permease
613159 2	C → T	L169F	PA5443 (<i>uvrD</i>)	DNA helicase II
617400 9	G → A	R221H	PA5483 (<i>algB</i>)	two-component response regulator
623303 9	G → A	S162N	PA5540	hypothetical protein

* - observed in PAO1DSM / MPAO-1⁷

[#] - observed in PAO1DSM / MPAO-1 and validated experimentally⁷

References

1. Klockgether, J. & Tümmler, B. Recent advances in understanding *Pseudomonas aeruginosa* as a pathogen. *F1000Res* **6**, (2017).
2. Azam, M. W. & Khan, A. U. Updates on the pathogenicity status of *Pseudomonas aeruginosa*. *Drug Discovery Today* (2018). doi:10.1016/j.drudis.2018.07.003
3. Holloway, B. W. Genetic recombination in *Pseudomonas aeruginosa*. *J. Gen. Microbiol.* **13**, 572–581 (1955).
4. Stanisich, V. & Holloway, B. W. Conjugation in *PSEUDOMONAS AERUGINOSA*. *Genetics* **61**, 327–339 (1969).
5. Stover, C. K. *et al.* Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* **406**, 959–964 (2000).
6. Winsor, G. L. *et al.* Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. *Nucleic Acids Res* **44**, D646–D653 (2016).
7. Klockgether, J. *et al.* Genome diversity of *Pseudomonas aeruginosa* PAO1 laboratory strains. *J. Bacteriol.* **192**, 1113–1121 (2010).
8. Bhowmik, B. K., Clevenger, A. L., Zhao, H. & Rybenkov, V. V. Segregation but Not Replication of the *Pseudomonas aeruginosa* Chromosome Terminates at Dif. *mBio* **9**, (2018).

9. Sidorenko, J., Jatsenko, T. & Kivisaar, M. Ongoing evolution of *Pseudomonas aeruginosa* PAO1 sublines complicates studies of DNA damage repair and tolerance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **797–799**, 26–37 (2017).
10. Luong, P. M. *et al.* Emergence of the P2 Phenotype in *Pseudomonas aeruginosa* PAO1 Strains Involves Various Mutations in *mexT* or *mexF*. *Journal of Bacteriology* **196**, 504–513 (2014).
11. Dunn, N. W. & Holloway, B. W. Pleiotropy of p-fluorophenylalanine-resistant and antibiotic hypersensitive mutants of *Pseudomonas aeruginosa*. *Genetics Research* **18**, 185–197 (1971).
12. Finger, J. & Krishnapillai, V. Host range, entry exclusion, and incompatibility of *Pseudomonas aeruginosa* FP plasmids. *Plasmid* **3**, 332–342 (1980).
13. Shintani, M., Sanchez, Z. K. & Kimbara, K. Genomics of microbial plasmids: classification and identification based on replication and transfer systems and host taxonomy. *Front. Microbiol.* **6**, (2015).
14. Holloway, B. W. & Fargie, B. Fertility factors and genetic linkage in *Pseudomonas aeruginosa*. *J. Bacteriol.* **80**, 362–368 (1960).
15. Stanisich, V. A. & Holloway, B. W. A mutant sex factor of *Pseudomonas aeruginosa*. *Genet. Res.* **19**, 91–108 (1972).
16. Kung, A. H. C. & Lee, B. T. O. Genetic analysis of radiation sensitive and chemical-mutagen sensitive mutants of *Pseudomonas aeruginosa*. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **27**, 191–199 (1975).
17. Rolfe, B. & Holloway, B. W. Genetic control of DNA specificity in *Pseudomonas aeruginosa*. *Genetics Research* **12**, 99–102 (1968).
18. Lasocki, K., Bartosik, A. A., Mierzejewska, J., Thomas, C. M. & Jagura-Burdzy, G. Deletion of the *parA* (*soj*) homologue in *Pseudomonas aeruginosa* causes *ParB* instability and affects growth rate, chromosome segregation, and motility. *J. Bacteriol.* **189**, 5762–5772 (2007).
19. Kawalek, A., Bartosik, A. A., Glabski, K. & Jagura-Burdzy, G. *Pseudomonas aeruginosa* partitioning protein *ParB* acts as a nucleoid-associated protein binding to multiple copies of a *parS*-related motif. *Nucleic Acids Res.* **46**, 4592–4606 (2018).
20. Bartosik, A. A. *et al.* Transcriptional profiling of *ParA* and *ParB* mutants in actively dividing cells of an opportunistic human pathogen *Pseudomonas aeruginosa*. *PLoS ONE* **9**, e87276 (2014).
21. Kawalek, A., Glabski, K., Bartosik, A. A., Fogtman, A. & Jagura-Burdzy, G. Increased *ParB* level affects expression of stress response, adaptation and virulence operons and potentiates repression of promoters adjacent to the high affinity binding sites *parS3* and *parS4* in *Pseudomonas aeruginosa*. *PLOS ONE* **12**, e0181726 (2017).

22. Ciok, A. *et al.* Plasmids of Psychrotolerant Polaromonas spp. Isolated From Arctic and Antarctic Glaciers – Diversity and Role in Adaptation to Polar Environments. *Front. Microbiol.* **9**, (2018).
23. Komatsu, T., Moriya, K. & Horikoshi, K. Preparation of Organic Solvent-Tolerant Mutants from Pseudomonas aeruginosa Strain PAO1161. *Bioscience, Biotechnology, and Biochemistry* **58**, 1754–1755 (1994).
24. Manavathi, B., Pakala, S. B., Gorla, P., Merrick, M. & Siddavattam, D. Influence of zinc and cobalt on expression and activity of parathion hydrolase from Flavobacterium sp. ATCC27551. *Pesticide Biochemistry and Physiology* **83**, 37–45 (2005).
25. Kujawa, M. *et al.* Nudix-type RNA pyrophosphohydrolase provides homeostasis of virulence factor pyocyanin and functions as a global regulator in Pseudomonas aeruginosa. *Molecular Microbiology* **106**, 381–394 (2017).
26. Papagiannitsis, C. C., Studentova, V., Ruzicka, F., Tejkalova, R. & Hrabak, J. Molecular characterization of metallo- β -lactamase-producing Pseudomonas aeruginosa in a Czech hospital (2009-2011). *J. Med. Microbiol.* **62**, 945–947 (2013).
27. Laudy, A. E., Osińska, P., Namysłowska, A., Zajac, O. & Tyski, S. Modification of the Susceptibility of Gram-Negative Rods Producing ES β LS to β -Lactams by the Efflux Phenomenon. *PLOS ONE* **10**, e0119997 (2015).
28. Romaniuk, K., Krucon, T., Decewicz, P., Gorecki, A. & Dziewit, L. Molecular characterization of the pA3J1 plasmid from the psychrotolerant Antarctic bacterium Pseudomonas sp. ANT_J3. *Plasmid* **92**, 49–56 (2017).
29. Liu, M. *et al.* ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res* doi:10.1093/nar/gky1123
30. Qiu, X., Gurkar, A. U. & Lory, S. Interstrain transfer of the large pathogenicity island (PAPI-1) of Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 19830–19835 (2006).
31. Khelik, K., Lagesen, K., Sandve, G. K., Rognes, T. & Nederbragt, A. J. NucDiff: in-depth characterization and annotation of differences between two sets of DNA sequences. *BMC Bioinformatics* **18**, 338 (2017).
32. Doberenz, S. *et al.* Identification of a Pseudomonas aeruginosa PAO1 DNA Methyltransferase, Its Targets, and Physiological Roles. *mBio* **8**, e02312-16 (2017).
33. Aravind, L., Leipe, D. D. & Koonin, E. V. Toprim—a conserved catalytic domain in type IA and II topoisomerases, DnaG-type primases, OLD family nucleases and RecR proteins. *Nucleic Acids Res* **26**, 4205–4213 (1998).
34. Cianciotto, N. P. & White, R. C. Expanding Role of Type II Secretion in Bacterial Pathogenesis and Beyond. *Infection and Immunity* **85**, e00014-17 (2017).

35. Ball, G., Durand, É., Lazdunski, A. & Filloux, A. A novel type II secretion system in *Pseudomonas aeruginosa*. *Molecular Microbiology* **43**, 475–485 (2002).
36. Ball, G., Viarre, V., Garvis, S., Voulhoux, R. & Filloux, A. Type II-dependent secretion of a *Pseudomonas aeruginosa* DING protein. *Res. Microbiol.* **163**, 457–469 (2012).
37. Xu, X. *et al.* Role of ppGpp in *Pseudomonas aeruginosa* acute pulmonary infection and virulence regulation. *Microbiol. Res.* **192**, 84–95 (2016).
38. Malinowska, A. *et al.* Diffprot - software for non-parametric statistical analysis of differential proteomics data. *J Proteomics* **75**, 4062–4073 (2012).
39. Köhler, T., Epp, S. F., Curty, L. K. & Pechère, J.-C. Characterization of MexT, the Regulator of the MexE-MexF-OprN Multidrug Efflux System of *Pseudomonas aeruginosa*. *Journal of Bacteriology* **181**, 6300–6305 (1999).
40. Uwate, M. *et al.* Two routes of MexS-MexT-mediated regulation of MexEF-OprN and MexAB-OprM efflux pump expression in *Pseudomonas aeruginosa*. *Microbiol. Immunol.* **57**, 263–272 (2013).
41. Köhler, T., Delden, C. van, Curty, L. K., Hamzehpour, M. M. & Pechere, J.-C. Overexpression of the MexEF-OprN Multidrug Efflux System Affects Cell-to-Cell Signaling in *Pseudomonas aeruginosa*. *Journal of Bacteriology* **183**, 5213–5222 (2001).
42. Lamarche, M. G. & Déziel, E. MexEF-OprN Efflux Pump Exports the *Pseudomonas* Quinolone Signal (PQS) Precursor HHQ (4-hydroxy-2-heptylquinoline). *PLOS ONE* **6**, e24310 (2011).
43. Maseda, H., Saito, K., Nakajima, A. & Nakae, T. Variation of the mexT gene, a regulator of the MexEF-OprN efflux pump expression in wild-type strains of *Pseudomonas aeruginosa*. *FEMS Microbiology Letters* **192**, 107–112 (2000).
44. Sobel, M. L., Neshat, S. & Poole, K. Mutations in PA2491 (mexS) Promote MexT-Dependent mexEF-oprN Expression and Multidrug Resistance in a Clinical Strain of *Pseudomonas aeruginosa*. *J Bacteriol* **187**, 1246–1253 (2005).
45. Fargier, E. *et al.* MexT Functions as a Redox-Responsive Regulator Modulating Disulfide Stress Resistance in *Pseudomonas aeruginosa*. *Journal of Bacteriology* **194**, 3502–3511 (2012).
46. Withers, T. R., Johnson, S. L. & Yu, H. D. Draft Genome Sequence for *Pseudomonas aeruginosa* Strain PAO579, a Mucoïd Derivative of PAO381. *J Bacteriol* **194**, 6617 (2012).
47. Ryan Withers, T., Heath Damron, F., Yin, Y. & Yu, H. D. Truncation of type IV pilin induces mucoidy in *Pseudomonas aeruginosa* strain PAO579. *Microbiologyopen* **2**, 459–470 (2013).
48. Yin, Y., Withers, T. R., Govan, J. R. W., Johnson, S. L. & Yu, H. D. Draft Genome Sequence of a Stable Mucoïd Strain of *Pseudomonas aeruginosa* PAO581 with a mucA25 Mutation. *Genome Announc* **1**, (2013).

49. Campbell, E. A. *et al.* Structural mechanism for rifampicin inhibition of bacterial RNA polymerase. *Cell* **104**, 901–912 (2001).
50. Severinov, K., Soushko, M., Goldfarb, A. & Nikiforov, V. Rifampicin region revisited. New rifampicin-resistant and streptolydigin-resistant mutants in the beta subunit of Escherichia coli RNA polymerase. *J. Biol. Chem.* **268**, 14820–14825 (1993).
51. Hall, A. R., Iles, J. C. & MacLean, R. C. The Fitness Cost of Rifampicin Resistance in Pseudomonas aeruginosa Depends on Demand for RNA Polymerase. *Genetics* **187**, 817–822 (2011).
52. Jatsenko, T., Tover, A., Tegova, R. & Kivisaar, M. Molecular characterization of Rifr mutations in Pseudomonas aeruginosa and Pseudomonas putida. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **683**, 106–114 (2010).
53. Binieda, A. *et al.* Purification, characterization, DNA sequence and cloning of a pimeloyl-CoA synthetase from Pseudomonas mendocina 35. *Biochem. J.* **340** (Pt 3), 793–801 (1999).
54. Ma, S., Wozniak, D. J. & Ohman, D. E. Identification of the histidine protein kinase KinB in Pseudomonas aeruginosa and its phosphorylation of the alginate regulator algB. *J. Biol. Chem.* **272**, 17952–17960 (1997).
55. Damron, F. H., Qiu, D. & Yu, H. D. The Pseudomonas aeruginosa sensor kinase KinB negatively controls alginate production through AlgW-dependent MucA proteolysis. *J. Bacteriol.* **191**, 2285–2295 (2009).
56. Chand, N. S., Clatworthy, A. E. & Hung, D. T. The two-component sensor KinB acts as a phosphatase to regulate Pseudomonas aeruginosa Virulence. *J. Bacteriol.* **194**, 6537–6547 (2012).
57. Balasubramanian, D., Schneper, L., Kumari, H. & Mathee, K. A dynamic and intricate regulatory network determines Pseudomonas aeruginosa virulence. *Nucleic Acids Res.* **41**, 1–20 (2013).
58. Madhuri Indurthi, S., Chou, H.-T. & Lu, C.-D. Molecular characterization of lysR-lysXE, gcdR-gcdHG and amaR-amaAB operons for lysine export and catabolism: a comprehensive lysine catabolic network in Pseudomonas aeruginosa PAO1. *Microbiology* **162**, 876–888 (2016).
59. Schreiber, K. *et al.* Anaerobic survival of Pseudomonas aeruginosa by pyruvate fermentation requires an Usp-type stress protein. *J. Bacteriol.* **188**, 659–668 (2006).
60. Kushner, S. R. mRNA decay in prokaryotes and eukaryotes: different approaches to a similar problem. *IUBMB Life* **56**, 585–594 (2004).
61. Whitney, J. C. *et al.* Genetically distinct pathways guide effector export through the type VI secretion system. *Mol. Microbiol.* **92**, 529–542 (2014).

62. Johnson, C. M. & Grossman, A. D. Integrative and Conjugative Elements (ICEs): What They Do and How They Work. *Annu. Rev. Genet.* **49**, 577–601 (2015).
63. Delavat, F., Miyazaki, R., Carraro, N., Pradervand, N. & van der Meer, J. R. The hidden life of integrative and conjugative elements. *FEMS Microbiol. Rev.* **41**, 512–537 (2017).
64. Wozniak, R. A. F. & Waldor, M. K. Integrative and conjugative elements: mosaic mobile genetic elements enabling dynamic lateral gene flow. *Nature Reviews Microbiology* **8**, 552–563 (2010).
65. He, J. *et al.* The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 2530–2535 (2004).
66. Grindley, N. D. F., Whiteson, K. L. & Rice, P. A. Mechanisms of site-specific recombination. *Annu. Rev. Biochem.* **75**, 567–605 (2006).
67. El-Sayed, A. K., Hothersall, J. & Thomas, C. M. Quorum-sensing-dependent regulation of biosynthesis of the polyketide antibiotic mupirocin in *Pseudomonas fluorescens* NCIMB 10586. *Microbiology (Reading, Engl.)* **147**, 2127–2139 (2001).
68. Molecular cloning□: a laboratory manual / J. Sambrook, E.F. Fritsch, T. Maniatis. *ResearchGate*
69. Kahn, M. *et al.* Plasmid cloning vehicles derived from plasmids ColE1, F, R6K, and RK2. *Meth. Enzymol.* **68**, 268–280 (1979).
70. Fellay, R., Frey, J. & Krisch, H. Interposon mutagenesis of soil and water bacteria: a family of DNA fragments designed for in vitro insertional mutagenesis of gram-negative bacteria. *Gene* **52**, 147–154 (1987).
71. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).
72. De Coster, W., D’Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669 (2018).
73. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput. Biol.* **13**, e1005595 (2017).
74. Tatusova, T. *et al.* NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res.* **44**, 6614–6624 (2016).
75. Sullivan, M. J., Petty, N. K. & Beatson, S. A. Easyfig: a genome comparison visualizer. *Bioinformatics* **27**, 1009–1010 (2011).
76. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)* **6**, 80–92 (2012).

77. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).
78. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief. Bioinformatics* **14**, 178–192 (2013).
79. Darling, A. C. E., Mau, B., Blattner, F. R. & Perna, N. T. Mauve: Multiple Alignment of Conserved Genomic Sequence With Rearrangements. *Genome Res* **14**, 1394–1403 (2004).
80. Guy, L., Kultima, J. R. & Andersson, S. G. E. genoPlotR: comparative gene and genome visualization in R. *Bioinformatics* **26**, 2334–2335 (2010).
81. Battle, S. E., Meyer, F., Rello, J., Kung, V. L. & Hauser, A. R. Hybrid pathogenicity island PAGI-5 contributes to the highly virulent phenotype of a *Pseudomonas aeruginosa* isolate in mammals. *J. Bacteriol.* **190**, 7130–7140 (2008).
82. Mavrodi, D. V., Loper, J. E., Paulsen, I. T. & Thomashow, L. S. Mobile genetic elements in the genome of the beneficial rhizobacterium *Pseudomonas fluorescens* Pf-5. *BMC Microbiol.* **9**, 8 (2009).
83. Klockgether, J., Reva, O., Larbig, K. & Tümmler, B. Sequence analysis of the mobile genome island pKLC102 of *Pseudomonas aeruginosa* C. *J. Bacteriol.* **186**, 518–534 (2004).