

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Fecal microbiota dysbiosis in macaques and humans within a shared environment

Erica Grant¹, Randall C. Kyes², Pensri Kyes², Pauline Trinh¹, Vickie Ramirez¹, Tawatchai Tanee³ Porntip Pinlaor⁴, Rungtiwa Dangtakot⁵, and Peter M. Rabinowitz^{1*}

¹ Center for One Health Research, Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, Washington, United States of America

² Department of Psychology and Washington National Primate Research Center, University of Washington, Seattle, Washington, United States of America

³ Faculty of Environment and Resource Studies, Mahasarakham University, Maha Sarakham and Genetics and Environmental Toxicology Group, Khon Kaen University, Khon Kaen, Thailand

⁴ Centre for Research and Development of Medical Diagnostic Laboratories, Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, Thailand

⁵ Biomedical Science Program, Graduate School, Khon Kaen University, Khon Kaen, Thailand

* Corresponding author

Email: peterr7@uw.edu

24 **Abstract**

25 Traditional zoonotic disease research focuses on detection of recognized pathogens and may
26 miss opportunities to understand broader microbial transmission dynamics between humans,
27 animals, and the environment. We studied human-macaque microbiome overlap in Kosum Phisai
28 District, Maha Sarakham Province, Thailand, where a growing population of long-tailed
29 macaques (*Macaca fascicularis*) in Kosumpee Forest Park interact with humans from an adjacent
30 village. We surveyed workers in or near the park with elevated exposure to macaques to
31 characterize tasks resulting in exposure to macaque feces in addition to dietary and lifestyle
32 factors that influence gut microbiome composition. Fecal samples were collected from 12
33 exposed workers and 6 controls without macaque exposure, as well as 8 macaques from
34 Kosumpee Forest Park and 4 from an isolated forest patch with minimal human contact. The V4
35 region of the 16S rRNA gene from fecal sample extracted DNA was amplified and sequenced
36 using Illumina HiSeq to characterize the microbial community. A permuted betadisper test on the
37 weighted UniFrac distances revealed significant differences in the dispersion patterns of gut
38 microbiota from exposed and control macaques ($p=0.03$). The high variance in gut microbiota
39 composition of macaques in contact with humans has potential implications for gut microbiome
40 stability and susceptibility to disease, described by the Anna Karenina principle (AKP). Human
41 samples had homogenous variance in beta diversity but different spatial medians between groups
42 ($p=0.02$), indicating a shift in microbial composition that may be explained by fundamental
43 lifestyle differences between the groups unrelated to exposure status. SourceTracker was used to
44 estimate the percent of gut taxa in exposed humans that was contributed by macaques. While one
45 worker showed evidence of elevated contribution, the overall trend was not significant. Task
46 observations among workers revealed opportunities to employ protective measures or training to

47 reduce exposure to occupational hazards. These results suggest the potential for hygiene
48 measures to mitigate negative aspects of contact between humans and macaques in order to
49 optimize the health of both populations.

50

51 **Introduction**

52 **Background**

53 Habitat fragmentation and human encroachment results in a patchwork of isolated non-
54 human primate populations across Thailand with potential for increased human-macaque contact
55 [1]. Supplemental feeding for religious reasons or tourism contributes to a growing macaque
56 population unconstrained by natural food resources. Amidst these changes, the high level of
57 human-macaque conflict has led researchers to call for improved management plans and
58 conservation strategies [1, 2]. The reasons for concern are two-fold: 1) human-macaque
59 conflict—such as crop-raiding—can disrupt or damage livelihoods, resulting in negative
60 perception of macaques and impairing conservation efforts and 2) the increased level of contact
61 can provide opportunities for transmission of zoonotic diseases into either macaque or human
62 populations.

63 Due to their genetic similarity, humans and macaques are susceptible to many of the same
64 infectious diseases including tuberculosis and hepatitis [3]. Parasitic infections from soil-
65 transmitted helminths capable of infecting humans and macaques, such as *Strongyloides*
66 *fuelleborni* and *S. stercoralis* have also been documented in this region [4, 5]. These pathogens
67 may (e.g. Ebola virus) or may not (e.g. Herpes B virus) cause overt disease in macaques,
68 however cross-species pathogen transmission from wildlife reservoirs to human hosts is a
69 recognized factor in the emergence of novel diseases [6]. These spillover events can readily

70 occur where natural and urban spaces meet. An example of such an interface is the Kosumpee
71 Forest Park (KFP), a small fragmented forest in northeastern Thailand that is home to over 700
72 long-tailed macaques (*Macaca fascicularis*) and lies adjacent to the Kosum Phisai community of
73 4,235 persons [7, 8]. Unlike other popular tourist sites in Southeast Asia, macaques in KFP
74 rarely climb onto people and feeding by the people often involves simply throwing food on the
75 ground [9]. However, workers in and around the park regularly feed macaques and sweep
76 macaque excrement from public spaces, elevating their risk of zoonotic disease transmission
77 relative to other members of the Kosum Phisai community. While these workers represent a
78 control point for broader spread of zoonoses, little is known about their knowledge, attitudes, and
79 practices surrounding macaque exposure [10]. Based on task observations, it may be possible to
80 identify intervention strategies to reduce exposure to macaque biological material. Such
81 strategies could include the use of personal protective equipment (PPE), training in basic hand
82 hygiene, or other measures to mitigate the risk of disease transmission. These measures would
83 also promote responsible wildlife conservation by protecting macaques from pathogens that the
84 workers could transmit through reverse zoonotic transmission.

85 The zoonotic spillover potential of certain pathogens, such as simian foamy virus, can be
86 investigated through blood sample collection, however this method can be logistically
87 challenging since it requires trapping and immobilization of wild macaques. Additionally,
88 previous surveys among this worker population indicated that scratches or bites are infrequent,
89 and ingestion of aerosolized fecal matter may be a more common route of exposure to zoonoses.
90 Therefore, in this setting, we chose to analyze the microbial communities of fecal samples in
91 humans and macaques with close contact. Advantages of analyzing the fecal microbiota include
92 the fact that, compared to the skin microbiota, it is better characterized in literature, more

93 temporally stable, and yield higher read counts [11]. Recent studies have demonstrated that the
94 community composition of human microbiota is influenced by our environment and the animals
95 sharing that environment. The degree of contribution from these sources can be quantified using
96 Bayesian approaches like SourceTracker [12, 13]. We performed a pilot study of fecal
97 microbiota of workers and macaques in a shared environment to test the hypothesis that workers
98 exposed to macaques will exhibit microbiota profiles that contain a greater percentage of
99 microbes found in macaque feces compared to unexposed individuals. The goals of this study
100 were to provide a baseline assessment of the risk of zoonotic disease transmission between
101 macaques and workers and guide prevention recommendations.

102

103 **Materials and methods**

104 **Study design**

105 This pilot study was a cross-sectional sampling of humans and macaques, comparing
106 humans with occupational contact with macaques (exposed humans) to humans without such
107 contact (human controls), and macaques in close contact with humans (exposed macaques)
108 compared to macaques without significant human contact (control macaques).

109

110 **Human participants**

111 Eligible workers (n=12) were defined as members of the community who contact
112 macaques or macaque bodily fluids (blood, feces, urine) as a component of their paid work at
113 least once per week. Workers were excluded if they had not worked at that site for a minimum of
114 three months. Human controls (n=6) were recruited from a convenience sampling of adults at

115 Maharakham University and were eligible if they were over 18 years of age and reported no
116 contact with macaques. Recruited participants were informed of study objectives and their rights
117 as participants and offered 100 Thai baht as compensation for their time.

118

119 **Animal participants**

120 Exposed macaques (n=8) were sampled at Kosumpee Forest Park, with an effort to
121 collect samples from macaques belonging to each of the social groups and age/sex distribution
122 representative of the overall population. These macaques are individually identifiable by facial
123 features or other unique characteristics by RCK. Control macaques (n=4) were sampled from a
124 nearby forest in Phon Ngam in the same manner as exposed macaques, and age/sex were
125 recorded.

126

127 **Environment**

128 Sites were selected based on the level of interaction between humans and macaques. The
129 study site for exposed human participants was a village of approximately 4,235 individuals
130 adjacent to Kosumpee Forest Park (KFP), Kosum Phisai District, Maha Sarakham Province in
131 northeastern Thailand (16°15'19"N 103°04'06"E) [8]. The forest park is an isolated forest patch
132 of approximately 0.2 km², bordered on the east by the Chi River and to the south by the Kosum
133 Phisai village. The park contains over 700 long-tailed macaques, divided into five social groups
134 with largely overlapping ranges [7]. Control sites were Maharakham University for humans,
135 approximately 24 km E of KFP, and a small forest tract in Phon Ngam (16°21'01"N
136 102°56'54"E) for macaques, approximately 16 km NW of KFP, where there is minimal human-
137 macaque interaction.

138

139 **Measurement**

140 Interviews, task observation, and sample collection was conducted from Sept 24 – Oct 7,
141 2017. Survey data and sample metadata were collected and stored using the REDCap electronic
142 database [14].

143 Macaque workers were surveyed regarding practices, training (e.g. macaque behavior,
144 PPE use, wound care) and their knowledge of the principle that macaques and humans can share
145 diseases. We piloted the occupational risk factor survey used in this study for eight park workers
146 in October 2016 and revised it to address limitations that emerged during administration and
147 analysis. Additions included a dietary questionnaire based on a modified food frequency
148 questionnaire (FFQ). The full occupational questionnaire is available in supplemental materials.
149 Task observations of workers were recorded using a GoPro HERO5 video recorder (GoPro, Inc.,
150 San Mateo, CA, USA) in order to assess work activities and supplement characterization of
151 exposure opportunities identified in the survey. Task observations were performed at the job title
152 level (vendor, park worker, and janitor), not for each individual, for feasibility. The scoring
153 criteria was devised by authors based on probable routes for fecal microbe transmission to
154 humans. The video recordings were reviewed by two individuals to maintain consistency and
155 discrepancies were addressed by reexamining the video segment. During review of recorded
156 tasks, an exposure category was assigned by the reviewer at 5 minute intervals, based on
157 proximity of macaques (high=direct contact or within 3m, low=beyond 3m or not visible) and
158 behavior (aerosol generation or hand-to-mouth contact).

159 Fresh fecal samples were placed immediately into OMNIgene.GUT kits (DNA Genotek,
160 Ontario, Canada) to stabilize and preserve microbial community composition and stabilize DNA

161 in the absence of a cold chain. Workers were provided with sterile collection kits and instructions
162 in Issan Thai for proper specimen collection; macaque samples were similarly collected using a
163 sterile spatula from the center of fresh excrement. Samples stored at ambient temperature as per
164 OMNIgene.GUT kit instructions until they were processed at Khon Kaen University. QIAamp
165 PowerFecal DNA Isolation kit (Qiagen, Hilden, Germany) was used to extract genomic DNA,
166 following manufacturer protocols. DNA concentration was determined using a NanoDrop2000
167 spectrophotometer (NanoDrop Technologies Inc., DE, USA) and the integrity of DNA was
168 evaluated by running 5 ul of sample on a 0.8% agarose gel under 100 V for 30 min and assessing
169 bands. Extracted DNA samples were shipped overnight on blue ice to Genewiz Laboratories in
170 Suzhou, China. DNA quality was verified by Genewiz using NanoDrop, Qubit, and agarose
171 electrophoresis. The V4 region of the bacterial 16S genes were amplified using the 515F-806R
172 primers, based on the Earth Microbiome Project protocol [15]. Amplicons were sequenced on an
173 Illumina HiSeq platform by Genewiz Laboratories. Raw FASTQ files and metadata can be
174 accessed through the Qiita database (<https://qiita.ucsd.edu/>) (accession no. 11835) and the
175 European Bioinformatics Institute, European Nucleotide Archive (accession no. ERP111664).

176

177 **Analysis**

178 DNA sequences or reads in the form of FASTQ files were analyzed with QIIME2 version
179 2017.12.0 pipeline [16]. DADA2 version 2017.12.1 was used for sequence quality control and
180 feature table construction [17]. Forward reads were truncated to 280 bp and reverse reads to 260
181 bp. Alpha diversity metrics (observed OTUs, Shannon's diversity index, Faith's Phylogenetic
182 Diversity, and Pielou's Evenness) were calculated in QIIME2. In order to attain valid
183 comparisons of abundance and diversity across samples, we normalized to the lowest sample

184 depth of 12,466 reads per sample [18]. Sequences were assigned taxonomy using the SILVA 132
185 reference database [19]. Analysis of Composition of Microbiomes (ANCOM) was performed in
186 QIIME2 between species and exposure groups with significantly different abundance values
187 identified based on the W-statistic [20]. Principal Coordinate of Analysis (PCoA) plots and taxa
188 bar plots were generated using the phyloseq package (version 1.22.3) in R [21]. PCoA plots were
189 generated to visualize clustering patterns based on weighted UniFrac distance measures, which
190 describes the degree of similarity between sample compositions by measuring the fraction of
191 unique branch length from the phylogenetic tree of sample features and weights the distance by
192 the relative abundance of that taxa within a sample. Profile clustering patterns from weighted
193 UniFrac distance measures were analysed using adonis and betadisper tests from the vegan
194 package (version 2.5.1) [22]. All tests were performed using 999 permutations based on the
195 spatial median. To further characterize microbial sharing, SourceTracker [23] was applied to
196 feature tables with macaques as source and humans as the sink under the default settings at a
197 rarefaction depth of 1000 with 100 burn-ins and 10 re-starts.

198

199 **Study team**

200 The COHERE guidelines for reporting of One Health studies were followed in the
201 preparation of this manuscript [24]. Study members represented the following areas of expertise:
202 primatology (RCK, PK), human health (PR), anthropological medicine (VR), microbial ecology
203 (EG), molecular biology (PP, RD), computational biology (PT), and environment/resource
204 management (TT).

205

206 **Ethics statement**

207 The research in this study was approved through the University of Washington
208 Institutional Review Board (IRB) for human subjects research and Institutional Animal Care and
209 Use Committee (IACUC) for animal research (#51546 and #3143-04, respectively). The study
210 also received approval through Mahasarakham University for human and animal subjects
211 research (protocol numbers 037/2016 and 0009/2016, respectively). Written informed consent
212 was obtained from all human participants and they were informed that participation was
213 voluntary, they could withdraw at any time, and questionnaire responses, individual microbiota
214 results, and task observation videos would be kept confidential and de-identified. Macaque
215 samples were obtained from fresh defecations, therefore no direct macaque handling occurred as
216 part of this study. This study was part of a larger project approved by the National Research
217 Council of Thailand (NRCT project approval to RCK - Project ID: 2016/048; “Healthy
218 Coexistence between Human and Non-human Primates: A One Health Approach”).

219

220 **Results**

221 **Questionnaire**

222 Exposed workers included government employees of Kosumpee Forest Park (n=8),
223 janitors at a nearby school (n=3), and a vendor stationed near the park entrance (n=1). All study
224 participants were born in Thailand and lived in the Maha Sarakham province for over a year.
225 Demographic factors are summarized in Table 1.

226

227 **Table 1. Human metadata.**

Factor	Exposed (n=12)	Control (n=6)
Age, years (mean \pm SD)	47.17 \pm 11.36	27.5 \pm 9.44
Sex		
Male	75% (9)	50% (3)
Female	25% (3)	50% (3)
Education, years (mean \pm SD)	9.0 \pm 3.05	16.8 \pm 5.76 ¹
Household size		
1-3	25% (3)	67% (4)
4-6	58% (7)	33% (2)
7-9	17% (2)	0
Self-rated general health		
Fair	77% (8)	0
Good	33% (4)	83% (5)
Excellent	0	17% (1)
Smoker	75% (9)	0
Health problems in past year		
Fever	92% (11)	67% (4) ¹
Respiratory problems	58% (7)	67% (4) ¹
Gastrointestinal problems	33% (4)	67% (4)
Skin problems	25% (3)	0
Infectious diseases in lifetime		
Tuberculosis	8% (1)	0
Malaria	8% (1)	0
Dengue	17% (2)	0
Other parasites, hookworm	58% (7) ¹	0 ¹
Vaginal birth method	77% (8)	50% (3)
Breast-fed as infant	92% (11)	33% (2)
BMI	25.5 \pm 5.8	23.8 \pm 3.5
Antibiotic use in past month	17% (2) ⁴	33% (2) ²

228 Demographic, early life history, dietary, and other health factors for exposed and control
229 humans, which may influence gut microbiota or may be related to macaque exposure.
230 Superscripted numbers reflect the number of missing datapoints.

231
232 Occupational factors related to microbial transmission are presented in Table 2. More
233 than half of workers regularly wash hands without soap. All participants reported handwashing
234 before and after eating (not listed in table), however task observation footage suggested this was
235 not the case for at least four participants. PPE use as reported in the survey was low, which was
236 further confirmed by the video recorded task observations. Respondents did not report receiving
237 training relevant to safe animal handling or disease prevention before working around macaques.
238 In an assessment of zoonotic disease knowledge, one-third of workers thought a diseased animal
239 could transmit that agent to a human. Only one worker thought a human could make an animal
240 sick and remarked that this would be with a high degree of contact. Workers typically only have
241 direct physical contact with carcasses, but occasionally trap live macaques to move them from
242 private properties to the forest park or when helping researchers. In one instance, a janitor had to
243 remove a macaque from a classroom using a stick and grabbing it by hand. When around animals
244 that appear sick, workers' primary form of precaution was to avoid contact.

245

246 **Table 2. Occupational risk factors.**

Factor	Response
Years at current job (mean \pm SD)	18.40 \pm 11.79
Hrs/wk around macaques or their feces (mean \pm SD)	45.08 \pm 8.694
Handwashing	
Water only	58% (7)

Soap and water	50% (6)
Alcohol-based sanitizer	8% (1)
PPE	
Disposable gloves	8% (1)
Paper or cloth dust masks	17% (2)
Rubber boots	25% (3)
Received animal/disease safety training ^a	0% (0)
Change in macaque behavior	42% (5)
Knowledge of animal to human transmission	33% (4)
Concerned about diseases from animals at work	33% (4)
Knowledge of human to animal transmission	8% (1)
Take precautions around animals that look sick	83% (10)

247 Occupational risk factors related to macaque exposure among park workers, janitors and vendors
248 based on a questionnaire.

249 ^a Training topics included animal behavior, animal capture/restraint, infectious disease
250 prevention, PPE use, or wound care.

251

252 Since starting their current job, workers noted that macaques seem “naughtier”, wait for
253 provisioning or do not look for natural food, and eat more human food (e.g. chicken, meatballs,
254 soda). All workers reported finding macaques that looked sick or had died. Carcasses were
255 typically buried or burned. One janitor remarked that, “Last month 3 monkeys die, pick them up
256 by broom into plastic bag and then threw them into the forest.”

257 Workers were asked what diseases they were primarily concerned about getting in
258 general, not necessarily from macaques. Responses included leptospirosis (n=3), cancer (n=2),
259 the common cold (n=2), cirrhosis (n=1), allergies (n=1), and an airborne infectious disease (n=1)
260 (Table 1). One worker was concerned about a “disease that come with monkey poo because I

261 have to sweep it every day.” In contrast, non-communicable diseases like high blood pressure
262 (n=3), cancer (n=1), diabetes (n=1) and hemorrhoids or constipation (n=1) were the primary
263 disease concerns among controls.

264 An abbreviated food frequency questionnaire revealed dietary differences in the type of
265 animal protein consumed. Control group members consumed more pork (p=0.04) and snail
266 (p=0.03), whereas exposed workers typically ate more frog (p=0.04). There were no significant
267 differences found in other dietary categories, including raw meat consumption. All respondents
268 reported that they pass normal formed stool (Type 3/4 on Bristol stool scale), except one, from
269 the exposed group, who reported Type 1/2.

270

271 **Task observation of workers**

272 Park workers engaged in the highest exposure activities based on recorded task
273 observations, followed by individuals working as school janitors, then vendors (Table 3). Using
274 the number of exposure events divided time observed to calculate relative risk (RR), a park
275 worker is 1.78 times more likely than a school janitor and 2.84 times more likely than a vendor
276 to work within 3m of macaques or engage in risk elevating activities (e.g. aerosol generation,
277 hand-mouth contact) during the task observation.

278

279 **Table 3. Potential exposure to macaque feces based on video-recorded task observation**

Occupation	Proximity to macaques or macaque feces				Risk elevation ^c	Time observed	RR (95% CI)
	Not visible ^a	Beyond 3m ^a	Within 3m ^b	Contact ^b			
Vendor	0	16	3	0	4	95 min	Ref
Janitor	4	7	2	0	11	65 min	1.78 (1.67-1.90)

Park worker	0	3	6	4	9	65 min	2.84 (2.67-3.02)
-------------	---	---	---	---	---	--------	------------------

280 Task observation assessment for the three worker groups (vendor, janitor, and park worker)
 281 reveals variation in proximity to macaques and behaviors that elevate risk of potential exposure.
 282 Risk of a potential exposure event during the task observation is presented relative to the lowest
 283 exposure occupation (vendor).

284 ^a No exposure

285 ^b Exposure

286 ^c Exposure, defined as engagement in activities that generate aerosols and/or eating, drinking, or
 287 smoking without prior handwashing

288

289 Macaque demographics

290 Macaques were age and sex identified according to Table 4. Among exposed macaques,
 291 members of four of the five social groups within Kosumpee Forest Park are represented in this
 292 study. Two individuals were sampled from each group (i.e. Red Dot, Stump Tail, Droop Lip and
 293 Hare Lip). At the control site, all four members were from the same social group.

294

295 **Table 4. Macaque metadata.**

Macaques	Exposed (n=8)	Control (n=4)
Age		
Juvenile	0	75% (3)
Subadult	37.5% (3)	0
Adult	62.5% (5)	25% (1)
Sex		
Male	50% (4)	50% (2) ¹
Female	50% (4)	25% (1) ¹

296 Age and sex of sampled macaques. Superscripted numbers reflect the number of missing
297 datapoints where age or sex could not be determined.

298

299 **Fecal microbiota analysis**

300 **Phylum-level abundance**

301 A total of 3,307 amplicon sequence variants (ASVs) were generated from 628,623 total
302 read counts. There was an average of 20,954 reads per sample (range: 12,466-35,318). Fig. 1
303 shows the relative abundance of bacterial phyla in each sample, after rarefaction to minimum
304 sample size. All sample profiles were dominated by Bacteroidetes, Firmicutes, and
305 Proteobacteria.

306

307 **Figure 1. Phylum-level abundance bar plot.** Relative abundance of bacterial phyla in macaque
308 control (n=4), macaque exposed (n=8), human exposed (n=12), and human control (n=6)
309 samples following rarefaction to the minimum library size. Samples are labelled with their record
310 ID, group (HC, Human Control; HE, Human Exposed; ME, Macaque Exposed; MC, Macaque
311 Control), and, if applicable, level of exposure (L, Low; M, Medium; H, High) and social group
312 (R, S, D, H).

313 Using ANCOM analysis, no taxonomic features were significantly different in abundance
314 between exposed and unexposed humans. Statistically significant differences in abundance
315 between humans and macaques are listed in Table 5.

316

317 **Table 5. ANCOM analysis of differential abundance in humans and macaques**

Feature taxonomy	W	Enriched in
------------------	---	-------------

Cyanobacteria>Melainabacteria	25	Macaques
Kiritimatiellaeota>Kiritimatiellae>WCHB1-41	45	Macaques
Bacteroidetes> Bacteroidia>Bacteroidales>Marinifilaceae	85	Macaques
Fusobacteria>Fusobacteriia>Fusobacteriales>Fusobacteriaceae	86	Humans
Proteobacteria>Gammaproteobacteria>Enterobacteriales>Enterobacteriaceae	90	Macaques
Bacteroidetes>Bacteroidia>Bacteroidales>Bacteroidaceae>Bacteroides	237	Macaques
Bacteroidetes> Bacteroidia>Bacteroidales>Rikenellaceae>Rikenellaceae RC9 gut group	249	Macaques

318 Features (ASVs) that were differentially abundant in humans or macaques are reported at the
319 most resolved taxonomic level. W-statistic indicated is the number of other items from which a
320 single item is found to be significantly different using the default $\alpha=0.05$.

321

322 **Alpha and beta diversity**

323 Fig. 2 displays the 1) total number of observed features in each sample, 2) Shannon's
324 index, which accounts for abundance and evenness of the taxa present using a natural logarithm,
325 and 3) Simpson's index, which measures the relative abundance of the different species making
326 up the sample richness. For both humans and macaques, alpha diversity was consistently lower
327 in the exposed groups (Fig 2), though this difference was only statistically significant among
328 exposed humans compared to non-exposed controls.

329

330 **Figure 2. Alpha diversity.** Alpha diversity was significantly lower among human exposed (HE)
331 relative to human controls (HC) based on the observed features ($p=0.04$), Shannon's index
332 ($p=0.02$), and Simpson's index ($p=0.04$). While there was a trend toward lower alpha diversity in
333 exposed macaques, this was not statistically significant.

334

335 PCoA plots for human and macaque gut microbial communities are shown in Fig. 3.
336 Dispersion using the betadisper test was significant for macaques ($p=0.03$), but not humans
337 ($p=0.66$). Findings among humans were unchanged after excluding people who reported taking
338 antibiotics in the past month (2 participants from control and 2 from exposed). Adonis [22] was
339 used to test for location shift of the spatial median based on exposure status, and this statistic was
340 significant for macaques and humans ($p=0.04$ and 0.02 , respectively). Dispersion and location
341 tests were also performed for unweighted UniFrac, Bray-Curtis, and Jaccard distance measures,
342 with the same conclusions.

343

344 **Figure 3. PCoA plot on weighted UniFrac distances.** 2D PCoA plot based on weighted
345 UniFrac distances demonstrate clustering and dispersion patterns for exposed human/macaque
346 and control human/macaque samples.

347

348 **SourceTracker analysis**

349 SourceTracker analysis (Fig. 4) revealed a higher percentage of microbes potentially
350 sourced by macaque microbiota in the exposed human samples (mean=3.37%) compared to the
351 controls (mean=1.84%). However, this difference was not significant by Mann-Whitney test
352 ($p=0.95$) and was driven by higher proportions for one individual. Similarly, the reverse analysis,
353 with humans as the source and macaques as the sink, showed a difference in proportions
354 attributed to human samples for exposed and control macaques (mean=4.21% and 3.98%,
355 respectively) that was not significant ($p=0.83$).

356

357 **Figure 4. SourceTracker analysis.** (A) Percent of each human sample attributed to macaque
358 exposed or macaque control source, remainder is an unknown source. (B) Percent of each
359 macaque sample attributed to human exposed or human control source, remainder is an unknown
360 source.

361

362 **Discussion**

363 Our study of gut microbiota in humans and macaques in close contact found that the
364 degree of sharing between was not statistically significant. The gut microbiota of the exposed
365 workers was significantly different from the control humans, although demographic differences
366 could explain the shift. Exposed macaques in close contact with humans, compared to a less
367 exposed population, exhibited beta-diversity dispersion effects that may reflect a dysbiotic,
368 unstable gut microbiota composition, which may be tied human contact in an urban environment.

369 SourceTracker analysis revealed no significant difference in microbial sharing between
370 humans and macaques. However, one exposed worker had a greater proportion of their
371 microbiota sourced from macaques than the other workers, suggesting that microbial sharing
372 could be occurring and could depend largely on individual factors or behavior.

373 It is also worth noting that a common diet may play a role in the detected similarities,
374 instead of or in addition to a shared environment. A study among urban Saudi and Bedouin
375 populations compared to local baboons found that the shared environment and dietary overlap
376 between Bedouins and local baboons resulted in more similar gut microbiome compositions
377 relative to urban populations [25]. However, the relative importance of environment or diet was
378 not characterized in that study. It is not clear to what degree the diet of macaques and humans
379 overlap in our study setting. In addition to bananas, local residents and tourists bring a variety of

380 foods to the macaques. During observations conducted between Sept-Dec 2016 [7] much of the
381 provisioning consisted of fruits and vegetables, but also included chips, breads, and other foods
382 not traditionally found in a macaque diet. Some macaques routinely foraged in trash or
383 consumed more atypical food than others, so individual level dietary differences should be better
384 characterized in the future.

385 Our analysis revealed that workers exhibited a different composition of fecal microbial
386 communities than controls, as evidenced by significantly different spatial medians. This finding
387 may be due to a number of other exposure factors that warrant further investigation to determine
388 the consequences of this location effect, including differences in age, SES, smoking status,
389 delivery mode, and history of infectious diseases. Early life factors are believed to play an
390 important role in shaping the adult microbiome, and there were differences in delivery method
391 and infant diet between exposed and control groups. While there is a considerable difference in
392 age, all subjects were adults, so this factor alone is not expected to greatly influence results as
393 gut microbiota, which tends to be well-established in healthy adults. Healthy adults' gut
394 microbiomes are usually less sensitive to perturbations than infants, whose microbiota are
395 developing and have not reached a stable state and elderly (>75 years old), who tend to have
396 lower total bacterial levels [26]. However, the difference age may be related to other factors
397 (e.g., infectious disease history), which could shift their microbial composition. We also
398 emphasize the need to exercise caution when excluding participants based on antibiotics use.
399 Some respondents listed paracetamol or anthelmintic medication when asked about antibiotics
400 use, or indicated they are unsure whether a drug they took was an antibiotic. Future studies
401 should ask participants about antibiotic use by referencing specific drugs based on locally used
402 names and example pills or obtain packaging from the medications used, if possible. A sub-

403 analysis excluding participants who reported antibiotic use did not alter the general conclusions
404 of this study.

405 The macaques in the park have a high level of gut microbiota dispersion relative to the
406 macaques with minimal human contact. Dispersion essentially reflects variation of microbiota
407 composition, that is the taxa present and their abundance differs from sample to sample among
408 exposed macaques, whereas the control macaques are composed of similar taxa at a similar
409 abundance, and therefore cluster tightly together, with minimal dispersion. This significant
410 dispersion pattern on exposed macaques is suggestive of the “Anna Karenina principle,” a
411 signature of dysbiosis characterized by increased variation in profiles of individuals in a disease
412 state [27]. This dysbiosis may be due to environmental stressors or diseases that perturb a stable
413 state in an unpredictable manner. We cannot definitively determine whether AKP effects are
414 occurring without longitudinal sampling, however the initial findings are suggestive of these
415 effects. In the KFP population, this dysbiosis could be a result of increased stress and
416 competition among macaques, an increased disease burden, or may be attributable to their
417 atypical diet. When asked if they noticed any changes in macaque behavior, workers reported
418 that the macaques drank more Coca-Cola and ate more chicken than they used to. While most of
419 the provisioned food appears to be fruits and vegetables, according to author RCK, who has
420 observed this population extensively, the more extreme dietary changes like foraging in trash,
421 might explain the high variation in composition among macaques at KFP. The population density
422 of the macaques in KFP also is approximately 3,670 individuals/km² which is considerably
423 higher than found in more natural settings [7]. This likely results in elevated stress and
424 aggression among macaques, which may ultimately facilitate pathogen spread. Since their
425 microbiota appear to be in a dysbiotic state relative to macaques with low levels of human

426 contact, a condition that may predispose them to gut-related diseases, they might be expected to
427 present a greater health threat to humans than wild macaques with typical gut flora [28, 29].

428 Given that AKP effects are associated with growths of opportunistic pathogens, we
429 expect to find lower evenness among the exposed macaques. While evenness based on Shannon
430 index, Pielou evenness, and Simpson evenness was marginally lower among exposed macaques,
431 this difference was not statistically significant. Our small sample size limited power and
432 increased the risk of beta error, in which a study may fail to reject the null hypothesis due to
433 insufficient power. This is a limitation that should be addressed in any future studies. By further
434 characterizing changes in susceptibility to pathogens related to gut dysbiosis, we can improve
435 understanding of the consequences of human activities such as diet supplementation or habitat
436 encroachment on wild macaque populations.

437 We also found a location shift in the spatial medians of control and exposed macaque
438 sample. It should be noted that, since the assumption of equal group variances is violated among
439 macaques, the test used is not technically valid, however, since the group larger sample size is
440 the same that exhibits greater dispersion, the test is liable to be too conservative, therefore the
441 detected shift in spatial medians likely represents a statistically significant finding [30].

442 Another limitation of the study was the choice of human controls, who differed in many
443 aspects from the exposed human population. As a result, microbiota differences between the
444 groups could be due to demographic differences rather than factors related to macaque contact or
445 occupation.

446 Further research should 1) investigate temporal trends and the stability of the dysbiosis
447 described in this study, 2) recruit well matched controls (e.g. matched age, SES, gender) in
448 Kosum Phisai to minimize the number of confounding factors in microbiota comparisons, and 3)

449 incorporate testing for GI parasitism since many members of this community take anti-
450 helminthic medication prophylactically and both factors have been shown to alter gut microbiota.
451 The cross-sectional study design employed is practical as a baseline assessment that could be
452 repeated in the future for continued, longitudinal surveillance of high risk worker populations
453 and matched controls.

454 While the threat of acquiring an infectious disease shed through macaque feces from their
455 work tasks appears low, we recommend that basic PPE be used, such as closed-toe shoes, to
456 reduce the risk of acquiring environmentally transmitted parasites shed in macaque feces, which
457 can enter through the skin. The high number of hand-to-mouth activities and work without
458 respiratory protection represents a pathway for transmission of microbes that may be present in
459 aerosolized macaque feces and offers insight into possible risk mitigating interventions. Due to
460 the small sample size, we elected not to use scores from task observation videos in microbial
461 composition analysis, instead treating all workers as exposed, however it is worth noting that the
462 degree of exposure does indeed vary within this group. One worker noted that they experienced
463 respiratory issues, which they attributed to the sweeping of macaque feces. Even if there were no
464 microbial hazards from this exposure, the dust particles or endotoxin from Gram negative
465 bacteria can cause irritation to the lungs. Therefore, future studies could assess lung function in
466 the workers to help determine whether this workplace exposure contributes to decreased lung
467 function or increased inflammation, respectively. Use of a mask during such tasks or misting of
468 the ground prior to sweeping may reduce exposure to aerosolized macaque feces and protect
469 worker health.

470 The use of a One Health approach, involving researchers from a range of disciplines,
471 allowed us to compare the microbial status for both humans and the macaques of Kosumpee

472 Forest Park. A traditional approach concerning only the human health impacts of macaque
473 exposure may not have yielded an overtly apparent risk. Our finding of dysbiosis in the gut flora
474 of macaques with close human contact may indicate increased susceptibility to pathogens. While
475 further investigation is needed to determine the implications of this finding, a microbiome-based
476 approach considering human and animal health in parallel may provide a more complete picture
477 of health in an ecosystem.

478

479 **Conclusions**

480 This study draws on a One Health approach to reduce human-animal conflict in a setting
481 modified by habitat encroachment and fragmentation. Characterizing shifts in gut microbial
482 communities allows for improved understanding of whether health changes are occurring due to
483 increased human-macaque contact in a shared environment. Based on our analysis, workers' gut
484 microbiota may under certain circumstances be influenced by their exposure to macaques, but
485 this would need to be confirmed in larger studies. The dispersion effects seen in macaques of
486 Kosumpee Forest Park suggest that their altered diet and/or interaction with an urban
487 environment may contribute to gut dysbiosis with unknown health consequences. Given the risk
488 of transmission or respiratory irritation from ingesting or inhaling fecal microbes, exposed
489 individuals and their employers should consider greater use of basic PPE and infection
490 prevention methods, particularly proper hand hygiene. Targeted health protection and disease
491 awareness promotion among forest park workers could limit opportunities for disease spillover
492 from macaque populations into the broader community.

493

494 **Acknowledgements**

495 We extend our gratitude to the study participants, Mr. Apichat Karaket, Director of
496 Kosumpee Forest Park and park staff, and the staff of the Mattayom Watklangkosum School
497 who helped facilitate this work. Many thanks to Janna Schurer and Gemina Garland-Lewis of the
498 University of Washington Center for One Health Research for their guidance and support in
499 manuscript revisions and IRB approval, respectively.

500

501 **References**

- 502 1. Malaivijitnond S, Hamada Y. Current situation and status of long-tailed macaques
503 (*Macaca fascicularis*) in Thailand. *Nat Hist J Chulalongkorn Univ.* 2008;8:185-204.
- 504 2. Kyes R, Jones-Engel L, Iskandar E, Onibala J, Lapin B, Chalise M, et al. Primate
505 conservation biology in the 21st century: Global partnerships in research, training and outreach.
506 *American Journal of Primatology.* 2006;68:144-. PubMed PMID: WOS:000239456400232.
- 507 3. Hankenson FC, Johnston NA, Weigler BJ, Di Giacomo RF. Zoonoses of occupational
508 health importance in contemporary laboratory animal research. *Comparative medicine.*
509 2003;53(6):579-601.
- 510 4. Thanchomnang T, Intapan PM, Sanpool O, Rodpai R, Tourtip S, Yahom S, et al. First
511 molecular identification and genetic diversity of *Strongyloides stercoralis* and *Strongyloides*
512 *fuelleborni* in human communities having contact with long-tailed macaques in Thailand.
513 *Parasitology Research.* 2017:1-7. doi: 10.1007/s00436-017-5469-z.

- 514 5. Wenz-Mücke A, Sithithaworn P, Petney TN, Taraschewski H. Human contact influences
515 the foraging behaviour and parasite community in long-tailed macaques. *Parasitology*.
516 2013;140(6):709-18. doi: 10.1017/S003118201200203X. PubMed PMID: 23363557.
- 517 6. Gao F, Bailes E, Robertson DL, Chen Y, Rodenburg CM, Michael SF, et al. Origin of
518 HIV-1 in the chimpanzee *Pan troglodytes troglodytes*. *Nature*. 1999;397(6718):436.
- 519 7. Kyes R, Tanee T, Thamsenanupap P, Karaket A, Kyes P. Population status of the long-
520 tailed macaques (*Macaca fascicularis*) at Kosumpee Forest Park, Maha Sarakham, Thailand.
521 *American Journal of Primatology*. 2017.
- 522 8. Village Health Volunteer Database Ministry of Public Health Thailand; 2014 [2017 May
523 19]. Available from: <http://www.thaiphc.net/>.
- 524 9. Ramirez V, Rabinowitz P, Kyes RC, Schurer JM, Grant ET, Trufan S, et al. Long-tailed
525 macaques (*Macaca fascicularis*) in urban landscapes: Gastrointestinal parasitism and barriers for
526 healthy co-existence in northeast Thailand. *American Journal of Tropical Medicine & Hygiene*.
527 2018;**Manuscript submitted for publication**.
- 528 10. Gregory AE, Lisa J-E, Michael S, Komang Gde S, Artha P, Agustin F, et al. Human
529 Exposure to Herpesvirus B–Seropositive Macaques, Bali, Indonesia. *Emerging Infectious
530 Disease journal*. 2002;8(8):789. doi: 10.3201/eid0808.010467.
- 531 11. Kim D, Hofstaedter C, Zhao C, Mattei L, Tanes C, Clarke E, et al. Optimizing methods
532 and dodging pitfalls in microbiome research. *Microbiome*. 2017;5. doi: 10.1186/s40168-017-
533 0267-5. PubMed PMID: WOS:000400644500002.
- 534 12. Song SJ, Lauber C, Costello EK, Lozupone CA, Humphrey G, Berg-Lyons D, et al.
535 Cohabiting family members share microbiota with one another and with their dogs. *elife*. 2013;2.

- 536 13. Mosites E, Sammons M, Otiang E, Eng A, Noecker C, Manor O, et al. Microbiome
537 sharing between children, livestock and household surfaces in western Kenya. Plos One.
538 2017;12(2). doi: 10.1371/journal.pone.0171017. PubMed PMID: WOS:000396161200073.
- 539 14. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic
540 data capture (REDCap)—a metadata-driven methodology and workflow process for providing
541 translational research informatics support. Journal of biomedical informatics. 2009;42(2):377-81.
- 542 15. Gilbert JA, Meyer F, Antonopoulos D, Balaji P, Brown CT, Brown CT, et al. Meeting
543 report: the terabase metagenomics workshop and the vision of an Earth microbiome project.
544 Standards in genomic sciences. 2010;3(3):243.
- 545 16. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al.
546 QIIME allows analysis of high-throughput community sequencing data. Nature methods.
547 2010;7(5):335.
- 548 17. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2:
549 High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581-3.
550 Epub 2016/05/23. doi: 10.1038/nmeth.3869. PubMed PMID: 27214047; PubMed Central
551 PMCID: PMC4927377.
- 552 18. Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, et al. Normalization and
553 microbial differential abundance strategies depend upon data characteristics. Microbiome.
554 2017;5(1):27. Epub 2017/03/03. doi: 10.1186/s40168-017-0237-y. PubMed PMID: 28253908;
555 PubMed Central PMCID: PMC5335496.
- 556 19. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
557 ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic


- 558 Acids Res. 2013;41(Database issue):D590-6. Epub 2012/11/28. doi: 10.1093/nar/gks1219.
559 PubMed PMID: 23193283; PubMed Central PMCID: PMCPMC3531112.
- 560 20. Mandal S, Van Treuren W, White RA, Eggesbø M, Knight R, Peddada SD. Analysis of
561 composition of microbiomes: a novel method for studying microbial composition. *Microb Ecol*
562 *Health Dis.* 2015;26:27663. Epub 2015/05/29. PubMed PMID: 26028277; PubMed Central
563 PMCID: PMCPMC4450248.
- 564 21. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis
565 and graphics of microbiome census data. *PLoS One.* 2013;8(4):e61217. Epub 2013/04/22. doi:
566 10.1371/journal.pone.0061217. PubMed PMID: 23630581; PubMed Central PMCID:
567 PMCPMC3632530.
- 568 22. Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ, et al. The vegan
569 package. *Community ecology package.* 2007;10:631-7.
- 570 23. Knights D, Kuczynski J, Charlson ES, Zaneveld J, Mozer MC, Collman RG, et al.
571 Bayesian community-wide culture-independent microbial source tracking. *Nat Methods.*
572 2011;8(9):761-3. Epub 2011/07/17. doi: 10.1038/nmeth.1650. PubMed PMID: 21765408;
573 PubMed Central PMCID: PMCPMC3791591.
- 574 24. Davis MF, Rankin SC, Schurer JM, Cole S, Conti L, Rabinowitz P, et al. Checklist for
575 one health epidemiological reporting of evidence (COHERE). *One Health.* 2017;4:14-21.
- 576 25. Angelakis E, Yasir M, Bachar D, Azhar EI, Lagier J-C, Bibi F, et al. Gut microbiome and
577 dietary patterns in different Saudi populations and monkeys. *Scientific Reports.* 2016;6:32191.
- 578 26. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and
579 resilience of the human gut microbiota. *Nature.* 2012;489(7415):220-30. doi:
580 10.1038/nature11550. PubMed PMID: 22972295; PubMed Central PMCID: PMCPMC3577372.




- 581 27. Zaneveld JR, McMinds R, Thurber RV. Stress and stability: applying the Anna Karenina
582 principle to animal microbiomes. *Nature microbiology*. 2017;2(9):17121.
- 583 28. Petersen C, Round JL. Defining dysbiosis and its influence on host immunity and disease.
584 *Cellular microbiology*. 2014;16(7):1024-33.
- 585 29. Van Den Elsen LW, Poyntz HC, Weyrich LS, Young W, Forbes-Blom EE. Embracing
586 the gut microbiota: the new frontier for inflammatory and infectious diseases. *Clinical &*
587 *translational immunology*. 2017;6(1).
- 588 30. Anderson MJ, Walsh DC. PERMANOVA, ANOSIM, and the Mantel test in the face of
589 heterogeneous dispersions: what null hypothesis are you testing? *Ecological monographs*.
590 2013;83(4):557-74.

591

592 **Supporting information**

593 **S1 File. Occupational exposure to macaques survey.** Questionnaire administered to workers to
594 assess demographic, life history, diet, and general health, in addition to knowledge, attitudes and
595 practices surrounding macaque exposure and zoonoses. Control surveys contained demographic,
596 life history, diet, and general health sections only.





PCoA on Weighted UniFrac Distance

