Short title: Bacterial diversity in feces of wild birds from the Pacific Northwest

RESEARCH ARTICLE

Bacterial Diversity in Feces of Wild Bald Eagles, Turkey Vultures and Common Ravens from the Pacific Northwest Coast, U.S.A.

Rocio Crespo,^{1#**¶} Scot E Dowd², Daniel E. Varland^{3¶}, Scott Ford⁴ and Thomas E. Hamer⁵

¹ Avian Health and Food Safety Laboratory, Washington Animal Disease Diagnostic Laboratory, Washington State University, Puyallup, Washington, United States of America.

² MR DNA (Molecular Research), 503 Clovis Road Shallowater, Texas, United States of America

³ Coastal Raptors, 90 Westview Drive, Hoquiam, Washington, United States of America

⁴ Avian Specialty Veterinary Services, 519 S 68th St, Milwaukee, Wisconsin, United States of America

⁵ Hamer Environmental, L.P., P.O. Box 2561, Mount Vernon, Washington, United States of America

[#]Current Address: Department of Population Health and Pathobiology, College of Veterinary Medicine, North Carolina State University, Raleigh, North Carolina, United States of America

*Corresponding author

E-mail: rcrespo@ncsu.edu (RC)

[¶]These authors contributed equally to this work.

Abstract

Birds harbor diverse microorganisms in their guts, which collectively fulfill important roles in providing their hosts with nutrition and protection from pathogens. Although numerous studies have investigated the presence of certain pathogenic bacteria in the feces of wild birds, only a few have attempted to investigate the microbiota of the gut. This study analyzed the avian bacteria present in the cloaca of avian scavengers captured on coastal beaches of Washington and Oregon between 2013 and 2015: 10 turkey vultures (*Cathartes aura*), 9 bald eagles (*Haliaeetus leucocephalus*), and 2 common ravens (*Corvus corax*). We used illumina sequencing based on the V4 region of the 16s gene was to characterize the bacterial diversity. Our investigation revealed phylum-level differences in the microbiome of turkey vultures, compared with bald eagles and common ravens. Substantial microbiome differences were found between bald eagles and ravens below the phylum level. Although little is known about the possible relations among these microorganisms, our analyses provides the first integrated look at the composition of the avian microbiota and serves as a foundation for future studies in this area.

Keywords: avian scavenger, turkey vulture, bald eagle, common raven, bacteria, coast, microbiota, microbiome, raptor.

Abbreviations: OTU = operational taxonomic unit

Introduction

Birds harbor diverse microorganisms in their gut, which collectively fulfill important roles in providing their hosts with nutrition and protection from pathogens [1]. Gut microorganisms have been isolated and characterized by a variety of methods, such as culture-based assays, culture-independent DNA sequencing, and fluorescence *in situ* hybridization targeting the 16S rRNA gene [2]. Most of these studies are limited in scope and are focused on pathogenic bacteria such as *Salmonella*, *Escherichia coli*, *Campylobacter*, and *Clostridium perfringens* [3-5].

In recent years, metagenomics using next-generation sequencing has emerged as a way to analyze complex microbial communities and their functions [6]. This approach is capable of sequencing thousands or millions of amplified DNA molecules in a single run and, for bacteria, does not require conventional cloning and amplification. Additionally, metagenomic studies offer a powerful tool for the comprehensive and unbiased assessment of microbial diversity within the complex gut ecosystem by allowing examination of organisms not easily cultured in the laboratory [6, 7].

Few studies have used genetic sequencing to explore the intestinal microbiota of wild birds [8]. Furthermore, these studies reported gene sequences to phylum or class levels only, with no attempt to understand extant microorganism populations at the species level. In this study, we analyze the avian microbiome present in the cloaca of bald eagles, turkey vultures, and common ravens collected in coastal Washington and Oregon, U.S.A. Because research has demonstrated that intestinal microorganisms can influence the lifecycle and reproduction of their hosts [9], data presented in this paper can be important for a better understanding of the health

and nutrition of these apex scavengers that comprise key components of the food web of the Pacific Northwest coast.

Materials and methods

We captured avian scavengers in coastal Washington and Oregon with a carcass-baited net launcher [10]. We captured turkey vultures and common ravens from May through August and bald eagles from February through June. Once in hand, we marked birds for identification with visual identification leg bands (eagles and ravens) or patagial tags (vultures) [11].

Aging. We assigned captured birds to two age classes, adult and immature. We classified bald eagles as adults (≥ 4 years old) with attainment of fully white head and tail plumage, yellow beak and yellow iris [12]. We classified turkey vultures as adults (≥ 2 years old) with attainment of a bright ivory bill [13]. Common ravens were ranked as adults (≥ 1 years old) by having glossy black wing and tail feathers and as immatures by having dull black or brownish wing and tail feathers [14].

Sexing. When possible, we determined the sex of individuals using measurements. We sexed bald eagles using bill depth and hallux length measurements [15] and common ravens using foot pad length and body mass [16]. Using this approach, the sex of one raven captured was equivocal. In this case, we classified the individual as female because we observed it with another raven that we assumed was male based on its demonstrably larger size [17]. We did not sex captured turkey vultures because it is not possible to differentiate gender using plumage characteristics or measurements [13].

Fecal sampling. We extracted up to 2 ml of fecal material directly from the cloaca of each bird using a disposable 3 ml pipette. In the field, we transferred fecal samples to cryovials

and then placed them in a pre-frozen Golden Hour Thermal Container (www.credothermal.com). With no more than 8 hours of elapsed time, samples were stored at -20°C for up to one month and then at -70°C until processed in 2017.

Genetic sequencing of all 21 cloaca flora samples was conducted by MR DNA, Molecular Research, LP (www.MrDNAlab.com; Shallowater, Texas). In summary, the 16S rRNA gene V4 variable region PCR primers 515/806 were used in a single-step 30 cycle PCR using the HotStarTaq Plus Master Mix Kit (Qiagen, USA) under the following conditions: 94°C for 3 minutes, followed by 30 cycles of 94°C for 30 seconds, 53°C for 40 seconds and 72°C for 1 minute, after which a final, 5-minute elongation step at 72°C was performed on a MiSeq system (Illumina Inc, San Diego, California). After amplification, PCR products were checked in 2% agarose gel to determine the success of amplification and the relative intensity of bands. Samples were pooled and purified using calibrated Ampure XP beads (Beckman Coulter, Life Sciences, Indianapolis, Indiana). This purified PCR product was then used to prepare an Illumina DNA library (<https://www.illumina.com/techniques/sequencing/ngs-library-prep.html>).

Sequence paired ends were processed using MR DNA taxonomic analysis pipeline (www.MrDNAlab.com). In summary, the process included depletion of barcodes and primers from genetic sequences, paired sequences were merged, followed by removal of sequences <150bp, and then removal of sequences with ambiguous base calls and homopolymer runs exceeding 6bp. Following these steps, operational taxonomic units (OTUs) were generated at 97% homology and chimeras were removed. Final OTUs were taxonomically classified using BLASTn (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) against a database derived from Ribosomal Database Project II (<http://rdp.cme.msu.edu>) and the National Center for Biotechnical Information (www.ncbi.nlm.nih.gov). When a given sequence matched < 77% of a reference sequence, no taxonomic identity was assigned. To designate a sequence in a phylum the

sequence was at least 77% identical to the reference sequence, for the class >80%, for the order > 85%, > 90% for the family, > 95% for the genus, and for the species >97%. We acknowledge that use of the V4 region of the 16s does not give more than a tentative evaluation of the lower phylogenetic levels (e.g. genus and species), yet we feel that some discussion of the findings in terms of genus and species provided interesting insight as part of this unique survey. Normalized to 20,000 sequences per sample, Alpha and Beta diversity analyses were completed using the Qiime2 [18] microbiome analysis pipeline (<https://qiime2.org/>).

Results

We captured and sampled 10 turkey vultures, 9 bald eagles and 2 common ravens. Except for one turkey vulture captured on an Oregon coastal beach, all individuals were captured on southwest Washington coastal beaches or adjacent to Grays Harbor (Fig 1). Eight of the turkey vultures were adults and two were immatures. Three of the bald eagles were adults and six were immature; six were male and four were female. One common raven was an adult female and the other was an immature male. We collected one fecal sample from each individual for a total of 21 samples.

Fig 1. Avian scavenger sampling sites.

Locations where fecal samples were collected from turkey vultures, bald eagles and common ravens in coastal Washington and Oregon. Numbers next to squares show the number of avian scavengers trapped at that location.

Genetic sequencing resulted in 1,905,510 sequences from the 21 samples we collected. Bacterial OTUs comprised 99.7% of the sequences found; 0.006% were Archaea. The other 0.25% constituted sparse cross-reactions of the primers with eukaryotes [19].

The total number of bacterial taxonomic OTUs detected among the 21 samples was 1065. Turkey vultures presented the largest diversity of bacterial fecal microbiome (1022 OTUs; $n = 10$), followed by common ravens (454 OTUs; $n = 2$), and bald eagles (604 OTUs; $n = 9$). In turkey vultures, 183 bacteria genera were identified, 141 in bald eagles, and 131 in common ravens (Table 1). Rarefaction curves of observed OTUs and Shannon diversity indices were generated based on a rarefaction level of 20,000 sequences per sample (Fig 2). Non-parametric pairwise comparisons made between turkey vultures and bald eagles (Fig 3) revealed significant differences in the number of observed OTUs ($p < 0.001$) as well as Shannon diversity indices between the two species ($p < 0.001$). The alpha diversity of turkey vultures was found to be significantly greater than the alpha diversity of bald eagles with regards to both measured alpha diversity metrics. Corresponding with alpha diversity analyses, based on ANOSIM pairwise comparisons (S1 Table), beta diversity between the turkey vultures and both the bald eagles as well as the common raven is shown. Beta diversity comparisons were based on the weighted unifrac distance matrix values which can be better visualized via principal coordinate analysis (PCoA) plot (Fig 4).

Table 1. Number of bacteria taxonomic groups.

Avian Species	Bacteria Taxonomy				
	Phylum	Class	Order	Family	Tentative Genus
Turkey vulture	13	23	49	94	183
Bald eagle	12	22	43	78	141
Common raven	12	19	39	76	131

Number of bacteria taxonomic groups identified in fecal samples collected from turkey vultures (n = 10), bald eagles (n = 9) and common ravens (n = 2) captured on coastal beaches in Washington and Oregon.

Fig 2. OTU Rarefaction and Shannon Diversity Index Curve.

16s rRNA sequence data was rarefied to 20,000 sequences per sample. (A) OTUs and (B) Shannon diversity indices were plotted for three avian species; Bald Eagle (n=9), Common Raven (n=2), and Turkey Vulture (n=10). OTUs were clustered at 97% identity.

Fig 3. Kruskal-Wallis Pairwise Results (Observed OTUs).

Significant differences present in alpha diversity metrics: (A) Observed OTUs and (B) Shannon diversity indices; were identified using the non-parametric Kruskal-Wallis H Test and differences between groups identified using multiple pairwise comparisons. Comparisons involving the Common Raven were not reported due to limited sample size (n=2).

Fig 4. Weighted UniFrac PCoA.

Principal coordinate plot of weighted UniFrac [18] data with colors keyed on the three avian species sampled. Bald Eagle (n=9; Red), Common Raven (n=2; blue), and Turkey Vulture (n=10; orange). The phylogenetic assemblage of the Turkey Vultures is clearly distinct from the remaining two sampled species.

The proportions of bacterial OTUs varied by phylum for each bird in the study (Fig 5). Thirteen phyla were represented among bacteria detected in the fecal samples (Fig 5, S1 Figure,

and S2 Table). Overall, more than 80% of the bacterial sequences detected belonged to two phyla, Firmicutes and Proteobacteria. In turkey vultures, the two most common phyla were Firmicutes and Fusobacteria and in bald eagles and common ravens Firmicutes and Proteobacteria were most common.

Fig 5. Mean proportions of bacterial operational taxonomic units by phylum.

Mean proportions of bacterial operational taxonomic units by phylum detected in fecal samples collected from turkey vultures (n = 10) bald eagles (n = 9), and common ravens (n = 2) captured on coastal beaches in Washington and Oregon. Proportions < 1% are not shown.

Twenty-three classes of bacteria were represented in the fecal samples. In turkey vultures, class Clostridia contained the highest proportions of OTUs by individual, while in bald eagles class Betaprotobacteria held this distinction (Fig 6). The two common raven fecal samples also included OTUs in class Betaproteobacteria, with a larger proportion of OTUs in the sample from the adult.

Fig 6. Proportions of bacterial operational taxonomic units by class.

Proportions of bacterial operational taxonomic units by class detected in fecal samples from turkey vultures (n=10), bald eagles (n = 9) and common ravens (n = 2) captured on coastal beaches in Washington and Oregon. Letters below columns are the visual identification codes on bands (eagles and ravens) or wing-tags (vultures) applied to birds at capture. Proportions < 1% are not shown.

A total of 188 tentative bacterial genera were represented in the feces of the three avian species. Of the 183 different genera detected in the 10 turkey vulture samples, 18 genera accounted for 90% of the bacterial sequences, with two genera, *Clostridium* and *Fusobacterium*, comprising 48% of the total on average (Table 2). A total of 141 genera were detected in the nine bald eagle samples and 14 of these genera represented 90% of all sequences detected. The two most common, *Burkholderia* and *Pseudomonas*, comprised 36% of the total on average (Table 2). While only two common ravens were sampled, 131 bacterial genera were detected with 9 of these comprising 90% of the OTUs (Table 2). Based on hierarchical clustering of the top 35 genera detected amongst the three species, there is a clear distinction between the fecal microbiome of turkey vultures and that of the bald eagle (Fig 4).

Table 2. Proportions of bacterial operational taxonomic units by genus.

Percentile scale

A) Turkey Vulture

Genus	Individual Patagial Tag ID Codes										Average
	CA	AR	HE	AY	BV	AV	AL	BY	AK	AX	
Enterococcus	0.16	0.03	15.08	0.05	0.06	0.55	0.16	0.1	0.03	0.03	1.63
Vagococcus	0.04	0.03	9.72	0.04	0.03	0.2	0.14	0.58	0.02	0.07	1.09
Lactobacillus	1.01	0.13	1.41	2.36	0.12	7.28	18.01	1.34	0.08	12.4	4.42
Clostridium	22.69	4.25	34.99	18.06	9.86	49.1	50.35	28.14	66.49	33.31	31.66
Peptostreptococcus	7.31	5.94	0.48	8.47	16.18	18.03	2.45	4.86	1.55	1.3	6.66
Ruminococcus	5.34	34.66	0.06	0.44	0.08	0.06	0.1	12.09	0.07	2.68	5.56
Anaerovorax	0.42	8.4	0.02	0.41	1.37	0.04	0.27	0.45	0.14	1.33	1.28
Eubacterium	3.12	3.01	0.19	3.43	0.08	5.34	0.4	3.32	0.61	0.11	1.96
Cellulosilyticum	1.99	0.05	0.14	0.53	0.04	0.14	0.33	0.06	8.46	2.94	1.47
Peptoclostridium	0.42	0.74	1.5	1.58	0.53	0.93	14.42	1.04	0.42	0.18	2.14
Holdemania	0.06	7.72	0.02	1.92	0.01	0.02	0.02	0.01	0.32	0.01	1.01
Peptoniphilus	1	8.99	0.09	1.65	9.97	0.69	0.22	18.07	0.11	0.05	4.08
Atopobium	3.74	6.71	0.07	10.67	0.04	0.12	0.32	6.94	2.84	6.46	3.79
Fusobacterium	38.98	9.96	4.04	33.3	36.35	8.08	10.68	13.8	0.19	14.2	16.96
Bradyrhizobium	0.04	0.01	0.02	0.04	1.51	0.03	0.04	0.51	0.75	6.89	0.98
Salmonella	5.03	0.64	7.46	0.56	0.1	0.14	0.19	0.09	0.3	0.59	1.51
Pseudomonas	0.13	0.1	0.32	1.43	9.87	0.08	0.09	0.09	1.19	4.67	1.8
Chlamydia	0.53	0.01	13.43	0.01	0.01	0.01	0.02	0.01	< 0.01	0.01	1.4

226 B) Bald Eagle

Genus	Individual Leg Band ID Codes									Average
	R/O	M/C	P/O	N/D	M/E	N/D	K/2	N/K	D/O	
Streptococcus	0.62	0.49	0.04	0.01	0.04	15.49	19.42	0.02	28.22	7.15
Clostridium	23.29	0.41	0.37	0.33	0.42	0.33	1.29	0.48	0.32	3.03
Bifidobacterium	0.01	0.03	0.02	0.01	0.01	0.02	5.52	19.87	0.01	2.83
Sphaerotilus	0.34	0.05	17.17	0.07	0.11	0.16	0.03	0.07	0.17	2.02
Bacteroides	2.23	4.39	0.01	0.01	0.00	9.21	4.23	0.02	0.00	2.23
Paraprevotella	>0.01	0.07	>0.01	>0.01	nd*	0.01	0.10	12.71	nd	1.43
Prevotella	2.13	nd	nd	>0.01	nd	nd	>0.01	nd	nd	1.58
Bradyrhizobium	0.82	0.13	0.02	0.14	0.01	5.99	0.02	5.14	0.01	1.37
Burkholderia	2.14	22.11	27.06	12.86	32.28	19.69	8.71	12.59	39.10	19.62
Delftia	6.62	1.06	11.01	0.35	41.61	14.51	0.31	8.70	4.07	9.80
Methylophilus	0.01	14.64	0.05	0.02	0.05	0.02	0.03	0.02	0.03	1.65
Salmonella	12.10	0.18	0.33	2.20	0.09	0.62	0.05	0.09	0.28	1.77
Pseudomonas	17.17	12.19	29.52	48.94	15.95	2.21	1.50	14.29	6.61	16.49
Stenotrophomonas	21.07	0.30	0.17	20.30	0.08	18.71	0.25	0.07	15.36	8.48

*nd = not detected

230 C) Common Raven

Genus	Individual Leg Band ID Codes		Average
	RSYLOG	ROGLSG	
Psychrobacter	0.01	12.47	6.24
Atopostipes	7.11	0.04	3.58
Catellibacter	18.12	0.05	9.09
Enterococcus	0.21	14.53	7.37
Vagococcus	0.15	11.81	5.98
Peptoclostridium	0.15	11.81	4.45
Erysipelothrix	0.02	12.64	6.33
Afipia	9.39	1.48	5.44
Daeguia	8.85	>0.01	4.43
Burkholderia	12.02	0.36	6.19
Delftia	14.27	1.23	7.75
Stenotrophomonas	11.14	5.11	8.12

232 Top 90th percentiles, expressed as a percentage of total, bacterial operational taxonomic units, by
 233 genus in fecal samples collected from A) turkey vultures (n = 10), B) bald eagles (n = 9), and C)
 234 common ravens (n = 2) captured on coastal beaches in Washington and Oregon. Letters above

columns are the visual identification codes on bands (eagles and ravens) or wing-tags (vultures) applied to birds at capture. Color denotes percentile category within each column.

As with genera, most of the tentatively classified bacteria species detected across all three avian species we studied comprised relatively small proportions of the overall fecal bacterial diversity (Table 3). Moreover, bacteria that were relatively abundant in one avian species in our study were less abundant or missing altogether in the other two. In turkey vulture samples, only *Fusobacterium mortiferum*, *Clostridium perfringens*, and *Clostridium ruminantium* represented over 10% each of the total bacteria species population. In bald eagles, *Burkholderia ubonensis* comprised more than 10% of all bacterial species and *Delftia tsuruhatensis* comprised nearly 10%. In contrast, no bacteria species represented $\geq 10\%$ of the total bacteria in common ravens.

246 **Table 3. Top 10th percentiles of bacterial operational taxonomic units tentatively classified to the closest species match in fecal**
247 **samples of turkey vultures, bald eagles and common ravens.**

Turkey Vulture (n = 10)	Number of OTUs (% of total)	Bald Eagle (n = 9)	Number of OTUs (% of total)	Common Raven (n = 2)	Number of OTUs (% of total)
<i>Fusobacterium mortiferum</i>	25,100 (17.0)	<i>Burkholderia ubonensis</i>	5,870 (17.8)	<i>Catelicoccus marimammalium</i>	5,632 (9.1)
<i>Clostridium perfringens</i>	21,764 (14.7)	<i>Delftia tsuruhatensis</i>	3,229 (9.8)	<i>Delftia tsuruhatensis</i>	4,801 (7.7)
<i>Clostridium ruminantium</i>	16,684 (11.3)	<i>Stenotrophomonas spp.</i>	2,598 (7.9)	<i>Stenotrophomonas spp.</i>	4,748 (7.7)
<i>Ruminococcus spp.</i>	7,943 (5.4)	<i>Streptococcus sanguinis</i>	2,229 (6.8)	<i>Enterococcus faecalis</i>	4,476 (7.2)
<i>Peptostreptococcus spp.</i>	7,526 (5.1)	<i>Pseudomonas spp.</i>	2,212 (6.7)	<i>Erysipelothrix tonsillarum</i>	3,924 (6.3)
<i>Peptoniphilus methionivorax</i>	6,046 (4.1)	<i>Pseudomonas veronii</i>	1,706 (5.2)	<i>Burkholderia ubonensis</i>	3,493 (5.6)
<i>Atopobium minutum</i>	5,560 (3.8)	<i>Pseudomonas plecoglossicida</i>	1,240 (3.8)	<i>Afipia spp.</i>	3,369 (5.4)
<i>Peptoclostridium clostridium sordellii</i>	2,771 (1.9)	<i>Bifidobacterium thermophilum</i>	933 (2.8)	<i>Peptoclostridium clostridium sordellii</i>	2,754 (4.4)
<i>Clostridium septicum</i>	2,374 (1.6)	<i>Clostridium spp.</i>	728 (2.2)	<i>Daeguia caeni</i>	2,745 (4.4)
<i>Lactobacillus hayakitensis</i>	2,336 (1.6)	<i>Sphaerotilus spp.</i>	665 (2.0)	<i>Psychrobacter proteolyticus</i>	2,394 (3.9)
<i>Peptostreptococcus anaerobius</i>	2,323 (1.6)	<i>Salmonella enterica</i>	584 (1.8)	<i>Atopostipes spp.</i>	2,202 (3.6)
<i>Enterococcus faecalis</i>	2,299 (1.6)	<i>Burkholderia spp.</i>	575 (1.7)	<i>Vagococcus carniphilus</i>	2,133 (3.4)
<i>Salmonella enterica</i>	2,235 (1.5)	<i>Bacteroides spp.</i>	575 (1.7)	<i>Vagococcus teuberi</i>	1,534 (2.5)
<i>Lactobacillus aviarius</i>	2,178 (1.5)	<i>Methylophilus spp.</i>	544 (1.7)	<i>Psychrobacter pulmonis</i>	1,431 (2.3)
<i>Cellulosilyticum spp.</i>	2,174 (1.5)	<i>Proteiniphilum acetatigenes</i>	472 (1.4)	<i>Ignatzschineria larvae</i>	1,226 (2.0)
<i>Chlamydia chlamydophila psittaci</i>	2,071 (1.4)	<i>Afipia spp.</i>	444 (1.3)	<i>Gardnerella vaginalis</i>	974 (1.6)
<i>Anaerovorax spp.</i>	1,901 (1.3)	<i>Bradyrhizobium spp.</i>	437 (1.3)	<i>Enhydrobacter aerosaccus</i>	965 (1.6)
<i>Holdemania spp.</i>	1,495 (1.0)	<i>Methylotenera spp.</i>	429 (1.3)	<i>Pelomonas aquatica</i>	893 (1.4)
<i>Bradyrhizobium spp.</i>	1,407 (1.0)	<i>Prevotella spp.</i>	423 (1.3)	<i>Pseudomonas spp.</i>	869 (1.4)
<i>Pseudomonas spp.</i>	1,140 (0.8)	<i>Sphingobacterium spp.</i>	360 (1.1)	Other ^a	
<i>Campylobacter spp.</i>	1,100 (0.7)	<i>Ralstonia spp.</i>	319 (1.0)		
<i>Burkholderia ubonensis</i>	1,065 (0.7)	<i>Cytophaga hutchinsonii</i>	315 (1.0)		
<i>Stenotrophomonas spp.</i>	929 (0.6)	<i>Tepidimonas spp.</i>	314 (1.0)		
<i>Vagococcus carniphilus</i>	859 (0.6)	<i>Paenibacillus barengoltzii</i>	246 (0.7)		

*Thioalkalivibrio
denitrificans*

241 (0.7)

Sum of top 10 Percentile	121,280 (82.3) ^b	27,447 (84.0) ^b	54,276 (81.5) ^b
--------------------------	-----------------------------	----------------------------	----------------------------

248 Top 10th percentiles of bacterial operational taxonomic units sequenced from fecal samples of bald eagles, turkey vultures and
 249 common ravens captured on coastal beaches in Washington and Oregon. Sequences detected in at least two avian species are
 250 identified in bold (includes two in the footnote).

251 ^aBacteria not in top 10 percentile, nevertheless comprising 1.1 - 1.3% of the total: *Acinetobacter johnsonii*, *Cosenzaea proteus*
 252 *myxofaciens*, ***Salmonella enterica***, ***Clostridium spp.***, *Staphylococcus epidermidis*. Bacteria shown in bold were also found in turkey
 253 vulture and bald eagle samples.

254 ^bPercent of total bacterial operational taxonomic units included in the top 10th percentile.

Archaea were identified in one turkey vulture sample and in two bald eagle samples. We tentatively classified these as uncultured *Candidatus nitrosocaldus*. No Archaea were identified in common ravens.

Discussion

Our analyses showed substantial microbiome diversity among turkey vultures, bald eagles, and common ravens. These differences probably represent their diverse anatomies, physiologies, and also to an extent, differences in diet. Turkey vultures, bald eagles and common ravens are among the many terrestrial vertebrates that scavenge opportunistically on the Pacific Northwest coast [20] and also inland from the coast. It should be noted that ceca, which act as sites of anaerobic activity in some species of birds, are vestigial in all three of our study species [21].

The authors are aware that bacteria species classification based upon V4 is tentative. The classification of ribosomal RNA genes has been the gold standard for molecular taxonomic research for decades, but standard primers for bacterial species identification does not exist yet [22]. On the other hand, sequences of the most commonly identified bacterial genera and species (those that are the 10 top percentile) in this study exist in the databases. These bacteria were not environmental or novel bacteria. For this reason, we discuss not only bacteria class, order, and phylum, but also genus and species.

Two classes, Fusobacteria and Clostridia, which are comprised exclusively of anaerobes, predominated the cloacal flora in turkey vultures. Roggnbuck et al. [23] also reported these two classes as most prevalent in turkey vultures. *Fusobacterium mortiferum*, which on average

comprised 17% of all bacterial species detected in Turkey Vulture feces, is a normal inhabitant of the alimentary tract in chickens where it protects against pathogenic bacteria [24].

Many clostridia species are normal gut flora and play a crucial role in gut homeostasis, however under stressful gut conditions (i.e. inflammation, parasitism, etc) these bacteria may release toxins that can cause serious diseases in humans and other animals, including birds [25, 26]. *Clostridium perfringens* and *C. ruminantium* accounted for 26% of the bacterial species we detected in turkey vulture feces in our study. *C. perfringens* is known to be a normal component of intestinal flora, but can cause necrotic enteritis in birds when intestinal dysbiosis occurs [27]. *C. ruminantium* is reported to contribute to rumen fermentation in cattle and it is also a food-spoiling bacterium, but it has not been directly linked to animal diseases [28].

Bacterial toxins can also be present in carrion. Turkey vultures have a remarkable tolerance for some of these toxins, such as botulinum [29, 30] and anthrax [23, 31]. Only 0.3% of sequences assembled from turkey vultures feces were identified as *C. botulinum* and no *Bacillus anthracis* was detected in these birds. Neither of the aforementioned species was present in bald eagles or common raven samples.

Most *Enterococcus* (phylum Firmicutes) are commensal bacteria; however there are a few species in this genus, including *E. faecalis*, that are associated with significant mortality in humans and other animals [32, 33]. *Enterococcus faecalis* comprised 7.2 % of the bacterial OTUs in the two common ravens we sampled and 1.6% in the turkey vultures. While this bacterium was detected in all nine bald eagle fecal samples, they comprised < 0.1% of the assembled sequences for that species. In Illinois, 25 raptors were tested for *Enterococcus* due to their potential to develop resistance to antibiotics; 53 of 56 cloacal samples collected contained *E. faecalis* [34]. *Streptococcus sanguinis* (phylum Firmicutes) comprised nearly 7% of all

bacteria in bald eagles but $\leq 0.3\%$ in turkey vultures and common ravens, Although these bacteria are considered normal inhabitants of mammals and humans [35], there is no information about the commonality of these bacteria or their significance in the digestive tracts of avian species. *Erysipelothrix tonsillarum* comprised 6% of the OTUs in the two common raven fecal samples and $< 0.1\%$ in turkey vultures and bald eagles. These non-pathogenic bacteria are commonly isolated from healthy swine and poultry as well as poultry litter. *E. tonsillarum* is closely related to the pathogenic species *E. rhusiopathae*, causative agent of erysipeloid in mammals and erysipelas in poultry [36, 37].

Almost 70% of fecal bacteria in the bald eagles we sampled ($n = 9$) and 55% in the two common ravens belonged to the phylum Proteobacteria. In contrast, Proteobacteria accounted for less than 10% of the turkey vulture microbiome ($n = 10$). In bald eagles, the most common species belonging to this phylum were tentatively classified as *Burkholderia ubonensis* and *Delftia tsuruhatensis*, comprising 18% and 10% of all fecal bacteria detected respectively. Both are common environmental bacteria that have been occasionally linked to infections in humans [38, 39]. *B. ubonensis* was also detected in the two common ravens sampled, ranking seventh most common and averaging 5.6%. *Catellibacterium marimammali* was the most common bacteria in the common ravens sampled, averaging 9% of the total. *C. marimammali* was the most common species in the feces of gulls in Wisconsin, U.S.A. [40].

Other notable bacteria detected from the Proteobacteria phylum included bacteria in *Escherichia*, *Campylobacter*, *Pseudomonas*, and *Salmonella*. *E. coli* inhabits the large intestine and distal ileum of most vertebrate species, including birds; the organism is shed in the feces. Generally-speaking, the vast majority of the strains are components of normal intestinal flora and are non-pathogenic. Some strains are of low virulence and may cause opportunistic infections. A

few are highly pathogenic and may cause septicemia. *E. coli* sequences were detected in the cloacal samples from eight of the ten turkey vultures, eight of the nine bald eagles, and both ravens. However, *E. coli* sequences were sparse and, on average, accounted for less than 0.01% of all the bacterial OTUs per bird. No further analysis was done on these *E. coli* sequences to determine if they had any pathogenic genes or not. *Campylobacter* sp. were detected in all the fecal samples from turkey vultures and accounted for 0.7% of the bacterial OTUs (Table 3). In bald eagles and common ravens, *Campylobacter* sp. were also detected in all samples, but the OTU frequency was less than 0.01%. Members of *Pseudomonas* are ubiquitous and can be opportunistic pathogens [41]. Bacteria from the *Pseudomonas* genus were detected in all fecal samples. They constituted 17% of the fecal flora in bald eagles, but < 2% of all bacterial OTUs in turkey vultures and common ravens.

Avian predators may become infected with *Salmonella* when they consume infected animals [42]. *S. enterica* colonize the intestinal tract and have been isolated from wild birds with or without clinical signs of disease [43]. In our study, *S. enterica* comprised 1-2% of bacterial species detected in all three avian scavengers. In other research, *S. enterica* sampled from the feces of scavenging birds ranged from 1% to 20% by isolation [44-46]. Differences in prevalence between isolation and detection by meta-analysis may be due to the stability of the DNA molecule that can be amplified even when the target organism is no longer viable. *S. enterica* sequence was performed only to the species level, and subspecies or serotype were not determined.

Archaea are single-celled prokaryotes that are difficult to culture [47]. They have been found in the digestive tracts of some animals where they produce methane [48]; the role these organisms play in the gut of birds is still unknown [1]. *Candidatus nitrosocaldus* is ubiquitous in

water environments being present in plankton, sediments, and surrounding soils of lakes and oceans [49]. It has a particular affinity for hot spring environments. It is also common in waste water treatment plants and fertilized soils. Its presence in the feces of two bald eagles and one turkey vulture in our study may be a reflection of their contact with marine carrion.

In summary, this is the first study utilizing metagenomics to investigate the fecal microbiome to the species level. This is also the first meta-analysis performed in bald eagles and common ravens. We demonstrated that the fecal flora varies among the three scavenger species even though they live in the same region and consume some of the same foods (D. Varland, unpub. data). We discovered that over 50% of the microbiota in the cloaca belongs to a single phylum in all three species, Proteobacteria for bald eagles and common ravens and Firmicutes for turkey vultures. As expected numerous species of microorganisms were tentatively identified; however, only about 10% of these tentative species contained > 1% of the OTUs. Finally, this method for identification and characterization of microorganism populations can be used in epidemiological, pathogen detection, and microbial diversity studies.

Acknowledgements

We thank the Washington State Parks and Recreation Commission and the Oregon Parks and Recreation Department for allowing vehicle access to coastal beaches, which was essential to our work. We also greatly appreciate the field assistance of many, including Nathalie Denis, Glenn Marquardt, Larry Warwick, Sandra Miller, Pam McCauley, Suzy Whittey, Tom Rowley and Dale Larson. D. Varland color marked birds in the study under federal banding permit 21417 and under state Scientific Collection Permits 13-014, 14-084 and 15-093 in Washington and under Scientific Taking Permit 049-13 in Oregon.

References

1. Waite DW, Taylor MW. Exploring the avian gut microbiota: current trends and future directions. *Front Microbiol.* 2015;6:673. Epub 3 July 2015.
2. Gomez-Alvarez V, Revetta RP, Domingo JWS. Metagenomic analyses of drinking water receiving different disinfection treatments. *Appl Environ Microb.* 2012;78(17):6095-102.
3. Tizard IR. Salmonellosis in wild birds. *Seminars in Avian and Exotic Pet Medicine.* 2004;13(2):50-66.
4. Tarello W. Etiologic agents and diseases found associated with clinical aspergillosis in falcons. *Intern J Micro.* 2011;2011:176963. doi: 10.1155/2011/176963.
5. Hernández J, Lindberg P, Waldenström J, Drobni M, Olsen B. A novel *Salmonella* serovar isolated from Peregrine Falcon (*Falco peregrinus*) nestlings in Sweden: *Salmonella enterica enterica* serovar Pajala (*Salmonella* Pajala). *Infection Ecology and Epidemiology.* 2012;2:10.3402/iee.v2i0.7373. Epub 2012 Aug 3. doi: 10.3402/iee.v2i0.7373. PubMed Central PMCID: PMC3432136.
6. Maeda I, Siddiki MSR, Nozawa-Takeda T, Tsukahara N, Tani Y, Naito T, et al. Population abundance of potentially pathogenic organisms in intestinal microbiome of jungle crow (*Corvus macrorhynchos*) shown with 16s rRNA gene-based microbial community analysis. *BioMed Res Int.* 2013;2013:438956, 5 pages.
7. Kovatcheva-Datchary P, Zoetendal EG, Venema K, Vos WMd, Smidt H. Tools for the tract: understanding the functionality of the gastrointestinal tract. *Ther Adv Gastroenter.* 2009;2(4 Supplement):9-22. doi: 10.1177/1756283X09337646.

- 391 8. Colston TJ, Jackson CR. Microbiome evolution along divergent branches of the
392 vertebrate tree of life: what is known and unknown. *Mol Ecol.* 2016;25(16):3776-800. doi:
393 <https://doi.org/10.1111/mec.13730>.
- 394 9. Waite DW, Deines P, Taylor MW. Gut Microbiome of the Critically Endangered New
395 Zealand Parrot, the Kakapo (*Strigops habroptilus*). *PLoS One.* 2012;7(4):e35803. doi:
396 10.1371/journal.pone.0035803.
- 397 10. Varland DE. A capture to remember: Monitoring the health of avian scavengers on the
398 pacific coast. *J Av Med Surg.* 2015;29(3):250-6.
- 399 11. Varland DE, Smallwood JA, Young LS, Kochert MN. Marking techniques. In: Bird DM,
400 Bildstein KL, editors. *Raptor Research and management Techniques*. Blaine, WA U.S.A. and
401 Surrey, B.C. Canada: Hancock House Publishers; 2007. p. 221-36.
- 402 12. McCollough MA. Molting sequence and aging of bald eagles. *Wilson Bull.*
403 1989;101(1):1-10.
- 404 13. Pyle P, editor. *Identification guide to North American birds. Part II. Point Reyes Station,*
405 *California: Slate Creek Press; 2008.*
- 406 14. Marzluff JM. Common Raven. In: del Hoyo J, Elliott A, Christie DA, editors. *Handbook*
407 *of the Birds of the World. 14, Bush-Shrikes to Old World Sparrows.* Barcelona, Spain: Lynx;
408 2009.
- 409 15. Bortolotti GR. Sexual size dimorphism and age-related size variation in Bald Eagles *J*
410 *Wild Managem.* 1984;48(1):72-81.
- 411 16. Bedrosian B, Loutsch J, Craighead D. Using morphometrics to determine the sex of
412 common ravens. *Northwestern Naturalist.* 2008;89(Spring):46-52.

- 413 17. Boarman WI, Heinrich B. Common raven. In: Poole AF, Gill FB, editors. The Birds of
414 North America. version 2.0 ed. Ithaca, NY, USA: Cornell lab of Ornithology; 1999.
- 415 18. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet C, Al-Ghalith GA, et al. QIIME
416 2: Reproducible, interactive, scalable, and extensible microbiome data science. PeerJ Preprints.
417 2018;6(1):e27295v1.
- 418 19. Yadav S, Kumar A, Gupta M, Maitra SS. Cross-reactivity of prokaryotic 16S rDNA-
419 specific primers to eukaryotic DNA: Mistaken microbial community profiling in environmental
420 samples. Curr Microbiol. 2018;75(8):1038-45. Epub 2018 Apr 2. doi: 10.1007/s00284-018-1482-
421 4.
- 422 20. Varland DE, Buchanan JB, Fleming TL, Kenney MK, Vanier C. Scavenging as a food-
423 acquisition strategy by peregrine falcons. J Raptor Res. 2018;52(3):291-308.
- 424 21. Clench M, Mathias J. The avian cecum: A review. Wilson Bull. 1995;107(1):93-121.
- 425 22. Mizrahi-Man O, Davenport ER, Gilad Y. Taxonomic classification of bacterial 16s rna
426 genes using short sequencing reads: Evaluation of effective study designs. PLoS One.
427 2013;8(1):e53608.
- 428 23. Roggenbuck M, Schnell IB, Blom N, Bælum J, Bertelsen MF, Sicheritz-Pontén T, et al.
429 The microbiome of New World vultures. Nature Communications. 2014;5:5498. doi:
430 10.1038/ncomms6498.
- 431 24. Portrait V, Cottenceau G, Pons AM. A *Fusobacterium mortiferum* strain produces a
432 bacteriocin-like substance(s) inhibiting *Salmonella* Enteritidis. Lett Appl Microbiol.
433 2000;31(2):115-7.
- 434 25. Popoff MR, Bouvet P. Clostridial Toxins. Future Microbiol. 2009;4(8):1021-64. doi:
435 <https://doi.org/10.2217/fmb.09.72>.

26. Lopetuso LR, Scaldaferri F, Petito V, Gasbarrini A. Commensal Clostridia: leading players in the maintenance of gut homeostasis. Gut Pathog. 2013;5:23. doi: <https://doi.org/10.1186/1757-4749-5-23>.
27. Cooper KK, Songer JG, Uzal FA. Diagnosing clostridial enteric disease in poultry. JVDI. 2013;25(3):314–27.
28. Reida R, Burgessa CM, McCabea E, Fanningb S, Whyteb P, Kerry J, et al. Real-time PCR methods for the detection of blown pack spoilage causing Clostridium species; *C. estertheticum*, *C. gasigenes* and *C. ruminantium*. Meat Sci. 2017;133:56-60. doi: <http://dx.doi.org/10.1016/j.meatsci.2017.06.001>.
29. Kalmbach ER. American vultures and the toxin of clostridium botulinum. JAVMA. 1939;94:187-91.
30. Ohishi I, Sakaguchi G, Riemann H, Behymer D, Hurvell B. Antibodies to Clostridium botulinum toxins in free-living birds and mammals. J Wild Dis. 1979 15(1):3-9.
31. Darling ST, Bates LB. Anthrax of animals in Panama, with a note on its probable mode of transmission by buzzards. Am Vet Rev. 1912;42:70-5.
32. Ellerbroek LK, Mac N, Peters J, Hultquist L. Hazard potential from antibiotic-resistant commensals like Enterococci. J Med Vet A. 2004;51:393-9.
33. Silva N, Igrejas G, Gonçalves A, Poeta P. Commensal gut bacteria: distribution of *Enterococcus* species and prevalence of *Escherichia coli* phylogenetic groups in animals and humans in Portugal. Annals Microbiol. 2012;62(2):449-59.
34. Marrow J, Whittington JK, Mitchell M, Hoyer LL, Maddox C. Prevalence and antibiotic-resistance characteristics of *Enterococcus* spp. isolated from free-living and captive raptors in central Illinois. J Wild Dis. 2009;45(2):302-13.

- 459 35. Caufield PW, Dasanayake AP, Yihong Li YP, Hsu J, Hardin JM. Natural history of
460 *Streptococcus sanguinis* in the oral cavity of infants: Evidence for a discrete window of
461 infectivity. Infect Immun. 2000;68(7):4018-23. doi: 10.1128/IAI.68.7.4018-4023.2000.
- 462 36. Takahashi T, Fujisawa T, Benn Y, Tamura Y, Sawada T, Suzuk S, et al. *Erysipelothrix*
463 *tonsillarum* sp. nov. isolated from tonsils of apparently healthy pigs. Int J Syst Bacteriol.
464 1987;37(2):166-8.
- 465 37. Eriksson H, Bagge E, Båverud V, Fellström C, Jansson DS. *Erysipelothrix rhusiopathiae*
466 contamination in the poultry house environment during erysipelas outbreaks in organic laying
467 hen flocks. Avian Pathol. 2014;43(3):231-7. doi:
468 <http://dx.doi.org/10.1080/03079457.2014.907485>.
- 469 38. Price EP, Sarovich DS, Webb JR, Hall CM, Jaramillo SA, Sahl JW, et al.
470 Phylogeographic, genomic, and meropenem susceptibility analysis of *Burkholderia ubonensis*.
471 PLoS Neglected Tropical Diseases. 2017;11(9):e0005928. doi:
472 <https://doi.org/10.1371/journal.pntd.0005928>.
- 473 39. Ranc A, Dubourg G, Fournier PE, Raoult D, Fenollar F. *Delftia tsuruhatensis*, an
474 Emergent Opportunistic Healthcare-Associated Pathogen. Emerg Infect Dis. 2018;24(3):594-6.
475 doi: <https://doi.org/10.3201/eid2403.160939>.
- 476 40. Koskey AM, Fisher JC, Traudt MF, Newton RJ, McLellan SL. Analysis of gull fecal
477 microbial community reveals dominance of *Catelliboccus marimammali* in relation to
478 culturable enterococci. Appl Environ Microb. 2014;80:757-65. Epub 15 November 2013. doi:
479 10.1128/AEM.02414-13.

41. Abdul-Aziz T, Barnes HJ. Miscellaneous and Sporadic Bacteria Infections. In: Swayne DE, Glisson JR, McDougald LR, Nolan LK, Suarez DL, Nair V, editors. Diseases of Poultry. 13th ed. Ames, IA: Wiley-Blackwell; 2013. p. 1017-27.
42. Benskin CM, Wilson K, Jones K, Hartley IR. Bacterial pathogens in wild birds: a review of the frequency and effects of infection. Biol Rev Camb Philos. 2009;84:349-73.
43. Henderson SC, Bounous DI, Lee MD. Early events in the pathogenesis of avian salmonellosis. Infect Immun. 1999;67:3580-6.
44. Janecko N, Čížek A, Halová D, Karpíšková R, Myšková P, Literák I. Prevalence, characterization and antibiotic resistance of *Salmonella* isolates in large corvid species of Europe and North America between 2010 and 2013. Zoonoses Public Hlth. 2015; 62(4):292-300. Epub 2014 Aug 7. doi: 10.1111/zph.12149.
45. Reche MP, Jiménez PA, Alvarez F, Ríos JEGdl, Rojas AM, Pedro Pd. Incidence of *Salmonellae* in captive and wild free-living raptorial birds in central Spain. J Med Vet A. 2003;50(1):42-4.
46. Sulzner K, Kelly T, Smith W, Johnson CK. Enteric pathogens and antimicrobial resistance in turkey vultures (*Cathartes aura*) feeding at the wildlife-livestock interface. J Zoo Wild Med. 2014 45(4):931-4. doi: <http://dx.doi.org/10.1638/2012-0217.1>.
47. Childers R, Reno P, Olson R. Prevalence and geographic range of *Nadelspora canceri* (Microspora) in Dungeness crab Cancer magister. Dis Aquat Organ. 1996;24:135-42.
48. Danielsson R, Dicksved J, Sun L, Gonda H, Müller B, Schnürer A, et al. Methane production in dairy cows correlates with rumen methanogenic and bacterial community structure. Front Microbiol. 2017;8:226. Epub 17 February 2017. doi: 10.3389/fmicb.2017.00226.

502 49. Abby SS, Melcher M, Kerou M, Krupovic M, Stieglmeier M, Rossel C, et al. *Candidatus*
503 *Nitrosocaldus cavascurens*, an ammonia oxidizing, extremely thermophilic archaeon with a
504 highly mobile genome. *Front Microbiol.* 2018;9(January):Article 28. doi:
505 <https://doi.org/10.3389/fmicb.2018.00028>.
506

Supporting information

S1 Figure. Proportions of bacterial operational taxonomic units by phylum. Proportions of bacterial operational taxonomic units by phylum detected in fecal samples from turkey vultures (n = 10), bald eagles (n = 9), and common ravens (n = 2) captured in coastal Washington and Oregon. Letters below columns are the visual identification codes on bands (eagles and ravens) or wing-tags (vultures) applied to birds at capture. Proportions < 1% are not shown.

S2 Figure. Hierarchal Clustering of Taxonomic Data. Evaluation of the taxonomic classification data using a dual hierarchal dendrogram. Each sample is clustered on the X-axis labeled based on the sampled species. Samples with more similar microbial populations are mathematically clustered closer together. The heatmap represents the relative percentages of each genus. The predominant genera are represented along the right Y-axis. The legend for the heatmap is provided in the upper left corner. The microbial population of Turkey Vultures (n=10) is distinct from the population identified in the Bald Eagle (n=9).

S1 Table. Pairwise ANOSIM Results (Weighted UniFrac)

Significant differences present in the weighted UniFrac distance matrix values were identified using the Analysis of Similarities (ANOSIM) Test and differences between groups identified using multiple pairwise comparisons over 999 permutations.

Group 1	Group 2	Sample size	R ^a	p-value	q-value
BALDEAGLE	COMMONRAVEN	11	0.204204	0.239	0.239
BALDEAGLE	TURKEYVULTURE	19	0.928121	0.001	0.003
COMMONRAVEN	TURKEYVULTURE	12	0.81087	0.015	0.0225

^aR scales from +1 to -1. A value of +1 indicating the most similar samples are in the same group. A value equal to 0 indicating there is no relationship observed between similar and dissimilar samples.

S2 Table. Mean number (and percent) of bacterial operational taxonomic units by phylum.

Mean number (and percent) of bacterial operational taxonomic units by phylum detected in fecal samples from turkey vultures (n = 10), bald eagles (n = 9), and common ravens (n = 2) captured on coastal beaches in Washington and Oregon. For each species, phyla with the largest number of operational taxonomic units are identified in bold.

Phylum	Turkey Vulture		Bald Eagle		Common Raven	
	Mean	(Percent)	Mean	(Percent)	Mean	(Percent)
Acidobacteria	5.7	(< 0.1)	0.2	(< 0.1)	ND ^a	ND
Actinobacteria	5441	(4.3)	2,033.8	(5.4)	951	(2.1)
Bacteroidetes	261.4	(0.2)	2,912.2	(8.3)	336.5	(0.4)
Chlamydiae	2,000	(1.4)	6.9	(< 0.1)	5.5	(< 0.1)
Cyanobacteria	7.3	(< 0.1)	195.3	(0.6)	150.5	(0.2)
Deferribacteres	19.2	(< 0.1)	0.1	(< 0.1)	ND	ND
Firmicutes	10,6044.8	(67.1)	4,937.4	(13.6)	27,467	(40.9)
Fusobacteria	22,873.3	(17.3)	81.2	(0.2)	61.5	(0.1)
Gemmatimonadetes	4.1	(< 0.1)	111.4	(0.3)	1	(< 0.1)
Ignavibacteriae	7.9	(< 0.1)	208.3	(0.6)	1	(< 0.1)
Proteobacteria	11,345.5	(9.7)	22,445.3	(70.9)	32,631	(55.5)
Spirochaetes	2.4	(< 0.1)	0.3	(< 0.1)	372.5	(0.9)
Tenericutes	2.1	(< 0.1)	ND	ND	ND	ND

^a ND = Not detected.

Figure 1

Figure 2a

Figure 2b

Groups				
Group 1	Group 2	H	p-value	q-value
Bald eagle (n=9)	Turkey vulture (n=10)	13.5	0.000238563	0.00071569

Figure 3a

Groups				
Group 1	Group 2	H	p-value	q-value
Bald eagle (n=9)	Turkey vulture (n=10)	11.76	0.000605172	0.001815517

Figure 3b

Figure 4

Figure 5

Figure 6