1 Neofunctionalisation of basic helix loop helix proteins occurred when plants colonised the 2 land

3
4 Clémence Bonnot¹², Alexander J. Hetherington¹, Clément Champion¹, Holger Breuninger¹³,
5 Steven Kelly¹, Liam Dolan^{1#}

6
7 ¹ Department of Plant Sciences, University of Oxford, South Parks Road, Oxford, OX1 3RB,
8 UK

9 ² Present address: Labex ARBRE, UMR 1136 INRA-Université de Lorraine (IAM), INRA-Grand
10 Est-Nancy, Champenoux, France

11 ³ Present address: ZMBP, Entwicklungsgenetik, 72076 Tübingen, Germany

12 **# Author for correspondence:** Tel: +44 (0) 1865 275147

13 Email: Liam.Dolan@plants.ox.ac.uk

14

15 **Abstract word count: 200**

16 **Total word count: 6277**

17 **Detail: Introduction (1045), Materials and Methods (2211), Results (1982), Discussion**
18 **(967), Acknowledgements (72)**

19 **Figures: 8 (Coloured figures: Figures 1, 2, 3, 5, 6 and 7)**

20 **Tables: 2**

21 **Supporting information: 10 (4 Sup Data and 6 Sup Figures)**

22

23 **ABSTRACT**

24 *ROOT HAIR DEFECTIVE SIX-LIKE (RSL)* genes control the development of structures –
25 rhizoids, root hairs, gemmae, mucilage papillae – that develop from single cells at the
26 surface of diverse groups of land plants. RSL proteins constitute a subclass (VIIIc) of the
27 basic helix loop helix (bHLH) class VIII transcription factor family. We set out to determine if
28 the function of RSL genes in the control of cell differentiation in land plants was inherited
29 from streptophyte algal ancestor. The Charophyceae are a monophyletic class of
30 streptophyte algae with tissue-like structures and rhizoids. We identified the single class VIII
31 bHLH gene from the charophyceae alga *Chara braunii* (*CbbHLHVIII*). Phylogenetic analysis
32 suggests that this protein is sister to the RSL (bHLH subclass VIIIc) proteins and together

they constitute a monophyletic group. Expression of *CbbHLHVIII* does not compensate for loss of the RSL function in either *Marchantia polymorpha* or *Arabidopsis thaliana*. Furthermore, *CbbHLHVIII* is expressed at sites of morphogenesis in *C. braunii* – the apices, nodes and gametangia – but not in rhizoids. This indicates that *C. braunii* class VIII protein is functionally different from land plant RSL proteins; they control rhizoid development in land plants but not in the charophycean algae. These data are consistent with the hypothesis that RSL proteins and their function in the differentiation of cells at the plant surface evolved in the lineage leading to land plants after the divergence of the land plants and *C. braunii* from their last common ancestor. This may have occurred by neofunctionalisation at or before the colonisation of the land by streptophytes.

Key words: basic helix loop helix (bHLH), *Chara braunii*, *Coleochate nitellarum*, Land plants, ROOT HAIR DEFECTIVE SIX-LIKE (RSL), Filamentous rooting cells, Streptophyte algae, Neofunctionalisation

INTRODUCTION

The colonisation of the land by streptophytes and the subsequent radiation of morphological diversity among the land plants was a major transition in Earth history (Kenrick & Crane, 1997; Pires & Dolan, 2012). The simple body plans of extant streptophyte algae has led to the hypothesis that the body plan of the common ancestor of the land plants was simple. It is hypothesized that plants then underwent morphological diversification during and after the colonisation of the dry continental surfaces (Graham *et al.*, 2000; Delaux *et al.*, 2012; Harrison, 2017). These adaptations to life on the land contributed to the establishment of the first complex terrestrial ecosystems by 407 Ma and led to the terrestrial ecosystems that exist today (Gibling & Davies, 2012; Lenton *et al.*, 2012).

The sequencing of land plant (Rensing *et al.*, 2008; Banks *et al.*, 2011; Bowman *et al.*, 2017) and streptophyte algae genomes (Hori *et al.*, 2014; Ju *et al.*, 2015; Nishiyama *et al.*, 2018) has led to the hypothesis that the evolution of the morphological diversity resulted from an increase in the number of regulatory genes in gene families (Floyd & Bowman, 2007; Lang *et al.*, 2010; Pires & Dolan, 2010a,b; Bowman *et al.*, 2017; Lehti-Shiu *et al.*, 2017). For example, transcription factor family number is higher in land plants than in

streptophyte algae (Tanabe *et al.*, 2005; Navaud *et al.*, 2007; Chanderbali *et al.*, 2015; Catarino *et al.*, 2016; Nishiyama *et al.*, 2018). It is also likely that the function of regulatory genes will have changed during the course of the transition to land. Such changes could result from genes assuming new functions (neofunctionalisation) or dividing their functions among their descendants (sub-functionalisation) (Prince & Pickett, 2002; Rensing, 2014).

The development of a diversity of morphological structures that form from single cells in the surface cell layer of organs is regulated by RSL class 1 genes (also known as subclass VIIIc1 basic helix loop helix) in a diversity of land plants (Honkanen & Dolan, 2016). For example, RSL class 1 genes positively regulate the development of rhizoids, mucilage papillae and gemmae (asexual propagules) in the liverwort *Marchantia polymorpha* (Proust *et al.*, 2016). Orthologs positively regulate the development of rhizoids and mucilage papillae in the moss *Physcomitrella patens* (Menand *et al.*, 2007; Jang *et al.*, 2011; Proust *et al.*, 2016). Class 1 RSL genes positively regulate the development of root hairs in diverse groups of angiosperms including *Arabidopsis thaliana* (Masucci & Schiefelbein, 1994; Menand *et al.*, 2007) and the grasses *Oryza sativa* and *Brachypodium distachyon* (Kim & Dolan, 2016; Kim *et al.*, 2017). Expression of RSL class 1 genes from one taxa of land plant can compensate for the loss of function in another taxa (Menand *et al.*, 2007; Kim & Dolan, 2016; Proust *et al.*, 2016; Kim *et al.*, 2017). For example, expression of *M. polymorpha* MpRSL1 gene using the cauliflower mosaic virus 35S promoter (35S) in the root hairless *Atrhd6 Atrsl1* mutants of *A. thaliana* restores root hair development (Proust *et al.*, 2016). This demonstrates that not only do RSL class 1 genes control the development of these structures in diverse land plants, but also that the function of the proteins has been conserved during the course of land plant evolution.

RSL class 1 genes (subclass VIIIc1) are members of class VIII basic helix loop helix (bHLH) transcription factors (Menand *et al.*, 2007; Pires & Dolan, 2010a; Proust *et al.*, 2016). Class VIII bHLH proteins includes two other subclasses, subclass VIIla and subclass VIIlb (Heim *et al.*, 2003; Pires & Dolan, 2010a; Catarino *et al.*, 2016). The functions of subclass VIIla proteins are unknown (Pires & Dolan, 2010a). Subclass VIIlb includes the HECATE-related transcription factors that control fruit development in *A. thaliana* (Liljegren *et al.*, 2004; Gremski *et al.*, 2007; Kay *et al.*, 2013). Together the subclasses VIIla, VIIlb and VIIIc

constitute a monophyletic group of proteins that is conserved among land plants. While the function of RSL class 1 genes (subclass VIIIc) have been shown to be conserved among diverse groups of land plants (Menand *et al.*, 2007; Jang *et al.*, 2011; Kim & Dolan, 2016, p. 20; Proust *et al.*, 2016; Kim *et al.*, 2017), the functions of subclass VIIIb have only been defined in *A. thaliana* to date (Liljegren *et al.*, 2004; Gremski *et al.*, 2007; Kay *et al.*, 2013).

Since RSL class 1 genes control the development of rhizoids and root hairs in the mosses, liverworts and angiosperms, it is hypothesized that RSL class 1 genes control the development of the rhizoidal rooting structure in the common ancestor of the land plants (Menand *et al.*, 2007; Proust *et al.*, 2016). This was one of a suite of new functions that evolved early in land plant evolution, which was a key adaptation to life on the continental surfaces of the planet (Kenrick & Crane, 1997; Delaux *et al.*, 2012). To investigate the origin of this regulatory mechanism we searched the genome (Nishiyama *et al.*, 2018) and transcriptomes of the streptophyte alga *Chara braunii* for genes encoding class VIII bHLH proteins. *C. braunii* is a member of the Charophyceae, the only class of streptophyte algae in which tissue-like structures and rhizoids develop (Smith & Allen, 1955; Pickett-Heaps, 1975; Graham & Wilcox, 2000; Nishiyama *et al.*, 2018). We identified a single class VIII gene (*CbbHLHVIII*) that is sister to the land plant RSL (subclass VIIIc) transcription factors. The expression of *CbbHLHVIII* in *M. polymorpha* or *A. thaliana* mutants does not compensate for the loss of endogenous RSL function, indicating that *C. braunii* class VIII protein is functionally different from the land plant RSL proteins. Furthermore, *CbbHLHVIII* is expressed in regions of organogenesis - apices and young nodes of the thallus, and gametangia - but not in rhizoids or in rhizoids morphogenesis zones. This is consistent with the hypothesis that *CbbHLHVIII* regulates development at *C. braunii* morphogenetic centres, but does not control rhizoid differentiation. We conclude that class VIII proteins evolved the ability to control the differentiation of cells such as rhizoids in the land plant lineage after the divergence of *C. braunii* and land plants from their last common ancestor. This is consistent with a model in which neofunctionalisation of class VIII proteins occurred during the increase in morphological diversity that occurred during the transition to land.

MATERIALS AND METHODS

Sequence identification

To identify class VIII sequences in *Chara braunii* and *Coleochaete nitellarum*, we used the land plants bHLH class VIII protein sequences of *Arabidopsis thaliana*, *Selaginella moelendorffii*, *Physcomitrella patens* and *Marchantia polymorpha* published previously (Catarino *et al.*, 2016) as queries in TBlastN searches (Altschul *et al.*, 1990) of *Coleochaete nitellarum* transcriptome (Bonnot *et al.*, 2017) and *C. braunii* genome (Nishiyama *et al.*, 2018) and transcriptome (see “Construction of the *C. braunii* transcriptome” hereafter). No E-value thresholds were used. All hits were manually investigated. Transcripts and genomic sequences were aligned. Amino acid sequences were predicted from transcripts using ExPASy translate (ExPASy, University of Geneva, Geneva, Switzerland) and analysed with SMART (<http://smart.embl-heidelberg.de>) and Pfam (<http://pfam.sanger.ac.uk/search>) to determine if a bHLH domain was coded in each transcript. Reciprocal TBlastN searches (Altschul *et al.*, 1990) were conducted on NCBI (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) to verify if the hits containing a bHLH domain belonged to the bHLH class VIII family.

Unique bHLH class VIII gene candidates (later named CnbHHLHVIII and CbbHHLHVIII) were found in the *C. nitellarum* transcriptome (transcript_12969), and in the *C. braunii* transcriptome (Cb_Transcript_119934) and genome (CHBRA233g00280) respectively. We verified the sequence of CnbHHLHVIII and CbbHHLHVIII transcripts by PCR (primers listed in Table 1) using cDNA from whole plant total RNA and Sanger sequencing (Source Bioscience, Nottingham, UK). The transcript encoding CnbHHLHVIII was 2769 nucleotides long and contained a 1725 nucleotide CDS coding for a 574 amino acid protein (Supplementary Data 1). The transcript encoding CbbHHLHVIII was 6103 nucleotides long and contained a 3468 nucleotides CDS coding for a 1155 amino acids protein (Supplementary Data 1). The CDS and protein sequences of CnbHHLHVIII and CbbHHLHVIII were uploaded on Genbank under the accession numbers MK292332 and MK292331 respectively.

Phylogeny and protein domain analysis

We used the iterative refinement method L-INS-i (Katoh *et al.*, 2005) on MAFFT--add (Katoh & Frith, 2012; Katoh & Standley, 2013) to add the full length protein sequences of CnbHHLHVIII and CbbHHLHVIII to the previously published alignment of full length bHLH proteins from the land plants *A. thaliana*, *Oryza sativa*, *S. moelendorffii*, *P. patens* and *M. polymorpha*, the streptophyte alga *Klebsormidium flaccidum*, the chlorophyte algae *Volvox*

carteri, *Chlamydomonas reinhardtii*, *Chlorella variabilis* and *Ostreococcus tauri* and, the rodophyte alga *Cyanidioschyzon merolae* (Supplementary Data 2) (Catarino *et al.*, 2016). The alignment was trimmed manually with BioEdit v.7.1 (Hall, 1999) to remove the non-conserved regions (Supplementary Data 3). Maximum-likelihood analysis was carried-out with PhyML 3.0 (Guindon *et al.*, 2010), with the Jones Taylor and Thornton (JTT) amino acid substitution model (Jones *et al.*, 1992), on the complete set of aligned and trimmed proteins (Archaeplastida bHLH phylogeny; Figure 1.A and Supplementary Figure S1) and on a reduced set (Supplementary Data 4) containing only the members of the bHLH class VIII family, CnbHLHVIII and CbbHLHVIII (bHLH class VIII phylogeny; Figure 1.B and Supplementary Figure S2). The statistical branch support values were calculated with an approximate likelihood ratio test (aLRT) (Guindon *et al.*, 2010). Trees were visualised using FigTree (<http://tree.bio.ed.ac.uk/software/figtree/>). The presence of the bHLH class VIII conserved motifs was assessed using the amino acid alignment (Supplementary Figure S3) and MEME (Multiple Em for Motif Elicitation) (Bailey & Elkan, 1994) with a constraint of 1 to 4 conserved motifs on members of each subclasses (VIIIa, HECATE, RSL class 1, RSL class 2 and RSL class 3) associated with CnbHLHVIII and CbbHLHVIII. The LOGO representation of the amino-acid sequence of each bHLH class VIII conserved motif was produced using WebLogo (<https://weblogo.berkeley.edu>).

Plant growth conditions

C. nitellarum (strain provided by the Skidmore Algal collection of Professor David Domozych) were grown in cell culture flasks (cell star, 250 mL, filter cap; Greiner Bio-One, Kremsmunster, Austria) in 75 mL of liquid NaNO_3^- -Bold Basal Medium (Nichols, 1973) supplemented with 1g.L^{-1} of NaNO_3^- , renewed every six weeks, under a cycle of 8h:16h dark:light ($38\text{ }\mu\text{mol.m}^{-2}.\text{s}^{-1}$) at 23°C at 45 rpm.

C. braunii (strain S276) (Nishiyama *et al.*, 2018) was grown under axenic conditions on a metallic net immersed in modified Forsberg liquid medium ($0.56\text{ mg.L}^{-1}\text{ Na}_2\text{HPO}_4$, $0.112\text{ g.L}^{-1}\text{ Ca}(\text{NO}_3)_2.3\text{H}_2\text{O}$, $0.1\text{ g.L}^{-1}\text{ MgSO}_4.7\text{H}_2\text{O}$, $0.0174\text{ g.L}^{-1}\text{ Na}_2\text{SiO}_3.5\text{H}_2\text{O}$, $0.03\text{ g.L}^{-1}\text{ KCl}$, $0.02\text{ g.L}^{-1}\text{ Na}_2\text{CO}_3$, $2\text{ }\mu\text{g.L}^{-1}\text{ MnCl}_2$, $2\text{ }\mu\text{g.L}^{-1}\text{ CoCl}_2$, $4\text{ }\mu\text{g.L}^{-1}\text{ CuCl}_2$, $0.4\text{ mg.L}^{-1}\text{ FeCl}_2$, $0.1\text{ mg.L}^{-1}\text{ ZnCl}_2$, $0.1\text{ mg.L}^{-1}\text{ NaMoO}_4$, $0.4\text{ mg.L}^{-1}\text{ H}_3\text{BO}_3$, $0.5\text{ g.L}^{-1}\text{ TRIS}$, 0.02 mg.L^{-1} Nitrilo triacetic acid (NTA)) (Forsberg, 1965) pH 7.8 supplemented with Kao and Michayluk vitamin solution (100X)

(#K3129; Sigma, St-Louis, MO, USA) at 23°C under a light cycle of 8 h : 16 h, dark : light (38 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$).

M. polymorpha wild type accessions Takaragaike-1 (Tak-1) male and Takaragaike-2 (Tak-2) female (Ishizaki *et al.*, 2008) and the loss of function *Mprsl1-1* mutant (Proust *et al.*, 2016) were used in this study. Meristem-containing thallus fragments of axenic *Mprsl1-1* and wild type and, surface sterile spores produced from the cross of Tak-1 and Tak-2 were transformed with binary vectors (see vectors construction hereafter) using agrobacterium (strain GV3101) following published co-cultivation protocols (Ishizaki *et al.*, 2008; Kubota *et al.*, 2013). After transformation, regenerated thalli or sporelings were selected on ½ Johnson's medium 1% agar supplemented with 100 $\mu\text{g.mL}^{-1}$ of cefotaxime and 10 $\mu\text{g.mL}^{-1}$ of hygromycin or 10 μM of chlorsulfuron depending on the vector (see vector construction hereafter). For phenotypic and expression analyses, Tak-1, Tak-2 and *Mprsl1-1* thalli, Tak-1 and Tak-2 sporelings, transformed sporelings or thalli were grown in axenic conditions on ½ Johnson's medium 0.7% agar at 23°C under continuous illumination (38 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$).

Wild type *A. thaliana* Columbia (Col-0) and the loss of function *Atrhd6-3 Atrsl1-1* double mutant (Menand *et al.*, 2007) were used in this study. Plants were transformed by floral deep (Zhang *et al.*, 2006) with binary vectors (see vectors construction hereafter) using agrobacterium (strain GV3101) cultures. Before, in vitro culture seeds were surface sterilized ten minutes with a solution of 70% Ethanol and 0.1% Triton X-100 and ten minutes with a solution of 99% ethanol. After transformation seeds were selected on MS medium 1% agar supplemented with 50 $\mu\text{g.mL}^{-1}$ of hygromycin. For phenotypic and expression analyses, seeds of wild type, mutants and of three T2 transformant lines for each construct were grown as previously described (Breuninger *et al.*, 2016).

Phenotypic analysis and image acquisition

Transmitted light microscopy images of *C. braunii*, *M. polymorpha* and *A. thaliana* were captured using a camera (Leica DFC310 FX; Leica, Wetzlar, Germany) mounted on a dissecting microscope (Leica M165 FC; Leica, Wetzlar, Germany). For *A. thaliana*, 15 plants per line were phenotyped 10 days after germination. For each line of *M. polymorpha* over-expressing 35S:*AtRHD6*, *MpRSL1* and *CbbHLHVIII* in the wild type background, 5 plants were

phenotyped four weeks after spore transformation. 20 plants were phenotyped for each line generated by transforming the vector into the *M. polymorpha* *Mprsl1* knockout mutant background, seven and ten weeks after thallus regeneration.

C. braunii rhizoid nodal growth pattern was analysed on 107 nodes (32 N₁, 26 N₂, 22 N₃, 15 N₄ and 12 N_{older}) from 13 plants. The stage (N₁, N₂, N₃, N₄ or N_{older}) and the presence or absence of initiating, unicellular elongating, multicellular and branched rhizoids were noted for each node. The percentage of node of each stage (N₁, N₂, N₃, N₄ and N_{older}) displaying initiating, unicellular elongating, multicellular and branched rhizoids were calculated. Most nodes displayed rhizoids at different stages and were therefore used in the calculation of the percentage of node for each of the rhizoid categories they displayed.

RNA extraction, cDNA synthesis

Total RNA was extracted with a mirTMVana Kit (#AM1560; Thermo Fisher scientific, Waltham, MA, USA) from frozen *C. braunii* whole plants for cDNA synthesis and cloning. Total RNA for transcriptomes was isolated using the same protocol from frozen rhizoids, green thalli with attached gametangia and whole plants. mRNA was extracted with a Dynabeads[®]mRNA DIRECTTM Kit (#61011; Thermo Fisher scientific, Waltham, MA, USA) from ground frozen tissues collected as follows. Two weeks after propagation, rhizoids grown below a metallic net were cut and flash frozen in liquid nitrogen for total RNA and mRNA extraction. *C. braunii* thallus cleaned from any remaining rhizoids with gametangia (for total RNA extraction) or without (for mRNA extraction) and, thallus parts (apices, nodes, branches, gametangia and zygotes) were collected by hand under a dissecting microscope (Leica M165 FC; Leica, Wetzlar, Germany) and flash frozen in liquid nitrogen. Total RNA and mRNA were DNase treated using the Turbo DNaseTM Kit (#AM2238; Thermo Fisher scientific, Waltham, MA, USA) according manufacturer's instructions. cDNAs were synthesized from 5 µg of total RNA or 20 ng of mRNA in 20 µL reaction volume using the SuperScript III First-strand synthesis System (#18080051; Thermo Fisher scientific, Waltham, MA, USA) with the oligo (dT₂₀) provided.

Total RNA was extracted with Direct-Zol RNA Miniprep Kits (#R2060; Zymo research, Irvine, CA, USA) from ground frozen whole plant for *C. nitellarum* and *M. polymorpha* and, from roots for *A. thaliana*. *C. nitellarum*, *M. polymorpha* and *A. thaliana* samples of total

RNA were DNase treated using the Turbo DNase-free™ Kit (#AM1907; Thermo Fisher scientific, Waltham, MA, USA). cDNAs were synthesised from 1 µg of total RNA in a 20 µL reaction volume using 200 U of Protoscript II reverse transcriptase (#M0368S; NEB, Ipswich, MA, USA), oligo (dT₁₇) at 2 µM and the NEB Murine RNase inhibitor (#M0314; NEB, Ipswich, MA, USA).

Construction of the *C. braunii* transcriptome

The whole plant *C. braunii* transcriptome was produced by the sequencing of ten strand-specific cDNA libraries (two whole plants libraries, two green thallus libraries and four rhizoid libraries) with an insert size of 500 bp, using a paired-end read length of 2 X 100 bp on Illumina Hiseq2000. GATC Biotech Ltd (Eurofingonomics, Konstanz, Germany) performed the preparation of the libraries from 2 µg of total RNA and sequencing.

Raw reads were quality trimmed with Trimmomatic-0.32 (Bolger *et al.*, 2014), to remove Illumina adaptors and low quality tails. Ribosomal RNA was filtered out with Sortmerna-1.9 (Kopylova *et al.*, 2012). Reads were further error corrected using Allpaths-LG-4832 (Butler *et al.*, 2008) (with setting PAIRED_SEP=-20 and ploidy = 1). Trimmed and corrected reads were normalised using Khmer-0.7.1 with a khmer size of 31. Before assembly, paired end reads were stitched together using Allpaths-LG-4832 (Butler *et al.*, 2008). A de novo transcriptome assembly was made with the cleaned, stitched reads using SGA (Simpson & Durbin, 2012), SSPACE-v3 (Boetzer *et al.*, 2011), and CAP3 (Huang & Madan, 1999). Finally assembled scaffolds were corrected using Pilon-1.6 (Walker *et al.*, 2014). The transcriptome assembly of *C. braunii* consisted of 117,611 transcripts with a mean sequence length of 749 bp. 21,917 of the transcripts were over 1kb in length. This Transcriptome Shotgun Assembly project has been deposited at DDBJ/EMBL/GenBank under the accession GGXX00000000. The version described in this paper is the first version, GGXX01000000.

The expression level of each transcript was next quantified. Raw reads were quality trimmed with Trimmomatic-0.32 (Bolger *et al.*, 2014), Sortmerna-1.9 (Kopylova *et al.*, 2012) and BAYESHAMMER (SPADES-3.5.0) (Nikolenko *et al.*, 2013). Corrected reads were mapped to the transcriptome using Salmon 0.9.1 (Patro *et al.*, 2017).

Cloning and vector construction

The CDSs encoding CbbHLHVIII, CnbHLHVIII, MpRSL1 and AtRHD6 were amplified respectively from *C. braunii*, *C. nitellarum*, *M. polymorpha* and *A. thaliana* cDNA made from total RNA, using the high-fidelity polymerase Phusion (#F530L; Thermo Fisher Scientific, Waltham, MA, USA) with the primers listed in Table 1. For *C. braunii* and *C. nitellarum* we used a slow elongation rate of 70s.Kb⁻¹. The CDSs of CbbHLHVIII, MpRSL1 and AtRHD6 were then cloned into pCR8-GW-TOPO vectors (#K2500-20; Thermo Fisher Scientific) to establish Gateway™ (GW) entry vectors. For *A. thaliana* transformations, the pCR8 entry vectors containing the CDSs encoding CbbHLHVIII and AtRHD6 were recombined with the destination vector pCambia p35S:GW:T (Breuninger *et al.*, 2016) by LR reactions (#12538-200; Thermo Fisher Scientific). For *M. polymorpha* transformations, the entry vectors containing the CDSs encoding CbbHLHVIII and MpRSL1 were recombined by LR reactions (#12538-200; Thermo Fisher scientific, Waltham, MA, USA) with the destination vectors pCambia p35S:GW:T (carrying the resistance to hygromycin) (Breuninger *et al.*, 2016) and pCambia pMpEF1α:GW:T carrying a mutated version of the MpALSm gene driven by the p35S promoter and conferring plant resistance to chlorsulfuron. This vector was generated by splicing PCR products of the MpALSm gene with the 35S promoter together with the MpEF1α promoter into a XhoI/PstI-digested pCambia backbone using the Clontech In-Fusion kit (Clontech Cat. #: 638909) and the primers indicated in Table 1. Subsequently, this vector was converted into a Gateway™ destination vector using the inserted SmaI site next to the MpEF1α promoter using the Gateway™ Vector Conversion System (Thermo Fisher Cat. #: 11828029).

Expression analysis

Quantitative PCR analyses were done on a 7300 Applied Biosystems thermocycler (Thermo Fisher scientific, Waltham, MA, USA) using the SensiFAST™ Sybr® Hi-ROX Kit (#BIO-92020; Bioline, London, UK). Amplification was performed in a reaction volume of 10 µL containing 500 nM of each of the forward and reverse primers and 4 µL of a two-fold dilution of *C. braunii* cDNA synthesized from mRNA or of ten-fold dilutions of *M. polymorpha* and *A.*

thaliana cDNA synthesized from total RNA. Data analysis were carried out as previously described (Saint-Marcoux *et al.*, 2015) using the reference genes *AtUBQ10* and *AtPDF2* (Czechowski *et al.*, 2005), *MpACT* and *MpCUL* (Saint-Marcoux *et al.*, 2015) and, *CbELF5a* (CHBRA130g00470). All primer used are listed in Table 1.

RESULTS

There is a single class VIII basic helix loop helix transcription factor in the *Chara braunii* genome

To determine if genes encoding class VIII bHLH transcription factors control rhizoid development in charophyceae algae, we searched a *C. braunii* transcriptome for similar sequences. We isolated RNA from whole *C. braunii* plants, green parts and from rhizoids. We sequenced the RNA and assembled a transcriptome with an N50 value of 1229 bp and a mean length of 749 bp (Table 2). Using the TBLASTN algorithm and the class VIII protein sequences of *A. thaliana*, *Selaginella moelendorffi*, *Physcomitrella patens* and *M. polymorpha* (Catarino *et al.*, 2016) as queries, we identified a single nucleotide sequence encoding a predicted protein of 1155 amino acids similar to the land plant class VIII proteins (*Cb_Transcript_119934*) (Supplementary Data 1). We further searched the *C. braunii* genome and transcriptome assemblies (Nishiyama *et al.*, 2018), using the same queries and the complete sequence of the putative *C. braunii* class VIII bHLH protein as a query but did not identify further new class VIII sequences. We also identified a putative class VIII bHLH protein from a *Coleochaete nitellarum* transcriptome (Bonnot *et al.*, 2017) (Supplementary Data 1). No putative class VIII bHLH sequence was found in *Klebsormidium flaccidum* (Hori *et al.*, 2014; Catarino *et al.*, 2016).

To test if the putative *C. braunii* and *C. nitellarum* class VIII bHLH proteins belong to the class VIII bHLH family, we aligned the sequences around their bHLH domain with bHLH sequences from 11 Archaeplastida taxa and generated a phylogenetic tree using the maximum likelihood statistics (Figure 1 and Supplementary Figure S1 and S2) (Catarino *et al.*, 2016). The topology of the tree indicated that the *C. braunii* putative class VIII bHLH protein was sister to the RSL (subclass VIIIc) proteins while *C. nitellarum* putative class VIII bHLH protein was sister to the RSL class 1 (subclass VIIIc 1) and class 2 (subclass VIIIc 2). Since the *C. braunii* and *C. nitellarum* proteins constituted a monophyletic group with the other class VIII bHLH proteins, we conclude that the *C. braunii* and *C. nitellarum* proteins are

class VIII members, and designated them respectively CbbHLHVIII and CnbHLHVIII. These data indicate that a single gene encoding a class VIII bHLH protein existed in the last common ancestor of *C. braunii*, *C. nitellarum* and the land plants. They also demonstrate that a single copy class VIII bHLH gene is present in both *C. braunii* that develops rhizoids and in *C. nitellarum* that does not develop rhizoids.

CbbHLHVIII and CnbHLHVIII lack the conserved motifs present in the land plant class VIII proteins

Land plant class VIII bHLH proteins are characterised by distinct, conserved motifs near and in the bHLH domain (Figure 2). They possess a very conserved bHLH domain (Figure 2.A and Supplementary Figure S3) containing an atypical basic domain characterised by the presence of a conserved Alanine (A²¹⁰ in AtRHD6) at position 9 of the bHLH domain (Figure 2.A and Supplementary Figure S3) that is not found in the other bHLH classes (Supplementary Figure S4 and Data S2) (Heim *et al.*, 2003; Liljegren *et al.*, 2004; Gremski *et al.*, 2007). There is an atypical basic domain, typical of class VIII bHLH proteins, in both CbbHLHVIII and CnbHLHVIII (Figure 2.C and Supplementary Figure S3 and S4).

There are conserved characteristic amino acid motifs in RSL (subclass VIIIc) and HECATE (subclass VIIIb) proteins (Pires & Dolan, 2010a). A conserved motif is located just after the bHLH domain in members of the RSL subfamily (VIIIc) (Figure 2.C and Supplementary Figure S3). The precise sequence of this motif is characteristic of each of the three monophyletic RSL subclasses. KVLATDEFWPAQGGKAPDISQVKDALDAI is found in members of the RSL subclass 1 (VIIIc1), APIAYNGMDIG in members of the RSL subclass 2 (VIIIc2) and NKDSASEVKCEKWKEFIDAQT in members of the RSL subclass 3 (VIIIc3) (Figure 2.B). Similarly, a conserved sequence (DPIAVSRPKRRNVRI) is located just before the N-terminal of the bHLH domain in members of the HECATE (VIIIb) subclass (Figure 2.B and C). None of the conserved amino-acid motifs of the RSL and HECATE proteins are present in the CbbHLHVIII or CnbHLHVIII protein sequences (Figure 2.C and Supplementary Figure S3). This suggests that these land plant specific motifs evolved after the divergence of *C. braunii* and *C. nitellarum* from the last common ancestors with the land plants.

CbbHLHVIII protein cannot replace the RSL proteins that positively regulate root hair development in *Arabidopsis thaliana*

To assess if *CbbHLHVIII* controls the development of filamentous rooting cells (rhizoids and root hairs) in land plants, we tested if *CbbHLHVIII* could substitute for the loss of RSL function in *A. thaliana* mutants. *Atrhd6 Atrsl1* double mutants are devoid of RSL class 1 function and do not develop root hairs (Figure 3) (Menand *et al.*, 2007). To test the ability of *CbbHLHVIII* to restore root hair development, *Atrhd6 Atrsl1* double mutants were transformed with a gene construct driving the expression of *CbbHLHVIII* under the control of the cauliflower mosaic virus 35S constitutive promoter (*p35S:CbbHLHVIII:T*). We identified three lines that expressed *CbbHLHVIII* at high levels (Figure 3.A and B). None of these *Atrhd6 Atrsl1 p35S:CbbHLHVIII:T* plants developed root hairs (Figure 3.C). As a control, *Atrhd6 Atrsl1* double mutants were transformed with either a *p35S:AtRHD6:T* or a *p35S:MpRSL1:T* gene constructs which overexpressed the *A. thaliana* and *M. polymorpha* RSL subclass 1 genes respectively (Figure 3.A and B). Both the *Atrhd6 Atrsl1 p35S:AtRHD6:T* and the *Atrhd6 Atrsl1 p35S:MpRSL1:T* plants developed root hairs (Figure 3.C). These data indicate that expression of a class VIII protein from *C. braunii* cannot compensate for loss of *AtRHD6* and *AtRSL1* function. This suggests *CbbHLHVIII* does not function in rhizoid development and indicates that *CbbHLHVIII* is functionally different from the RSL subclass 1 proteins *AtRHD6* and *AtRSL1*.

CbbHLHVIII* protein cannot replace the RSL protein that positively regulates rhizoid development in *Marchantia polymorpha

To independently determine if the function of *CbbHLHVIII* could substitute for RSL function in land plants, we tested the ability of *CbbHLHVIII* to restore rhizoid development in the rhizoidless *Mprsl1* loss of function mutant of *M. polymorpha* (Figure 4 and 5). We transformed the *Mprsl1* mutant with a gene construct in which the *CbbHLHVIII* gene was under the control of the constitutive promoter of *M. polymorpha* *EF1α* (*pMpEF1α:CbbHLHVIII:T*) (Althoff *et al.*, 2014). Five *Mprsl1 pMpEF1α:CbbHLHVIII:T* lines with high steady state levels of the *C. braunii* transgene were identified (Figure 4). None of these *Mprsl1 pMpEF1α:CbbHLHVIII:T* plants developed rhizoids (Figure 5). As a control, we transformed the *Mprsl1* loss of function mutant with a *pMpEF1α:MpRSL1:T* gene construct. *Mprsl1* plants transformed with *pMpEF1α:MpRSL1:T* developed rhizoids indicating that overexpression of the *MpRSL1* class VIII protein driven by the promoter of *M. polymorpha* *EF1α* compensate for the loss of *MpRSL1* function (Figure 5). The lack of rhizoid

development on *Mprs1* pMpEF1 α :CbbHLHVIII:T plants demonstrates that class VIII bHLH protein from *Chara braunii* (CbbHLHVIII) cannot substitute for MpRSL1 function during rhizoid development in *Mprs1* mutants. These data suggest that CbbHLHVIII and MpRSL1 are functionally different.

Expression of CbbHLHVIII in wild type *Marchantia polymorpha* does not induce supernumerary rhizoid development

To verify that CbbHLHVIII could not promote rhizoid development in *M. polymorpha* using a different experimental approach, we ectopically overexpressed CbbHLHVIII in wild type and compared the phenotypes to the phenotypes of plants that ectopically overexpress the endogenous MpRSL1 gene (Figure 6). Ectopic overexpression of MpRSL1 using the 35S promoter (p35S:MpRSL1:T) induced the development of supernumerary rhizoids in wild type. However, expression of CbbHLHVIII from the 35S promoter (p35S:CbbHLHVIII:T) in wild type *M. polymorpha* did not induce supernumerary rhizoid development. This verifies that CbbHLHVIII cannot function during rhizoid development in *M. polymorpha*. This is consistent with the hypothesis that CbbHLHVIII does not control rhizoid development and that CbbHLHVIII and MpRSL1 proteins are functionally different.

CbbHLHVIII protein cannot replace the RSL protein that positively regulates gemma development in *Marchantia polymorpha*

To independently verify that CbbHLHVIII cannot substitute for MpRSL1 in *M. polymorpha*, we tested the ability of CbbHLHVIII to restore gemma development in *Mprs1* mutants (Figure 5). *Mprs1* mutants rarely develop gemmae; the gemma cups of *Mprs1* mutants are empty while gemmae can fill gemma cups of wild type plants. None of the *Mprs1* plants transformed with the pMpEF1 α :CbbHLHVIII:T gene construct developed gemmae and gemma cups were empty. The control *Mprs1* plants transformed with the pMpEF1 α :MpRSL1:T gene construct developed gemmae and gemma cups were full. Together these data indicate that overexpression of the CbbHLHVIII protein cannot compensate for loss of MpRSL1 function during gemma development in *Mprs1* mutants. This is consistent with the hypothesis that the function of CbbHLHVIII is different from MpRSL1.

***Chara braunii* develop rhizoids from multicellular nodes**

The green thallus of *C. braunii* comprises several axes, each consisting of alternating nodes and internodes (Figure 7.A) (Smith & Allen, 1955; Pickett-Heaps, 1975; Graham & Wilcox, 2000; Nishiyama *et al.*, 2018). The internodes are composed of a single elongated cell, while the nodes are multi-cellular and are the sites from which branches (determinate structures) and new axes (indeterminate structures) initiate. To define where *C. braunii* rhizoids develop, we grew individual *C. braunii* algae in axenic liquid culture and found that rhizoids developed from the nodal complexes (Figure 7). No rhizoids were present on nodes 1 and 2 where node 1 is the node nearest the apex (Figure 7.B). After initiation from a nodal cell, the rhizoid elongates by tip growth (Supplementary Figure S5) and cell division. Growing rhizoids can branch (Supplementary Figure S5). Rhizoids of all developmental stages – initiating rhizoids, elongating rhizoids, multicellular rhizoids and branched rhizoids – are present on node 3 and older nodes (Figure 7.B). These observations indicate that, in our growth condition, rhizoids develop from the nodal complexes of the thallus. Nodal initiation of rhizoids begins from the nodes in 3rd position from the apical meristem and initiation continues in older nodes (Figure 7.B and C).

CbbHLHVIII* is expressed at the apex and gametangia of *Chara braunii

The demonstration that expression of *CbbHLHVIII* does not restore rhizoid development in *Atrhd6 Atrs1* double mutants or *Atrhd6 Atrs1* double mutants suggests that this protein is not involved in rhizoid development in *C. braunii*. In land plants, RSL Class 1 encoding genes (subclass VIIIc) are expressed in filamentous rooting cells – rhizoid cells and root hair cells – and the cells from which rhizoid cells and root hair cells develop (Menand *et al.*, 2007; Jang *et al.*, 2011; Kim & Dolan, 2016; Proust *et al.*, 2016; Kim *et al.*, 2017). To determine where *CbbHLHVIII* is expressed during *C. braunii* development, we measured the steady state levels of *CbbHLHVIII* mRNA using quantitative RT-PCR (Figure 8). In the first experiment, RNA was isolated from thallus, rhizoids, gametangiophores (antheridia and archegonia) and zygotes (Figure 7 and Supplementary Figure S6). The highest steady state levels of *CbbHLHVIII* mRNA were detected in the thallus and gametangiophores (Figure 8.A). Low steady state levels of *CbbHLHVIII* mRNA were detected in rhizoids and zygotes. These results were consistent with the levels of expression of *CbbHLHVIII* (Cb_Transcript_119934) detected in the thallus and rhizoid transcriptomes (Supplementary Table 1). To identify where in the thallus *CbbHLHVIII*

mRNA accumulated, we isolated RNA from different thallus structures (Figure 8.B). Steady state levels of *CbbHLHVIII* mRNA were highest at the apex (apical meristem and node 1) and levels were progressively lower in nodes 2 and nodes 3. Relatively high steady state levels of *CbbHLHVIII* mRNA were observed in gametangia-bearing branches, while *CbbHLHVIII* mRNA levels were lower in branches without gametangia. This suggests that *CbbHLHVIII* mRNA accumulates in apices and gametangia. These data are consistent with *CbbHLHVIII* being expressed in the morphogenetic centres of *C. braunii*, the apex, node 1 and gametangia. The low steady state levels of *CbbHLHVIII* mRNA in rhizoids and in nodes bearing initiating rhizoids (node 3 and older) suggest that *CbbHLHVIII* expression is not involved in rhizoid development. Taken together these data indicate that *CbbHLHVIII* is expressed in the *C. braunii* morphogenetic centres but suggest that this transcription factor not involved in rhizoid development.

DISCUSSION

Charophycean algae develop complex bodies from morphogenetic centres located at the apices of their axes, like land plants (Smith & Allen, 1955; Pickett-Heaps, 1975; Graham & Wilcox, 2000). These axes develop a diversity of cell types including tip-growing cells called rhizoids that are involved in nutrient uptake (Box, 1986, 1987; Andrews, 1987; Vermeer *et al.*, 2003; Wuestenberg *et al.*, 2011) and anchorage (Graham & Wilcox, 2000). Tip-growing cells involved in nutrient uptake and anchorage also evolved among the land plants (Jones & Dolan, 2012; Bonnot *et al.*, 2017): the tip-growing rhizoids of bryophytes and root hairs of euphyllophytes. No other streptophyte algae form morphogenetic centres or develop the same level of cellular complexity that is characteristic of the charophycean algae (Pickett-Heaps, 1975; Graham & Wilcox, 2000). The distribution of morphological traits on the most recent streptophyte phylogenies suggests that increased body plan complexity evolved independently in the charophycean lineage and in the lineage leading to the land plants (embryophytes) (Wodniok *et al.*, 2011; Wickett *et al.*, 2014; Nishiyama *et al.*, 2018). If rhizoid evolved independently in the land plants and charophycean lineages, we might predict that different mechanisms regulating rhizoid differentiation evolved in the two lineages.

Two broad conclusions can be made from our results. First, we conclude that different mechanisms control the initiation of rhizoid development in Charophycean algae

and land plants. Second, we conclude that the acquisition of the rhizoid development function by class VIII bHLH evolved in the land plant lineage after the divergence of extant streptophyte algae and the land plants from their last common ancestor.

Class VIII bHLH proteins are required for the formation of tip-growing rooting cells in land plants (Masucci & Schiefelbein, 1994; Menand *et al.*, 2007; Jang *et al.*, 2011; Kim & Dolan, 2016; Proust *et al.*, 2016; Kim *et al.*, 2017). Here, we identified a class VIII bHLH protein (*CbbHLHVIII*) gene from the charophycean alga *C. braunii* that is sister to land plant RSL genes on protein trees. *CbbHLHVIII* transcripts accumulate at high levels in parts of the plant undergoing morphogenesis, including the apex, the nodes and the gametangia. However, the expression of *CbbHLHVIII* was very low in rhizoids and the transcript was hardly detectable in regions of the plant from which rhizoids develop. This suggests that *CbbHLHVIII* does not function in rhizoid development in *C. braunii*. This conclusion is supported by the observation that a single class VIII bHLH protein is also present in the genome of *C. nitellarum*, which does not develop rhizoids (Pickett-Heaps, 1975; Graham & Wilcox, 2000). These data support the hypothesis that class VIIIc bHLH proteins carry out different functions in streptophyte algae and land plants.

Further evidence supporting the hypothesis that *CbbHLHVIII* does not control rhizoid development is its inability to promote rhizoid development when expressed in land plants. *CbbHLHVIII* does not substitute for loss of RSL class 1 function in land plant *rs1* loss of function mutants. This indicates that *CbbHLHVIII* is functionally different from land plant RSL class 1 genes. This finding is further supported by the inability of *CbbHLHVIII* expression to restore gemma development in *rs1* loss of function mutants of *M. polymorpha*. These results suggest that RSL class 1 proteins acquired the ability to promote the development of surface structures from single epidermal cells, including rhizoids, after *C. braunii* and land plants last shared a common ancestor. If correct, it would suggest that class VIII proteins evolved novel functions during or after the transition to land. It is possible that these novel functions are conferred by the RSL motifs that are conserved among all land plant RSL proteins but are not present in the streptophyte alga (*C. braunii*, *C. nitellarum*) RSL proteins.

An alternative, though less parsimonious, hypothesis is that the ancestral function of class VIII bHLH transcription factors was to control the development of structures from single cells as it is in land plants, but that this function was lost in both the *C. braunii* and *C. nitellarum* lineages. Ultimately, discovering the function of class VIII bHLH proteins in *Chara*

species and *Coleochaete* species requires the functional characterisation of their class VIII bHLH genes. While such characterisation is not currently possible in these algae it would demonstrate the ancestral function of these proteins in the last common ancestor using the comparative approach. The function of RSL class 1 genes in land plants is to control the development of structures that develop from single surface cells. If class VIII bHLH transcription factors controlled the development of a similar process in streptophyte algae, it would suggest that this function was ancestral and likely acted in the last common ancestor of these organism and land plants. However, the current data suggest that function of family VIII bHLH transcription factors in the development of streptophyte algae is entirely different from their function in land plants.

Taken together, the data support a model in which there was a single class VIII bHLH protein in the genome of a streptophyte algal ancestor of the land plants. This gene did not function in rhizoid development. Neofunctionalization occurred in the lineage leading to the land plants and was followed by multiple rounds of gene duplication (Pires & Dolan, 2010a,b; Catarino *et al.*, 2016). These duplicated genes control the development of diverse structures, such as specialised cells and organs derived from single epidermal cells including rhizoids and root hairs (*RSL* genes) (Masucci & Schiefelbein, 1994; Menand *et al.*, 2007; Jang *et al.*, 2011; Kim & Dolan, 2016; Proust *et al.*, 2016; Kim *et al.*, 2017) as well as complex organ development (*HECATE* genes) (Liljegren *et al.*, 2004; Gremski *et al.*, 2007; Kay *et al.*, 2013). This is consistent with the hypothesis that the increase in morphological diversity that accompanied the colonisation of the land resulted from gene duplication and neofunctionalisation of ancestral regulatory genes.

ACKNOWLEDGMENTS

This work was supported by a European Research Council Advanced award (EVO500 project no. 25028) and a Marie Skłodowska-Curie award (PLANTORIGINS project no. 238640) to L.D. A.J.H. was supported by a Biotechnology and Biological Sciences Research Council Doctoral Training Partnership Scholarship (Grant BB/J014427/1). We are grateful to Pr D. Domozych (Skidmore College, NY, USA) for providing *C. nitellarum* and, to Pr H. Sakayama and Pr T. Nishiyama for providing *C. braunii*.

AUTHOR CONTRIBUTIONS

L.D. and C.B. designed the research and wrote the paper. C.B. and H.B. produced the RNA samples for *C. braunii* transcriptome RNAseq. A.J.H. and S.K. performed *C. braunii* transcriptome assembly. H.B. built the vectors backbones. C.B. and C.C. performed *M. polymorpha* transformations. C.B. performed all other experiments and analysed all the data.

BIBLIOGRAPHY

Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T, Zachgo S. 2014. Comparison of the MpEF1 α and CaMV35 promoters for application in *Marchantia polymorpha* overexpression studies. *Transgenic research* **23**: 235–244.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of molecular biology* **215**: 403–410.

Andrews M. 1987. Phosphate uptake by the component parts of *Chara hispida*. *British Phycological Journal* **22**: 49–53.

Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to discover motifs in bipolymers. *Proceedings of the second international Conference on Intelligent systems for molecular biology*. Menlo Park, CA, USA: AAAI Press, 28–36.

Banks JA, Nishiyama T, Hasebe M, Bowman JL, Gribskov M, dePamphilis C, Albert VA, Aono N, Aoyama T, Ambrose BA. 2011. The *Selaginella* genome identifies genetic changes associated with the evolution of vascular plants. *Science* **332**: 960–963.

Boetzer M, Henkel C V, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27**: 578–579.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

- 594 **Bonnot C, Proust H, Pinson B, Colbalchini FP, Lesly-Veillard A, Breuninger H, Champion C,**
 595 **Hetherington AJ, Kelly S, Dolan L. 2017.** Functional PTB phosphate transporters are present
 596 in streptophyte algae and early diverging land plants. *New Phytologist* **214**: 1158–1171.
- 597 **Bowman JL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S, Nishihama R,**
 598 **Nakamura Y, Berger F. 2017.** Insights into land plant evolution garnered from the
 599 *Marchantia polymorpha* genome. *Cell* **171**: 287–304. e15.
- 600 **Box RJ. 1986.** Quantitative short-term uptake of inorganic phosphate by the *Chara hispida*
 601 rhizoid. *Plant, Cell & Environment* **9**: 501–506.
- 602 **Box RJ. 1987.** The uptake of nitrate and ammonium nitrogen in *Chara hispida* L.: the
 603 contribution of the rhizoid. *Plant, Cell & Environment* **10**: 169–176.
- 604 **Breuninger H, Thamm A, Streubel S, Sakayama H, Nishiyama T, Dolan L. 2016.**
 605 Diversification of a Transcription Factor Family Led to the Evolution of Antagonistically
 606 Acting Genetic Regulators of Root Hair Growth. *Current Biology* **26**: 1622–1628.
- 607 **Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, Nusbaum C, Jaffe**
 608 **DB. 2008.** ALLPATHS: de novo assembly of whole-genome shotgun microreads. *Genome*
 609 *research* **18**: 810–20.
- 610 **Catarino B, Hetherington AJ, Emms DM, Kelly S, Dolan L. 2016.** The stepwise increase in the
 611 number of transcription factor families in the Precambrian predated the diversification of
 612 plants on land. *Molecular biology and evolution* **33**: 2815–2819.
- 613 **Chanderbali AS, He F, Soltis PS, Soltis DE. 2015.** Out of the water: origin and diversification
 614 of the LBD gene family. *Molecular biology and evolution* **32**: 1996–2000.
- 615 **Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible W-R. 2005.** Genome-wide
 616 identification and testing of superior reference genes for transcript normalization in
 617 *Arabidopsis*. *Plant physiology* **139**: 5–17.
- 618 **Datta S, Prescott H, Dolan L. 2015.** Intensity of a pulse of RSL4 transcription factor synthesis
 619 determines *Arabidopsis* root hair cell size. *Nature Plants* **1**: 15138.

- 620 **Delaux P-M, Nanda AK, Mathé C, Sejalon-Delmas N, Dunand C. 2012.** Molecular and
621 biochemical aspects of plant terrestrialization. *Perspectives in Plant Ecology, Evolution and*
622 *Systematics* **14**: 49–59.
- 623 **Floyd SK, Bowman JL. 2007.** The ancestral developmental tool kit of land plants.
624 *International journal of plant sciences* **168**: 1–35.
- 625 **Forsberg C. 1965.** Nutritional studies of Chara in axenic cultures. *Physiologia Plantarum* **18**:
626 275–290.
- 627 **Gibling MR, Davies NS. 2012.** Palaeozoic landscapes shaped by plant evolution. *Nature*
628 *Geoscience* **5**: 99.
- 629 **Graham LE, Cook ME, Busse JS. 2000.** The origin of plants: body plan changes contributing
630 to a major evolutionary radiation. *Proc Natl Acad Sci USA* **97**.
- 631 **Graham LE, Wilcox LW. 2000.** *Algae*. Prentice-Hall, Inc.
- 632 **Gremski K, Ditta G, Yanofsky MF. 2007.** The HECATE genes regulate female reproductive
633 tract development in Arabidopsis thaliana. *Development* **134**: 3593–3601.
- 634 **Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010.** New
635 algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
636 performance of PhyML 3.0. *Systematic biology* **59**: 307–321.
- 637 **Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, Bailey PC. 2003.** The basic helix–
638 loop–helix transcription factor family in plants: a genome-wide study of protein structure
639 and functional diversity. *Molecular biology and evolution* **20**: 735–747.
- 640 **Honkanen S, Dolan L. 2016.** Growth regulation in tip-growing cells that develop on the
641 epidermis. *Current opinion in plant biology* **34**: 77–83.
- 642 **Hori K, Maruyama F, Fujisawa T, Togashi T, Yamamoto N, Seo M, Sato S, Yamada T, Mori**
643 **H, Tajima N. 2014.** Klebsormidium flaccidum genome reveals primary factors for plant
644 terrestrial adaptation. *Nature communications* **5**.

645 **Huang X, Madan A. 1999.** CAP3: A DNA sequence assembly program. *Genome research* **9**:
646 868–877.

647 **Ishizaki K, Chiyoda S, Yamato KT, Kohchi T. 2008.** Agrobacterium-mediated transformation
648 of the haploid liverwort *Marchantia polymorpha* L., an emerging model for plant biology.
649 *Plant and cell physiology* **49**: 1084–1091.

650 **Jang G, Yi K, Pires ND, Menand B, Dolan L. 2011.** RSL genes are sufficient for rhizoid system
651 development in early diverging land plants. *Development* **138**: 2273–2281.

652 **Harrison JC. 2017.** Development and genetics in the evolution of land plant body plans.
653 *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**: 20150490.

654 **Jones VAS, Dolan L. 2012.** The evolution of root hairs and rhizoids. *Annals of botany* **110**:
655 205–212.

656 **Jones DT, Taylor WR, Thornton JM. 1992.** The rapid generation of mutation data matrices
657 from protein sequences. *Bioinformatics* **8**: 275–282.

658 **Ju C, Van de Poel B, Cooper ED, Thierer JH, Gibbons TR, Delwiche CF, Chang C. 2015.**
659 Conservation of ethylene as a plant hormone over 450 million years of evolution. *Nature*
660 *Plants* **1**: 14004.

661 **Katoh K, Frith MC. 2012.** Adding unaligned sequences into an existing alignment using
662 MAFFT and LAST. *Bioinformatics* **28**: 3144.

663 **Katoh K, Kuma K, Toh H, Miyata T. 2005.** MAFFT version 5: improvement in accuracy of
664 multiple sequence alignment. *Nucleic acids research* **33**: 511–518.

665 **Katoh K, Standley DM. 2013.** MAFFT multiple sequence alignment software version 7:
666 improvements in performance and usability. *Molecular biology and evolution* **30**: 772–780.

667 **Kay P, Groszmann M, Ross JJ, Parish RW, Swain SM. 2013.** Modifications of a conserved
668 regulatory network involving INDEHISCENT controls multiple aspects of reproductive tissue
669 development in *Arabidopsis*. *New Phytologist* **197**: 73–87.

- 670 **Kenrick P, Crane PR. 1997.** *The origin and early diversification of land plants. Nature* **389**:
671 33-39.
- 672 **Kim CM, Dolan L. 2016.** ROOT HAIR DEFECTIVE SIX-LIKE class I genes promote root hair
673 development in the grass *Brachypodium distachyon*. *PLoS genetics* **12**: e1006211.
- 674 **Kim CM, Han C, Dolan L. 2017.** RSL class I genes positively regulate root hair development in
675 *Oryza sativa*. *New Phytologist* **213**: 314–323.
- 676 **Kopylova E, Noe L, Touzet H. 2012.** SortMeRNA: fast and accurate filtering of ribosomal
677 RNAs in metatranscriptomic data. *Bioinformatics* **28**: 3211–3217.
- 678 **Kubota A, Ishizaki K, Hosaka M, Kohchi T. 2013.** Efficient *Agrobacterium*-mediated
679 transformation of the liverwort *Marchantia polymorpha* using regenerating thalli.
680 *Bioscience, biotechnology, and biochemistry* **77**: 167–172.
- 681 **Lang D, Weiche B, Timmerhaus G, Richardt S, Riaño-Pachón DM, Corrêa LG, Reski R,**
682 **Mueller-Roeber B, Rensing SA. 2010.** Genome-wide phylogenetic comparative analysis of
683 plant transcriptional regulation: a timeline of loss, gain, expansion, and correlation with
684 complexity. *Genome Biology and Evolution* **2**: 488–503.
- 685 **Lehti-Shiu MD, Panchy N, Wang P, Uygün S, Shiu S-H. 2017.** Diversity, expansion, and
686 evolutionary novelty of plant DNA-binding transcription factor families. *Biochimica et*
687 *Biophysica Acta (BBA)-Gene Regulatory Mechanisms* **1860**: 3–20.
- 688 **Lenton TM, Crouch M, Johnson M, Pires N, Dolan L. 2012.** First plants cooled the
689 Ordovician. *Nature Geoscience* **5**: 86–89.
- 690 **Liljegren SJ, Roeder AH, Kempin SA, Gremski K, Østergaard L, Guimil S, Reyes DK, Yanofsky**
691 **MF. 2004.** Control of Fruit Patterning in *Arabidopsis* by INDEHISCENT. *Cell* **116**: 843–853.
- 692 **Masucci JD, Schiefelbein JW. 1994.** The *rh6* mutation of *Arabidopsis thaliana* alters root-
693 hair initiation through an auxin-and ethylene-associated process. *Plant Physiology* **106**:
694 1335–1346.

- 695 **Menand B, Yi K, Jouannic S, Hoffmann L, Ryan E, Linstead P, Schaefer DG, Dolan L. 2007.**
 696 An ancient mechanism controls the development of cells with a rooting function in land
 697 plants. *Science* **316**: 1477.
- 698 **Navaud O, Dabos P, Carnus E, Tremousaygue D, Hervé C. 2007.** TCP transcription factors
 699 predate the emergence of land plants. *Journal of molecular evolution* **65**: 23–33.
- 700 **Nichols HW. 1973.** Growth media-freshwater. *Handbook of phycological methods: culture*
 701 *methods and growth measurements* **1**: 39–78.
- 702 **Nikolenko SI, Korobeynikov AI, Alekseyev MA. 2013.** BayesHammer: Bayesian clustering for
 703 error correction in single-cell sequencing. *BMC Genomics* **14**: S7.
- 704 **Nishiyama T, Sakayama H, de Vries J, Buschmann H, Saint-Marcoux D, Ullrich KK, Haas FB,**
 705 **Vanderstraeten L, Becker D, Lang D. 2018.** The chara genome: Secondary complexity and
 706 implications for plant terrestrialization. *Cell* **174**: 448–464. e24.
- 707 **Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017.** Salmon provides fast and bias-
 708 aware quantification of transcript expression. *Nature Methods* **14**: 417–419.
- 709 **Pickett-Heaps JD. 1975.** *Green algae: structure, reproduction and evolution in selected*
 710 *genera*. Mass., Sinauer Associates.
- 711 **Pires N, Dolan L. 2010a.** Origin and diversification of basic-helix-loop-helix proteins in
 712 plants. *Molecular biology and evolution* **27**: 862–874.
- 713 **Pires N, Dolan L. 2010b.** Early evolution of bHLH proteins in plants. *Plant signaling &*
 714 *behavior* **5**: 911–912.
- 715 **Pires ND, Dolan L. 2012.** Morphological evolution in land plants: new designs with old
 716 genes. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**: 508–518.
- 717 **Prince VE, Pickett FB. 2002.** Splitting pairs: the diverging fates of duplicated genes. *Nature*
 718 *Reviews Genetics* **3**: 827–837.

719 **Proust H, Honkanen S, Jones VAS, Morieri G, Prescott H, Kelly S, Ishizaki K, Kohchi T, Dolan**
720 **L. 2016.** RSL Class I Genes Controlled the Development of Epidermal Structures in the
721 Common Ancestor of Land Plants. *Current Biology* **26**: 93–99.

722 **Rensing SA. 2014.** Gene duplication as a driver of plant morphogenetic evolution. *Current*
723 *Opinion in Plant Biology* **17**: 43–48.

724 **Rensing SA, Lang D, Zimmer AD, Terry A, Salamov A, Shapiro H, Nishiyama T, Perroud P-F,**
725 **Lindquist EA, Kamisugi Y. 2008.** The Physcomitrella genome reveals evolutionary insights
726 into the conquest of land by plants. *Science* **319**: 64–69.

727 **Saint-Marcoux D, Proust H, Dolan L, Langdale JA. 2015.** Identification of Reference Genes
728 for Real-Time Quantitative PCR Experiments in the Liverwort *Marchantia polymorpha*. *PloS*
729 *one* **10**: e0118678.

730 **Simpson JT, Durbin R. 2012.** Efficient de novo assembly of large genomes using compressed
731 data structures. *Genome Research* **22**: 549–556.

732 **Smith DM, Allen SJ. 1955.** *Cryptogamic botany*. McGraw-Hill Book company.

733 **Tanabe Y, Hasebe M, Sekimoto H, Nishiyama T, Kitani M, Henschel K, Münster T, Theissen**
734 **G, Nozaki H, Ito M. 2005.** Characterization of MADS-box genes in charophycean green algae
735 and its implication for the evolution of MADS-box genes. *Proceedings of the National*
736 *Academy of Sciences of the United States of America* **102**: 2436-2441.

737 **Vermeer CP, Escher M, Portielje R, de Klein JJ. 2003.** Nitrogen uptake and translocation by
738 *Chara*. *Aquatic Botany* **76**: 245–258.

739 **Vijayakumar P, Datta S, Dolan L. 2016.** ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes
740 root hair elongation by transcriptionally regulating the expression of genes required for cell
741 growth. *New Phytologist* **212**: 944–953.

742 **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,**
743 **Wortman J, Young SK, et al. 2014.** Pilon: an integrated tool for comprehensive microbial
744 variant detection and genome assembly improvement. *PloS one* **9**: e112963.

Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S, Barker MS, Burleigh JG, Gitzendanner MA. 2014. Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proceedings of the National Academy of Sciences* **111**: E4859–E4868.

Wodniok S, Brinkmann H, Glöckner G, Heidel A, Philippe H, Melkonian M, Becker B. 2011. Origin of land plants: Do conjugating green algae hold the key? *BMC Evolutionary Biology* **11**: 104-114.

Wuestenberg A, POeRS Y, Ehwald R. 2011. Culturing of stoneworts and submersed angiosperms with phosphate uptake exclusively from an artificial sediment. *Freshwater Biology* **56**: 1531–1539.

Zhang X, Henriques R, Lin S-S, Niu Q-W, Chua N-H. 2006. Agrobacterium-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nature protocols* **1**: 641-646.

769

770

771

772 **Table 1. List of primers for cloning, vector construction and expression analysis**

Experiment	Gene Name	Forward (F) & Reverse (R) primer sequences
CDS verification	<i>CnbHLHVIII</i>	F: CTGAGTACCGTATCTGTCTCACTTTGG
		R: GTAGTCCCTATCAAGCGTCGGTGC
	<i>CbbHLHVIII</i>	F: GAGAGAAGCCTAGAGGATGCTTCC
		R: CGTTTGATCAAGCGACGTTGAGC
Cloning	MpALS _m	F: AAATCTATCTCTCTCGAGCATGGCGTCAATTAGTGGATGC
		R: ACATTATTATGGAGAAACAAGTTAGTATATAGTCCTTCCATC
	MpEF1 α	F: ATGATACGAACGAAAGCTCCGGGGCACATCAATACTGTTGAGGCAACACC
		R: TGACGCCATGCTCGAGAGAGATAGATTGTAGAGAGAG
	<i>CnbHLHVIII</i>	F: AGCCAAGATGTCAACACAGCAGG
		R: GTAGTCCCTATCAAGCGTCGGTGC
	<i>CbbHLHVIII</i>	F: ATGAACAGTGTGGTGGCGGAGG
		R: CTAAACGATGCGTTCTTGGGGATG
	MpRSL1	F: ATGATGAGCTCGAGAACGCTGA
		R: TCAGGACGAGTTGTTGTCTGCTTG
	AtRHD6	F: ATGGCACTCGTTAATGACCATCC
		R: TTAATTGGTGATCAGATTCGAATTCC
Quantitative-PCR analysis	<i>CbbHLHVIII</i> *	F: CAGAAGCAGAGCCAAGAGGAGC
		R: CTAAACGATGCGTTCTTGGGGATG
	CbELF5A	F: CTAGCCAATCAGCTGAAGGAAGG
		R: TCAGCAAACCCTCTACATGGTCG
	MpRSL1	F: AGATGAGTCTGGGGCAACC
		R: GGATGAGCGCTTTAGAGTGG
	MpCUL [#]	F: AGGATGTGGACAAGGATAGACG
		R: GTTGATGTGGCAACACCTTG
	MpACT	F: AGGCATCTGGTATCCACGAG
		R: ACATGGTCGTTCTCCAGAC
	AtRHD6	F: CCTAAATCCGCTGGAAACAA
		R: CTCTTCGATTCTTGGCTGCT
RT-PCR analysis	AtUBQ10 [#]	F: GGCCTTGTATAATCCCTGATGAATAAG
		R: AAAGAGATAACAGGAACGGAAACATAG
	AtPDF2	F: TAACGTGGCCAAAATGATGC
		R: GTTCTCCACAACCGCTTGGT
	MpRSL1	F: ACTCTCGGGAATGACACTTCCAGG
		R: CCTTTTCAAGCATGGTGACCAAGTC
	AtRHD6	F: TCAACCGTCGAAGAACTGAG
		R: TTAATTGGTGATCAGATTCGAATTCC
	AtRSL1	F: CCCTAAACTGGCTGGCAATA

	R: TTATTCTGCTATACTTGTTTCTCTAGTTGAG
--	------------------------------------

* Annealing step at 65°C instead of 60°C; # Also used for RT-PCR analysis.

Table 2. Transcriptome parameters

Number of sequences	117611
Median sequence length	395 nucleotides
Mean sequence length	749 nucleotides
Max sequence length	21672 nucleotides
Min sequence length	200 nucleotides
No. sequences > 1kb	21917
No. sequences > 10kb	63
No. gaps	660
N50	1229
Combined sequence length	88124568

Figure 1. CbbHLHVIII is sister to land plant class VIII bHLH transcription factors.

a. Unrooted maximum-likelihood tree of the Archaeplastida bHLH transcription factors. The, the complete class VIII bHLH family including the RSL (subclass VIILc; in red) and related bHLH families (class X, XV, XIII and XIV) are marked with ellipses. CbbHLHVIII and CnbHLHVIII are marked with red and a blue triangle respectively. **b.** Maximum-likelihood tree of the class VIII bHLH transcription factors. Tree rooted with the bHLH class XIII and XIV sequences. The approximate likelihood ratio test (aLRT) support values are given for the major nodes of the tree and marked by a red circle. The red and the blue triangles mark CbbHLHVIII and CnbHLHVIII respectively. *A. thaliana* (At), *O. sativa* (Os), *S. moelendorfii* (Sm), *P. patens* (Pp) and *M. polymorpha* (Mp).

Figure 2. Conserved amino acid motifs of the class VIII bHLH family.

a. LOGO representation of the amino acid sequence of the atypical bHLH domain of class VIII proteins. The red star marks the position of the conserved Alanine (A²¹⁰ in AtRHD6) specific of the class VIII bHLH proteins. **b.** LOGO representations of the amino acid sequences of the conserved motifs of the class VIII bHLH subclasses. **c.** Position of the conserved class VIII bHLH amino acid domains in the land plant class VIII proteins and CbbHLHVIII and CnbHLHVIII. Class VIII bHLH domain (green box), HECATE domain (purple box), RSL domains

(blue boxes). The sequence of the *C. braunii* and *C. nitellarum* bHLH domains are given. Red stars mark the position of the conserved Alanine (A²¹⁰ in AtRHD6).

Figure 3. CbbHLHVIII expression does not restore root hair development on root hairless *Arabidopsis thaliana* Atrhd6 Atrsl1 mutants.

a. Histograms showing the mean steady state levels ($n=3$) of AtRHD6, MpRSL1 and CbbHLHVIII mRNA in *Arabidopsis thaliana* wild type (WT), Atrhd6 Atrsl1 double mutants and Atrhd6 Atrsl1 double mutants transformed with p35S:AtRHD6:T (AtRHD6), p35S:MpRSL1:T (MpRSL1) or p35S:CbbHLHVIII:T (CbbHLHVIII). L1, L2 and L3 are three independently transformed lines for each transgene. Transcripts levels were normalized to AtUBQ10 and AtPDF2 mRNA levels. Each biological replicate is represented by a square, a triangle and a diamond for replicate 1, replicate 2 and replicate 3, respectively. In some lines the expression of CbbHLHVIII was not detected (nd). Different letters refers to statistically different groups ($P<0,05$, Kruskal-Wallis test). **b.** Analysis of the presence or absence of AtRHD6 and Atrsl1 transcripts by RT-PCR in *Arabidopsis thaliana* wild type (WT), Atrhd6 Atrsl1 double mutants and Atrhd6 Atrsl1 double mutants transformed with p35S:AtRHD6:T (AtRHD6), p35S:MpRSL1:T (MpRSL1) or p35S:CbbHLHVIII:T (CbbHLHVIII) constructs. L1, L2 and L3 are three independently transformed lines for each transgene. AtUBQ10 is used as a reference gene. **c.** Root hair phenotypes of *Arabidopsis thaliana* wild type (WT), Atrhd6 Atrsl1 double mutant and three independent lines (L1, L2, L3) for each of the Atrhd6 Atrsl1 double mutants transformed with p35S:AtRHD6:T, p35S:MpRSL1:T or p35S:CbbHLHVIII:T constructs. Scale bars : 1 mm.

Figure 4. CbbHLHVIII expression in *Marchantia polymorpha* Mprsl1 mutants.

a. Histograms showing the mean steady state levels ($n=3$) of AtRHD6, MpRSL1 and CbbHLHVIII mRNA in *M. polymorpha* wild type male (Tak-1) and female (Tak-2), Mprsl1 mutant (m) and Mprsl1 mutants transformed with pMpEF1 α :MpRSL1:T (MpRSL1) or pMpEF1 α :CbbHLHVIII:T (CbbHLHVIII). For each construct, L1, L2 and L3 are three independently transformed lines. Transcripts levels were normalized to MpACT and MpCUL mRNA levels. Each biological replicate is represented by a square, a triangle and a diamond for replicate 1, replicate 2 and replicate 3, respectively. CbbHLHVIII was not detected (nd) in some lines. Different letters refers to statistically different groups ($P<0,05$, Kruskal-Wallis

test). **b.** Analysis of the presence or absence of *MpRSL1* transcript by RT-PCR in *Marchantia polymorpha* wild type male (Tak-1) and female (Tak-2), *Mprs1* mutant (m) and *Mprs1* mutants transformed with pMpEF1α:*MpRSL1:T* (*MpRSL1*) and pMpEF1α:*CbbHLHVIII:T* (*CbbHLHVIII*). *MpCUL* is the reference gene. L1, L2 and L3 are three independently transformed lines.

Figure 5. *CbbHLHVIII* expression in *M. polymorpha* *Mprs1* mutants does not restore rhizoid and gemmae development.

Rhizoid and gemma cup phenotypes of wild type male (Tak-1) and female (Tak-2) *M. polymorpha*, *Mprs1* mutants and three independent lines (L1, L2, L3) of *Mprs1* mutants transformed with pMpEF1α:*MpRSL1:T* or pMpEF1α:*CbbHLHVIII:T*. For each genotype the top image represents the rhizoid phenotype of regenerated thalli seven weeks after transformation. Scale bars: 3 mm. The bottom images are the gemma cups of regenerated thalli ten weeks after transformation. White arrows indicate gemma cups full of gemmae. # indicates empty gemma cups. Scale bars: 2 mm.

Figure 6. Expression of *CbbHLHVIII* in *Marchantia polymorpha* does not induce supernumerary rhizoid development.

a. Histograms showing the steady state level of *MpRSL1* and *CbbHLHVIII* mRNA in *M. polymorpha* wild type male (Tak-1) and female (Tak-2) and wild type *M. polymorpha* transformed with p35S:*CbbHLHVIII:T* (*CbbHLHVIII*) or p35S:*MpRSL1:T* (*MpRSL1*). For each construct, five lines (L1, L2, L3, L4 and L5) were independently transformed. Transcripts levels were normalized with the geometrical mean of *MpACT* and *MpCUL* expression levels. In some lines the expression of *CbbHLHVIII* was not detected (nd). Different letters refers to statistically different groups (P<0,05, Kruskal-Wallis test). **b.** Rhizoid phenotypes of *M. polymorpha* WT male (Tak-1) and female (Tak-2) and *M. polymorpha* wild type transformed with p35S:*MpRSL1:T* or p35S:*CbbHLHVIII:T*. Plants are four weeks old thalli grown from spores that were transformed with transgenes. Scale bars: 3 mm.

Figure 7. Rhizoids develop on *Chara braunii* nodal complexes.

a. *C. braunii* thallus. (a) Nodal complexes on the main axis of the thallus are framed in white. A star indicates the unicellular internodes of the main axis of the thallus. White arrowheads

indicate the new lateral axes that developed from nodal complexes. (b) *C. braunii* rhizoids (rh) growing from a nodal complexes. (c),(d) and (e) White stars mark the internodal cells on both sides of the multicellular nodes. Branches (br), new axes (white arrowheads) and rhizoids (rh) develop from the nodal complexes. Red arrowheads indicate where each rhizoid is attached to the nodal complex. Scale bars: 1 mm (a) and (b), 250 μ m (c), (d) and (e). Branches were removed in (d) and (e) to reveal the site where rhizoids and branches attach to the node. **b.** Percentages of nodes of different developmental stages bearing rhizoids at different developmental stage (Initiating, elongating, multicellular, branched). N_1 are the first nodes below the apical meristem, N_2 are the second, N_3 are the third, N_4 are the fourth and N_{older} are the fifth and older. **c.** Schematic of a *C. braunii* axis showing branches attached to the nodes, rhizoids (in brown), new axes (in blue) and apical meristems (in red).

Figure 8. CbbHLHVIII mRNA is expressed in morphogenetic centres.

a. Histograms showing the mean steady state levels ($n=3$) of CbbHLHVIII mRNA in *C. braunii* thallus (th), rhizoids (rh), gametangia (ga) and zygotes (zy). Transcripts levels were normalized with CbELF5a expression. Each biological replicate is represented by a square, a triangle and a diamond for replicate 1, replicate 2 and replicate 3, respectively. Different letters refer to statistically different groups ($P<0,05$, Kruskal-Wallis test). **b.** Histogram showing the mean steady state level of CbbHLHVIII mRNA in different regions of the *C. braunii* thallus: the apex (apical cells and first nodal complex; ap), nodes 2 (N_2), nodes 3 (N_3), older nodes (N_{older}), branches (br) and branches bearing gametangia (br+ga). Transcript levels were normalized to CbELF5a mRNA levels. Each biological replicate is represented by a square, a triangle and a diamond for replicate 1, replicate 2 and replicate 3, respectively. Different letters refers to statistically different groups ($P<0,05$, Kruskal-Wallis test).

Supplementary Table 1. Expression of Cb_Transcript_119934 (CbbHLHVIII) in the transcriptome.

Libraries	Expression level (TPM)							
	Th1	Th 2	Rh1	Rh2	Rh3	Rh4	WP1	WP2
Cb_Transcript_119934	0.95	2.10	0.04	0.00	0.15	0.01	0.39	1.51

Expression level in Transcript Per Million (TPM) of the CbbHLHVIII transcript (Cb_Transcript_119934) in the different libraries used for the transcriptome (two green thallus (Th) libraries, four rhizoids libraries (Rh) and two whole plants libraries (WP)).

Supplementary Data 1. Transcript, CDS and amino acid sequences of *CbbHLHVIII* and *CnbHLHVIII*.

Supplementary Data 2. Amino acid alignment of the Archaeplastida bHLH transcription factors.

Supplementary Data 3. Trimmed amino acid alignment used for the phylogenetic analysis of the Archaeplastida bHLH transcription factors.

Supplementary Data 4. Trimmed amino acid alignment used for the phylogenetic analysis of the class VIII bHLH transcription factors.

Supplementary Figure S1. Unrooted maximum-likelihood tree of Archeplastida bHLH transcription factors.

CbbHLHVIII and *CnbHLHVIII* in the class VIII bHLH family are highlighted respectively with a red and a blue triangle. The approximate likelihood ratio test (aLRT) support values are included at nodes. The tree includes sequences from the land plants *Arabidopsis thaliana* (At), *Oryza sativa* (Os), *Selaginella moelendorffii* (Sm), *Phycomitrella patens* (Pp) and *Marchantia polymorpha* (Mp); the streptophyte algae *C. braunii*, *C. nitellarum* and *Klebsormidium flaccidum* (Kf), the chlorophyte algae *Volvox carteri* (Vc), *Chlamydomonas reinhardtii* (Cr), *Chlorella variabilis* (Cv), *Ostreococcus tauri* (Ot); the rodophyte alga *Cyanidioschyzon merolae* (Cm).

Supplementary Figure S2. Maximum-likelihood tree of the class VIII bHLH transcription factors.

The class VIII bHLH tree is rooted with the bHLH class XIII and XIV families. It includes sequences from the land plants *Arabidopsis thaliana* (At), *Oryza sativa* (Os), *Selaginella moelendorffii* (Sm), *Phycomitrella patens* (Pp) and *Marchantia polymorpha* (Mp); streptophyte algae *C. braunii*, *C. nitellarum* and *Klebsormidium flaccidum* (Kf); the chlorophyte algae *Volvox carteri* (Vc), *Chlamydomonas reinhardtii* (Cr), *Chlorella variabilis* (Cv) and *Ostreococcus tauri* (Ot); and the rodophyte alga *Cyanidioschyzon merolae* (Cm).

The positions of *CbbHLHVIII* and *CnbHLHVIII* in the class VIII bHLH family are indicated

respectively with a blue and a red triangle. The approximate likelihood ratio test (aLRT) support values are included at nodes.

Supplementary Figure S3. Conserved domains of the bHLH family VIII proteins.

Location and amino acid sequences of the conserved domains in the class VIII bHLH proteins. The conserved domains are framed in black (bHLH domain), purple (HECATE domain), light blue (RSL class 1/VIIIc1), grey blue (RSL class 2/VIIIc2) and navy blue (RSL class 3/VIIIc3). The red star marks the position of the conserved Alanine (A²¹⁰ in AtRHD6).

Supplementary Figure S4. LOGO representation of the conserved amino acid sequences of the bHLH domain of class X, XIII, XIV and XV proteins.

Supplementary Figure S5. Rhizoid development on *C. braunii*.

Rhizoid developmental stages. (a) Initiating rhizoid indicated with a red arrowhead. (b) Elongating rhizoid. (c) Multicellular rhizoid with cell walls indicated with an arrowhead. (d) Branched rhizoid. Scale bars: 250 µm.

Supplementary Figure S6. *C. braunii* gametangia and zygotes.

(a) and (b) *C. braunii* gametangia (archegonia (ar) and anteridia (an)) develop on the branches (br). (c) and (d) Zygote (zy) retained on the branches (br). Scale bars : 500 µm (a) and (d) and 250 µm (b) and (c).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.