

Vitamin D: marker, cause or consequence of depression? An exploration using genomics

Yuri Milaneschi¹, Wouter J Peyrot¹, Michel G Nivard², Hamdi Mbarek², Dorret I Boomsma², Brenda WJH Penninx¹

¹*Department of Psychiatry, Amsterdam Public Health and Amsterdam Neuroscience, Amsterdam University Medical Center/GGZ inGeest, Amsterdam, NL;*

²*Department of Biological Psychology, VU University Amsterdam, Amsterdam, NL*

Address for correspondence:

Yuri Milaneschi, PhD
Department of Psychiatry, Amsterdam UMC / GGZ inGeest
Oldenaller 1
1081 HJ Amsterdam
The Netherlands
Email: y.milaneschi@ggzingeest.nl

Running head: vitamin D and depression

Keywords: vitamin D, major depression, genomics, polygenic risk, etiology, causality, mendelian randomization

Word count: Abstract 250 words, Text 1,200 words, 15 References, 2 Tables, 1 Figure and online supplemental materials

ABSTRACT

Background: Observational studies suggest an association between circulating vitamin D and depression. Trials testing the effect of vitamin D supplementation on depression reported inconclusive findings. It remains unknown whether the vitamin D-depression association stems from shared etiology or from a direct causal relationship. We explored the nature of the association between 25-hydroxyvitamin D (25-OH-D) and major depressive disorder (MDD) exploiting data and statistical tools from genomics.

Methods: Results from the two largest GWAS on 25-OH-D (79,366 samples) and major depressive disorder (MDD; 135,458 cases and 344,901 controls) were applied to individual-level data (>2,000 subjects with measures of genotype, circulating 25-OH-D and DSM-IV lifetime MDD) and summary-level data analyses.

A genetic association between 25-OH-D and MDD was tested by polygenic risk scores (PRS) and by estimating genetic correlation between traits. Two-sample Mendelian Randomization (2SMR) analyses tested the potential bidirectional causality between 25-OH-D and depression.

Results: In individual-level data, the 25-OH-D PRS was associated ($p=1.4e-20$) with 25-OH-D level, but not with lifetime MDD. Conversely, the MDD PRS was associated with MDD ($p=2.3e-5$), but not with 25-OH-D. In summary-level data analyses, the rg between the traits was low and not significant (-0.06 , $p=0.11$). 2SMR analyses provided no evidence of a significant causal role of 25-OH-D for MD and vice versa.

Conclusions: The use of genomics tools indicated that shared etiology or direct causality between vitamin D concentrations and depression is unlikely: vitamin D may represent a marker rather than a cause, or consequence, of depression.

INTRODUCTION

Meta-analyses of large observational studies provided evidence of cross-sectional and longitudinal associations between circulating vitamin D and depression.¹ In >2,300 psychiatrically well-characterized participants from the Netherlands Study of Depression and Anxiety (NESDA), we previously showed² that concentrations of 25-hydroxyvitamin D (25-OH-D, the body reserve of vitamin D) were lower in patients with remitted and current depressive disorders as compared to healthy controls. Furthermore, in patients with current depression low 25-OH-D was associated with symptoms severity and chronicity of depressive disorders after 2 years. It is not trivial to disentangle whether this observational correlation stems from shared underlying aetiological roots or a direct causal relationship between vitamin D and depression. Alternatively, as highlighted by Ioannidis,³ observational associations emerging in nutritional epidemiology may be the product of unresolved confounding. Nevertheless, several hypotheses based on preclinical data emerged about the potential impact of vitamin D on brain structure (vitamin D receptor is expressed in prefrontal cortex, amygdala and hippocampus) and pathophysiological processes relevant for mood.⁴ Meta-analyses⁵⁻⁷ pooling results from trials testing the effect of vitamin D supplementation on depression reported substantially inconclusive findings.

To move forward it is crucial to obtain a clearer knowledge of the exact nature of the vitamin D-depression relationship. Do both traits emerge from a common genetic liability? Does vitamin D truly exert a potential causal effect on depression, or is the effect in the reverse direction?

Genetic research has made enormous progress over the last years, and now provide unique opportunities to investigate shared risk and causality between traits applying new statistical tools and results from genome-wide association studies (GWAS).⁸ In the latest GWAS⁹ on 25-OH-D serum concentrations, the estimate of genetic correlation with depressive symptoms

was not-significant, indicating lack of shared genetic risk. Nevertheless this result may have been affected by the different genetic architectures of depression (highly polygenic with many genetic variants with small effects) and 25-OH-D (few genetic loci with relatively larger effect sizes), or by the use of data for depression from GWAS¹⁰ relatively underpowered as compared to those currently available.

In the present study, we leveraged on results from the two largest GWAS on 25-OH-D (SUNLIGHT consortium: 79,366 samples)⁹ and MDD (Psychiatric Genomics Consortium (PGC): 135,458 cases and 344,901 controls)¹¹. We estimated the degree of genetic overlap between the two traits using both individual-level data (in NESDA²) and summary-level data analyses. Using summary data, Mendelian randomization analyses exploring the potential bidirectional causality between 25-OH-D and depression were also performed.

METHODS

All methods reported hereby are extensively described in eMaterials. Individual-level data analyses were based on up to 2,376 NESDA participants with measures of genotype, circulating 25-OH-D and DSM-IV lifetime diagnoses of MDD (1,750 cases). Polygenic risk scores (PRS) for 25-OH-D and MDD were built based on GWAS summary statistics from, respectively, SUNLIGHT consortium (79,366 samples)⁹ and PGC (135,458 cases and 344,901 controls).¹¹ The PRS for increased 25-OH-D included the six independent genome-wide significant SNPs, jointly explaining an appreciable proportion (38%) of SNP-heritability in the discovery GWAS.⁹ PRS for MDD was built based on the full polygenic signal from the PGC GWAS using LDpred method.¹² Same- and cross-trait associations of PRS with 25-OH-D concentrations and lifetime MDD diagnosis were estimated using regression models adjusted for sex and 10 ancestry-informative genetic principal components. The proportion of variance explained by GPRS was additionally estimated. In additional analyses focusing on MDD cases, the association between the two PRS and symptom severity measured by Inventory of Depressive Symptoms (IDS-SR₃₀)¹³ was also estimated adjusting for sex and principal components.

Finally, summary-level data analyses were run using results from the GWAS of the two traits. Firstly, LD-score regression¹⁴ was applied to GWAS summary statistics to estimate genome-wide genetic correlation between 25-OH-D and MD. Then, the six genome-wide significant SNPs for increased 25-OH-D were used as instrument in two-sample MR (2SMR)¹⁵ testing the potential causal role of vitamin D on MDD using inverse-variance weighted fixed effects-meta-analysis (sensitivity analyses included median- and mode-mode base estimators). Opposite MR analyses testing the causal role of depression for 25-OH-D levels were additionally run using 37 independent genome-wide significant SNPs selected from the discovery GWAS¹¹ as instrument for MDD.

RESULTS

In analyses based on NESDA data, each SD increase in 25-OH-D levels was associated with lower odds of lifetime MDD (sex-adjusted OR=0.77, 95% CIs=0.70-0.87, $p=5.8e-6$). The PRS for increased 25-OH-D was strongly associated (per SD increase: $\beta=5.28$, $p=1.4e-20$; full results in Table 1) with 25-OH-D level, explaining 3.5% of trait variance (figure 1). No association was found with MDD; furthermore, when focusing on 1,750 MDD cases, PRS for 25-OH-D was not associated ($\beta=-0.17$, 95% CIs=-0.80-0.46, $p=0.60$) with symptoms severity measured with IDS-SR₃₀. In contrast, the PRS for MDD was associated with MDD (per SD increase: OR=1.27, $p=2.3e-5$, explaining 1.7% of MDD liability variance, but not with 25-OH-D levels. Within cases, MDD PRS was associated with symptoms severity ($\beta=1.29$, 95% CIs=0.66-1.93, $p=6.5e-5$).

These results were complemented with summary-level data analyses. The estimate of genome-wide genetic correlation between 25-OH-D levels and MD was not significantly different from 0 ($r_g=-0.06$, $se=0.04$, $p=0.11$). Table 2 report the main results from 2SMR analyses; the genetic instrument for 25-OH-D was not causally related to MDD risk (OR=0.96, $p=0.50$). Similarly, no significant evidence of a causal role of MDD on 25-OH-D levels was found ($\beta=-0.02$, $p=0.60$).

DISCUSSION

We examined the nature of the association between vitamin D and depression using data and statistical tools from genomics. Evidence of a common genetic base between the two traits, emerging from shared etiology or through a direct causal mechanism, that could potentially explain their phenotypic association was not found. Even in data from the NESDA cohort, showing a strong phenotypic association between 25-OH-D and MDD,² the underlying polygenic risk for each trait was not associated with the phenotype of the other trait, indicating the lack of genetic overlap. In line with these findings, different MR analyses provided no evidence of a causal effect of 25-OH-D levels on major depression risk, despite the use of a strong instrument for 25-OH-D (explaining ~4% of variance) and input data drawn from large GWAS studies. Also, a reverse causal association from MDD to 25-OH-D was not detected. Based on the present findings, unresolved confounding³ should be considered at this stage the most likely explanation for the association reported by observational studies. The relationship between blood concentrations of 25-OH-D levels may be indeed affected by several factors such health-related lifestyle, habits relates to sun exposure, physical activity, diet and comorbidity. The present findings represent a cautionary tale for further research testing the potential therapeutic effect of vitamin D supplementation on depression, as the expectations of a direct causal effect of vitamin D on mood should be substantially reconsidered.

Nevertheless, hypovitaminosis D in depression remains an important issue. As we previously demonstrated² one third of patients with established psychiatric diagnoses had 25-OH-D levels considered insufficient and risk for musco-skeletal complications. Normalization of vitamin D in depressed patients may therefore be important to prevent related complications and disability which may, in turn, indirectly impact on depression itself in particular during late-life. Nevertheless, applying genomics tools we found that that shared etiology or direct

causality between vitamin D concentrations and depression are unlikely: vitamin D may represent a health marker rather than a cause, or consequence, of depression.

Table 1. Same- and cross-trait associations of polygenic risk scores with circulating 25-hydroxyvitamin D and Major Depressive Disorder in >2,000 participants from NESDA.

PRS	25-OH-D			MDD		
	β	95%CIs	<i>p</i>	OR	95%CIs	<i>p</i>
25(OH)D	5.28	4.18 - 6.39	1.4E-20	0.93	8.29 - 1.04	0.20
MDD	-0.57	-1.70 - 0.55	0.32	1.27	1.14 - 1.43	2.3E-05

Results from linear (outcome: 25-OH-D) and binary logistic (outcome: MDD) regression analyses adjusted for sex and ten ancestry-informative genetic principal components.

Unit of measure 25(OH)D: nmol/l (multiply by 0.4 to obtain ng/ml).

Table 2. Two-samples Mendelian Randomization analyses based on GWAS summary statistics estimating causal effects between circulating 25-hydroxyvitamin D and Major Depressive Disorder

Exposure	Outcome	N SNPs	Inverse Variance Weighted			Weighted Median			Weighted Mode		
			β /OR	95%CI _s	<i>p</i>	β /OR	95%CI _s	<i>p</i>	β /OR	95%CI _s	<i>p</i>
25-OH-D	MD	6^a	0.96	0.90 - 1.10	0.50	0.98	0.89 - 1.12	0.68	0.98	0.88 - 1.09	0.71
MD	25(OH)D	37	-0.02	-0.04 - 0.01	0.25	-0.002	-0.04 - 0.03	0.90	0.01	-0.05 - 0.08	0.65

Unit of measures: 25-OH-D, 1 unit increase in (log)concentrations; MDD, 1 log-unit increase in risk.

SNP effects heterogeneity assessment:

- 25-OH-D causal for MDD, Cochran's Q $p=0.87$; MDD causal for 25-OH-D, Cochran's Q $p=0.18$
- leave-one-out SNP IVW analyses (eFigure1b and eFigure2b) indicated that the estimates were substantially not influenced by individual SNPs

^a eTable2 shows the estimated power for each of the 6 vitamin D-associated SNP to be found significantly ($\alpha=0.05$) associated with MDD in PGC GWAS assuming a true causal link between 25-OH-D and MDD. None of the 6 SNPs, for which adequate statistical power was established, was found to be associated with MDD, additionally confirming that that the phenotypic link between Vitamin-D and MDD is not attributable to a causal effect of Vitamin-D on MDD liability.

Acknowledgments

For NESDA, funding was obtained from the Netherlands Organization for Scientific Research (Geestkracht program grant 10-000-1002); the Center for Medical Systems Biology (CSMB, NWO Genomics), Biobanking and Biomolecular Resources Research Infrastructure (BBMRI-NL), VU University's Institutes for Health and Care Research (EMGO+) and Neuroscience Campus Amsterdam, University Medical Center Groningen, Leiden University Medical Center, National Institutes of Health (NIH, R01D0042157-01A, MH081802, Grand Opportunity grants 1RC2 MH089951 and 1RC2 MH089995). Part of the genotyping and analyses were funded by the Genetic Association Information Network (GAIN) of the Foundation for the National Institutes of Health. Computing was supported by BiG Grid, the Dutch e-Science Grid, which is financially supported by NWO.

We would like to thank the investigators from the Psychiatric Genomics Consortium and the SUNLIGHT Consortium for sharing summary statistics of their GWAS.

In order to perform the present analyses we received summary statistics of the GWAS by Hyde et al. (Nat Genet. 2016; 48:1031-6) from 23andMe. We would like therefore to thank research participants and employees of 23andMe for making this work possible.

References

1. Anglin RES, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *Br J Psychiatry*. 2013;202(2):100-107. doi:10.1192/bjp.bp.111.106666.
2. Milaneschi Y, Hoogendijk W, Lips P, et al. The association between low vitamin D and depressive disorders. *Mol Psychiatry*. 2014;1936:444-451. doi:10.1038/mp.2013.36.
3. Ioannidis JPA. The Challenge of Reforming Nutritional Epidemiologic Research. *JAMA - J Am Med Assoc*. 2018. doi:doi:10.1001/jama.2018.11025.
4. McCann J, Ames B. Is there convincing biological or behavioral evidence linking vitamin D deficiency to brain dysfunction? *FASEB J*. 2008;22(4):982-1001.
5. Gowda U, Mutowo MP, Smith BJ, Wluka AE, Renzaho AMN. Vitamin D supplementation to reduce depression in adults: Meta-analysis of randomized controlled trials. *Nutrition*. 2015;31(3):421-429. doi:10.1016/j.nut.2014.06.017.
6. Shaffer JA, Edmondson D, Wasson LT, et al. Vitamin d supplementation for depressive symptoms: A systematic review and meta-analysis of randomized controlled trials. *Psychosom Med*. 2014;76(3):190-196. doi:10.1097/PSY.000000000000044.
7. Li G, Mbuagbaw L, Samaan Z, et al. Efficacy of Vitamin D Supplementation in Depression in Adults: A Systematic Review. *J Clin Endocrinol Metab*. 2014;99(3):757-767. doi:10.1210/jc.2013-3450.
8. Maier RM, Visscher PM, Robinson MR, Wray NR. Embracing polygenicity: a review of methods and tools for psychiatric genetics research. *Psychol Med*. 2017:1-19. doi:10.1017/S0033291717002318.
9. Jiang X, O'Reilly PF, Aschard H, et al. Genome-wide association study in 79,366

- European-ancestry individuals informs the genetic architecture of 25-hydroxyvitamin D levels. *Nat Commun.* 2018;9(1):260. doi:10.1038/s41467-017-02662-2.
10. Okbay A, Baselmans BML, De Neve J-E, et al. Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nat Genet.* 2016;48(6):1-13. doi:10.1038/ng.3552.
 11. Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depressive disorder. *Nat Genet.* 2018. doi:<https://doi.org/10.1101/167577>.
 12. Vilhjálmsón BJ, Yang J, Finucane HK, et al. Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. *Am J Hum Genet.* 2015;97(4):576-592. doi:10.1016/j.ajhg.2015.09.001.
 13. Rush A, Gullion C, Basco M, Jarrett R, Trivedi M. The Inventory of Depressive Symptomatology (IDS): psychometric properties. *Psychol Med.* 1996;26(3):477-486.
 14. Bulik-Sullivan BK, Loh P-R, Finucane HK, et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet.* 2015;47(3):291-295. doi:10.1038/ng.3211.
 15. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife.* 2018;7:e34408. doi:10.7554/eLife.34408.

Figure legend

Figure 1. Proportion of trait variance explained by polygenic risk scores for increased 25-hydroxyvitamin D and MDD in > 2,000 participants from NESDA.

See eMaterials section 2.3 for estimation of variance explained by PRS.

P-values from Table 1

25-OH-D
N=2,376

MDD
1,771 cases + 370 controls





Figure 1