

1 The bat influenza H17N10 can be neutralized by broadly-
2 neutralizing monoclonal antibodies and its neuraminidase can
3 facilitate viral egress.

4 George Carnell^{a,b,1}, Efsthios S Giotis^{c,1}, Keith Grehan^{a,d,1}, Francesca Ferrara^a,
5 Stuart Mather^a, Eleonora Molesti^a, Simon Scott^a, Antonello Pessi^e, Krzysztof Lacek^{e,f},
6 & Nigel Temperton^a

7 ^a Viral Pseudotype Unit, Medway School of Pharmacy, University of Kent and University of Greenwich, Chatham, United Kingdom

8 ^b Laboratory of Viral Zoonotics, Department of Veterinary Medicine, University of Cambridge, United Kingdom

9 ^c Section of Virology, Faculty of Medicine, Imperial College London, St Mary's Campus, United Kingdom

10 ^d School of Molecular and Cellular Biology, University of Leeds, United Kingdom

11 ^e Ceinge Biotechnologie Avanzate S. C.R.L., Napoli, Italy

12 ^f The Laboratory of Virus Molecular Biology, University of Gdansk, 80-822 Gdansk, Poland

13 ¹ Authors contributed equally

14
15 Corresponding author: Nigel Temperton, n.temperton@kent.ac.uk, ORCID 0000-0002-7978-3815

16

17 Key words: Bat Influenza virus, broadly neutralizing antibody, Hemagglutinin, Neuraminidase

18 **Abstract**

19 The diversity of subtypes within the Influenza A virus genus has recently expanded with the
20 identification of H17N10 and H18N11 from bats. In order to further study the tropism and zoonotic
21 potential of these viruses, we have successfully produced lentiviral pseudotypes bearing both
22 haemagglutinin H17 and neuraminidase N10. These pseudotypes were shown to be efficiently
23 neutralized by the broadly-neutralizing monoclonal antibodies CR9114 and FI6. Our studies also
24 confirm previous reports that H17 does not use sialic acid as its cellular receptor, as pseudotypes
25 bearing the H17 envelope glycoprotein are released into the cell supernatant in the absence of NA.
26 However, we demonstrate that N10 facilitates heterosubtypic (H5 and H7) influenza HA-bearing
27 pseudotype release in the absence of another source of NA, significantly increasing luciferase
28 pseudotype production titres. Despite this, N10 shows no activity in the enzyme-linked lectin assay
29 used for traditional sialidases. These findings suggest that this protein plays an important role in viral
30 egress, but is perhaps involved in further accessory roles in the bat influenza lifecycle that are yet to
31 be discovered. Thus we show the lentiviral pseudotype system is a useful research tool, and
32 amenable for investigation of bat influenza tropism, restriction and sero-epidemiology, without the
33 constraints or safety issues with producing a replication-competent virus, to which the human
34 population is naïve.

35 Significance statement

36 Influenza virus is responsible for mortality and morbidity across the globe; human populations are
37 constantly at risk of newly emerging strains from the aquatic bird reservoir which harbors most of
38 the subtypes of influenza A (H1-H16). Recently identified subtypes (H17N10, H18N11) from bats
39 have broadened the reservoir from which potential pandemic strains of influenza can emerge. To
40 evaluate the potential for these novel subtypes to cross over into human populations, their ability to
41 establish an infection, in addition to the extent of cross-reactive immunity established by human
42 seasonal strains needs to be investigated. This study highlights a novel platform for the study of the
43 bat H17 and N10 envelope glycoproteins, using a lentiviral pseudotype system. Following the
44 generation of this pseudotype it was employed in cell entry and microneutralization assays. These
45 showed that two well-characterised monoclonal antibodies (mAb) which target avian and human
46 influenza subtypes will also neutralize H17. Furthermore the data presented in this study show a
47 novel aspect of the N10 glycoprotein in its ability to facilitate the budding of pseudotypes bearing
48 different influenza HAs.

49 Introduction

50 Influenza A virus is the principal causative agent of influenza, which is a substantial burden to global
51 economies and represents a significant public health risk worldwide (Ma et al. 2009; Reperant et al.
52 2012). The first stages of the viral life cycle are mediated by the influenza glycoprotein
53 hemagglutinin (HA), which is located in the lipid outer membrane of the virus. This protein mediates
54 both the attachment of the influenza virus to the target cell and the fusion process which allows the
55 virus to infect its host cell. Through its activity, HA is a key determinant of virus tropism (Dos Reis et
56 al. 2011; Sahini et al. 2010). Furthermore, HA is a highly variable protein and its features are often
57 used as methods for distinguishing different viral strains (Ebrahimi et al. 2014; Wilson and Cox
58 1990).

59 While the association between influenza viruses and wild birds has long been established (Gelfond
60 et al. 2009; Webster et al. 1992), the discovery of novel influenza-like viruses in New World bats
61 (H17N10, H18N11) represents a possible challenge to the notion that avian species are its sole
62 reservoir (Freidl et al. 2015; Sun et al. 2013; Tong et al. 2012, 2013). Recently, Egyptian *Rousettus*
63 *aegyptiacus* bats have been reported to harbor a third bat influenza subtype (Kandeil et al. 2018). It
64 is highly likely that further subtypes will be discovered in future years, warranting further research
65 on existing strains of bat influenza.

66 The H17N10 and H18N11 subtypes of influenza A were discovered in bat species in Peru and
67 Guatemala respectively. They are significantly diverged from other known influenza strains, in
68 respect to all eight gene segments (Tong et al. 2012, 2013). The HA of these novel strains contain a
69 number of apparently unique structural features and exhibit receptor-binding activities that differ
70 from other influenza viruses (Sun et al. 2013). Notably, it has been shown that H17 and H18
71 hemagglutinins do not bind to sialic acid receptors, and the true receptors are currently still
72 unknown (Tong et al. 2013).

73 The neuraminidase (NA) of these viruses is also divergent from other known NAs, but the overall
74 structure is preserved, despite differences in the active site (Juozapaitis et al. 2014; Ma et al. 2015).
75 Lack of sialidase activity has been reported previously for these NAs (García-Sastre 2012; Li et al.
76 2012; Tong et al. 2013; Zhu et al. 2012), suggesting they utilize a different substrate altogether. The
77 combination of atypical HA binding profiles alongside a lack of NA activity suggests that this virus

78 functions differently to previously discovered influenza A subtypes, despite relative phylogenetic
79 relatedness.

80 Although the receptor remains unknown, H17 and H18 sequences have been extensively analyzed
81 and compared to those of other HA gene sequences, to investigate the potential for these bat
82 viruses becoming zoonotic (Freidl et al. 2015; Mänz et al. 2013).

83 At present, the degree to which the H17N10 and H18N11 isolates are capable of infecting non-bat
84 hosts is unknown, as attempts to isolate wildtype virus have not been successful. Sequence analysis
85 indicates that there is significant potential for a spillover occurrence, but further research is required
86 to assess the true potential of these viruses as pandemic threats (Freidl et al. 2015; Juozapaitis et al.
87 2014).

88 To study the zoonotic potential of H17N10 and H18N11 viruses, attempts to integrate bat HAs into
89 influenza A reverse genetics systems were made but none yielded infectious virus (Juozapaitis et al.
90 2014; Zhou et al. 2014). More recently, progress has been achieved by Moreira et al. 2016 to this
91 end, using a pseudotyped VSV platform.

92 Influenza pseudotyped viruses (PVs) are useful tools to study both viral entry mechanisms and the
93 antibody response directed against the influenza HA and NA. Furthermore, when use of PVs is
94 coupled with detailed sequence analysis and phylogenetic inference, they offer the potential to
95 establish safe and effective assays to inform epidemiological and public health models of viral spread
96 and risk. This is especially the case with later generation, single cycle lentiviral vectors, allowing
97 experimentation on functional glycoproteins with the use of a reporter incorporated in the lentivirus
98 genome.

99 In this study, we report generation of the first H17 and H17N10 pseudotyped lentiviruses, their use
100 in virus neutralization assays using broadly neutralizing monoclonal antibodies (bnmAbs) and
101 investigation into the debated substrate specificity of the putative N10 neuraminidase.

102 Results

103 Generation of H17- and N10- pseudotype viruses

104 Generation of H17- (A/little yellow shouldered bat/060/2010) and H5- (A/Vietnam/1194/2004)
105 lentiviral PVs was achieved after transfection of HEK293T/17 cells with lentiviral pseudotype
106 production plasmids (Figure 1), followed by transduction of target human U87 MG (glioblastoma)
107 cells with PV supernatant. Generation of H17- PVs was only possible in the presence of HAT or
108 TMPRSS2 proteases in HEK293T/17 cells (Figure 2), as has been seen with other subtypes of
109 influenza (Bertram et al. 2010; Böttcher et al. 2006). Subsequent experiments were carried out using
110 canine MDCK II and RIE1495 target cells which were previously reported to allow production of VSV-
111 H17 (Moreira et al. 2016). In initial experiments, attempts were made to generate functional H17
112 and H17N10 PV utilizing different protease-expressing plasmids to promote HA maturation. PV
113 luciferase-based titres increased significantly when HAT-, TMPRSS2- and TMPRSS4- expression
114 plasmids were used in the transfection mix when compared to controls, indicating activation of the
115 bat HA and fusion competence. Other proteases tested (furin, KLK5, and TMPRSS3) did not yield
116 significant titre increases. MDCK II and RIE1495 cells were notably transduced, whereas HEK293T/17
117 cells were not. In the absence of co-transfected, protease-encoding plasmids, no significant PV titre
118 was measurable (Figure 3).

119 **H17 entry of target cells is inhibited by broadly neutralizing monoclonal antibodies**

120 Treatment of RIE1495 cells with the acidifying agent ammonium chloride resulted in significantly
121 lower levels of luciferase activity, demonstrating that H17 requires a low pH for membrane fusion
122 similar to conventional influenza viruses, such as H5 (Figure 4). Several studies have demonstrated
123 that the HA trimers of conventional influenza viruses must simultaneously coordinate their
124 conformational changes to complete membrane fusion (Otterstrom et al. 2014). HA-binding broadly
125 neutralising antibodies (bnmAbs) typically recognize loop regions surrounding the receptor binding
126 site or conserved regions of the stem and their induced-inhibition is serotype-specific (Dreyfus et al.
127 2013; Ekiert et al. 2009; Sui et al. 2009). In order to test whether cross reactive antibody responses
128 would affect the H17 glycoprotein, several characterised bnmAbs were employed. Neutralization
129 potency was measured as IC₅₀ (concentration or serum dilution required to neutralise 50% of input
130 virus). Neutralization tests were conducted using both the permissible target cell lines identified
131 above (MDCK II and RIE1495). The first bnmAb, CR9114, binds to a conserved epitope in the HA stalk
132 of group 1 and 2 influenza A viruses, and has also been shown to protect against lethal influenza
133 challenge in a mouse model against both lineages of influenza B (Dreyfus et al. 2012). The second,
134 CR6261, has been shown to neutralize H1 and H5 subtypes (Friesen et al. 2010). The third, F16-nt,
135 can neutralize H1 to H16 subtypes of influenza A. A fourth, F16-chol is a derivative of F16-nt,
136 conjugated to cholesterol (Corti et al. 2011; Lacey et al. 2014). H17 is effectively neutralized by
137 CR9114, at a concentration of 0.05 µg/ml. F16-nt and F16-chol also neutralize H17N10- PV, however,
138 concentrations required exceed those of CR9114. In contrast, CR6261 was not able to neutralize
139 H17 using the H17 PV assay (data not shown). Neutralization data is shown in Figure 5.

140 **N10 facilitates production of conventional H5 and H7 bearing PV.**

141 Pretreatment of target cells with sialidase, which removes cell surface sialic acids, did not affect the
142 entry of H17 and H17N10 PVs, which supports previous studies that H17 does not bind sialic acids
143 for infection (Maruyama et al. 2016), see Figure 6. N10-bearing PVs were titrated alongside H17 and
144 H17N10 PV, in order to measure their ability to transduce cells. Results showed that N10 had no
145 effect on H17 mediated viral entry and did not improve transduction when co-expressed with H17
146 (data not shown). However, addition of N10 in the generation of H5 or H7 PVs (H5N10, H7N10)
147 produced high titre PV in the absence of any other neuraminidase source (Figure 7), indicating that
148 N10 is facilitating release of viral particles bearing sialic acid-binding glycoproteins. H5N10 luciferase
149 titres were one log lower than parental H5 PV produced with exogenous neuraminidase, but two
150 logs higher than N10, ΔEnvelope (ΔEnv) or cell only controls (Figure 7). Despite this, the same
151 H5N10 PV preparation did not show any detectable NA activity when titrated using the enzyme
152 linked lectin assay (ELLA), confirming previous reports that this protein is not a sialidase (García-
153 Sastre 2012; Li et al. 2012; Tong et al. 2013; Zhu et al. 2012). Similarly, multiple different
154 preparations of H17N10 or of N10 bearing PV produced negative results in ELLA (data not shown).

155 **Discussion**

156 We have successfully produced H17 and H17N10 bearing lentiviral pseudotypes, and shown that
157 N10 is not required for H17 pseudotype budding, but that it can mediate release of a heterologous,
158 sialic acid-binding HA bearing PVs. These bat influenza PVs were neutralized by cross-reactive
159 bnmAbs, suggesting that the stalk region of the H17 glycoprotein retains conserved epitopes present
160 in group 1 and 2 influenza HAs (Sun et al. 2013).

161 Of the target cell lines tested, H17 and H17N10 lentiviral PV were able to transduce U87 MG, MDCK
162 II and RIE1495 cells to varying degrees when activated by proteolytic cleavage. Reports differ on the
163 permissibility of the MDCK I type cells (Hoffmann et al. 2016; Moreira et al. 2016). Due to the nature
164 of the pseudotype based microneutralisation (pMN) assay used in this study, the addition of cells in
165 suspension into PV containing supernatant may allow infection/transduction to occur before cell

166 adhesion and thus by a different route than *in vivo*. This may explain our results in the context of
167 Moriera and colleagues' findings that the bat influenza H18 VSV pseudotypes enter at the
168 basolateral membrane and inefficiently infected confluent cell monolayers (Moreira et al. 2016).
169 MDCK I and II differ in passage number (low and high, respectively), differences between them
170 include the mucin-type transmembrane protein podoplanin, which is expressed only in MDCK I, and
171 the Forssman glycosphingolipid, which is expressed only in MDCK II cells (Hansson et al. 1986;
172 Zimmer et al. 1997). The susceptibility of MDCK cells to bat influenza viruses is unclear, and is partly
173 compounded by the availability of various lineages with different characteristics (Dukes et al. 2011).
174 The widespread use of MDCK cells in influenza research is at odds with a previous study that showed
175 H17 failed to bind to MDCK cells (Sun et al. 2013). While the reasons for this are currently unclear, it
176 has been hypothesized that the level of HA binding is below the threshold of detection for the assays
177 used in this earlier study (Maruyama et al. 2016). Alternatively, only a subset of MDCK lineages may
178 be susceptible to these viruses (Table 2). RIE1495 cells are morphologically similar to MDCK II cells
179 and express the Forssman antigen, a possible factor in their susceptibility to H17 bearing PVs
180 (Moreira et al. 2016). The cell tropism data reported in the Moreira study raise some interesting
181 ideas concerning the H17 receptor, as only three out of eight bat cell lines were transduced by the
182 H17-VSV pseudotypes, originating from two *Miniopterus* and one *Pteropus* species. This suggests
183 that these viruses may be restricted to a particular set of closely related species belonging to the
184 *Miniopteridae* and *Pteropidae* families, which are closely related to the *Phyllostomidae* family, from
185 which the original H17 and H18 samples were isolated (Agnarsson et al. 2011; Tong et al. 2012).
186 However, this raises further questions, as the cell lines derived from *Pteropus dasymallus*
187 *yayeyamae* and *Rousettus leschenaultii* were not permissive, indicating a complex pattern of
188 susceptibility of species to these new influenza viruses. Nevertheless, as these cell lines have only
189 recently been isolated and immortalised (Maeda et al. 2008; Maruyama et al. 2014), detailed
190 characterisation of their surface proteins and expressed proteases are not yet available, requiring
191 additional research before conclusions can be drawn. Further investigation is required to determine
192 the mechanisms involved for the transduction of canine cells but not HEK293T/17 cells, particularly
193 relating to the expression of putative receptors on permissive cell lines.

194 Our results, contrary to previous studies, indicate that N10 is performing a similar function to other
195 NAs in enabling release of new influenza virus particles, which indicate its ability to form VLPs rather
196 than releasing H17 from its cellular receptor (Yondola et al. 2011). PVs bearing H5/H7 and N10
197 envelope glycoproteins successfully budded from producer cells into the surrounding medium in the
198 absence of a sialic acid cleaving neuraminidase, resulting in significantly increased titres when
199 compared to the same glycoproteins generated without the addition of N10 (Figure 7). This increase
200 in budding PV may be due to the action of N10 on an unknown substrate, or perhaps action of the
201 protein itself in virus morphogenesis and budding (Barman et al. 2004; Yondola et al. 2011).
202 Sialidase activity was however not detected using the enzyme-linked lectin assay, suggesting either a
203 lack of sensitivity of this assay or another mechanism for the removal of surface sialic acids by N10
204 (Juozapaitis et al. 2014; Sun et al. 2013; Wu et al. 2014). Similarly, PVs bearing solely the N10
205 glycoprotein did not show sialidase activity via ELLA, or successfully transduce cells to give a
206 significant luciferase reading. Further investigation is required on the role of the bat influenza
207 neuraminidase in its lifecycle, the combination of the findings described in this article, with the fact
208 that the N10 enzymatic structure remains conserved and NA-like, suggests that it is involved in
209 accessory functions other than simply aiding in viral egress. Our results highlight the distinct
210 difference between bat and traditional influenza A viruses where a delicate balance is in place
211 between HA and NA. It may be the case that such a balance exists between bat HA and NAs which
212 will be made clear once the substrate(s) of the bat NA is discovered.

213 In a previous study, it was demonstrated that the TMPRSS2 protease was capable of inducing HA
214 maturation of H17 through cleavage from HA0 to HA1 and HA2 (Hoffmann et al. 2016). In our study,
215 we demonstrated that this maturation can also be facilitated by TMPRSS4 and HAT. This is of
216 particular interest as the expression of specific proteases is a known limiting factor in viral tropism
217 for a number of different viruses (Böttcher-Friebertshäuser et al. 2010; Ferrara et al. 2012; Millet
218 and Whittaker 2015). This, coupled with the observed susceptibility of a canine-derived cell line to
219 the H17 PV, will need to be factored into future analysis of the potential for zoonotic spillover from
220 bat origin influenza viruses.

221 The fact that bnmAbs were able to neutralize the virus particles via the H17 hemagglutinin, and their
222 requirement for proteolytic activation, reinforces that we have only scratched the surface in terms
223 of their characterization. Further exploration is required to establish whether these viruses are
224 endogenous or capable of forming infectious particles *in vivo*. Research has revealed that all eight
225 segments of the genome encode functional proteins (Juozapaitis et al. 2014; Moreira et al. 2016; Wu
226 et al. 2014; Zhu et al. 2012), despite the inability to isolate wildtype virus to date. Nevertheless, the
227 development of tools, such as those described here, which can be used in H17 receptor
228 identification studies may ultimately aid discovery of wildtype virus samples from the bat reservoir.

229 **Methods**

230 **Plasmids:**

231 The H17 HA and N10 NA genes from A/little yellow shouldered bat/Guatamala/060/2010 were
232 synthesised commercially by Genscript (Genscript, USA) and subcloned into vector pl.18 (Cox et al.
233 2002). Lentiviral packaging plasmids p8.91 (Zufferey et al. 1997) and pCSFLW (Demaison et al. 2002)
234 containing a firefly luciferase reporter were used to produce PV. Protease encoding plasmids
235 pHCMV-Tag3 (TMPRSS4-myc) and pcDNA3.1-hTMPRSS3 were kindly provided by Prof. Stefan
236 Pöhlmann, Infection Biology Unit, German Primate Center, Germany. pCAGGS-TMPRSS2 and
237 pCAGGS-HAT were kindly provided by Eva Böttcher-Friebertshäuser, Philipps University of Marburg,
238 Germany. The plasmid bearing the Vesicular stomatitis virus envelope protein (VSV-G), pMD.G was
239 obtained from Dr Yasu Takeuchi, University College London, United Kingdom. All the plasmids that
240 were used in this study are listed in Table 3.

241 **Cell lines:**

242 Multiple cell lines were used for titration of PV containing supernatants. HEK293T/17 cells were
243 kindly provided by Dr Edward Wright (University of Westminster, UK). MDCK II and RIE1495 were
244 kindly provided by Dr. Gert Zimmer (Institute of Virology and Immunology, Switzerland). U-87 cells
245 were provided by Dr Simon Scott (University of Kent, UK). Madin-Darby Canine Kidney (MDCK) cells
246 were kindly provided by Prof. Sarah Gilbert (Jenner Institute, University of Oxford, UK). All cells were
247 cultured in Dulbecco's modified eagle medium supplemented with 10% fetal bovine serum.

248 **Antibodies:**

249 mAbs CR9114 and CR6261 were produced by Crucell (Janssen Vaccines AG, Bern, Switzerland). FI6-nt
250 and cholesterol conjugated FI6 (FI6-Chol) were produced by Alfredo Pesci and Krzysztof Lacey from
251 sequence information derived from Corti et al. 2011.

252 **Production and quantification of H17N10 and H17 bearing lentiviral pseudotypes:**

253 PV were produced as described previously (Ferrara et al. 2012; Temperton et al. 2007) and as shown
254 in Figure 1. Briefly, transfection of HEK293T/17 cells was performed using a variety of combinations

255 of plasmids pl.18-H17, pl.18-N10, p8.91 and pCSFLW using polyethylenimine transfection reagent
256 (Sigma Aldrich, UK). Protease-encoding expression plasmids were also included (Table 3). Medium
257 was replaced 12h post-transfection. Supernatants were harvested 48h post-transfection and passed
258 through a 0.45µm filter (Millipore, UK). PV-containing supernatants were titrated using the firefly
259 luciferase Bright-Glo™ system (Promega, UK). Serial dilution of 100µl of PV-containing supernatant
260 was performed across a white flat bottom 96-well Nunclon® plate (Thermo Fisher Scientific, UK).
261 Subsequently, approximately 1×10^4 cells per well were added per well in 50µl of medium, plates
262 were incubated in a humidified incubator at 37°C 5% CO₂ for 48h, after which 50µl of Bright-Glo™
263 substrate was added and luciferase reading recorded in relative luminescence units (RLU) after a 5
264 minute incubation period. Further sets of N10-bearing PV were produced by transfection of
265 HEK293T/17 cells with 500ng p8.91, 750ng pCSFLW and various amounts of pl.18-N10 plasmid.
266 Medium was replenished after 12h; PV containing supernatants were collected 72h later and passed
267 through a 0.45µm filter. Transfections were carried out in 6-well Nunclon® plates (Thermo Fisher
268 Scientific, UK).

269 GFP-expressing pseudotypes were produced by substituting the pCSFLW firefly luciferase lentiviral
270 vector mentioned previously with the GFP expressing vector pCSGW. PV-containing supernatants
271 were titrated down clear 96-well Nunclon® plates (Thermo Fisher Scientific, UK) with the addition of
272 1×10^4 cells per well. Plates were incubated for 72h and visualized by Nikon Eclipse 50i
273 epifluorescence microscope with a charge-coupled QICAM Fast 1394 camera (QImaging) at x200
274 magnification.

275 **Screening of cell lines**

276 Two-fold serial dilutions of PV-containing supernatant were performed as previously described using
277 white 96-well Nunclon® plates (Thermo Fisher Scientific, UK). Subsequently, approximately 1×10^4 of
278 each cell line was added in 50µl of medium per well. Plates were incubated in a humidified incubator
279 at 37°C and 5% CO₂ for 48h, after which 50µl of Bright-Glo™ substrate was added and luciferase
280 reading recorded in relative luminescence units (RLU), following a 5 minute incubation period.
281 Control wells were used to measure cell populations for each cell line.

282 **Pseudotype based microneutralization assay (pMN) using H17N10 and H17 bearing lentiviral 283 pseudotypes:**

284 Monoclonal antibodies (mAbs) were serially diluted 1:2 across white 96-well Nunclon® plates
285 (Thermo Fisher Scientific, UK) in 50 µl of DMEM. PV-containing supernatants were diluted and added
286 to each well to give an approximate RLU value of 1×10^6 per well in 50 µl of DMEM. After 1h
287 incubation at 37°C and 5% CO₂, approximately 1×10^4 cells were added per well in a volume of 50µl.
288 Plates were incubated for 48h at 37°C and 5% CO₂, then 50 µl of Bright-Glo™ was added and
289 luminescence read after a 5 minute incubation at room temperature. Results were analyzed with
290 Graphpad (Prism 7), using nonlinear regression on luminescence values normalized to cell only and
291 virus only control thresholds (100% and 0% neutralization equivalent, respectively). IC₅₀ values
292 represent the concentration (ng/ml) required for each antibody to neutralize 50% of functional
293 pseudotyped virus, based on luciferase activity.

294 **Cell treatment**

295 Overnight treatment of RIE1495 cells with the endosomal acidification inhibitor
296 ammonium chloride (SIGMA) and pretreatment for 2h with neuraminidase (Roche) was done as
297 previously described (Maruyama et.al, 2016). Treated cells were washed three times with serum
298 free medium, and then incubated for 24h with H17 PV. PV titration and luciferase activity was
299 monitored with a luminometer as described previously.

300 Pseudotype based enzyme linked lectin assay (pELLA) utilizing N10:

301 ELLA was performed as described by Couzens et al. 2014, but adapted in order to allow use of
302 lentiviral pseudotypes as a source of NA (Biuso et al. 2018; Prevato et al. 2015). Briefly, PV
303 containing supernatant was serially diluted (1:2) across a standard clear microtitre plate in sample
304 diluent (PBS, 1% BSA, 0.5% Tween 20). 50 μ l of the resulting dilution series was transferred in
305 duplicate to Nunclon[®] Maxisorp 96-well plates (Thermo Fisher Scientific, UK) previously coated with
306 Fetuin (Sigma Aldrich, UK) and containing 50 μ l of sample diluent per well. Plates were then
307 incubated for 18h and washed, prior to addition of conjugate diluent (PBS, 1% BSA) containing a
308 1:500 concentration of peanut lectin conjugated to horseradish peroxidase (PNA-HRPO, Sigma
309 Aldrich, UK). Plates were then incubated for 2h in the dark, whereupon OPD-based substrate in
310 citrate buffer was added (Sigma Aldrich, UK). Reactions were stopped after 10 minutes using 50 μ l
311 1M H₂SO₄, and readings recorded using a standard ELISA plate reader at 492nm. Exogenous
312 neuraminidase from *Clostridium perfringens* (Sigma Aldrich, UK) was used as a positive benchmark
313 control and samples were assayed alongside PV bearing neuraminidases from other influenza
314 subtypes.

315 Statistical analysis

316 Where possible, statistical analysis was carried out to determine whether differences in PV titre
317 were significant. One-way ANOVA t-tests were performed using fold change scores with a Tukey's
318 multiple comparisons test. P-values were set at 0.05 ($P \leq 0.05$) unless indicated otherwise.
319 Significantly different data are denoted with asteriks representing $p = <0.01$ (*), <0.001 (**), <0.0001
320 (***) and <0.00001 (****).


321

322

323

324 Figures and Tables


325



326


327 Figure 1. Cartoon showing the production of H17N10 PV via plasmid transfection.

328



329

330 Figure 2. Transduction of U87 MG cells by influenza GFP PV (x100 magnification). A) H17 produced
331 using pCAGGS-HAT. B) H17 PV produced using pCAGGS-TMPRSS2. C) Cell only control. D) H5 (A/Viet
332 nam/1194/2004) PV positive control.



333

334 Figure 3. Transduction of various target cell lines. A) HEK293T/17, B) RIE1495 and C) MDCK II with
335 H17 and H17N10 pseudotyped viruses carrying the luciferase reporter gene. Results given in Relative
336 Luminescence Units per ml; RLU/ml). Average cell only luminescence shown as a red line.


337

338

339

340

341




342

343

344 Figure 4. The effect of ammonium chloride based acidification on the transduction of RIE1495 cells
345 by H17 PV. Increases in acidification of RIE1495 cell cultures prior to transduction by H17 resulted in
346 lower luciferase based titres, indicating that the pH sensitivity of H17 is similar to that of H5.
347 Significant differences in transduction denoted by asteriks, ** represents $p = <0.001$ and *** $p =$
348 <0.0001 respectively.

349



350

351


352 Figure 5. Neutralization curves and IC₅₀ values for mAbs CR9114, F16-nt and F16-Chol against
353 H17N10 bearing lentiviral pseudotypes on cell lines RIE1495 and MDCK II. Nonlinear regression
354 carried out using Graphpad (Prism 7) in order to provide IC₅₀ values for each graph. Each IC₅₀ is the
355 concentration of mAb required for reduction of 50% of the virus input in terms of luciferase activity.

356

357

358

359




360

361

362 Figure 6. The effect of neuraminidase pre-treatment on the transduction of RIE1495 cells by H17 and
363 H5 PV. No relationship is seen between pre-treatment of cells with neuraminidase for H17 bearing
364 PV, but H5-based transduction is significantly reduced when sialic acids are stripped from target
365 cells. Significant difference in transduction denoted by asteriks, * represents $p = <0.01$ and ** $p =$
366 <0.001 respectively.

367

368



369

370

371

372 Figure 7. Release of H5 and H7 bearing PV by N10. PV bearing the glycoproteins H5
373 (A/Vietnam/1194/2004) and H7 (A/Shanghai/1/2013) were produced with exogenous
374 neuraminidase (+NA), with no NA (-NA) or accompanied with an N10 glycoprotein (+N10). H5 and
375 H7 PVs produced in the presence of N10 have significantly higher luciferase titres than those
376 produced with no NA. Background luciferase titre control (Δ Env) control shown. Significant
377 differences in transduction denoted by asteriks, **** represents $p < 0.00001$.

378

Table 1. Cell lines used in the production of Bat influenza VSV pseudotypes.

Cell Line	Species	Tissue:
<i>Vero E6</i>	<i>Chlorocebus sp.</i>	Kidney
<i>HEK293</i>	<i>Homo sapiens</i>	Kidney
MDCK	<i>Canis lupus familiaris</i>	Kidney
<i>SK-L</i>	<i>Sus scrofa</i>	Kidney
<i>QT6</i>	<i>Coturnix japonica</i>	Muscle
Cell Line:	Bat species:	Tissue:
<i>BKT1</i>	<i>Rhinolophus ferrumequinum</i>	Kidney
<i>FBKT1</i>	<i>Pteropus dasymallus yayeyamae</i>	Kidney
<i>YubFKT1</i>	<i>Miniopterus fuliginosus</i>	Kidney
<i>IndFSPT1</i>	<i>Pteropus giganteus</i>	Spleen
<i>DemKT1</i>	<i>Rousettus leschenaultii</i>	Kidney
<i>ZFBK11-97</i>	<i>Epomophorus gambianus</i>	Kidney
<i>SuBK12-08</i>	<i>Miniopterus schreibersii</i>	Kidney
<i>ZFBS13-75A</i>	<i>Eidolon helvum</i>	Spleen

List of cell lines used in VSV based pseudotyping of H17 and H18 viruses. Bold entries are those cell lines which were found to be permissive for viral transduction by Maruyama et al. 2016.

379

380

Table 2: MDCK cell lines.

NBL-2 (ATCC® cat # CCL-34™)
MDCK I (EEACC cat # 00062106)
MDCK II (EEACC cat# 00062107)
MDCK.1 (ATCC® cat# CRL-2935™)
MDCK.2 (ATCC® cat# CRL- CRL-2936™)
super dome (ATCC® cat# CRL-2286™)
super tube (ATCC® cat# CRL-2285™)
Different MDCK cell lines available commercially

381

382

383

384

385

386

387

388

389 Table 3: Genes, plasmids and sources.

Gene	Plasmid	Source
Transmembrane protease, serine 2	pCAGGS-TMPRSS2	(Böttcher et al. 2006)
Transmembrane protease, serine 3	pcDNA3.1-hTMPRSS3	(Bertram et al. 2010)
Transmembrane protease, serine 4	phCMV-Tag3 (TMPRSS4-myc)	(Jung et al. 2008)
Human Airway Trypsin	pCAGGS-HAT	(Böttcher et al. 2006)
Kallikrein-related peptidase 5	pl.18-KLK5	Synthesised, Sigma
Furin	pFurin	Kind gift from H. D. Klenk
H17 (A/little yellow shouldered bat/Guatamala/060/2010)	pl.18-H17	Synthesised, Genscript
N10 (A/little yellow shouldered bat/Guatamala/060/2010)	pl.18-N10	Synthesised, Genscript
H5 (A/Viet Nam/1194/2004)	pl.18-H5	(Temperton et al. 2007)
HIV-1 packaging plasmid	pCMVΔR8.91 (p8.91)	(Zufferey et al. 1997)
Lentiviral vector for expression of Firefly Luciferase	pHR-SIN-SE (pCSFLW)	(Demaison et al. 2002)
Vesicular Stomatitis Virus glycoprotein	pMD.G (VSV-G)	Yasu Takeuchi, University College London

390

391 **Acknowledgements**

392 EG is supported by the Biotechnology and Biological Sciences Research Council
393 (<http://www.bbsrc.ac.uk>) via Strategic LoLa grant BB/K002465/1 “Developing Rapid Responses to
394 Emerging Virus Infections of Poultry (DRREVIP)”.

395

396

397 **References**

- 398 Agnarsson, I., Zambrana-Torrel, C. M., Flores-Saldana, N. P., & May-Collado, L. J. (2011). A time-
399 calibrated species-level phylogeny of bats (chiroptera, Mammalia). *PLoS Currents*, 3, RRN1212.
400 doi:10.1371/currents.RRN1212
- 401 Barman, S., Adhikary, L., Chakrabarti, A. K., Bernas, C., Kawaoka, Y., Nayak, D. P., et al. (2004). Role of
402 transmembrane domain and cytoplasmic tail amino acid sequences of influenza A virus
403 neuraminidase in raft association and virus budding. *Journal of virology*, 78(10), 5258–5269.
404 doi:10.1128/JVI.78.10.5258
- 405 Bertram, S., Glowacka, I., Blazejewska, P., Soilleux, E. J., Allen, P., Danisch, S., et al. (2010). Tmprss2
406 and Tmprss4 facilitate trypsin-independent spread of influenza virus in Caco-2 cells. *J Virol*,
407 84(19), 10016–10025. doi:10.1128/JVI.00239-10
- 408 Biuso, F., Carnell, G. W., Montomoli, E., & Temperton, N. (2018). A Lentiviral Pseudotype ELLA for
409 the Measurement of Antibodies Against Influenza Neuraminidase. *Bio-Protocol*, 8(14), 1–15.
410 doi:10.21769/BioProtoc.2936
- 411 Böttcher-Friebertshäuser, E., Freuer, C., Sielaff, F., Schmidt, S., Eickmann, M., Uhlenhorff, J., et al.
412 (2010). Cleavage of influenza virus hemagglutinin by airway proteases Tmprss2 and HAT
413 differs in subcellular localization and susceptibility to protease inhibitors. *Journal of virology*,
414 84(11), 5605–14. doi:10.1128/JVI.00140-10
- 415 Böttcher, E., Matrosovich, T. Y., Beyerle, M., Klenk, H.-D., Garten, W., & Matrosovich, M. N. (2006).
416 Proteolytic activation of influenza viruses by serine proteases Tmprss2 and HAT from human
417 airway epithelium. *Journal of virology*, 80(19), 9896–8. doi:10.1128/JVI.01118-06
- 418 Corti, D., Voss, J., Gambelin, S. J., Codoni, G., Macagno, A., Jarrossay, D., et al. (2011). A Neutralizing
419 Antibody Selected from Plasma Cells That Binds to Group 1 and Group 2 Influenza A
420 Hemagglutinins. *Science*, 333(6044), 850–6. doi:10.1126/science.1205669
- 421 Couzens, L., Gao, J., Westgeest, K. B., Sandbulte, M. R., Lugovtsev, V. Y., Fouchier, R. A. M., &
422 Eichelberger, M. C. (2014). An optimized enzyme-linked lectin assay to measure influenza A
423 virus neuraminidase inhibition antibody titers in human sera. *Journal of Virological Methods*,
424 210, 7–14. doi:10.1016/j.jviromet.2014.09.003
- 425 Cox, R. J., Mykkeltvedt, E., Robertson, J. S., & Haaheim, L. R. (2002). Non-lethal viral challenge of
426 influenza haemagglutinin and nucleoprotein DNA vaccinated mice results in reduced viral
427 replication. *Scandinavian Journal of Immunology*, 55(1), 14–23. doi:10.1046/j.1365-
428 3083.2002.01015.x
- 429 Demaison, C., Parsley, K., Brouns, G., Scherr, M., Battmer, K., Kinnon, C., et al. (2002). High-level
430 transduction and gene expression in hematopoietic repopulating cells using a human
431 immunodeficiency [correction of immunodeficiency] virus type 1-based lentiviral vector
432 containing an internal spleen focus forming virus promoter. *Human gene therapy*, 13(7), 803–
433 813. doi:10.1089/10430340252898984
- 434 Dos Reis, M., Tamuri, A. U., Hay, A. J., & Goldstein, R. A. (2011). Charting the host adaptation of
435 influenza viruses. *Molecular Biology and Evolution*, 28(6), 1755–1767.
436 doi:10.1093/molbev/msq317
- 437 Dreyfus, C., Ekiert, D. C., & Wilson, I. A. (2013). Structure of a Classical Broadly Neutralizing Stem
438 Antibody in Complex with a Pandemic H2 Influenza Virus Hemagglutinin. *Journal of Virology*,
439 87(12), 7149–7154. doi:10.1128/JVI.02975-12

- 440 Dreyfus, C., Laursen, N. S., Kwaks, T., Zuijdggeest, D., Khayat, R., Ekiert, D. C., et al. (2012). Highly
441 Conserved Protective Epitopes on Influenza B Viruses. *Science*, 337(6100), 1343–1348.
442 doi:10.1126/science.1222908
- 443 Dukes, J. D., Whitley, P., Chalmers, A. D., Gaush, C., Hard, W., Smith, T., et al. (2011). The MDCK
444 variety pack: choosing the right strain. *BMC Cell Biology*, 12(1), 43. doi:10.1186/1471-2121-12-
445 43
- 446 Ebrahimi, M., Aghagolzadeh, P., Shamabadi, N., Tahmasebi, A., Alsharifi, M., Adelson, D. L., et al.
447 (2014). Understanding the undelaying mechanism of HASubtyping in the level of physic-
448 chemical characteristics of protein. *PLoS ONE*, 9(5), 1–14. doi:10.1371/journal.pone.0096984
- 449 Ekiert, D. C., Bhabha, G., Elsliger, M., Friesen, R. H. E., Jongeneelen, M., Throsby, M., et al. (2009).
450 Antibody recognition of a highly conserved influenza virus epitope: implications for universal
451 prevention and therapy. *Science*, 324(5924), 246–251. doi:10.1126/science.1171491.Antibody
- 452 Ferrara, F., Molesti, E., Böttcher-Friebertshäuser, E., Cattoli, G., Corti, D., Scott, S. D., & Temperton,
453 N. J. (2012). The human Transmembrane Protease Serine 2 is necessary for the production of
454 Group 2 influenza A virus pseudotypes. *Journal of molecular and genetic medicine: an
455 international journal of biomedical research*, 7, 309–14.
- 456 Freidl, G. S., Binger, T., Müller, M. A., De Bruin, E., Van Beek, J., Corman, V. M., et al. (2015).
457 Serological evidence of influenza a viruses in frugivorous bats from Africa. *PLoS ONE*, 10(5), 1–
458 7. doi:10.1371/journal.pone.0127035
- 459 Friesen, R. H. E., Koudstaal, W., Koldijk, M. H., Weverling, G. J., Brakenhoff, J. P. J., Lenting, P. J., et al.
460 (2010). New class of monoclonal antibodies against severe influenza: Prophylactic and
461 therapeutic efficacy in ferrets. *PLoS ONE*, 5(2), 1–7. doi:10.1371/journal.pone.0009106
- 462 García-Sastre, A. (2012). The neuraminidase of bat influenza viruses is not a neuraminidase. *Proc
463 Natl Acad Sci U S A*, 109(46), 18635–18636. doi:10.1073/pnas.1215857109
- 464 Gelfond, J. A. L., Gupta, M., & Ibrahim, J. G. (2009). A bayesian hidden markov model for motif
465 discovery through joint modeling of genomic sequence and ChIP-chip data. *Biometrics*, 65(4),
466 1087–1095. doi:10.1111/j.1541-0420.2008.01180.x
- 467 Hansson, G. C., Simons, K., & Meer, G. Van. (1986). Two strains of the Madin- Darby canine kidney (
468 NDCK) cell line have distinct glycosphingolipid compositions. *Proceedings of the National
469 Academy of Sciences*, 5(3), 483–489.
- 470 Hoffmann, M., Krüger, N., Zmora, P., Wrensch, F., Herrler, G., & Pöhlmann, S. (2016). The
471 hemagglutinin of bat-associated influenza viruses is activated by TMPRSS2 for pH-dependent
472 entry into bat but not human cells. *PLoS ONE*, 11(3), 1–18. doi:10.1371/journal.pone.0152134
- 473 Jung, H., Lee, K. P., Park, S. J., Park, J. H., Jang, Y. S., Choi, S. Y., et al. (2008). TMPRSS4 promotes
474 invasion, migration and metastasis of human tumor cells by facilitating an epithelial-
475 mesenchymal transition. *Oncogene*, 27(18), 2635–2647. doi:10.1038/sj.onc.1210914
- 476 Juozapaitis, M., Aguiar Moreira, E., Mena, I., Giese, S., Riegger, D., Pohlmann, A., et al. (2014). An
477 infectious bat-derived chimeric influenza virus harbouring the entry machinery of an influenza
478 A virus. *Nat Commun*, 5, 4448. doi:10.1038/ncomms5448
- 479 Kandeil, A., Gomaa, M. R., Shehata, M. M., El Taweel, A. N., Mahmoud, S. H., Bagato, O., et al.
480 (2018). Isolation and characterization of a distinct influenza A virus from Egyptian bats. *Journal
481 of virology*, JVI.01059-18. doi:10.1128/JVI.01059-18
- 482 Lacek, K., Urbanowicz, R. A., Troise, F., De Lorenzo, C., Severino, V., Di Maro, A., et al. (2014).

- 483 Dramatic potentiation of the antiviral activity of HIV antibodies by cholesterol conjugation.
484 *Journal of Biological Chemistry*, 289(50), 35015–35028. doi:10.1074/jbc.M114.591826
- 485 Li, Q., Sun, X., Li, Z., Liu, Y., Vavricka, C. J., Qi, J., & Gao, G. F. (2012). Structural and functional
486 characterization of neuraminidase-like molecule N10 derived from bat influenza A virus.
487 *Proceedings of the National Academy of Sciences of the United States of America*, 109(46),
488 18897–18902. doi:10.1073/pnas.1211037109
- 489 Ma, W., García-Sastre, A., & Schwemmler, M. (2015). Expected and Unexpected Features of the
490 Newly Discovered Bat Influenza A-like Viruses. *PLoS Pathog*, 11(6), e1004819.
491 doi:10.1371/journal.ppat.1004819
- 492 Ma, W., Lager, K. M., Vincent, A. L., Janke, B. H., Gramer, M. R., & Richt, J. A. (2009). The role of
493 swine in the generation of novel influenza viruses. *Zoonoses and Public Health*, 56(6–7), 326–
494 337. doi:10.1111/j.1863-2378.2008.01217.x
- 495 Maeda, K., Hondo, E., Terakawa, J., Kiso, Y., Nakaichi, N., Endoh, D., et al. (2008). Isolation of novel
496 adenovirus from fruit bat (*Pteropus dasymallus yayeyamae*) [7]. *Emerging Infectious Diseases*,
497 14(2), 347–349. doi:10.3201/eid1402.070932
- 498 Mänz, B., Schwemmler, M., & Brunotte, L. (2013). Adaptation of avian influenza A virus polymerase in
499 mammals to overcome the host species barrier. *Journal of virology*, 87(April), 7200–9.
500 doi:10.1128/JVI.00980-13
- 501 Maruyama, J., Miyamoto, H., Kajihara, M., Ogawa, H., Maeda, K., Sakoda, Y., et al. (2014).
502 Characterization of the Envelope Glycoprotein of a Novel Filovirus, Lloviu Virus. *Journal of*
503 *Virology*, 88(1), 99–109. doi:10.1128/JVI.02265-13
- 504 Maruyama, J., Nao, N., Miyamoto, H., Maeda, K., Ogawa, H., Yoshida, R., et al. (2016).
505 Characterization of the glycoproteins of bat-derived influenza viruses. *Virology*, 488, 43–50.
506 doi:10.1016/j.virol.2015.11.002
- 507 Millet, J. K., & Whittaker, G. R. (2015). Host cell proteases: Critical determinants of coronavirus
508 tropism and pathogenesis. *Virus research*, 202, 120–34. doi:10.1016/j.virusres.2014.11.021
- 509 Moreira, É. A., Locher, S., Kolesnikova, L., Bolte, H., Aydillo, T., García-Sastre, A., et al. (2016).
510 Synthetically derived bat influenza A-like viruses reveal a cell type- but not species-specific
511 tropism. *Proceedings of the National Academy of Sciences*, 201608821.
512 doi:10.1073/pnas.1608821113
- 513 Otterstrom, J. J., Brandenburg, B., Koldijk, M. H., Juraszek, J., Tang, C., Mashaghi, S., et al. (2014).
514 Relating influenza virus membrane fusion kinetics to stoichiometry of neutralizing antibodies at
515 the single-particle level. *Proceedings of the National Academy of Sciences*, 111(48), E5143–
516 E5148. doi:10.1073/pnas.1411755111
- 517 Prevato, M., Cozzi, R., Pezzicoli, A., Taddei, A. R., Ferlenghi, I., Nandi, A., et al. (2015). An innovative
518 pseudotypes-based Enzyme-Linked Lectin Assay for the measurement of functional anti-
519 neuraminidase antibodies. *PLoS ONE*, 10(8), e0135383. doi:10.1371/journal.pone.0135383
- 520 Reperant, L. A., Kuiken, T., & Osterhaus, A. D. (2012). Adaptive pathways of zoonotic influenza
521 viruses: From exposure to establishment in humans. *Vaccine*, 30(30), 4419–4434.
522 doi:10.1016/j.vaccine.2012.04.049
- 523 Sahini, L., Tempczyk-Russell, A., & Agarwal, R. (2010). Large-scale sequence analysis of hemagglutinin
524 of influenza A virus identifies conserved regions suitable for targeting an anti-viral response.
525 *PLoS ONE*, 5(2), 1–9. doi:10.1371/journal.pone.0009268

- 526 Sui, J., Hwang, W. C., Perez, S., Wei, G., Aird, D., Chen, L., et al. (2009). Structural and functional
527 bases for broad-spectrum neutralization of avian and human influenza A viruses. *Nature*
528 *structural & molecular biology*, 16(3), 265–273. doi:10.1038/nsmb.1566
- 529 Sun, X., Shi, Y., Lu, X., He, J., Gao, F., Yan, J., et al. (2013). Bat-Derived Influenza Hemagglutinin H17
530 Does Not Bind Canonical Avian or Human Receptors and Most Likely Uses a Unique Entry
531 Mechanism. *Cell Reports*, 3(3), 769–778. doi:10.1016/j.celrep.2013.01.025
- 532 Temperton, N. J., Hoschler, K., Major, D., Nicolson, C., Manvell, R., Hien, V. M., et al. (2007). A
533 sensitive retroviral pseudotype assay for influenza H5N1-neutralizing antibodies. *Influenza and*
534 *other respiratory viruses*, 1(3), 105–112. doi:10.1111/j.1750-2659.2007.00016.x
- 535 Tong, S., Li, Y., Rivallier, P., Conrardy, C., Castillo, D. A. A., Chen, L.-M., et al. (2012). A distinct lineage
536 of influenza A virus from bats. *Proceedings of the National Academy of Sciences of the United*
537 *States of America*, 109(11), 4269–4274. doi:10.1073/pnas.1116200109
- 538 Tong, S., Zhu, X., Li, Y., Shi, M., Zhang, J., Bourgeois, M., et al. (2013). New World Bats Harbor Diverse
539 Influenza A Viruses. *PLoS Pathogens*, 9(10). doi:10.1371/journal.ppat.1003657
- 540 Webster, R. G., Bean, W. J., Gorman, O. T., Chambers, T. M., & Kawaoka, Y. (1992). *Evolution and*
541 *Ecology of Influenza A Viruses. MICROBIOLOGICAL REVIEWS* (Vol. 56).
542 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC372859/pdf/microrev00028-0170.pdf>.
543 Accessed 27 August 2018
- 544 Wilson, I. a, & Cox, N. J. (1990). Structural basis of immune recognition of influenza virus
545 hemagglutinin. *Annual review of immunology*, 8(1), 737–771.
546 doi:10.1146/annurev.iy.08.040190.003513
- 547 Wu, Y., Wu, Y., Tefsen, B., Shi, Y., & Gao, G. F. (2014). Bat-derived influenza-like viruses H17N10 and
548 H18N11. *Trends in Microbiology*, 22(4), 183–191. doi:10.1016/j.tim.2014.01.010
- 549 Yondola, M. A., Fernandes, F., Belicha-Villanueva, A., Uccellini, M., Gao, Q., Carter, C., & Palese, P.
550 (2011). Budding capability of the influenza virus neuraminidase can be modulated by tetherin.
551 *Journal of virology*, 85(6), 2480–91. doi:10.1128/JVI.02188-10
- 552 Zhou, B., Ma, J., Liu, Q., Bawa, B., Wang, W., Shabman, R. S., et al. (2014). Characterization of
553 Uncultivable Bat Influenza Virus Using a Replicative Synthetic Virus. *PLoS Pathogens*, 10(10),
554 e1004420. doi:10.1371/journal.ppat.1004420
- 555 Zhu, X., Yang, H., Guo, Z., Yu, W., Carney, P. J., Li, Y., et al. (2012). Crystal structures of two subtype
556 N10 neuraminidase-like proteins from bat influenza A viruses reveal a diverged putative active
557 site. *Proceedings of the National Academy of Sciences of the United States of America*, 109(46),
558 18903–18908. doi:10.1073/pnas.1212579109
- 559 Zimmer, G., Lottspeich, F., Maisner, A., Klenk, H. D., & Herrler, G. (1997). Molecular characterization
560 of gp40, a mucin-type glycoprotein from the apical plasma membrane of Madin-Darby canine
561 kidney cells (type I). *The Biochemical journal*, 326, 99–108. doi:10.1042/bj3260099
- 562 Zufferey, R., Nagy, D., Mandel, R. J., Naldini, L., & Trono, D. (1997). Multiply attenuated lentiviral
563 vector achieves efficient gene delivery in vivo. *Nat Biotechnol*, 15(9), 871–875.
564 doi:10.1038/nbt0997-871
- 565