Genetic control of pod dehiscence in domesticated common bean: Associations with range expansion and local aridity conditions

Travis A. Parker, Jorge C. Berny Mier y Teran, Antonia Palkovic, Judy Jernstedt, and Paul Gepts¹

University of California, Department of Plant Sciences, 1 Shields Avenue, Davis, CA 95616-8780.

Classification

Major category: BIOLOGICAL SCIENCES

Minor category: Agricultural Sciences

Keywords: dirigent | aridity tolerance | GWAS | local adaptation | pod shattering

Author contributions: T.A.P. prepared the manuscript and conducted laboratory phenotyping, QTL mapping, GWAS, microscopy, and sequencing. J.C.B.M.T. genotyped the IxS population, gathered field phenotypes, co-conducted QTL mapping, and provided guidance for other procedures. A.P. assisted with field and greenhouse trials. J.J. led the sectioning and microscopy studies. P.G. conceived the initial project and provided guidance. All authors edited the manuscript.

The authors declare no conflict of interest.

Data deposition: The coding DNA sequence of *PvPdh1* for ICA Bunsí and SXB 405 will be deposited in the NCBI database.

¹To whom correspondence should be addressed. Email: plgepts@ucdavis.edu. Phone: 1 (530) 752-7743.

Significance

Plant domestication has radically modified crop morphology and development. Nevertheless, many crops continue to display some atavistic characteristics that were advantageous to their wild ancestors, such as pod dehiscence (PD). Domesticated common bean (*Phaseolus vulgaris*), a nutritional staple for millions of people globally, shows considerable variation in PD. Here, we identified multiple genetic regions controlling PD in common bean grown throughout geographically distributed lineages. For example, on chromosome Pv03, *PvPdh1* shows a single base-pair substitution that is strongly associated with decreased PD and expansion of the crop into northern Mexico, where the arid conditions promote PD. The environmental dependency and genetic redundancy explain the maintenance of atavistic traits under domestication. Knowledge of PD genetics will assist in developing aridity-adapted varieties.

Abstract

A reduction in pod dehiscence (PD) is an important part of the domestication syndrome in legumes, including common bean. Despite this, many modern dry bean varieties continue to suffer yield reductions due to dehiscence, an atavistic trait, which is particularly problematic in hot, dry environments. To date, the genetic control of this important trait has been only partially resolved. Using QTL mapping and GWAS, we identified major PD QTLs in dry beans on chromosomes Pv03, Pv05, Pv08, and Pv09, three of which had not been described previously. We further determined that the QTL on chromosome Pv03, which is strongly associated with PD in Middle American beans, includes a dirigent-like candidate gene orthologous to *Pod dehiscence 1 (Pdh1)* of soybean. In this gene, we identified a substitution in a highly conserved amino acid that is unique to PD-resistant varieties. This allele is associated with the expansion of Middle American domesticated common beans into the arid environments of northern Mexico, resulting in a high allelic frequency in the domesticated ecogeographic race Durango. The polygenic redundancy and environmental dependency of PD resistance may explain the maintenance of this atavistic characteristic after domestication. Use of these alleles in breeding will reduce yield losses in arid growing conditions, which are predicted to become more widespread in coming decades.

dirigent | aridity tolerance | GWAS | local adaptation | pod shattering

\body

Introduction

Effective seed dispersal is vital for spermatophytes. In the Fabaceae, the third largest family of flowering plants (1), seed dispersal is typically mediated by the explosive dehiscence (“shattering”) of pods at maturity. This mechanism is effective for the propagation of plants in the wild, but is associated with reduced yield and harvest constraints for cultivated crops. As such, there has been selection against pod dehiscence (PD), which continues to this day. A reduction in PD has been a central part of the domestication syndrome of many domesticated pulses. Anatomical differences are associated with some, but not all variation in PD in these species (2, 3). Reviews of the developmental genetics related to PD are available (4, 5).

In soybean (*Glycine max* (L.) Merr.), a domestication-related reduction in PD is mediated by the *NAC* family transcription factor *SHAT1-5* (6). A further reduction in PD is controlled by *Pod dehiscence 1* (*Pdh1*) (7). *Pdh1* encodes a dirigent-like gene related to lignin synthesis. This mutation is associated with the expansion of soybeans into arid regions. *Pdh1* is highly expressed in obliquely oriented fibers lining the soybean pod walls and has a minimal effect on gross pod anatomy (8).

Common bean (*Phaseolus vulgaris* L.) is the foremost grain legume for direct human consumption and is a dietary staple for hundreds of millions of people worldwide (9). Common bean diverged into distinct Middle American and Andean gene pools approximately 87,000 years before present, well before the first human migrations into the Americas (10). Subsequently, human populations independently domesticated members of each gene pool, making up two of at least seven domestication events in the genus *Phaseolus* (11) and 41 domestication events in the Fabaceae (12). Each of the two major domesticated gene pools of common bean is divided into several ecogeographic races. In the Middle American domesticated gene pool, it is important to single out race Durango, which includes varieties from arid, higher altitude regions of northern Mexico, and race Mesoamerica, from the warmer, humid lowlands of southern Mexico and Central America (13).

Phaseolus vulgaris can be separated into two economic groups: snap beans, grown for pods as a vegetable, and dry beans, grown for grain. Dry beans produce fibrous pods, which can be easily separated from seeds during threshing. In snap beans, selections in the 19th century led to “stringless” varieties with extreme PD resistance and very little pod suture fiber deposition

(14). Stringless varieties now dominate the snap bean market, but stringlessness is absent in dry beans. Using a recombinant inbred (RI) population derived from stringless cv. ‘Midas’ and wild accession G12873, Koinange *et al.* (15) identified a major pod fiber QTL on linkage group Pv02 (16). This gene, called *Stringless* (*St*), maps near the common bean ortholog of *INDEHISCENT* (*PvIND*), but there is a lack of complete co-segregation between the loci and no causal polymorphism is known to exist in the *PvIND* sequence (17). Rau *et al.* (18) used QTL mapping to identify a segregating locus on Pv05 in the Midas x G12873 genetic background. Despite this, a comprehensive evaluation of the genetic basis of PD in diverse germplasm has not yet been conducted and no molecular polymorphisms with a potential causal effect on PD have been described.

In the research reported here, we used high-precision phenotyping techniques, both in an RI population and diversity panels, to identify PD QTLs in common bean. A locus underlying one of the major QTLs was sequenced to identify possible causal polymorphisms. We found orthologous mechanisms that regulate pod dehiscence in this species. We were further able to identify associations between PD and local environmental conditions. Alleles identified in this study will be valuable for developing common bean varieties suited to the increasingly arid climatic conditions of coming decades.

Results

Anatomical analysis of developing pods. Clear differences in pod anatomy were found between domesticated snap bean, domesticated dry bean, and wild common bean (Fig. 1). Wild beans produce a lignified wall fiber layer (LFL) in the pods that is thicker than the bundle cap layer, while the LFL is greatly reduced in domesticated varieties. Stringless snap beans have weakly lignified vascular bundle sheaths (VS) at the suture, with a reduction in the number of lignified cells and the extent of secondary cell wall deposition in each cell. In stringless beans, the LFL is typically absent. In contrast to the clear anatomical differences between these three groups, no variation between PD-resistant and PD-susceptible domesticated dry bean pods was observed (Fig. 1B, 1C).

Variation in the ICA Bunsu/SXB 405 (IxS) population. Segregation for PD was determined in an RI population derived from PD-susceptible cv. ‘ICA Bunsu’ and PD-resistant

cv. ‘SXB 405’. Both parental genotypes belong to the Middle American gene pool. Three phenotyping approaches were used to evaluate PD (Fig. S1) and each had a unique distribution pattern (Fig. S2). These phenotypes were strongly correlated (Fig. S3). Varieties that dehiscence in the field had higher rates of PD after desiccation at 65°C (two-tailed t-test, $p=3.1 \times 10^{-8}$) and required lower levels of force to induce fracture at the sutures (two-tailed t-test, $p=1.2 \times 10^{-9}$). Similarly, the proportion dehiscing in the desiccator and force required to cause PD were negatively correlated ($r^2 = 0.71$ simple linear model, $p < 2 \times 10^{-16}$).

QTL mapping identified a major, PD-related QTL peak located in the same position on linkage group Pv03 using each of the three phenotyping methods (Fig. 2). The QTL mapped between SNP markers ss715639553 and ss715639323, which are separated by approximately 900 kb of physical distance (Table 1). Force measurement produced the most significant results (LOD score 53.3), followed by desiccation (LOD score 42.7), and field notes (LOD score 8.9). All methods produced results that were statistically significant based on 1000 randomized permutations of the data. The allele at the most significant SNP explained 17% of the variation in PD based on field notes, 59% of the variation based on desiccation, and 64% of the variation in force required for fracture in the population. Analyses to find additional PD QTLs failed to identify other regions of interest in the IxS population.

Syntenic mapping and expression. Due to the close phylogenetic relationship and extensive microsynteny between *P. vulgaris* and *G. max* (19, 20), gene families known to affect PD in soybean were primary candidates for control of the trait in common bean. These families include the *NAC*-domain transcription factors and dirigent-like genes.

No *NAC*-domain transcription factors exist in the Pv03 QTL mapping interval. In contrast, the LegumeIP 2.0 synteny tool indicated that strong synteny exists between the soybean region surrounding *GmPdh1* and the common bean QTL (Table S3). This is in agreement with previous synteny analyses (20, 21). An amino acid BLAST of GmPDH1 (cv. Toyosume) against the *P. vulgaris* G19833 proteome (21) indicates that the most similar common bean protein is encoded by Phvul.003G252100, which maps between the two most significant Pv03 QTL SNP markers. A neighbor-joining tree of common bean and soybean dirigent proteins indicates that GmPDH1 and the protein product of Phvul.003G252100 cluster together (Fig. S4). Furthermore, the common bean gene’s expression is limited to developing pods, with no detectable expression

in any other tissues (Fig. S5; data from (22)). This is comparable to the expression of soybean *PDHI* (7), and indicates that the gene serves a function unique to pods. Together, these results suggest that Phvul.003G252100 is orthologous to *GmPDHI*. Phvul.003G252100 is hereafter referred to as *PvPdhI*.

Sequencing of *PvPdhI*. Sequencing of *PvPdhI* in ICA Bunsu and SXB 405 revealed a non-synonymous single-base-pair substitution at position 485 of the gene's coding sequence (Fig. S6A). This substitution leads to a threonine/asparagine polymorphism (T162N) in the protein product (Fig. S6B). The 11 RILs with recombination between the most significant markers from QTL mapping showed complete co-segregation between the threonine/asparagine polymorphism and the PD phenotype (Table S1).

To investigate the functional importance of T162N, we evaluated the extent of its sequence conservation, surveyed literature related to this position in closely related dirigent proteins and used software tools to predict the effect of this substitution at the position. Sequencing of *PvPdhI* in several species of wild and domesticated *Phaseolus* from the USDA National Plant Germplasm System (NPGS) and UC Davis showed that the asparagine at this position was unique to the Middle American domesticated gene pool (Table S2). No polymorphism in the Andean gene pool was consistently associated with PD. In the Middle American gene pool, PD was significantly higher among genotypes with a threonine at position 162 than an asparagine (t-test: $p=0.0002$, $n=47$, Fig. S7). This threonine was strictly conserved in Andean domesticated common bean, Middle American and Andean wild common bean, and the closely related *P. lunatus* and *P. dumosus* (Table S2). The residue is present in 99 of the 100 most similar proteins in the NCBI database (Fig. S8A; see Discussion), indicating its functional importance. This threonine is also conserved in the 19 most similar proteins of *Selaginella moellendorffii* (Fig. S8B), a member of the first diverging group of lignin-containing plants. No comparable protein could be found in the proteome of *Physcomitrella patens*, a non-lignified moss.

Studies of closely related dirigent proteins indicate that this threonine is a component of one of the protein's active sites ("T163" in Fig. S8C; from (23)), and its substitution eliminates protein function (Fig. S8D; from (24)). An analysis with PROVEAN (25) predicted that the T162N mutation would have a deleterious effect (score: -4.587, cutoff = -2.5).

Validation through association mapping. The BeanCAP Middle American Diversity Panel (MDP) (26) was grown to determine, using the desiccation method, whether the Pv03 *PvPdh1* locus was related to PD in a broader population. A genome-wide association study (GWAS) indicated that the SNP closest to *PvPdh1* in physical distance was also the most significantly associated with PD (Fig. 3A, MAF threshold = 0.1). This SNP (S1_149243152) was 5.7 kb from the polymorphism in *PvPdh1*. Pv06 and Pv08 also included loci significantly associated with PD.

GWAS was also conducted in the Andean diversity panel (ADP) (27) to determine which loci control PD in this independently domesticated population. Chromosomes Pv03, Pv05, Pv08, and Pv09 all included major regions significantly associated with PD (Fig. 3B). The QTL on chromosome Pv08 was in an overlapping physical position with a QTL from the MDP (Fig. 3A).

In both the Andean and Middle American gene pools, PD varied greatly between market classes (Table S4). In Andean beans, PD after desiccation averaged 3% in the purple speck market class and 41% in the cranberry types. In Middle American beans, averages were below 1% for pinto types of race Durango, and as high as 18% in the black beans of race Mesoamerica. Members of Middle American race Mesoamerica displayed considerable variation in PD. GWAS using only members of this race (MDP with PC1 > 50) showed that the Pv08 QTL was most closely associated with PD in the population (Fig. S9). SNP S1_329543689, near the center of this interval of interest, was used for subsequent analyses. The region near *PvPdh1* did not include significant SNPs in this race, further indicating that races Durango and Mesoamerica rely on different genes for PD resistance.

To visualize the correlation between PD and population substructure in the MDP, PD was plotted against the first principal component of the genetic data. Each point was color-coded by its allele at the GWAS SNP peaks on Pv03 (S1_149243152, 5.7kb from *PvPdh1*) and Pv08 (SNP S1_329543689) (Fig. 4). Members of the MDP with the Pv03 PD resistance allele exhibited mean PD in the desiccator of 0.0067, with a maximum value of 0.14. In genotypes with no known resistance allele, the mean level of PD was 0.206 and ranged up to 0.63 (Fig. 4).

Discussion

A reduction in PD is a fundamental component of the domestication syndrome in common bean (15), and is important for future green and dry bean production and food security. In snap beans, a major gene - *St* - controls the presence or absence of pod strings and PD (Prakken, 1934; Koinange et al. 1996). For dry beans, we report here three novel QTLs mapped for the trait (on Pv03, Pv08, and Pv09), confirm a QTL on Pv05, identify a putative causal polymorphism in the *PvPdh1* gene underlying the major QTL on Pv03, and describe the association between Pv03, Pv05, and Pv08 QTLs and climate variables, especially precipitation.

Anatomical differences among wild and domesticated types. Our microscopy results are consistent with those of previous researchers, who noted that reduced lignification in the LFL is correlated with reduced PD among wild beans, domesticated dry beans, and domesticated snap beans (2, 3, 28). However, we were not able to identify anatomical differences responsible for the variation in dehiscence among dry bean cultivars. This is consistent with the pattern seen in *GmPdh1* in soybean (8) and the expected result of a mutation in *PvPdh1*. Indeed, *PvPdh1* is thought to modify the biochemical structure of lignin, rather than its total quantity or cell fate in the relevant pod structures. A loss of function mutation in this gene would therefore not lead to clear anatomical differences relative to the wild-type.

***PvPdh1* as a candidate gene for the Pv03 QTL identified in this study.** The strict conservation of the threonine at position 162 in *PvPdh1* highlights its functional importance. Of the 100 most similar known protein models, the one that lacks a threonine at this position is found in *Trifolium subterraneum*, a legume that produces pods that mature underground. PD is not relevant for seed dispersal in this species, and the gene may be undergoing pseudogenization. This threonine is maintained in the proteome of *Selaginella moellendorffii*, indicating that the residue has been conserved since before the lycophyte-euphyllophyte divergence. This coincides with the origin of lignin and lignans, and indicates the residue's functional role for members of the protein family. In a remarkable example of parallelism, independent loss of function mutations in this gene are found in certain domesticated populations in soybean (*G. max*) and in *P. vulgaris*, both species being subjected to selection for reduced dehiscence. This provides an additional example of strongly convergent phenotypic and molecular evolution (29). Similar examples of parallel evolution in common bean included the determinacy trait (*fin* or *PvTFL1*); (30, 31)) and absence of pigmentation (*P*; (32)).

Further research is needed to identify the biochemical and biophysical aspects responsible for differences in PD in domesticated dry beans. Notably, our results could shed light on the fundamental process of lignin synthesis. Dirigent-like genes, including *PvPdh1*, encode non-enzymatic proteins that guide the dimerization of lignin and lignan monomers (33). The role of these proteins in lignin synthesis has been debated, with suggestions that polymerization is guided (34) or unguided (35). Varieties of common bean with mutations in *Pdh1* could be used to elucidate the role of this protein family in lignin synthesis generally.

QTLs and Candidate Genes Identified by Association Mapping. Association mapping revealed several other dehiscence-related QTLs across the gene pools and races of common bean (Table 1). Our ADP association mapping identified significant Pv03 SNPs in an interval that is syntenic with a region controlling dehiscence in cowpea (36). *NAC* family and C2H2-type zinc finger transcription factors are found in this region (Table 1), and members of these families affect PD in soybean (6) and rapeseed (37), respectively. Orthologs of these genes may also affect dehiscence in cowpea (36). Interestingly, the QTL is large enough to include *PvPdh1*, although the QTLs discovered in Middle American beans and cowpeas are non-overlapping.

Another major QTL for PD in Andean beans maps to Pv05, as described recently (18), and several genes in this region are candidates for future study. Rau *et al.* (18) noted that an ortholog of *MYB26* exists in the qPD5.1-Pv region of interest on Pv05, which may be responsible for variation in PD. Significant Pv05 SNPs from our association mapping completely envelope the qPD5.1-Pv interval, supporting this result. Interestingly, our most significant Pv05 SNPs in the ADP are found just 22kb from *MYB46*. *MYB46* is involved in the same pathway as *MYB26* and the soybean PD resistance gene *SHAT1-5* (38, 6). *MYB46* also works redundantly with *MYB83*, a gene that may play a role in cowpea pod development (38, 39), making *MYB46* another potential subject of future study.

Several genes of interest exist near the middle of the ADP's Pv08 GWAS peak. These include a MYB family transcription factor with similarity to *A. thaliana* *MYB17*, three *WRKY* family transcription factors, which are related to genes involved in sorghum dehiscence (40), and a polygalacturonase, a group known to influence PD in *A. thaliana* (41) (Table 1).

The Pv09 GWAS peak found in the ADP included a gene predicted to be *cellulose synthase A7* (*CESA7*, Table 1). *CESA7* may play a role in fiber development in cowpea (39).

Similarly, two polygalacturonases are found in this interval, and members of this family are known to affect seed dispersal in *A. thaliana* (41). These genes may regulate dehiscence by altering the breakdown of cell wall material in developing pods. Identifying polymorphisms in PD candidate genes will be a promising area for future study.

Associations with environmental conditions. PD in common bean is correlated with environmental parameters. The PD-resistant allele of *PvPdh1* on Pv03 is found exclusively in genotypes with ancestry from race Durango (Table 1). Race Durango is adapted to higher elevations and lower humidity regions, particularly in the northern part of Mexico (13). The semi-arid conditions in these areas cause pods to become dry and brittle, which exacerbates PD. The non-functional *PvPdh1* allele rose to very high frequency in this ecogeographic race. In contrast, race Mesoamerica is adapted to humid lowlands, where environmental conditions mask PD and reduce selection pressure against it. In this race, the loss-of-function *PvPdh1* allele remains at low frequency and PD is widespread (Fig. 4A). Interestingly, this ecogeographic pattern closely parallels that of soybean, in which *Pdh1*-mediated resistance to PD is most common in arid regions (7). *PvPdh1* may also be responsible for the ease of threshing that has been noted in race Mesoamerica (13). In humid environments, the wild type *PvPdh1* allele may facilitate separation of seeds from pod material, while PD in the field remains low. In northern Mexico, the semi-arid climate facilitates threshing but increases PD in the field. Under these conditions, the PD-resistant allele is advantageous. Because of this trade-off, the polymorphism in *PvPdh1* appears to be related to local adaptation (Fig. 5). Alleles that prevent PD will be valuable in coming decades, which are predicted to be increasingly arid (42).

Redundancies in genetic control and maintenance of atavistic traits. Crosses between races have tremendous potential for crop improvement (for example, between races Durango and Mesoamerica (43)), but could also result in problematic gene complementation. Because several genes influence PD redundantly, cultivars descended from crosses between races could demonstrate atavistic transgressive segregation. This may be responsible for the high levels of dehiscence seen in some varieties of common bean. The interactions between these loci will also be of considerable importance for plant breeders.

Methods

Details regarding materials and methods can be found in SI Methods. Pods were sectioned for microscopy using a Vibratome. Lignified and hydrophobic structures were visualized using epifluorescence microscopy after staining with Auramine O.

The IxS population was genotyped using the Illumina Infinium II BARCBean6K_3 BeadChip. In Spring 2014, 238 RILs were grown in an unreplicated trial and visually phenotyped for the presence or absence of PD. In fall 2014, 191 RILs in a partially replicated trial were phenotyped based on proportion of pods dehiscing due to desiccation and force required to cause fracture. The maximum LOD score of 1000 random permutations of the data was used as a significance threshold.

Synteny mapping was conducted using LegumeIP 2.0 (44) and CoGe SynMap (45). Candidate genes were identified by NCBI BLAST and clustered through the NCBI portal. Gene expression data were accessed through the Common Bean Gene Expression Atlas (22). *PvPdh1* of ICA Bunsí, SXB 405, and RILs of interest was amplified by PCR and sequenced at the UC DNA Sequencing Facility. The CONstraint-Based multiple ALIGNment Tool (COBALT) (46) was used to align the PvPDH1 amino acid sequence to the most similar documented proteins of the NCBI database. PROVEAN (25) was used to estimate mutational effects.

PvPdh1 was sequenced in accessions with known pod shattering phenotypes from NPGS and UC Davis. Because members of the reproductively isolated Andean gene pool did not carry the T162N substitution, these individuals were filtered from subsequent analyses. For Middle American accessions, the categorical shattering scale used by USDA was translated into a simple numeric scale and PD between allele groups was compared using a student's t-test.

The ADP (27) consisted of 208 phenotyped accessions, and these were evaluated based on presence of PD in the field, proportion dehiscing in the desiccator, and force required to cause fracture. The MDP (26) included 278 phenotyped varieties that were evaluated by the desiccation method alone. GWAS in both populations were conducted using TASSEL (47) through SNIPlay (48). Manhattan plots were visualized using the qqman R package (49).

Worldclim2 precipitation data (50) were compiled with Natural Earth national boundary shapefiles (51) in QGIS to visualize precipitation patterns in the range of Middle American beans.

ACKNOWLEDGEMENTS. The authors would like to thank S. Beebe (CIAT, Cali, Colombia) for providing seeds of the IxS population. Seeds of the ADP and MDP were generously provided

by R. Lee and P. McClean (North Dakota State University). Undergraduates M. Rocha, P. Silva Rezende, G. Coelho Portilho, N. Hamada, E. Yang, A. Herrera, J. Pimentel, M. Bustamante, E. White, J. Gonzales, and P. Augello contributed to DNA extractions, pod phenotyping, and other lab protocols. Paola Hurtado, Andrea Ariani, and other members of the Gepts lab provided ideas for data analysis. Funding for T.A.P. was provided through a Clif Bar Family Foundation Seed Matters fellowship and Lundberg Family Farms research support.

References

1. LPWG. (2017). A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny The Legume Phylogeny Working Group (LPWG). *Taxon*, 66(1), 44-77.
2. Prakken R (1934). Inheritance of colours and pod characters in *Phaseolus vulgaris* L. *Genetica*, 16(3-4), 177-296.
3. Tiwari SP, Bhatia VS (1995). Characters of pod anatomy associated with resistance to pod-shattering in soybean. *Ann Bot*, 76(5), 483-485.
4. Dong Y, Wang YZ (2015). Seed shattering: from models to crops. *Front Plant Sci*, 6, 476.
5. Ballester P, Ferrandiz C (2017). Shattering fruits: variations on a dehiscent theme. *Curr Opin Plant Biol*, 35, 68-75.
6. Dong Y, et al. (2014). Pod shattering resistance associated with domestication is mediated by a NAC gene in soybean. *Nat Comm*, 5, 3352.
7. Funatsuki H, et al. (2014). Molecular basis of a shattering resistance boosting global dissemination of soybean. *Proc Natl Acad Sci*, 111(50), 17797-17802.
8. Suzuki M, Fujino K, Funatsuki H (2009). A major soybean QTL, qPDH1, controls pod dehiscence without marked morphological change. *Plant Prod Sci*, 12(2), 217-223.
9. Gepts, P et al. (2008). Genomics of *Phaseolus* beans, a major source of dietary protein and micronutrients in the tropics. *Genomics of tropical crop plants* eds. Moore PH, Ming R. (Springer, New York), 113-143.
10. Ariani A, Berny Mier y Teran JC, Gepts P (2018). Spatial and temporal scales of range expansion in wild *Phaseolus vulgaris*. *Mol Biol Evol*, 35(1), 119-131.
11. Bitocchi E, et al. (2017). Beans (*Phaseolus* spp.) as a model for understanding crop evolution. *Front Plant Sci*, 8, 722.
12. Harlan JR (1992). *Crops and man* (No. Ed. 2). (American Society of Agronomy, Madison, Wisconsin, USA).
13. Singh SP, Gepts P, Debouck DG (1991). Races of common bean (*Phaseolus vulgaris*, Fabaceae). *Econ Bot*, 45(3), 379-396.
14. Foster N, Cordell LS (Eds.). (1992). *Chilies to chocolate: Food the Americas gave the world*. University of Arizona Press.
15. Koinange EM, Singh SP, Gepts P. (1996). Genetic control of the domestication syndrome in common bean. *Crop Sci*, 36(4), 1037-1045.
16. Freyre R, et al. (1998). Towards an integrated linkage map of common bean. 4. Development of a core linkage map and alignment of RFLP maps. *Theor Appl Genet*, 97(5-6), 847-856.
17. Gioia T, Logozzo G, Kami J, Spagnoletti Zeuli P, Gepts P (2012). Identification and characterization of a homologue to the *Arabidopsis* *INDEHISCENT* gene in common bean. *J Hered*, 104(2), 273-286.

18. Rau D, *et al.* (2018). Genomic dissection of pod shattering in common bean: mutations at nonorthologous loci at the basis of convergent phenotypic evolution under domestication of leguminous species. *The Plant Journal*.
19. Gepts P, *et al.* (2005) Legumes as a model plant family. Genomics for food and feed Report of the Cross-Legume Advances through Genomics Conference. *Plant Physiol*, 137(4):1228-1235.
20. McClean PE, Mamidi S., McConnell M, Chikara S, Lee R (2010). Synteny mapping between common bean and soybean reveals extensive blocks of shared loci. *BMC Genomics*, 11(1), 184.
21. Schmutz J, *et al.* (2014). A reference genome for common bean and genome-wide analysis of dual domestications. *Nature genetics*, 46(7), 707.
22. O'Rourke JA *et al.* (2014). An RNA-Seq based gene expression atlas of the common bean. *BMC Genomics*, 15(1), 866.
23. Kim KW, *et al.* (2015). Trimeric structure of (+)-pinoresinol-forming dirigent protein at 1.95 Å resolution with three isolated active sites. *J Biol Chem*, 290(3), 1308-1318.
24. Gasper R, *et al.* (2016). Dirigent protein mode of action revealed by the crystal structure of AtDIR6. *Plant Physiol*, pp-01281.
25. Choi Y, Chan AP (2015). PROVEAN web server: a tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics* 31(16): 2745-2747.
26. Moghaddam SM, *et al.* (2016). Genome-wide association study identifies candidate loci underlying agronomic traits in a Middle American diversity panel of common bean. *Plant Genome*, 9(3).
27. Cichy KA *et al.* (2015). A *Phaseolus vulgaris* diversity panel for Andean bean improvement. *Crop Sci*, 55(5), 2149-2160.
28. Murgia ML *et al.* (2017). A comprehensive phenotypic investigation of the “pod-shattering syndrome” in common bean. *Front Plant Sci*, 8, 251.
29. Lenser T, Theißen G (2013). Molecular mechanisms involved in convergent crop domestication. *Trends in plant science*, 18(12), 704-714.
30. Repinski SL, Kwak M, Gepts P (2012). The common bean growth habit gene *PvTFL1y* is a functional homolog of *Arabidopsis TFL1*. *Theor Appl Genet*, 124(8), 1539-1547.
31. Kwak M, Toro O, Debouck DG, Gepts, P. (2012). Multiple origins of the determinate growth habit in domesticated common bean (*Phaseolus vulgaris*). *Ann Bot*, 110(8), 1573-1580.
32. McClean PE, *et al.* (2018). White seed color in common bean (*Phaseolus vulgaris*) results from convergent evolution in the *P (Pigment)* gene. *New Phytol*.
33. Davin LB, *et al.* (1997). Stereoselective bimolecular phenoxy radical coupling by an auxiliary (dirigent) protein without an active center. *Science*, 275(5298), 362-367.
34. Hosmani PS, *et al.* (2013). Dirigent domain-containing protein is part of the machinery required for formation of the lignin-based Casparian strip in the root. *Proc Natl Acad Sci*, 110(35), 14498-14503.
35. Ralph J, *et al.* (2008). Lignification: are lignins biosynthesized via simple combinatorial chemistry or via proteinaceous control and template replication. *Recent Advances in Polyphenol Research*, eds. Daayf F, Lattanzio, V. (Blackwell, Chichester, UK), 36-66.
36. Lo, S *et al.* (2018). Identification of QTL controlling domestication-related traits in cowpea (*Vigna unguiculata* L. Walp). *Sci Rep*, 8(1), 6261.
37. Tao Z, *et al.* (2017). BnLATE, a Cys2/His2-type zinc-finger protein, enhances silique shattering resistance by negatively regulating lignin accumulation in the silique walls of *Brassica napus*. *PLoS one*, 12(1), e0168046.
38. McCarthy RL, Zhong R, Ye ZH (2009). MYB83 is a direct target of SND1 and acts redundantly with MYB46 in the regulation of secondary cell wall biosynthesis in *Arabidopsis*. *Plant Cell Physiol*, 50(11), 1950-1964.

39. Suanum W, *et al.* (2016). Co-localization of QTLs for pod fiber content and pod shattering in F2 and backcross populations between yardlong bean and wild cowpea. *Mol Breed*, 36(6), 80.
40. Tang H *et al.* (2013). Seed shattering in a wild sorghum is conferred by a locus unrelated to domestication. *Proc Natl Acad Sci*, 201305213.
41. Ogawa M, Kay P, Wilson S, Swain SM (2009). ARABIDOPSIS DEHISCENCE ZONE POLYGALACTURONASE1 (ADPG1), ADPG2, and QUARTET2 are polygalacturonases required for cell separation during reproductive development in Arabidopsis. *Plant Cell*, 21(1), 216-233.
42. Sherwood S, Fu Q (2014) A drier future? *Science* 343.6172, 737-739.
43. Singh SP, Molina A, Urrea CA, Gutiérrez JA (1993). Use of interracial hybridization in breeding the race Durango common bean. *Canadian Journal Plant Science* 73, 785-793.
44. Li J, Dai X, Zhuang Z, Zhao PX (2016). LegumeIP 2.0—a platform for the study of gene function and genome evolution in legumes. *Nucleic Acids Res*, 44(D1), D1189-D1194.
45. Lyons E, Pedersen B, Kane J, Freeling M (2008). The value of nonmodel genomes and an example using SynMap within CoGe to dissect the hexaploidy that predates the rosids. *Trop Plant Biol*, 1(3-4), 181-190.
46. Papadopoulos JS, Agarwala R (2007). COBALT: constraint-based alignment tool for multiple protein sequences. *Bioinformatics*, 23(9), 1073-1079.
47. Bradbury PJ, *et al.* (2007). TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*, 23(19), 2633-2635.
48. Dereeper A, *et al.* (2011). SNIPlay: a web-based tool for detection, management and analysis of SNPs. Application to grapevine diversity projects. *BMC Bioinformatics*, 12(1), 134.
49. Turner SD (2018). qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *J Open Source Softw*, 3(25), 731.
50. Fick SE, Hijmans RJ (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Intl J Climatol*, 37(12), 4302-4315.
51. Kelso NV, Patterson T (2010). Introducing natural earth data-natureearthdata.com. *Geographia Technica*, 5, 82-89.
52. Hagerty CH, *et al.* (2016). Mapping snap bean pod and color traits, in a dry bean×snap bean recombinant inbred population. *J Am Soc Hortic Sci*, 141(2), 131-138.
53. Liljegren SJ, *et al.* (2004). Control of fruit patterning in *Arabidopsis* by INDEHISCENT. *Cell* 116, 843-853.

Figures

Fig. 1. Variation in PD-related structures in common bean. (A) Cross-section of the ventral suture of G12873, a wild Middle American bean. Wild beans show very high pod dehiscence (PD) and extensive lignified vascular sheath (VS) and fiber layer (LFL) deposition in pod walls. (B) In PD-susceptible domesticated dry beans (cv. ICA Bunsu shown), LFL deposition is reduced relative to wild types, indicating that these cells may be related to Middle American common bean domestication. (C) PD-resistant dry beans (cv. SXB 405 shown) are anatomically similar to PD-susceptible domesticated types (see B). (D) Stringless varieties (cv. Midas shown) display a reduction in VS lignification, including a reduction in secondary cell wall thickening. The LFL is absent in these varieties. Scale bars represent 100µm.

Fig. 2. Pod dehiscence (PD) QTL mapping based on three phenotyping methods. (A) Genome-wide and (B) Pv03-specific mapping results. All methods produced statistically significant results in the same region of chromosome Pv03. The significance threshold, determined by 1000 randomized permutations of the data, is shown as a black bar at LOD=5.80. The common bean ortholog of *Pdh1*, which regulates PD in soybean, is located between the most significant markers from QTL mapping (Table S1).

Fig. 3. GWAS of PD in independently domesticated common bean populations. (A) In the Middle American Diversity Panel (MDP), the most significant SNP is located 5.7kbp from the *PvPdh1* putative causal polymorphism. Pv06 and Pv08 also included loci of interest. (B) In the Andean Diversity Panel (ADP), chromosomes Pv03, Pv05, Pv08, and Pv09 include major regions of interest. SNPs located near *PvMYB26* (18) on Pv05 were highly significant. Horizontal red and blue lines indicate the Bonferroni-corrected significance threshold for an alpha of 0.01 and 0.05, respectively. Based on the proportion of pods dehiscing in a desiccator, with correction for population structure by PCA.

Fig. 4. The relationship between PD, ecogeographic race, and resistance alleles. (A) The first principal component of genetic data for the MDP separates race Durango (at left) from race Mesoamerica (at right). Members of race Durango have low susceptibility to PD relative to members of race Mesoamerica. Accessions are color coded by genotype at the GWAS peaks on Pv03 and Pv08. (B) A violin plot showing of the extent of PD by allele in the MDP. Accessions with these PD resistance loci have significantly lower levels of PD than accessions with neither allele. Letters “a” and “b” distinguish significantly different groups (Tukey HSD).

Fig. 5. *PvPdh1* is related to local adaptation and range expansion in common bean. PD is nearly absent in Race Durango, a group adapted to the hot, dry environments of northern Mexico, where environmental aridity exacerbates PD. The loss of function *PvPdh1* allele is nearly at fixation in this population. In contrast, race Mesoamerica is adapted to humid lowlands, where conditions mask PD susceptibility. PD has been selected against less strongly in this population, and the wild type *PvPdh1* predominates. For detailed information on the geographic distribution of these races, see Singh *et al.* (13).

Table 1. Summary of pod fiber or dehiscence QTLs, their genome locations, potential candidate genes, and homologies with other species

Chromosome or Linkage Group	Gene pool	Ecogeographic race, from 13; if available)	QTL location (bp, v1.0, from 8)	Potential candidate genes (when identified)	Source in <i>Phaseolus vulgaris</i>	Homologies in other species (when known)
Pv02	Andean	Nueva Granada	43,425,893-43,900,872	<i>PvIND</i>	(15, 52)	<i>Arabidopsis</i> : (53)
Pv03	Middle American	Durango	47,527,006-48,475,205	<i>PvPdh1</i> : dirigent family	This research	Soybean: (7)
Pv03	Andean		39,768,300-48,451,789	<i>NAC</i> family, C2H2 zinc finger	This research	Cowpea: (36)
Pv04	Middle American		42,310,662		(52)	
Pv05	Andean	Nueva Granada	35,000,893-39,497,309	<i>MYB26</i> , <i>MYB46</i>	(18); this research	Cowpea: (36, 39); <i>Arabidopsis</i> : (38)
Pv08	Andean & Middle American	Mesoamerica	330,345-9,215,942	<i>MYB</i> family, <i>WRKY</i> family, polygalacturonase	This research	Sorghum: (40); <i>Arabidopsis</i> : (41)
Pv09	Andean		29,587,741-37,450,759	<i>CESA7</i> , polygalacturonases	This research	Cowpea: (39)

Supporting Information

Genetic control of reduced pod dehiscence in domesticated common bean: Associations with range expansion and local aridity conditions

Travis A. Parker, Jorge C. Berny Mier y Teran, Antonia Palkovic, Judy Jernstedt, and
Paul Gepts

SI Methods

Microscopy. Pods of G12873 (wild, high dehiscence), ICA Bunsí (domesticated dry bean, dehiscence susceptible) SXB 405 (domesticated dry bean, dehiscence resistant), and Midas (domesticated snap bean, dehiscence susceptible) were Vibratome-sectioned to identify morphological differences that might be associated with PD. All sectioned pods were greenhouse-grown and harvested when pods were at full size with seeds filled, at the onset of pod color change. All sections were 100 micrometers thick and made in a transverse plane perpendicular to the fibers of interest. All sections were treated with Auramine O for at least 20 minutes. Fluorescence was visualized using an Olympus microscope.

RI population and phenotyping for pod dehiscence. A recombinant inbred (RI) population developed from a cross between ICA Bunsí (domesticated, PD-susceptible) and SXB 405 (domesticated, PD-resistant) was used for QTL mapping (1). The population (IxS) of 238 RILs was field-grown during the spring and summer of 2014. The spring planting was an un-replicated trial conducted in Coachella, California. At maturity, plots were visually evaluated for the presence or absence of PD, and the data were used as a phenotype for QTL mapping. During the summer of 2014, the RI population was grown in a replicated trial in Davis, California. At maturity, dried pods from 191 RILs were harvested from each plot; these were evaluated for susceptibility to PD by two methods. First, all pods were desiccated at 65°C for seven days, and then returned to room temperature for a minimum of seven additional days. The proportion of pods dehiscence in this process was recorded for each plot. Second, the amount of force required to induce pod fracture was measured using an Imada force measurement gauge (method modified from (2)). A bit mounted to the gauge was used to press the ventral side of each pod at the most apical seed, and the peak force required to cause fracture at the apical end of the pod beak was recorded. Force required for PD was normalized to account for small but significant differences between note-takers, and the standardized score was used for QTL mapping.

Genotyping. Genomic DNA was extracted from parents and RILs of the IxS population using a modified CTAB protocol. DNA quality was confirmed using a NanoDrop spectrophotometer. The IxS population was genotyped using the Illumina Infinium II BARCBean6K_3 BeadChip (3); 382 segregating SNPs were identified in the population. Primers spanning the transcribed sequence of Phvul.003G252100, a candidate gene underlying the major QTL identified in this study, were developed using the NCBI Primer-BLAST tool. Differences in the genomic sequence around

PvPDH1 exist between the Middle American and Andean gene pools, so variable PCR primers were used between the gene pools. *PvPdh1*ALL MA Forward (CATCTCCCCCATTTTCCCCC) and *PvPdh1*ALL Reverse (AACACGTGGAAGAGGAGGATT) were used for Middle American accessions, while *PvPdh1*ALL Andean Forward (CATCTCTCCCATTTTCTCCT) and *PvPdh1*ALL Reverse (AACACGTGGAAGAGGAGGATT) were used for Andean types. PCR conditions for this amplification included an initial denaturation at 95°C for 180s, 38 cycles of 95°C for 30s, 51°C for 30s, and 68°C for 60s, and a final elongation step of 68°C for 300s. PCR products were cleaned using a GeneJET PCR Purification Kit and sequenced at the UC DNA Sequencing Facility by Sanger sequencing.

QTL mapping. Composite interval mapping was conducted using the R package R/qtl (4). Field dehiscence, proportion dehiscing in a desiccator, and force measurements were separately used to identify PD QTLs marked by SNPs. The maximum LOD score of 1000 randomized permutations of the data was used as a significance threshold. Multiple QTL mapping was conducted using the scantwo function in R/qtl and by running the analysis with RILs subsetted by genotype at the most significant marker near *PvPdh1* on Pv03.

Syntenic mapping and expression. Candidate genes related to PD were identified in Phytozome 12 (5). Syntenic comparisons between common bean and soybean were made using the Legume Information System 2.0 (6); these were verified using available literature (7, 8). The CoGe SynMap (9) and LegumeIP 2.0 (6) syntenic tools were used to compare syntenic regions between *Arabidopsis* (Col-0, TAIR10), common bean (G19833, *Pvulgaris_V1.0_218*; (8)), and soybean (Williams 82, Release 1.1; (10)). For tree generation, the *PvPDH1* amino acid sequence was BLASTed against the *A. thaliana*, *G. max*, and *P. vulgaris* proteomes. Default Grishin settings were used to construct the distance matrix. A fast-minimum evolution tree (11) was generated based on a maximum sequence difference of 0.85. Gene expression from a variety of tissues and developmental stages were based on published data (12) and visualized in R.

Amino acid conservation analyses. The complete amino acid sequence of *PvPdh1* from accession G19833 was BLASTed against the NCBI proteome database. The CONstraint-Based multiple ALIGNment Tool (COBAL) (13) was used to align the most similar proteins known among several plant taxa and identify conserved residues. The Protein Variation Effect Analyzer (PROVEAN) software tool (14) was used to estimate the effect of mutations of interest.

Validation of the role of *PvPdh1* in a wider population. The Genetic Resources Information Network (GRIN) database of the National Plant Germplasm (NPGS) includes PD phenotype data for the genus *Phaseolus*. PD-susceptible and PD-resistant varieties from this pool were selected for validation of the role of *PvPdh1* in PD. A small number of varieties commonly grown at UC Davis with known PD phenotypes were also genotyped. Stringless snap bean varieties were specifically excluded from the analysis to avoid the epistatic effect of the *Stringless* (*St*) locus on PD. Genomic DNA was extracted using a modified CTAB method; amplification and Sanger sequencing of *PvPdh1* were conducted as described previously. Genotypes were separated into the Andean or Middle American gene pool based on an indel in the 3' UTR of

PvPdh1. This indel consistently predicted the gene pool in varieties of known ancestry. After sequencing, Middle American varieties were divided into groups based on amino acid at position 162 of *PvPdh1*. The degree of dehiscence between these groups was evaluated by student's t-test.

Validation of QTL mapping results using association mapping.

Two hundred and eight accessions of the Andean Diversity Panel (ADP) (15) were grown in Davis, CA during summer 2016. PD in the field, proportion dehiscing in a desiccator, and force required for fracture were recorded. Principal component analysis was conducted on SNP data for the population, and the results were used as covariates to account for population structure. Two hundred seventy-eight members of the BeanCAP Middle American Diversity Panel (MDP) (16) were phenotyped for PD by desiccation in 2017. Association mapping was conducted using GLM in TASSEL (17) via SNIPlay (18). All results were visualized using the qqman R package (19).

Precipitation map generation. Precipitation across the native range of Middle American beans was mapped in QGIS 2.18.19 using data from worldclim2 (20). National boundaries and coastlines were added using shapefiles available through Natural Earth (21). USGS topographical global raster data grids were also used to improve the visualization of coastlines (https://topotools.cr.usgs.gov/gmted_viewer/gmted2010_global_grids.php).

Supporting Information

Supplemental Figures

Fig. S1. Phenotyping methods. PD was evaluated by (A) visual inspection of PD in the field, (B) proportion of dehiscing pods in a desiccator (none dehiscing in this sample), and (C) force required to induce fracture with a force measurement gauge.

Fig. S2. Phenotyping distributions in the ICA Bunsu/SXB 405 RI population. (A) Presence/absence of PD in the field. (B) The proportion of pods dehiscing after desiccation. (C) The force required for pod fracture. Force measurements resulted in a bimodal distribution, indicating that a single large-effect gene was responsible for much of the population's variation. "Frequency" represents the number of RILs falling into each bin.

Fig. S3. Correlations between phenotyping methods in the IxS RI population. RI lines are color coded by genotype at the *PvPdh1* locus. The numbers in the lower left panels indicate the correlation coefficients between those methods. PDS=PD susceptible, PDR=PD resistant.

Fig. S4. A rooted neighbor joining tree based on sequence of GmPDH1, PHAVU_003G252100g, and the most similar dirigent proteins of *G. max* and *P. vulgaris* in the NCBI database. GmPDH1 and PHAVU_003G252100g form a clade among all the proteins of these species, supporting their orthology. Tree derived from a Grishin protein distance matrix and rooted using 12 distantly related dirigent-like proteins.

Fig. S5. Expression of Phvul.003G252100.1 (*PvPdh1*) is unique to pods in *P. vulgaris* cv. Negro Jamapa. This pattern is extremely unusual even among homologs of *Arabidopsis* genes affecting PD. YL- Fully expanded 2nd trifoliate leaf tissue from fertilized plants L5- Leaf tissue collected 5 days after plants were inoculated with effective rhizobium; LF- Leaf tissue from fertilized plants collected at the same time of LE and LI; LE- Leaf tissue collected 21 days after plants were inoculated with effective rhizobium LI- Leaf tissue collected 21 days after plants were inoculated with ineffective rhizobium; YS- All stem internodes above the cotyledon collected at the 2nd trifoliate stage; ST- Shoot tip, including the apical meristem, collected at the 2nd trifoliate stage; FY- Young flowers, collected prior to floral emergence; PY- Young pods, collected 1 to 4 days after floral senescence. Samples contain developing embryos at globular stage PH- Pods approximately 9cm long, associated with seeds at heart stage (pod only); P1- Pods between 10 and 11 cm long, associated with stage 1 seeds (pod only); P2- Pods between 12 and 13 cm long associated with stage 2 seeds (pod only); SH- Heart stage seeds, between 3 and 4 mm across and approximately 7 mg S1- Stage 1 seeds, between 6 and 7 mm across and approximately 50 mg; S2- Stage 2 seeds, between 8 and 10 mm across and between 140 and 150 mg; RT- Root tips, 0.5 cm of tissue, collected from fertilized plants at 2nd trifoliate stage of development; YR- Whole roots, including root tips, collected at the 2nd trifoliate stage of development; R5- Whole roots separated from 5 day old pre-fixing nodules; RF- Whole roots from fertilized plants collected at the same time as RE and RI; RE- Whole roots separated from fix+ nodules collected 21 days after inoculation; RI- Whole roots separated from fix- nodules collected 21 days after inoculation; N5- Pre-fixing (effective) nodules collected 5 days after inoculation ;NE- Effectively fixing nodules collected 21 days after inoculation; NI- Ineffectively fixing nodules collected 21 days after inoculation. From O'Rourke *et al.* (12).

A.

Bunsi 300	CATTTACGACACCAAGAACACTTACACTGCGTACCTCGGATTCAACTTTGCTCTCAATAG	359
SXB 405 455	CATTTACGACACCAAGAACACTTACACTGCGTACCTCGGATTCAACTTTGCTCTCAATAG	396
Bunsi 360	CGCAAATCATCAGGGAACCATCACCGTCGCTGGAGCTGACCCACCTTGAAGAAGACCAG	419
SXB 405 395	CGCAAATCATCAGGGAACCATCACCGTCGCTGGAGCTGACCCACCTTGAAGAAGACCAG	336
Bunsi 420	AGACATCTCAGTCACAGGTGGCACTGGAGATTTCTTCATGCATAGAGGAATCGCCACCAT	479
SXB 405 335	AGACATCTCAGTCACAGGTGGCACTGGAGATTTCTTCATGCATAGAGGAATCGCCACCAT	276
Bunsi 480	CATGACCGATGCTTTTGAAGGTGATGTTTATTTCCGCCTTCGTGTTGAAATCAAGTTTTA	539
SXB 405 275	CATGAACGATGCTTTTGAAGGTGATGTTTATTTCCGCCTTCGTGTTGAAATCAAGTTTTA	216
Bunsi 540	TGAGTGTGGTGatataataagtatatatatatGCACGCACACACAAAGTTACAGTAAATA	599
SXB 405 215	TGAGTGTGGTGATATATAAGTATATATATGCACGCACACACAAAGTTACAGTAAATA	156
Bunsi 600	AAGATGTCTGTTTCTGTCAACAGCCTCGTCTAACAGTTACTCTTATTACTATATTAATA	659
SXB 405 155	AAGATGTCTGTTTCTGTCAACAGCCTCGTCTAACAGTTACTCTTATTACTATATTAATA	96
Bunsi 660	AATATGGCTGTTTCTGTCTACTGCCTCTATAGTATGTGGCTTGCTTGTGTAATAACGTT	719
SXB 405 95	AATATGGCTGTTTCTGTCTACAGCCTCTATAGTATGTGGCTTGCTTGTGTAATAACGTT	36

B.

Bunsi 1	MGAKVTLFVFFTFALCSTFPLQRKQYAPCKHLVLFHDIYNGRNALNATSAIIAAPQG	60
SXB 405 1	MGAKVTLFVFFTFALCSTFPLQRKQYAPCKHLVLFHDIYNGRNALNATSAIIAAPQG	60
Bunsi 61	ANLTKLANNFHFGLNVFDDPVTLDNNLHSEPVGRAQGFYIYDTKNTYTAYLGFNFALNS	120
SXB 405 61	ANLTKLANNFHFGLNVFDDPVTLDNNLHSEPVGRAQGFYIYDTKNTYTAYLGFNFALNS	120
Bunsi 121	ANHQTITVAGADPTLKKTRDISVTGGTGDFFMHARGIATIMDAFEGDVYFRLRVEIKFY	180
SXB 405 121	ANHQTITVAGADPTLKKTRDISVTGGTGDFFMHARGIATIMDAFEGDVYFRLRVEIKFY	180
Bunsi 181	ECW 183	
SXB 405 181	ECW 183	

Fig. S6. A polymorphism exists in *PvPdh1* between the parents of the RI population. A) At position 485 of the CDS of *PvPdh1*, there is a C/A polymorphism between ICA Bunsi and SXB 405. This nonsynonymous substitution leads to B) a threonine/asparagine polymorphism at position 162 in the amino acid sequence of the protein products.

Fig. S7. Dehiscence in Middle American GRIN NPGS accessions. A) In individuals with an asparagine at position 162, dehiscence resistance predominates. B) In individuals with a wild-type threonine at position 162, dehiscence susceptibility predominates. Accessions were phenotyped by GRIN NPGS and genotyped by Sanger sequencing of *PvPdh1*.

A.

Query_150380 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [1] 184
 XP_007156024 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 BAP91518 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 XP_014508512 134 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 182
 XP_0176131817 134 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 182
 XP_004503651 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 181
 AFK40037 133 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 XP_013447143 137 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 GAU32639 136 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 XP_017441955 133 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [2] 183
 XP_019420874 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 XP_014634029 126 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 174
 XP_007160128 128 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 176
 XP_007040198 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 XP_002297990 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 XP_004298854 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_002297995 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 AC767169 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 XP_011022578 135 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 AF284548 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 AF284547 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 XP_009334474 137 TLKK-YRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 XP_011026952 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 XP_006439431 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 AC267121 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 XP_015806528 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 XP_008388884 137 TLKK-YRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 KXP58666 135 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 183
 AF284545 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 K0076393 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 AFK42402 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 XP_006476461 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 XP_012470512 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 XP_008238999 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 AC767120 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 XP_015577413 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 GAU32648 124 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 172
 GAU32641 124 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 172
 AAO25355 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 XP_016741331 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 AF284546 131 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 179
 P13248 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 XP_017234471 147 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 195
 K2086017 147 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [50] 245
 XP_002276430 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 XP_002276448 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 AAA33662 127 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 175
 KVM95339 134 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 182
 XP_008451985 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 XP_010162707 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 XP_011022577 150 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 198
 XP_012087199 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 OLV94014 103 TLKK-TRGSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 151
 XP_002308536 126 TLKK-SKDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 198
 K2086017 126 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 161
 XP_004164540 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 191
 GAU32643 124 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 173
 EE51229 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [1] 190
 XP_007209654 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 AB24_3 116 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 164
 A317752 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_012086792 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 KX111359 136 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 184
 A3177512 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_016741354 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 K0072356 117 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 165
 XP_017620779 139 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 XP_018055805 137 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 185
 XP_012470511 139 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 AAF25363 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 AAF25364 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 AAF25361 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 XP_010055806 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 A3177523 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_010251577 140 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 188
 KVM95347 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 XP_006439432 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 AAF25366 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 AAF25365 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 XP_010106579 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 XP_017972306 140 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [3] 191
 XP_018809828 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 XP_010106578 143 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 191
 AAF25368 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 XP_006343908 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 AAF25362 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 ACJ40439 151 TLKK-YRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [13] 212
 XP_004298853 143 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 191
 XP_018829049 142 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 190
 XP_016741587 139 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 XP_002297997 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 AB052129 152 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW [24] 224
 XP_006826467 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_002272144 144 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 192
 OAF59689 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 OAF59685 139 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 OAF59686 139 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 187
 XP_015085509 141 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 189
 XP_002508091 138 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 186
 XP_019225306 143 TLKK-TRDSVVGTTGDF FHHGATITDAFEGGVYRLRVEIKFYECW 191

B.

Query_274211 157 IATITTHDAFEGGVYRLRVEIKFYECW [5] 184
 XP_002981852 154 IARLSSTQSISGVTFVLKVVTLF [2] 175
 XP_002982822 165 IARLSSTQSISGVTFVLKVVTLF [2] 189
 XP_002976809 2467 TANITTHHSDGDTFTVMTSTY [4] 2493
 XP_002980538 120 IANLVTHHSDGDTFTVMTSTY [4] 138
 XP_002971526 138 IATITTHHSDGDTFTVMTSTY [2] 162
 XP_002991608 138 IATITTHHSDGDTFTVMTSTY [2] 162
 XP_002974933 120 YAIISSTKMPPLSVL-LDVQVFL 142
 XP_002993354 152 YAIIVTAVDHGLVYVLDVQVFL [2] 177
 XP_002977365 112 YALTSSTGAEPGLIATYEVHLVR [1] 136
 XP_002977900 121 YAAVVT----- 128
 XP_002974944 112 YALTSSTGAEPGLIATYEVHL-- 133
 XP_002991290 123 YAIIVT----- 128
 XP_002972740 119 YAIIVT----- 124
 XP_002988690 141 YATITTHASAGGSVILEIDVKVSH 164
 XP_002986379 141 YATITTHASAGGSVILEIDVKVSH 164
 XP_002968306 141 FAVITTSNTNRRATTVGFDVTI [QV] 164
 XP_002967407 141 FAVITTSNTNRRATTVGFDVTI [QV] 164
 XP_002966352 119 FAVITTSNVN SAVLHNVFTQH 142
 XP_002978209 119 FAVITTSNVN SAVLHNVFTQH 142
 XP_002985735 150 FASSSTVDITRTNTTYSVDI NVFN [4] 177
 XP_002974438 150 FASSSTVDITRTNTTYSVDI NVFN [4] 177

C.

D.

670
 671 **Fig. S8.** The threonine at position 162 is a highly conserved component of the active site for
 672 dirigent-like genes. (A) Of the 100 most similar proteins to PvPDH1 in the NCBI database, 99
 673 have a threonine at the aligned position, indicating it is vital for protein functionality. The one
 674 exception is a gene from *Trifolium subterraneum*, which places pods underground and the gene
 675 may be undergoing gene decay. (B) The 19 most similar dirigent-like genes from *Selaginella*
 676 *moellendorffii* have a threonine at this position, indicating that the residue has been very
 677 strongly conserved for over 400 million years (22, 23). (C) In the closely related protein PsDIR6,
 678 the homologous threonine (T163) is an important component of the active site (from (24)). (D)
 679 Targeted mutagenesis of the equivalent residue (T166) in a closely related *Arabidopsis* protein
 680 showed that substituting the threonine with a valine led to a complete loss of gene function
 681 (from (25)).

Fig. S9. GWAS of pod dehiscence (PD) in Race Mesoamerica (MDP, PC1>50) using GLM in SNIPlay/TASSEL. Pv08 was most significantly associated with variation in PD, although no SNPs achieved significance in this smaller population, the most significant SNPs were located in an overlapping interval on Pv08 as a major QTL of the ADP, indicating that the same gene may be responsible for the variation across populations. MAF threshold = 0.1.

Supplemental Tables

Table S1. Co-segregation between dehiscence phenotype and position 162 in *PvPdh1*. The 11 RILs with recombination between the flanking markers from QTL mapping showed perfect correspondence between phenotype and genotype at this position.

RIL	Phenotype	<i>PvPdh1</i> CDS residue 485	PvPDH1 peptide residue 162
013	Dehiscent	Cytosine	Threonine
036	Dehiscent	Cytosine	Threonine
052	Dehiscent	Cytosine	Threonine
094	Dehiscent	Cytosine	Threonine
096	Dehiscent	Cytosine	Threonine
006	Non-dehiscent	Adenine	Asparagine
033	Non-dehiscent	Adenine	Asparagine
065	Non-dehiscent	Adenine	Asparagine
101	Non-dehiscent	Adenine	Asparagine
117	Non-dehiscent	Adenine	Asparagine
154	Non-dehiscent	Adenine	Asparagine

Table S2. Sequencing of *PvPdh1* in several species of wild and domesticated *Phaseolus*

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 477040	<i>P.</i> <i>acutifolius</i>	PvPdh1ALL	C	NA	W6	High shattering		United States	Wild
PI 195359	<i>P.</i> <i>dumosus</i>	PvPdh1ALL	C	NA	W6	Shattering resistant		Guatemala	Landrace
PI 311194	<i>P.</i> <i>dumosus</i>	PvPdh1ALL	C	NA	W6	Shattering resistant		Guatemala	Landrace
PI 326055	<i>P.</i> <i>dumosus</i>	PvPdh1ALL	C	NA	W6	Shattering resistant		Venezuela	Landrace
PI 535379	<i>P. hintonii</i>	PvPdh1ALL	C	NA	W6	Full and easily shattered		Mexico	Wild
PI 257418	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		Argentina	Landrace
PI 264603	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		Costa Rica	Wild
PI 264606	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		Costa Rica	Wild
PI 264607	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		Costa Rica	Wild
PI 264610	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		Costa Rica	Wild
PI 347796	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Shattering resistant		United States	Cultivated
PI 347797	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		United States	Cultivated
PI 347811	<i>P. lunatus</i>	PvPdh1ALL	C	NA	W6	Moderate to high shattering		United States	Cultivated
Canario 707	<i>P. vulgaris</i>	PvPdh1Andes	C	Andean	Davis	Shattering resistant		United States	Cultivated
Etna	<i>P. vulgaris</i>	PvPdh1Andes	C	Andean	Davis	Moderate to high shatter		United States	Cultivated
PI 136701	<i>P. vulgaris</i>	PvPdh1ALL	C	Andean	W6	No shattering		Canada	Cultivated
PI 136745	<i>P. vulgaris</i>	PvPdh1ALL	C	Andean	W6	No shattering		Canada	Cultivar
PI 161952	<i>P. vulgaris</i>	PvPdh1ALL	C	Andean	W6	No shattering		Belgium	Cultivated
PI 167105	<i>P. vulgaris</i>	PvPdh1ALL	C	Andean	W6	No shattering		Turkey	Cultivated
PI 169804	<i>P. vulgaris</i>	PvPdh1ALL	C	Andean	W6	No shattering		Turkey	Cultivated

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 180734	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Germany	Cultivar
PI 226859	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Spain	Cultivar
PI 226879	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Ukraine	Cultivated
PI 226934	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Germany	Cultivated
PI 234257	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Turkey	Cultivated
PI 278689	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		United States	Cultivated
PI 279818	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Canada	Cultivar
PI 281596	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Italy	Cultivated
PI 289355	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Hungary	Cultivar
PI 304813	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shattering		United States	Breeding material
PI 304824	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		United States	Breeding material
PI 309766	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		Mexico	Landrace
PI 313270	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter	Full and easily shattered	Mexico	Landrace
PI 324628	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Hungary	Cultivar
PI 325630	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		Mexico	Landrace
PI 353500	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		China	Cultivated
PI 353531	P. vulgaris	PvPdh1ALL	C	Andean	W6	No shattering		China	Cultivated
PI 361284	P. vulgaris	PvPdh1ALL	C	Andean	W6	Full and easily shattered		India	Uncertain
PI 433622	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		United States	Cultivar

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 439555	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		Netherlands	Landrace
PI 439571	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		Netherlands	Landrace
PI 476686	P. vulgaris	PvPdh1ALL	C	Andean	W6	Full and easily shattered		Mexico	Cultivated
PI 533322	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		Mexico	Cultivated
PI 533323	P. vulgaris	PvPdh1ALL	C	Andean	W6	Moderate to high shatter		Mexico	Cultivated
PI 632356	P. vulgaris	PvPdh1ALL	C	Andean	W6	Full and easily shattered		United States	Cultivated
Tiger's Eye	P. vulgaris	PvPdh1Andes	C	Andean	Davis	Moderate to high shatter		United States	Landrace
UC 0801	P. vulgaris	PvPdh1Andes	C	Andean	Davis	No shattering		United States	Cultivated
UC Holstein	P. vulgaris	PvPdh1Andes	C	Andean	Davis	No shattering		United States	Cultivated
UC Jacob's Cattle	P. vulgaris	PvPdh1ALL	C	Andean	Davis	No shattering		United States	Cultivated
Matterhorn	P. vulgaris	PvPdh1ALL	A	Middle American	Davis	No shattering		United States	Cultivar
PI 136722	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Canada	Cultivated
PI 169722	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Turkey	Cultivated
PI 169725	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Turkey	Landrace
PI 169731	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Turkey	Cultivated

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 175866	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Turkey	Cultivated
PI 289341	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 289342	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 289358	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 291368	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		China	Cultivated
PI 309729	P. vulgaris	PvPdh1ALL	A	Middle American	W6	Moderate to high shatter		Mexico	Landrace
PI 311907	P. vulgaris	PvPdh1ALL	A	Middle American	W6	Moderate to high shattering		Mexico	Landrace
PI 313309	P. vulgaris	PvPdh1ALL	A	Middle American	W6	Full and easily shattered		Mexico	Landrace
PI 313313	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Mexico	Landrace
PI 324580	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 324593	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 324604	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Hungary	Cultivar
PI 353505	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		China	Cultivated
Pink 9634	P. vulgaris	PvPdh1ALL	A	Middle American	Davis	No shattering		United States	Cultivar

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
W6 27625	P. vulgaris	PvPdh1ALL	A	Middle American	W6	Full and easily shattered		Mexico	Labeled as wild, but has solid black seeds that weigh 38g/100
W6 9719	P. vulgaris	PvPdh1ALL	A	Middle American	W6	No shattering		Russian Federation	Cultivar
Zuni Gold	P. vulgaris	PvPdh1ALL	A	Middle American	Davis	No shattering		United States	Landrace
SXB 405	P. vulgaris	PvPdh1ALL	A	Middle American	Davis	Non-shattering			Cultivated
PI 194572	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Full and easily shattered		Guatemala	Landrace
PI 227115	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Australia	Cultivated
PI 262977	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Netherlands	Cultivated
PI 282057	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Chile	Cultivar
PI 307820	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shattering		El Salvador	Landrace
PI 310586	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Honduras	Cultivated
PI 311843	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Shattering resistant		Guatemala	Landrace
PI 313572	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Full and easily shattered		Colombia	Cultivated

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 324618	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Hungary	Cultivar
PI 358218	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Macedonia	Cultivar
PI 476681	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Mexico	Cultivated
PI 476688	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Mexico	Cultivated
PI 476694	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Mexico	Cultivated
PI 476695	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Mexico	Cultivated
PI 476701	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Mexico	Cultivated
PI 476703	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Full and easily shattered		Mexico	Cultivated
PI 476707	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Guatemala	Cultivated
PI 476709	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Guatemala	Cultivated
PI 476710	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Guatemala	Cultivated
PI 476730	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		Guatemala	Cultivated
PI 476737	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shatter		Guatemala	Cultivated
PI 642946	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Full and easily shattered		United States	Cultivated

Accession	Species	Primer	Position 485 C-->A; Threonine - -> Asparagine	Middle American or Andean based on duplication of repetitive element near position 648	Source/ station	Shattering	Shattering_2	Country of origin	Improvement status
PI 642947	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Moderate to high shattering	Full and easily shattered	United States	Cultivated
PI 661740	P. vulgaris	PvPdh1ALL	C	Middle American	W6	Full and easily shattered		Honduras	Landrace
W6 11336	P. vulgaris	PvPdh1ALL	C	Middle American	W6	No shattering		China	Cultivar
G12873	P. vulgaris	PvPdh1ALL	C	Middle American	Davis	Full and easily shattered		Mexico	Wild
ICA Bunsí	P. vulgaris	PvPdh1ALL	C	Middle American	Davis	Shattering		Colombia	Cultivated
PI 339544	P. vulgaris	PvPdh1ALL	"M" ambiguity code. Excluded from analysis	Middle American	W6	No shattering		Turkey	Landrace
PI 226900	P. vulgaris	PvPdh1ALL	C	NA	W6	No shattering		Spain	Cultivated
PI 433561	P. vulgaris	PvPdh1ALL	C	NA	W6	Moderate to high shatter		United States	Cultivar
G23584	P. debouckii	PvPdh1Andes	C	Andean	Davis	Shattering		Peru	Wild

692

Table S3. Synteny near *Pdh1* in *G. max* and *P. vulgaris* – sharing of gene models.

<i>Glycine max</i>		<i>Phaseolus vulgaris</i>	
Gene model	Predicted gene function	Gene model	Predicted gene function
Glyma16g25490.1	Proline-rich extensin-like receptor kinase 4	Phvul.003G252900.1	Proline-rich extensin-like receptor kinase 4
Glyma16g25500.1	Leucine-rich repeat (LRR) family protein	Phvul.003G252700.1	Leucine-rich repeat (LRR) family protein
Glyma16g25530.2	Plant invertase/pectin methylesterase inhibitor superfamily protein		
Glyma16g25540.1	Major facilitator superfamily protein	Phvul.003G252500.1	Major facilitator superfamily protein
Glyma16g25550.1	C2H2-like zinc finger protein	Phvul.003G252400.1	C2H2-like zinc finger protein
Glyma16g25560.1	Protein phosphatase 2C family protein	Phvul.003G252300.1	Protein phosphatase 2C family protein
Glyma16g25570.1	GRAS family transcription factor	Phvul.003G252200.1	GRAS family transcription factor
Glyma16g25580.1	Disease resistance-responsive (dirigent-like protein) family protein	Phvul.003G252100.1	Disease resistance-responsive (dirigent-like protein) family protein
Glyma16g25600.1	G-box binding factor 1		
Glyma16g25600.5	G-box binding factor 1	Phvul.003G252000.1	G-box binding factor 1
Glyma16g25600.4	G-box binding factor 1		
Glyma16g25600.6	G-box binding factor 1		
Glyma16g25600.7	G-box binding factor 1		
Glyma16g25611.1	Protein kinase superfamily protein	Phvul.003G251900.1	Protein kinase superfamily protein
Glyma16g25620.1	geranylgeranyl pyrophosphate synthase 1	Phvul.003G251500.1	geranylgeranyl pyrophosphate synthase 1
Glyma16g25650.1	Protein of unknown function (DUF677)		
Glyma16g25660.1			
Glyma16g25670.1			
Glyma16g25680.2	Protein of unknown function (DUF607)	Phvul.003G251400.1	Protein of unknown function (DUF607)

Glyma16g25690.1	GNS1/SUR4 membrane protein family	Phvul.003G251300.1	GNS1/SUR4 membrane protein family
Glyma16g25700.1	PQ-loop repeat family protein / transmembrane family protein	Phvul.003G251100.1	PQ-loop repeat family protein / transmembrane family protein
Glyma16g25710.2	Pectin lyase-like superfamily protein		
Glyma16g25720.1	calmodulin-like 41	Phvul.003G251000.1	calmodulin-like 41
Glyma16g25740.2	vacuolar ATP synthase subunit H family protein	Phvul.003G250900.1	vacuolar ATP synthase subunit H family protein

693

Table S4. PD after desiccation, by market class, gene pool, and ecogeographic race

Market class	Gene pool	Race	Mean PD (%)	Median PD (%)	Standard deviation (%)	n
Cranberry	Andean	Nueva Granada	41.43	46.29	29.86	24
Kidney	Andean	Nueva Granada	21.09	13.89	18.32	43
Purple speck/mottled	Andean	Nueva Granada	3.11	0	5.84	17
Great northern	Middle American	Durango	0.94	0	2.12	31
Pink	Middle American	Durango	2.48	0	6.37	23
Pinto	Middle American	Durango	0.74	0	2.38	93
Black	Middle American	Mesoamerica	17.63	19	13.22	43
Navy/small white	Middle American	Mesoamerica	15.2	8.5	16.62	46
Red/small red	Middle American	Variable	9.59	4	14.7	29

References

- Assefa T, et al. (2013). Pod harvest index as a selection criterion to improve drought resistance in white pea bean. *Field Crops Res*, 148, 24-33.
- Dong Y, et al. (2014). Pod shattering resistance associated with domestication is mediated by a NAC gene in soybean. *Nat Comm*, 5, 3352.
- Song Q, et al. (2015). SNP assay development for linkage map construction, anchoring whole-genome sequence, and other genetic and genomic applications in common bean. *G3 (Bethesda)*, 5(11), 2285-2290.
- Broman KW, Wu H, Sen S, Churchill, GA (2003). "R/qtl: QTL mapping in experimental crosses." *Bioinformatics* 19(7), 889-890.
- Goodstein DM, et al. (2011). Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res*, 40(D1), D1178-D1186.
- Li J, Dai X, Zhuang Z, Zhao PX (2016). LegumeIP 2.0—a platform for the study of gene function and genome evolution in legumes. *Nucleic Acids Res*, 44(D1), D1189-D1194.
- McClellan PE, Mamidi S., McConnell M, Chikara S, Lee R (2010). Synteny mapping between common bean and soybean reveals extensive blocks of shared loci. *BMC Genomics*, 11(1), 184.
- Schmutz J, et al. (2014). A reference genome for common bean and genome-wide analysis of dual domestications. *Nature Genetics*, 46(7), 707.
- Lyons E, Pedersen B, Kane J, Freeling M (2008). The value of nonmodel genomes and an example using SynMap within CoGe to dissect the hexaploidy that predates the rosids. *Trop Plant Biol*, 1(3-4), 181-190.
- Schmutz J, et al. (2010). Genome sequence of the palaeopolyploid soybean. *Nature*, 463(7278), 178.
- Desper R, Gascuel O (2004). Theoretical foundation of the balanced minimum evolution method of phylogenetic inference and its relationship to weighted least-squares tree fitting. *Mol Biol Evol*, 21(3), 587-598.
- O'Rourke JA et al. (2014). An RNA-Seq based gene expression atlas of the common bean. *BMC Genomics*, 15(1), 866.
- Papadopoulos JS, Agarwala R (2007). COBALT: constraint-based alignment tool for multiple protein sequences. *Bioinformatics*, 23(9), 1073-1079.

14. Choi Y, Chan AP. "PROVEAN web server: a tool to predict the functional effect of amino acid substitutions and indels." *Bioinformatics* 31.16 (2015): 2745-2747.
15. Cichy KA et al. (2015). A *Phaseolus vulgaris* diversity panel for Andean bean improvement. *Crop Sci*, 55(5), 2149-2160.
16. Moghaddam SM, et al. (2016). Genome-wide association study identifies candidate loci underlying agronomic traits in a Middle American diversity panel of common bean. *Plant Genome*, 9(3).
17. Bradbury PJ, et al. (2007). TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*, 23(19), 2633-2635.
18. Dereeper A, et al. (2011). SNIPlay: a web-based tool for detection, management and analysis of SNPs. Application to grapevine diversity projects. *BMC Bioinformatics*, 12(1), 134.
19. Turner SD (2018). qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *J Open Source Softw*, 3(25), 731.
20. Fick SE, Hijmans RJ (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Intl J Climatol*, 37(12), 4302-4315.
21. Kelso NV, Patterson T (2010). Introducing natural earth data-natureearthdata. com. *Geographia Technica*, 5, 82-89.
22. Soltis PS, Soltis DE, Savolainen V, Crane PR, Barraclough TG (2002). Rate heterogeneity among lineages of tracheophytes: integration of molecular and fossil data and evidence for molecular living fossils. *Proc Natl Acad Sci*, 99(7), 4430-4435.
23. Zimmer A, et al. (2007). Dating the early evolution of plants: detection and molecular clock analyses of orthologs. *Mol Genet Genomics*, 278(4), 393-402.
24. Kim KW, et al. (2015). Trimeric structure of (+)-pinorensinol-forming dirigent protein at 1.95 Å resolution with three isolated active sites. *J. Biol Chem*, 290(3), 1308-1318.
25. Gasper R, et al. (2016). Dirigent protein mode of action revealed by the crystal structure of AtDIR6. *Plant Physiol*, pp-01281.