1 Title: Risks to pollinators from different land-use transitions: bee species' responses to
2 agricultural expansion show strong phylogenetic signal
3 Running title: Phylogenetic signal in bee sensitivity
4
5 Adriana De Palma^{*1}, Michael Kuhlmann^{2,1}, William D. Pearse³, Emma Flynn⁴, Stuart P.M.
6 Roberts⁵, Simon G. Potts⁵, Andy Purvis^{1,4}
7 1 Department of Life Sciences, Natural History Museum, London, SW7 5BD, UK
8 2 Zoological Museum, University of Kiel, Kiel, Germany
9 3 Ecology Center and Department of Biology, Utah State University, Logan, UT, USA
10 4 Department of Life Sciences, Imperial College London, Silwood Park, Ascot, SL5 7PY, UK
11 5 Centre for Agri-Environmental Research, School of Agriculture, Policy and Development,
12 The University of Reading, Reading, UK
13
14 *Corresponding author: A.De-Palma@nhm.ac.uk
15

16 **Abstract**

17 Bee species worldwide are facing a future of further land-use change and
 18 intensification. Populations of closely-related species with similar ecological
 19 characteristics are likely to respond similarly to such pressures. Such phylogenetic
 20 signal in species' responses could undermine the stability of pollination services in
 21 agricultural and natural systems. We use abundance data from a global compilation of
 22 bee assemblages in different land uses to assess the sensitivity of 573 bee species to
 23 agricultural expansion, intensification and urbanization; and combine the results with the
 24 Bee Tree of Life to assess phylogenetic signal. In addition, we assess whether variation
 25 in species' sensitivity to land-use change is better explained by phylogenetic or
 26 available functional trait differences. Bee species show strong phylogenetic signal in
 27 sensitivity to agricultural land expansion but only a weak signal in sensitivity to
 28 agricultural intensification and urbanisation. Sensitivities were usually best explained
 29 by a combination of functional and phylogenetic distances. This finding suggests that
 30 the commonly-recorded traits, despite being meaningful as functional response traits,
 31 do not capture all important determinants of bee species' vulnerability or resistance.
 32 However, it also suggests that model-based predictions of the sensitivity of poorly
 33 known species may be sufficient to help guide conservation efforts.

34 **Keywords**

35 Land-use change, pollination services, correlates of risk, species decline,
 36 conservation

38 **Introduction**

39 Land-use change is the most important driver of present-day terrestrial biodiversity
 40 loss [1,2] and is predicted to cause continued damage in the future [3,4]. Most
 41 scenarios incorporate further loss of natural and semi-natural land, driven by
 42 agricultural expansion and urbanisation, as well as increased degradation due to
 43 agricultural intensification [5]. The effect such global change will have on bee
 44 populations could have serious consequences for crop pollination worldwide [6] and—
 45 as bee species are the most important pollinators of flowering plants globally [7]—for
 46 wild plant populations too [8].

47 Predicting likely effects of land-use change on ecosystem functions such as

pollination requires an understanding of whether and how species' sensitivities to land-use change are shaped by their functional traits and evolutionary history [9,10]. Traits are known to predict responses of European bee populations to land use and related pressures, but associations are inconsistent [11–13]. In bumblebees, long-term population declines in North America show phylogenetic signal (i.e., closer relatives tend to show more similar trends) [14], as does their global conservation status as assessed by the IUCN Red List [15], but information on other bees is limited. The relative usefulness of phylogeny and available functional trait data in predicting species' responses is an open question. Although phylogenetic relatedness is in this context only a proxy for similarity of species' functional responses [16], it may outperform functional trait data if responses are driven by a broader set of (phylogenetically patterned) ecological differences than are captured by available trait data.

A strong phylogenetic signal in species' responses to a particular pressure increases the risk that the pressure could impact ecosystem functioning [16], as the order of species losses can strongly influence pollination networks [17]. Phylogenetic diversity and redundancy of bee communities may also decline; although debate is ongoing [18,19], phylogenetic diversity can be important for ecosystem functioning and stability [20]. The protection of phylogenetic diversity (by conserving those clades that are most vulnerable) may also be important for maintaining robust pollination networks: high phylogenetic diversity can correlate with interaction diversity [21], as closely-related species also tend to share resources [22,23]. The strength of phylogenetic signal can vary among pressures [e.g. 24], meaning that different land-use transitions could carry different risks of pollination impairment that might not be apparent from the effects on species diversity or overall abundance. Nonetheless, so far there has been no exploration of the phylogenetic pattern of bee sensitivities to particular threats such as agricultural expansion, agricultural intensification and urbanization.

Understanding which disturbances prompt the most phylogenetically patterned response may therefore indicate where ecosystem services might be at greatest risk and so where conservation action will be most important. However, it is unclear whether responses to different pressures will all show phylogenetic signal [16] as the links between ecological traits and the response to human impacts are not always straightforward [11,16]. We use data from 86 studies and 2,599 sites to assess the sensitivity of 573 bee species (from 96 genera) to human-dominated land uses, including agriculture and urban areas, and to increasing agricultural intensity. We quantify the strength of phylogenetic signal

83 in these sensitivity estimates, accounting for uncertainty in sensitivities. Using trait data
84 for a subset of 143 bee species, we assess whether phylogenetic differences are better
85 able than functional traits to explain variation in species' sensitivities.

86 **Materials and methods**

87 Site-level data on bee abundance and occurrence were extracted from the PREDICTS
88 database [25] and De Palma et al. [26]. We refer to each survey of multiple sites that
89 used the same sampling method within the same season and the same country as a
90 'study'. Differences in sampling effort among sites within a study were corrected for
91 where necessary, assuming that recorded abundance increases linearly with sampling
92 effort [26]. Within each study, we recorded any blocked or split-plot design. See
93 Appendix Table S1.1 for a list of data sources (some containing multiple studies).

94 The major land use and land-use intensity at each site was recorded based on
95 information in the associated paper, supplementary files or other information provided
96 by the original authors, as described in [27]. Land use was classified as primary
97 vegetation, secondary vegetation, cropland, plantation forest, pasture or urban. The
98 use-intensity scale assesses human disturbance on a three-level qualitative scale within
99 each land use (minimal, light and intense) [28]. For instance, intensively-used cropland
100 includes monocultures with characteristic features of intensification (e.g., large fields,
101 high levels of external inputs, irrigation and mechanization); lightly-used cropland would
102 show some, but not all, or the same features; whereas minimally-used cropland would
103 include small mixed-cropping fields with little or no external input, irrigation or
104 mechanization.

105 *Species sensitivity*

106 We focus on four land-use transitions (Table 1). The first, with the largest sample size,
107 compares species abundances in natural/semi-natural land (i.e., primary or secondary
108 vegetation) with all human-dominated land uses combined (i.e., all other land-use
109 classes) as a recent synthesis showed that, in terms of species composition,
110 assemblages in human-dominated land uses are more similar to each other than to
111 those in natural or semi-natural land [3]. However, within this broad categorisation,
112 particular transitions may influence species in different ways, so we also explore
113 separately the impact of conversion to agricultural land, conversion to urban land, and
114 increases in agricultural intensity. The land-use classes within the dataset were

coarsened depending upon the land-use comparison of interest, allowing species abundances to be compared between intact (natural and semi-natural) land uses and converted (human-dominated, agricultural or urban, respectively) land uses, or with increasing agricultural intensity (Table 1).

We define the Species Sensitivity Index (SSI) as the log-response ratio:

$$SSI = \ln\left(\frac{n_c}{n_i}\right)$$

where n_c is the mean abundance of the species in converted land uses, and n_i is the mean abundance in intact land-uses. SSI was estimated as follows. For each species, abundance was modelled statistically as a function of the land-use category (intact vs converted) or agricultural intensity. Study identity was included as a factor to account for among-study differences in methodology, sampling effort and biogeography. Study identity would ideally be treated as a random variable, but many species were in too few studies to accurately estimate random-effect variances [29] (~80% of species in our dataset were represented in six or fewer studies). Bayesian generalized linear models were used [with weakly informative default priors, as described in 30, implemented in the arm package using the bayesglm function] with a Poisson error structure unless overdispersion required use of a quasi-Poisson structure [31]. This approach provides more realistic coefficient estimates and more conservative standard errors than frequentist generalized linear models given that there can be complete separation of the data (for example, when a species is always absent from converted sites but always present in intact sites). The land-use coefficient (the log response ratio) from the model was used as the estimate of SSI and the standard error as its associated uncertainty.

In all further analyses, we use data for species that have abundance values in at least six sites (three intact and three converted), aiming for a balance between maximizing numbers of species and having sufficient data for each one. Repeating the analyses with a more stringent threshold (at least 12 sites: six intact and six converted site) produced qualitatively similar results, so we show results with the more lenient threshold, which include more species.

Phylogenetic tree

We used the Bee Tree of Life [32], a recent phylogeny of over 1,300 bee species from

around the world, after rate-smoothing using PATHd8 [a computationally efficient method for large phylogenetic trees 33] with the root age constrained to one. Of the species present in our dataset, 141 were also in the phylogeny. Although there was no difference in mean SSI between these and the species absent from the phylogeny (two-tailed t-test: $t_{1,245.0} = -0.33$, n.s.), the latter tended to have higher uncertainty in SSI (two tailed t-test: $t_{1,230.1} = -4.29$, $p < 0.001$). Because of this non-randomness, we used the *pastis* package in R [a birth-death polytomy resolver, 34] to estimate—1000 times—possible placements for missing species, given their taxonomic affinities [35], to produce 1000 complete trees. See Appendix S2 for full details. Birth-death polytomy resolvers can bias trait-based analyses [36], so we perform all analyses on the rate-smoothed incomplete tree of 141 species as well as the 1000 complete trees.

Phylogenetic analyses

Phylogenetic signal was quantified using Pagel's [37] λ , which produces reliable estimates when sample sizes are large [38], as here. We used the implementation in the R package *phytools* [39,40], which accommodates uncertainty in the SSI estimates, though we also estimated λ without accounting for uncertainty in species sensitivity, for comparison. As bumblebees (*Bombus*) may respond differently from other species to human impacts [14,41,42], which could drive phylogenetic signal in the overall dataset, we also estimated λ for both bumblebees alone and all other species.

We used 'traitgrams' [43] to assess the relative power of phylogenetic and functional distances to explain variation in SSI. Phylogenetic and functional distances were combined into a single set of Euclidean distances [43] according to:

$$FPDist = \sqrt{(aPDist^2 + (1 - a)FDist^2)}$$

where $PDist$ is the phylogenetic (cophenetic) species distance [pez package: 47], $FDist$ is the trait pairwise distance [Gower's dissimilarity with transformation to provide Euclidean properties: 48,49], and a governs the relative weighting of $FDist$ and $PDist$ in $FPDist$. Setting $a = 0$ computes functional diversity and $a = 1$ computes phylogenetic diversity. We calculated 11 distance matrices using a values spaced evenly from 0 to 1, and compared their explanatory power to find which best explained our data in distance-based generalised linear models [dbstats package, 47,48] weighted by the inverse squared standard errors of the SSI. We also modelled SSI as a function of a randomly-generated distance matrix (with original values drawn from a normal distribution)

as a null model. Models were compared using an adapted version of Akaike's Information Criterion (AIC) [47,48]. We calculated models' pseudo R^2 as: $1 - \left(\frac{\text{Residual deviance}}{\text{Null deviance}} \right)$.

Ecological trait data were collated for 556 of the species in our dataset by SPMR, MK and EF by assessing peer-reviewed and grey literature and by measuring available museum specimens. Traits included inter-tegular distance (ITD) [a proxy for foraging range, 52], flight season duration, dietary breadth, nesting strategy, and reproductive phenology and strategy [11], and are generally phylogenetically conserved (see Appendix Table S3.2). However, because data for the complete set of seven traits were available for only 75 species in both our data set and the original phylogeny, we re-ran this analysis using a reduced set of traits (ITD, nesting strategy and reproductive strategy), enabling inclusion of 143 species. We also repeated the analysis using the completed trees (259 species with data for all traits; 537 species with the minimal set).

Results

SSI values

Species' sensitivities to different land-use transitions were relatively normally distributed (see Fig 1), with wide variation in species responses. For the comparison between natural and human-dominated land-uses, for instance, 40 species had $SSI < -2$ and 28 species had $SSI > 2$; 82 species had SSI estimates that were significantly non-zero (44 negative, 38 positive; coloured lines on Fig 1).

Phylogenetic signal

The strength of phylogenetic signal in SSI differed among land-use comparisons (Fig. 2a). The strongest signal was in comparisons between (semi-)natural and human-dominated land uses, with high and significantly non-zero λ values for all but one of the completed phylogenetic trees ($p < 0.05$; Fig. 2a and Fig. 3). SSIs comparing semi-natural and natural land to agriculture showed λ values nearly as high: λ was significantly non-zero for the rate-smoothed tree ($\lambda = 0.59$, $p < 0.05$), and for >99% of the completed trees. The signal in sensitivity to urbanization and to increasing agricultural intensity depended on which phylogeny was used, being very low except with the completed trees (Fig. 2a). Weighting the analyses by the standard error of species sensitivity was extremely important: λ values were always low when standard

errors were not accounted for (Fig. 2b).

There was no significant phylogenetic signal within the bumblebees. Phylogenetic signal was also lower among non-bumblebees, particularly when assessing species' sensitivity to all human-dominated land-uses, and particularly when analyzing only the species in the rate-smoothed tree (Table 2); analyses using the complete phylogenies often found significant, albeit reduced, phylogenetic signal in non-bumblebees for both agricultural expansion and loss of natural vegetation (Table 2).

Relative importance of traits and phylogeny

Species' sensitivities to human-dominated land uses were best explained by a combination of both phylogeny and functional distances: this was true when using the rate-smoothed tree and across the completed trees (with $\alpha \geq 0.7$ producing the highest AIC weights and high R^2 values; Figure 4 and Appendix Figure S5.1). However, phylogenetic distance was more important than functional distances for transitions whose SSIs showed higher phylogenetic signal (i.e., sensitivity to human dominated land and agricultural land), whereas sensitivity to agricultural intensification seemed to be more strongly influenced by functional distances ($\alpha < 0.4$ had the highest AIC weights and R^2 values; Figure 4 and Appendix Figure S5.1).

Discussion

Bee species show strong heterogeneity in their responses to land-use practices with, in our data set, roughly as many 'winners' (positive SSI) as 'losers' (negative SSI) (Fig 1). This is in line with the wide range of responses reported in the literature (e.g., urbanization has been reported as having positive [50,51], neutral [52] and negative [50,51,53] effects on bee populations). Clearly many bee species are able to benefit from the environmental changes (potentially including removal of competitors: [54]), while others struggle to persist.

The strong phylogenetic signal ($\lambda \sim 0.7$) seen in species' responses to human-dominated land uses is not surprising: SSI estimates a species' susceptibility to one particular driver (land-use change), which will be mediated by functional response traits [11,12,55,56] that are often phylogenetically conserved [57] (though estimated responses will also be influenced by the contexts of individual studies, probably reducing the phylogenetic signal).

The different focused land-use comparisons find different degrees of phylogenetic signal in species' responses (Fig 2), with conversion to agriculture and urbanization producing moderately strong signal ($\lambda \sim 0.6$, except for the analysis of urbanization using the rate-smoothed tree, where sample size is small) while agricultural intensification elicits negligible signal ($\lambda \sim 0$). Accounting for measurement error was crucial for estimating phylogenetic signal strength; omitting it led to much lower estimates of λ (Fig 2), as expected [39], though results were consistent. Results were also not strongly driven by bumblebees; clustering of sensitivity was seen throughout the tree.

Why does agricultural intensification not produce a stronger phylogenetic signal? We consider three possible biological explanations. First, species' responses to agricultural intensity may be mediated by features of landscapes—not considered in our models—more than by features of sites. Field diversity and landscape composition, for example, have been shown to shape the biotic effects of agricultural intensity [58]; because such attributes are not recorded in the PREDICTS database, we were not able to consider them, perhaps causing responses to appear idiosyncratic. Second, the low-intensity agricultural habitats to which higher-intensity agriculture is compared may have already filtered out the most vulnerable native bee species (i.e., those with the most negative SSI values), and perhaps had synanthropic (i.e., SSI > 0) non-natives added, by the original conversion from natural to agricultural land. Such SSI-biased turnover of species would tend to erode SSI's phylogenetic signal. This explanation—that the initial conversion of a landscape to an 'anthrome' [59] leaves a strong phylogenetic signal whereas subsequent intensification of use does not—requires that much the same traits mediate responses to agricultural expansion and agricultural intensification. Such trait-based extinction filters are likely why mammalian extinction risk shows no phylogenetic signal in regions with a long history of intense human activity [60]. If this explanation is correct, then even though low-intensity agricultural practices score better than more intensive systems for community-level measures of bee biodiversity [35,58], they may have already altered community composition irreversibly [42]. The third possibility is that the ability of species to persist in the face of agricultural intensification depends most strongly on traits that are not phylogenetically conserved. For instance, we found no strong phylogenetic signal in diet breadth (Appendix Table 3.2). Diet breadth can be flexible for some species (e.g. *Bombus terrestris* can increase diet breadth when faced with increased resource competition [61]), but on the other hand, even generalist species can have rigid host-plant preferences [62]. A fourth, non-

biological, explanation for the lack of phylogenetic signal in responses to agricultural intensification is that our use-intensity criteria mix multiple pressures (e.g. pesticide and fertilizer use); each component might on its own elicit phylogenetically patterned responses, but the signal may be lost by mixing them.

Although SSI values usually showed phylogenetic signal, in the subset of bee species with available trait information, SSIs were generally better explained by a combination of both phylogenetic and trait distances [43], highlighting the added value of ecological trait data for species. These results are in line with previous analyses revealing that functional traits can significantly influence species sensitivities to a number of land uses and land-use practices in a variety of systems [11,12,55]. However, such trait-environment relationships can have low explanatory power and results across studies can be contradictory [13]. Taken with our results, this suggests that species' sensitivities may be influenced by ecological differences that are not fully captured by the commonly measured traits, while phylogeny may provide a closer approximation to these unmeasured characteristics [63]. For instance, the phylogenetic relatedness of bee species can inform the structure of plant-pollinator networks [23,64]; as ecological interactions are lost more quickly than species from a system [65], a given species' network could have a strong influence on its resilience to disturbances. Responses to agricultural intensification do not fit this model: trait differences explained some variation in SSI but phylogenetic distance did not. This combination of results is consistent with the suggestion that trait-mediated competition may underlie the responses [66].

Although the importance of traits in mediating species' sensitivities to land-use changes is congruent with previous work [12], our dataset may not be representative; it is therefore possible that a more complete set of species with trait data would change the relative importance of traits and phylogeny. The species in our dataset capture significantly less phylogenetic diversity than expected from a random selection of species from the Bee Tree of Life (see Appendix 6). Local assemblages are often a non-random subset of the global phylogeny [67,68]; our analyses focus on such assemblages and so this is, in part, an inescapable consequence of our study's objectives.

The strong phylogenetic signal in bee species' sensitivities to human-dominated land uses like agriculture means that losses of diversity are likely to be concentrated within a subset of clades, where they will be correspondingly more severe; likewise, any gains in diversity will be restricted to groups of related species. Clustering of losses can greatly

reduce the phylogenetic and functional diversity of a system [16], potentially jeopardizing its ability to function under environmental change [20]. Crop pollination may be more robust than other ecosystem functions to species losses as a few dominant species are the main contributors at local scales [69]. However, higher species diversity may be necessary at larger spatial scales [70]. Furthermore, some wild plant species require specialist pollinators, potentially meaning that they face a double threat from conversion of land to agriculture: directly, through loss of habitat, and indirectly, through the decline in their pollinators.

Our results suggest that the phylogenetic risk assessment framework set out by Díaz et al. [16] could inform management practices and highlight gaps in knowledge. Even though bees are well studied relative to many other invertebrate groups, there are still uncertainties about their current status and vulnerability to human impacts [71] as well as gaps in trait data. The strong phylogenetic patterning in species' sensitivities to agricultural expansion could help to predict the sensitivity of understudied species, identifying those that are most vulnerable or resistant to guide conservation planning [72,73]. For instance, many species of bumblebees showed positive or neutral responses to agricultural land, but negative responses to increased agricultural intensity (Appendix Figure S4.2 and S4.4); this combination of results suggests that agricultural production can support many species of bumblebees, but only if intensity is low. However, our estimate of sensitivity to particular pressures—SSI—does not necessarily indicate the conservation status of a species, which is a product of both sensitivity to combined pressures and exposure to those pressures. This may explain why we found no phylogenetic signal in bumblebee SSIs, even though extinction risk in these species is significantly non-random [15]. Phylogenetically patterned responses may also open opportunities to monitor bee communities at higher taxonomic levels [74–76]: this would reduce the need for species identification by expert taxonomists and the need for destructive sampling of bees [77,78], save time and money, and facilitate citizen science.

Our results provide a basis from which to make, test, and inform predictions about bee species' sensitivities to land-use change, with potentially important benefits for monitoring and conservation prioritization, as well as identifying land-use pressures that may most affect pollination services to crops and wild species. Analyses of how these results scale up to changes in abundance-weighted phylogenetic diversity of communities are necessary to identify spatial patterns in diversity and potential areas of

pollination deficit [35,79].

Acknowledgements

We are grateful to all those who contributed data to this project, particularly members of the Greenveins project, and the PREDICTS team for assistance with data collation and curation, including Sara Contu, Sam Hill, Helen Phillips, Tim Newbold and Lawrence Hudson. We thank Gavin Thomas and Susy Echeverria-Londoño for assistance with phylogeny estimation. Much of the trait data used in this work was compiled by the EU FP7 project “Status and Trends of European Pollinators” (244 090, www.STEP-project.net). This work was supported by the BBSRC (grant BB/F017324/1 to ADP) and NERC (grant NE/M014533/1 to AP) and is a contribution from the Imperial College Grand Challenges in Ecosystems and the Environment Initiative. PREDICTS is endorsed by the GEO-BON.

Author Contributions

ADP and AP conceived the study. ADP carried out statistical analyses and wrote the first draft of the manuscript. MK assessed species taxonomy. EF, MK, SPMR and SP collated trait data. WP provided guidance on phylogenetic analyses. All authors contributed significantly to revisions of the manuscript. All authors gave final approval for publication.

References

1. Maxwell SL, Fuller RA, Brooks TM, Watson JEM. 2016 Biodiversity: The ravages of guns, nets and bulldozers. *Nature* **536**, 143–145. (doi:10.1038/536143a)
2. WWF. 2016 Living Planet Report 2016 Risk and resilience in a new era.
3. Newbold T *et al.* 2015 Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50. (doi:10.1038/nature14324)
4. Pereira HM *et al.* 2010 Scenarios for global biodiversity in the 21st century. *Science* **330**, 1496–501. (doi:10.1126/science.1196624)
5. Hurtt GC *et al.* 2011 Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Clim. Change* **109**, 117–161. (doi:10.1007/s10584-011-0153-2)
6. Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham S a, Kremen C, Tscharntke T. 2007 Importance of pollinators in changing landscapes for world crops. *Proc. Biol. Sci.* **274**, 303–13. (doi:10.1098/rspb.2006.3721)
7. Danforth BN, Sipes S, Fang J, Brady SG. 2006 The history of early bee diversification based on five genes plus morphology. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 15118–15123. (doi:10.1073/pnas.0604033103)
8. Ollerton J, Winfree R, Tarrant S. 2011 How many flowering plants are pollinated by animals? *Oikos* **120**, 321–326. (doi:10.1111/j.1600-0706.2010.18644.x)
9. Suding KN *et al.* 2008 Scaling environmental change through the community-level: a trait-based response-and-effect framework for plants. *Glob. Chang. Biol.* **14**, 1125–

1140. (doi:10.1111/j.1365-2486.2008.01557.x)
10. D'iaz S *et al.* 2013 Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol. Evol.* **3**, 2958–2975. (doi:10.1002/ece3.601)
11. De Palma A, Kuhlmann M, Roberts SPM, Potts SG, Börger L, Hudson LN, Lysenko I, Newbold T, Purvis A. 2015 Ecological traits affect the sensitivity of bees to land-use pressures in European agricultural landscapes. *J. Appl. Ecol.* **52**, 1567–1577. (doi:10.1111/1365-2664.12524)
12. Rader R, Bartomeus I, Tylianakis JM, Laliberté E. 2014 The winners and losers of land use intensification: pollinator community disassembly is non-random and alters functional diversity. *Divers. Distrib.* **20**, 908–917. (doi:10.1111/ddi.12221)
13. Bartomeus I, Cariveau DP, Harrison T, Winfree R. 2017 On the inconsistency of pollinator species traits for predicting either response to land-use change or functional contribution. *Oikos*, doi: 10.1111/oik.04507. (doi:10.1111/oik.04507)
14. Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R. 2013 Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 4656–60. (doi:10.1073/pnas.1218503110)
15. Arbetman MP, Gleiser G, Morales CL, Williams P, Aizen MA. 2017 Global decline of bumblebees is phylogenetically structured and inversely related to species range size and pathogen incidence. *Proceedings. Biol. Sci.* **284**, 20170204. (doi:10.1098/rspb.2017.0204)
16. D'iaz S *et al.* 2013 Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol. Evol.* **3**, 2958–2975. (doi:10.1002/ece3.601)
17. Kaiser-Bunbury CN, Muff S, Memmott J, Müller CB, Caflisch A. 2010 The robustness of pollination networks to the loss of species and interactions: a quantitative approach incorporating pollinator behaviour. *Ecol. Lett.* **13**, 442–452. (doi:10.1111/j.1461-0248.2009.01437.x)
18. Cardinale BJ *et al.* 2015 Further re-analyses looking for effects of phylogenetic diversity on community biomass and stability. *Funct. Ecol.* (doi:10.1111/1365-2435.12540)
19. Cadotte MW, Livingstone SW, Yasui S-LE, Dinnage RD, Li J-T, Marushia R, Santangelo J, Shu W. 2017 Explaining ecosystem multifunction with evolutionary models. *Ecology* **98**, 3175–3187. (doi:10.1002/ecy.2045)
20. Cadotte MW, Dinnage R, Tilman D. 2012 Phylogenetic diversity promotes ecosystem stability. *Ecology* **93**, S223–S233. (doi:10.1890/11-0426.1)
21. Miranda M, Parrini F. 2015 Congruence between species phylogenetic and trophic distinctiveness. *Biodivers. Conserv. Conserv.* **24**, 355–369. (doi:10.1007/s10531-014-0813-4)
22. Naisbit RE, Rohr RP, Rossberg a. G, Kehrli P, Bersier L-F. 2012 Phylogeny versus body size as determinants of food web structure. *Proc. R. Soc. B Biol. Sci.* **279**, 3291–3297. (doi:10.1098/rspb.2012.0327)
23. Rezende EL, Lavabre JE, Guimarães PR, Jordano P, Bascompte J. 2007 Non-random coextinctions in phylogenetically structured mutualistic networks. *Nature* **448**, 925–928. (doi:10.1038/nature05956)
24. Fritz SA, Purvis A. 2010 Selectivity in mammalian extinction risk and threat types: a new measure of phylogenetic signal strength in binary traits. *Conserv. Biol.* **24**, 1042–1051. (doi:10.1111/j.1523-1739.2010.01455.x)
25. Hudson LN *et al.* 2017 The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecol. Evol.* **7**, 145–188. (doi:10.1002/ece3.2579)
26. De Palma A *et al.* 2016 Predicting bee community responses to land-use changes: Effects of geographic and taxonomic biases. *Sci. Rep.* **6**, 31153. (doi:10.1038/srep31153)
27. Hudson LN *et al.* 2014 The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecol. Evol.* **4**, 4701–4735.

- (doi:10.1002/ece3.1303)
28. Hudson LN *et al.* 2014 The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecol. Evol.* **4**, 4701–4735. (doi:10.1002/ece3.1303)
29. Fox GA, Negrete-Yankelevich S, Sosa VS. 2015 *Ecological Statistics: Contemporary Theory and Application*. Oxford: Oxford University Press.
30. Gelman A, Jakulin A, Pittau MG, Su Y-S. 2008 A weakly informative default prior distribution for logistic and other regression models. *Ann. Appl. Stat.* **2**, 1360–1383. (doi:10.1214/08-AOAS191)
31. Crawley MJJ. 2007 *The R book*. Chichester: John Wiley & Sons Ltd. See http://books.google.com/books?hl=en&lr=&id=XYDI0mIH-moC&oi=fnd&pg=PA2008&dq=The+R+Book&ots=HYTI-mHeHF&sig=ut_4HwPVkTuZ-bg-gk7N2Zwrn1s.
32. Hedtke SM, Patiny S, Danforth BN. 2013 The bee tree of life: a supermatrix approach to apoid phylogeny and biogeography. *BMC Evol. Biol.* **13**, 138. (doi:10.1186/1471-2148-13-138)
33. Britton T, Anderson CL, Jacquet D, Lundqvist S, Bremer K. 2007 Estimating divergence times in large phylogenetic trees. *Syst. Biol.* **56**, 741–752. (doi:10.1080/10635150701613783)
34. Thomas GH, Hartmann K, Jetz W, Joy JB, Mimoto A, Mooers AO. 2013 PASTIS: an R package to facilitate phylogenetic assembly with soft taxonomic inferences. *Methods Ecol. Evol.* **4**, 1011–1017. (doi:10.1111/2041-210X.12117)
35. De Palma A, Kuhlmann M, Bugter R, Ferrier S, Hoskins AJ, Potts SG, Roberts SPM, Schweiger O, Purvis A. 2017 Dimensions of biodiversity loss: Spatial mismatch in land-use impacts on species, functional and phylogenetic diversity of European bees. *Divers. Distrib.* **23**, 1435–1446. (doi:10.1111/ddi.12638)
36. Rabosky DL. 2015 No substitute for real data: phylogenies from birth-death polytomy resolvers should not be used for many downstream comparative analyses. , 1–23. (doi:arXiv:1503.04978)
37. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884. (doi:10.1038/44766)
38. Münkemüller T, Lavergne S, Bzeznik B, Dray S, Jombart T, Schiffrers K, Thuiller W. 2012 How to measure and test phylogenetic signal. *Methods Ecol. Evol.* **3**, 743–756. (doi:10.1111/j.2041-210X.2012.00196.x)
39. Ives AR, Midford PE, Garland T. 2007 Within-species variation and measurement error in phylogenetic comparative methods. *Syst. Biol.* **56**, 252–270. (doi:10.1080/10635150701313830)
40. Revell LJ. 2012 phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223. (doi:10.1111/j.2041-210X.2011.00169.x)
41. De Palma A *et al.* 2016 Predicting bee community responses to land-use changes: Effects of geographic and taxonomic biases. *Sci. Rep.* **6**. (doi:10.1038/srep31153)
42. Carvalheiro LG *et al.* 2013 Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecol. Lett.* **16**, 870–878. (doi:10.1111/ele.12121)
43. Cadotte M, Albert CH, Walker SC. 2013 The ecology of differences: Assessing community assembly with trait and evolutionary distances. *Ecol. Lett.* **16**, 1234–1244. (doi:10.1111/ele.12161)
44. Pearse WD, Cadotte MW, Cavender-Bares J, Ives AR, Tucker C, Walker SC, Helmus MR. 2015 pez: Phylogenetics for the Environmental Sciences.
45. Gower JC. 1971 A General Coefficient of Similarity and Some of Its Properties. *Biometrics* **27**, 857. (doi:10.2307/2528823)
46. Debastiani VJ, Pillar VD. 2012 SYNCSEA--R tool for analysis of metacommunities based on functional traits and phylogeny of the community components. *Bioinformatics* **28**, 2067–8. (doi:10.1093/bioinformatics/bts325)

47. Boj E, Caballe A, Delicado P, Fortiana J. 2014 dbstats: Distance-Based Statistics.
48. Boj E, Caballé A, Delicado P, Esteve A, Fortiana J. 2015 Global and local distance-based generalized linear models. *TEST* (doi:10.1007/s11749-015-0447-1)
49. Greenleaf SS, Williams NM, Winfree R, Kremen C. 2007 Bee foraging ranges and their relationship to body size. *Oecologia* **153**, 589–596. (doi:10.1007/s00442-007-0752-9)
50. Hernandez JL, Frankie GW, Thorp RW. 2009 Ecology of Urban Bees: A Review of Current Knowledge and Directions for Future Study. *Cities Environ.* **2**, 1–15.
51. Fortel L, Henry M, Guilbaud L, Guirao AL, Kuhlmann M, Mouret H, Rollin O, Vaissière BE. 2014 Decreasing Abundance, Increasing Diversity and Changing Structure of the Wild Bee Community (Hymenoptera: Anthophila) along an Urbanization Gradient. *PLoS One* **9**, e104679. (doi:10.1371/journal.pone.0104679)
52. Baldock KCR *et al.* 2015 Where is the UK's pollinator biodiversity? The importance of urban areas for flower-visiting insects. *Proc. R. Soc. B Biol. Sci.* **282**. (doi:10.1098/rspb.2014.2849)
53. Bates AJ, Sadler JP, Fairbrass AJ, Falk SJ, Hale JD, Matthews TJ. 2011 Changing bee and hoverfly pollinator assemblages along an urban-rural gradient. *PLoS One* **6**, e23459. (doi:10.1371/journal.pone.0023459)
54. Tuck SL, Porter J, Rees M, Turnbull LA. 2018 Strong responses from weakly interacting species. *Ecol. Lett.* (doi:10.1111/ele.13163)
55. Williams NM, Crone EE, Roulston TH, Minckley RL, Packer L, Potts SG. 2010 Ecological and life-history traits predict bee species responses to environmental disturbances. *Biol. Conserv.* **143**, 2280–2291. (doi:10.1016/j.biocon.2010.03.024)
56. Violle C, Navas M-L, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E. 2007 Let the concept of trait be functional! *Oikos* **116**, 882–892. (doi:10.1111/j.2007.0030-1299.15559.x)
57. Srivastava DS, Cadotte MW, Macdonald a. AM, Marushia RG, Mirotchnick N. 2012 Phylogenetic diversity and the functioning of ecosystems. *Ecol. Lett.* **15**, 637–648. (doi:10.1111/j.1461-0248.2012.01795.x)
58. Kennedy CM *et al.* 2013 A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecol. Lett.* **16**, 584–599. (doi:10.1111/ele.12082)
59. Ellis EC, Ramankutty N. 2008 Putting people in the map: anthropogenic biomes of the world. *Front. Ecol. Environ.* **6**, 439–447. (doi:10.1890/070062)
60. Fritz S a, Bininda-Emonds ORP, Purvis A. 2009 Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecol. Lett.* **12**, 538–549. (doi:10.1111/j.1461-0248.2009.01307.x)
61. Fontaine C, Collin CL, Dajoz I. 2008 Generalist foraging of pollinators: diet expansion at high density. *J. Ecol.* **96**, 1002–1010. (doi:10.1111/j.1365-2745.2008.01405.x)
62. Kleijn D, Raemakers I. 2008 A retrospective analysis of pollen host plant use by stable and declining bumble bee species. *Ecology* **89**, 1811–1823. (doi:10.1890/07-1275.1)
63. Faith DP. 2013 Biodiversity and evolutionary history: useful extensions of the PD phylogenetic diversity assessment framework. *Ann. N. Y. Acad. Sci.* **1289**, 69–89. (doi:10.1111/nyas.12186)
64. Chamberlain S a, Cartar R V, Worley AC, Semmler SJ, Gielens G, Elwell S, Evans ME, Vamosi JC, Elle E. 2014 Traits and phylogenetic history contribute to network structure across Canadian plant – pollinator communities. , 545–556. (doi:10.1007/s00442-014-3035-2)
65. Valiente-Banuet A *et al.* 2015 Beyond species loss: the extinction of ecological interactions in a changing world. *Funct. Ecol.* **29**, 299–307. (doi:10.1111/1365-2435.12356)
66. Kraft NJB, Adler PB, Godoy O, James EC, Fuller S, Levine JM. 2015 Community assembly, coexistence and the environmental filtering metaphor. *Funct. Ecol.* **29**, 592–599. (doi:10.1111/1365-2435.12345)

67. Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW. 2009 The merging of community ecology and phylogenetic biology. *Ecol. Lett.* **12**, 693–715. (doi:10.1111/j.1461-0248.2009.01314.x)
68. Vamosi SM, Heard SB, Vamosi C, Webb CO. 2009 Emerging patterns in the comparative analysis of phylogenetic community structure. *Mol. Ecol.* **18**, 572–592. (doi:10.1111/j.1365-294X.2008.04001.x)
69. Kleijn D *et al.* 2015 Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nat. Commun.* **6**, 7414. (doi:10.1038/ncomms8414)
70. Winfree R, Reilly JR, Bartomeus I, Cariveau DP, Williams NM, Gibbs J. 2018 Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science (80-.)*. **359**, 791–793. (doi:10.1126/SCIENCE.AAO2117)
71. Nieto A *et al.* 2015 *European Red List of Bees*. Luxembourg: Publication Office of the European Union. (doi:10.2779/77003)
72. Mace GM, Gittleman JL, Purvis A. 2003 Preserving the tree of life. *Science (80-.)*. **300**, 1707–1709. (doi:10.1126/science.1085510)
73. Bland LM, Collen B, Orme CDL, Bielby J. 2015 Predicting the conservation status of data-deficient species. *Conserv. Biol.* **29**, 250–259. (doi:10.1111/cobi.12372)
74. Bevilacqua S, Terlizzi A, Claudet J, Frascchetti S, Boero F. 2012 Taxonomic relatedness does not matter for species surrogacy in the assessment of community responses to environmental drivers. *J. Appl. Ecol.* **49**, 357–366. (doi:10.1111/j.1365-2664.2011.02096.x)
75. Guénard G, Von Der Ohe PC, De Zwart D, Legendre P, Lek S. 2011 Using phylogenetic information to predict species tolerances to toxic chemicals. *Ecol. Appl.* **21**, 3178–3190. (doi:10.1890/10-2242.1)
76. Kallimanis AS, Mazaris AD, Tsakanikas D, Dimopoulos P, Pantis JD, Sgardelis SP. 2012 Efficient biodiversity monitoring: Which taxonomic level to study? *Ecol. Indic.* **15**, 100–104. (doi:10.1016/j.ecolind.2011.09.024)
77. Lebuhn G *et al.* 2013 Detecting Insect Pollinator Declines on Regional and Global Scales. *Conserv. Biol.* **27**, 113–120. (doi:10.1111/j.1523-1739.2012.01962.x)
78. Tepedino VJ, Durham S, Cameron SA, Goodell K. 2015 Documenting bee decline or squandering scarce resources. **29**, 280–282. (doi:10.1111/cobi.12439)
79. Thuiller W, Lavergne S, Roquet C, Boulangeat I, Lafourcade B, Araujo MB. 2011 Consequences of climate change on the tree of life in Europe. *Nature* **470**, 531–4. (doi:10.1038/nature09705)

Figure Legends

Figure 1: Spread of SSI values (points) and their standard error (lines) for each land-use transition, for all species in the dataset. Coloured points indicate that SSI values were significantly different from 0. Particularly vulnerable species (i.e., those with an SSI significantly below 0) are coloured in yellow to red, while species that strongly benefit from land-use change (i.e., those with SSI values significantly above 0) are coloured light to dark blue.

Figure 2: Phylogenetic signal in species' sensitivity to various land use pressures, when (a) standard errors are accounted for and (b) when they are ignored. Red triangles indicate the values for the rate-smoothed tree, where some species are missing from the tree. Numbers in red show the number of species included in these tests. Boxplots show the distribution of values across the 1,000 completed trees, with numbers in black showing the number of species included in these tests.

Figure 3: Phylogenetic signal in species' sensitivity to all human-dominated land uses using the rate-smoothed (incomplete) tree. Tips of the phylogeny are coloured according to the species' sensitivity: from blue to red indicate less to more sensitive. The right panel shows species sensitivity $\pm$ the standard error. See Appendix Figures S4.2 to S4.4 for similar figures for other land-use transitions.

Figure 4: Akaike's Weights of models assessing species sensitivities as a function of α value, where $\alpha = 0$ uses only functional traits to calculate the distance, $\alpha = 1$ uses only phylogeny to calculate the distance, and $\alpha = 0.5$ considers both traits and phylogeny equally. Model performance was also calculated where distances were randomly computed. Red triangles indicate the values for the rate-smoothed tree, where some species are missing from the tree. Numbers in red show the number of species included in these tests. Boxplots show the distribution of values across the 1,000 completed trees, with numbers in black showing the number of species included in these tests. For a similar figure assessing explanatory power of models, see Appendix Figure S5.5.

Tables

Table 1: Land-use transitions of interest and the component land-use classes. Dataset sample sizes are also given: number of studies, species and species that match with the recently-published Bee Tree of Life [32].

Land use comparison	Intact	Converted	Number of studies	Number of species	Number of species in the phylogeny
Loss of natural and semi-natural habitat	Primary & Secondary vegetation	Cropland, Pasture, Plantation forest & Urban	86	488	141
Agricultural expansion	Primary & Secondary vegetation	Cropland, Pasture & Plantation forest	84	437	121
Urbanisation	Primary & Secondary vegetation	Urban	54	146	56
Agricultural intensification	Agricultural sites (as defined above) with a continuous use-intensity scale from 1 (minimal use) to 3 (intense use)		53	268	87

Table 2: λ values of the sensitivity of all bee species, bumblebees only and other species (bumblebees excluded from the phylogenetic tree) to land-use change. For the completed trees, this is the mean $\lambda \pm$ standard deviation across the 1000 phylogenies.

Sensitivity to:	Tree	λ		
		All species	Bumblebees	Other species
Human-dominated land	Rate-smoothed	0.741	0	0
	Completed	0.672 ± 0.09	0 ± 0	0.626 ± 0.14
Agricultural land	Rate-smoothed	0.594	0	0
	Completed	0.64 ± 0.1	0 ± 0	0.586 ± 0.16
Urban land	Rate-smoothed	0	0	0.143
	Completed	0.633 ± 0.15	0.256 ± 0.44	0.2 ± 0.12
Higher-intensity agriculture	Rate-smoothed	0.046	0	0.041
	Completed	0.116 ± 0.03	0 ± 0	0.121 ± 0.04

Sensitivity to human-dominated land use

Sensitivity to agricultural land

Sensitivity to urban land

Sensitivity to agricultural intensification

bioRxiv preprint doi: <https://doi.org/10.1101/524546>; this version posted January 18, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Sensitivity Index

Sensitivity to Human dominated land

138 Species

55 Species

Sensitivity to Agricultural land

134 Species

52 Species

Sensitivity to Agricultural intensification

148 Species

50 Species

Sensitivity to Urban land

64 Species

29 Species