

Figure Supplementary 1 for Figure 1

wnt11f1 is only expressed by prim1-derived neuromasts





Figure Supplementary 2 for Figure 1

Analysis of hair cell orientation with respect to the nearest neighbor
(do hair cells show relative orientation with respect to each other?)



Hair cell orientation in double mutants




Figure Supplementary 3 for Figure 2

Analysis of angle difference between a given hair cell's orientation and the nearest fitting ellipse (concentricity)





Figure Supplementary 4 for Figure 2

Analysis of hair cell orientation with respect to the nearest neighbor




Figure Supplementary 5 for Figure 3




Figure Supplementary 6 for Figure 3

Vangl2 localization is absent in *vangl2*^{m209} mutants






Figure Supplementary 7 for Figure 4

Analyses of cell behaviors of hair cell precursors during development show no differences between Wild type and *MZwnt11f1* mutants

● Hair cell precursor
 ● Hair cells produced from precursor division
 ● Hair cells produced from precursor division
 Rearrangement



e Division angle with respect to the radius (XY)







Figure Supplementary 8 for Figure 4

Angle comparison with nearest neighbor of hair cells in primII-derived neuromasts





Figure Supplementary 9 for Figure 5

Pipeline for Segmentation of Phalloidin images for Support Cell orientation analysis


Step 1

Phalloidin staining




Step 2

Custom sharpening




Step 3

MorphoJ segmentation




Step 4

Image Clean-up



Step 5


Fitting ellipse




(3h after Neomycin)

Figure Supplementary 10 for Figure 6


wnt11f1 is expressed in the muscle



fzd7a



gpc4



scrib







Figure Supplementary 11 for Figure 6

Neuromasts derived from the Occipital prim and primD show a hair cell phenotype in *MZwnt11f1* mutants




Figure Supplementary 12

Distribution, but not ratio of expression, of Emx+ hair cells is affected in primII-derived neuromasts of *MZwnt11f1* mutants

