FabriEVES: A dedicated platform for endogenous viral elements in fishes, amphibians, birds, reptiles and invertebrates

Jixing Zhong^{1,2,3§}, Jiacheng Zhu^{2,3§}, Zaoxu Xu^{2,3}, Geng Zou⁴, JunWei Zhao⁵, SanJie Jiang², Wei Zhang⁶, Jun Xia^{2,3,7}, Lin Yang^{2,3}, Fang Li⁵, Ya Gao³, Fang Chen³, Yiquan Wu^{8*}, Dongsheng Chen^{3*}

1 School of Biological Sciences and Medical Engineering, Southeast University, Xuanwu District, Nanjing 210096, China

2 BGI Education Center, University of Chinese Academy of Sciences, Beishan Industrial Zone, Shenzhen, 518083, China.

3 BGI-Shenzhen, Beishan Industrial Zone, Shenzhen, 518083, China.

4 State Key Laboratory of Agricultural Microbiology, College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

5 Department of Gynaecology, Shanghai First Maternity and Infant Hospital, Tongji University School of Medicine, Shanghai, 200040, China.

6 Sigmovir Biosystems Inc., 9610 Medical Center Drive, Suite 100, Rockville, MD, 20850, USA

7 MGI, BGI-Shenzhen, Beishan Industrial Zone, Shenzhen, 518083, China.

8 Current address: HIV and AIDS Malignancy Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892; yiquanwuhzau@hotmail.com

§, These authors contribute equally to this work

*, Corresponding authors:

Dongsheng Chen, chendongsheng@genomics.cn

Yiquan Wu, yiquanwuhzau@hotmail.com

ABSTRACT: Endogenous viral elements (EVEs) are heritable viral deriving elements present in the genomes of other species. As DNA 'fossils' left by ancient viruses, EVEs were used to infer the characteristics of extinct viruses. EVEs in mammals have been well classified by several databases, however, EVEs in non-mammalian organisms are poorly documented. Here, we report FabriEVEs (<http://tfbsbank.co.uk/FabriEVEs>), the first dedicated and comprehensive online platform for the collection, classification and annotation of EVEs in fishes, amphibians, birds, reptiles and invertebrates. In total, nearly 1.5 million EVEs from 82 species deriving from class I (dsDNA), II (ssDNA), III (dsRNA), IV (positive ssRNA), V (negative ssRNA), VI (ssRNA-RT) and VII (dsDNA-RT) viruses were recorded in FabriEVEs, accompanying with comprehensive annotation including the species name, location, genomic features, virus family and associated literature. Flexible and powerful query options were provided to pinpoint desired EVEs. Furthermore, FabriEVEs provides free access to all EVEs data in case users need to download them for further analysis. Taken together, our database provided systematic classification and annotation of EVEs in non-mammal species, which paves the way for comparative analysis of EVEs and throws light upon the co-evolution of EVEs and their hosts.

INTRODUCTION:

Although endogenous viral elements (EVEs) mainly origin from retroviruses(1–4), increasing number of EVEs have been identified from non-retroviruses families including parvoviruses, filoviruses, bornaviruses and circoviruses(2). EVEs can be classified into partial EVEs or entire EVEs (proviruses) depending on whether they represent the whole genomes of extant viruses. EVEs provide a unique opportunity for researchers to investigate extinct viruses, which give rise to the emerging Paleovirology(5). Comparison EVEs with their extant counterparts give us a clue

regarding the evolutionary trajectories of viruses and the co-evolution or the arm race between viruses and hosts(2,6). EVEs have been found to play important roles in regulating the expression of host genes by introducing novel cis-regulatory elements including promoters and enhancers(1,7–9). Domesticated EVEs are found to be involved in several critical biological processes of hosts such as immunity(10) and the formation of placenta(11,12). The contribution of immunity by EVEs can be interpreted in two aspects. Firstly, EVEs were found to regulate several immune-related cellular genes and were essential for the innate immunity of host(13). Besides, Some endogenous viral genes can be domesticated and utilized by hosts to fight against exogenous viruses by blocking their cell entry and replication(10,14–18).The initial studies of EVEs mainly focused on mammals, probably attributed to the available integral and abundant whole genome sequences. Recently, increasingly more EVEs have been identified in non-mammals species, along with the progress of modENCODE(19), Genome10K program(20), Avian Phylogenomics Project(21) and B10K Project(22). In fishes, several cases of EVEs have been reported in spotted green pufferfish and elephant shark respectively(23,24). A systematic screen on 48 bird species revealed a variety of EVEs from both retroviruses and non-retroviruses, though the frequency of EVEs is much lower than that in mammals(4). In reptiles, EVEs deriving from hepadnaviruses, bornaviruses and circoviruses have been identified from several snake species and Chinese soft shell turtle(25,26). EVEs have also been found in amphibians such as western clawed frog (*Xenopus tropicalis*)(27). In invertebrates, 27 species were found to accommodate several virus origin fragments(6,26,28). Of particular interest, a remarkable diversity of EVEs was found to exist in a crustacean genome, pill bug (*Armadillidium vulgare*)(29). Up to now, there are two EVEs databases for mammals. HERVd, the first database for human endogenous retroviruses, was released in 2002 and updated in

2004(30,31). gEVE version 1.1 houses 774,172 EVEs identified from 20 mammalian genomes(32). Despite a large amount of EVEs have been identified and reported in non-mammal species, there is no specific database to accommodate these invaluable data sets. To fill the gap between increasing EVEs data in fishes, amphibians, birds, reptiles and invertebrates and our limited ability to interpret them, we developed a dedicated online platform for the storage, classification, annotation, querying and downloading of non-mammal animal EVEs (Figure 1).

The statistics of data in FabriEVEs

In total, 1,483,658 EVEs from 19, 1, 48, 4 and 28 species in fishes, amphibians, birds, reptiles and invertebrates were included in FabriEVEs (Table 1). Although the majority of the EVEs (>99.99%) derived from retroviruses as expected, we still observed a high diversity of EVE originating from non-retroviruses. For example, in birds, 249 EVEs from hepadnaviruses (double strand DNA viruses), bornaviruses (negative single strand RNA viruses), circoviruses (single strand DNA viruses) and parvoviruses (single strand DNA viruses) were presented in our database. It is of particular interest that some species accommodate much more EVEs than their close relatives. In birds, *Phalacrocorax carbo* (Cormorant) contains 68 copies of non-retroviruses EVEs, compared to 0-14 copies of non-retroviruses EVEs in remaining 47 bird species. Admittedly, this unusual high copy number of non-retroviruses EVEs in Cormorant might simply be due to differences in genome size or genome assembly among the 48 sequenced bird species. However, we ruled out this possibility based on following reasons: 1) All of the genome size range from 1.05 Gb in *Acanthisitta chloris* (Rifleman) to 1.26 Gb in *Aptenodytes forsteri* (Emperor penguin), which means that no bird genome is significantly larger than the others among the 48 sequenced genomes in the Avian phylogenomics project(33–36). 2) The

sequencing depth of Great cormorant is only 24X compared to the depth of 160X in *Melopsittacus undulatus* (Budgerigar); 3) Among those 48 species, *Gallus gallus* (chicken) is the most well-studied bird species with its draft genome released as early as 2004(37). However, no non-retroviruses EVEs have been reported in chicken genome according to Cui's systematic screen on the 48 bird species(4). In summary, the high copy number of non-retroviruses EVEs in *Phalacrocorax carbo* (Cormorant) was unlikely to be caused by genome size or genome assembly variance. Considering previous reports have shown that EVEs might play important roles in the evolution of animals(1–3), it is reasonable to speculate that those *Phalacrocorax carbo* (Cormorant)-specific non-retroviruses might contribute to its divergence from other species or facilitate its adaptation to its unique ecology niches.

EVEs data Search

As there are 1.5 million EVEs in FabriEVEs, we provided advanced methods for users to retrieve the data they are looking for (Figure 2a). Specifically, four key attributes of each EVE elements were used to build the query. To begin with, the first filtering method was based on the class and scientific name of species. Class names (fishes, amphibians, birds, reptiles and invertebrates) were used to initiate the classification. After choosing the specific class, all the species in this class will be shown in a selection list, from which users can choose what species they are interested in. In addition, EVEs can be classified into *hepadnaviridae*, *bornaviridae*, *circoviridae*, *parvoviridae*, *densovirinae*, *reoviridae*, *rhodoviridae*, *flaviviridae*, *bunyaviridae*, *orthomyxoviridae*, *retroviridae* or “unassigned” based on which family the corresponding extant counterpart viruses belong to. All those options were provided as check-boxes and the default status for them are 'checked'. Users can deselect singular or several of them, and the

corresponding types of EVEs will be included or excluded from the results. Similarly, EVEs can be divided into partial or entire depending on whether they are homologous to part or entire genome sequences of extant viruses. After choosing desired options for class, species, type and completeness of EVEs, all the selected values will be submitted to a PHP script, which will execute the query based on provided information. For example, if the class 'bird', species '*Acanthisitta chloris*', EVE type 'hepadnaviruses', completeness 'entire' were chosen, then only 1 record fulfilling all those criteria will be returned. That is 'Rifleman_hepadnaviruses_1' located on the scaffold27280 of *Acanthisitta chloris* (Rifleman). If multiple records were found, then they will be listed in a table (Figure 2e). Each row begins with a unique EVE ID such as Rifleman_hepadnaviruses_1, which contains the species name, EVE type and a number to indicate its rank in the database. To provide more detailed information for every EVE, the EVE ID was hyperlinked to a detail page.

EVEs detail demonstration

In the detail information page, the following information was provided (Figure 2f): 1) EVE ID indicating the unique identifier of the EVE in FabriEVEs database; 2) Class, common name and species name specifying the name of certain animal species; 3) EVEs type representing the type of EVEs; 4) Locus showing the location of EVE on animal chromosomes or contigs; 5) completeness indicating whether the EVE is complete or partial; 6) Reference showing from which literature the EVEs data is retrieved from;

EVE data downloading

In addition to comprehensive annotation, we provided free access to all the EVEs in our database (Figure 2c). In the downloading page, EVEs data from fishes, amphibians, birds, reptiles and invertebrates were compressed as zip format files and hyperlinked to a button with the information of host class and the number of EVEs in that corresponding class. Therefore, users can download data sets from any class by clicking that button. These datasets are also available in CNGBdb.

Analysis of functions of EVEs in *Drosophila melanogaster*

To explore the functional implications EVEs, we performed comprehensive analysis on the EVEs elements in *Drosophila melanogaster*. In total, 4183 *Drosophila melanogaster* EVEs elements were stored in FabriEVEs and but after filtering only 3594 of them were analyzed using ChIPseeker (references). The results suggest that the majority of EVEs were distributed on chromosome 2R, 3L and 3R (Figure 3a). To assess the genomic features of EVEs, we calculated the frequency of EVEs in different genomic regions including promoter ($\leq 1\text{kb}$), promoter (1–2kb), promoter (2–3kb), 5' UTR, 3' UTR, other exon, 1st Intron, other intron, downstream ($\leq 3\text{kb}$) and distal Intergenic. We noticed that fruit fly EVEs are most enriched in distal intergenic and intron region, implying that they might function as cis regulatory elements (Figure 3b). To infer the potential functions of those EVEs, we predicted their targets using the “annotatePeak” function from ChIPseeker that regard the nearest gene as the target of corresponding EVE, resulting in 313 putative genes. GO term analysis showed that those putative targets are enriched in GO terms associated with “regulation of organelle organization”, “intracellular signal transduction” and “regulation of vesicle fusion”, “SUMOylation of DNA damage response and repair proteins” and “Transmission across Chemical Synapses” (Figure

3c). As shown in Figure 3d, multiple EVEs elements were located around *step* and *cta* genes, associated with the GO term “regulation of organelle organization”. Further experiments need to be performed to investigate if those EVEs elements indeed regulate the expression patterns of nearby genes.

DISCUSSION

The classification of EVEs from 82 non-mammalian species makes it possible to dissect EVEs in a comprehensive and systematic way. The annotation of EVEs deriving from a variety of literature with a standard script makes the EVEs are comparable to each other which greatly facilitates the comparative analysis on them. Flexible searching options supplied users with powerful and efficient strategies to find EVEs they are interested in. The present of data in user-friendly web interfaces and offering the data for downloading accelerate the sharing and spreading of those valuable EVEs resources. Benefiting from those enormous conveniences provided by FabriEVEs, we observed several unique properties of EVEs in non-mammalian organisms, which give rise to following questions: 1) The copy number of non-retroviral EVEs are much higher in *Phalacrocorax carbo* (cormorant) compared to other bird species. Did it possibly contribute to the evolution of cormorant? 2) The non-retroviral EVEs in birds (225 out of 251) mainly origin from *hepadnaviridae*, while the non-retroviral EVEs reported in invertebrates (136 out of 330) are mainly from *densovirinae*. Is this related to the life styles of *hepadnaviridae* and *densovirinae*? 3) Hundreds of non-retroviruses deriving EVEs were found in fishes, amphibians, birds, reptiles and invertebrates. Of particular interest, high diversity of EVEs were reported in birds and invertebrates. Unlike retroviruses, most non-retroviruses viruses could not integrate into host genome and some RNA viruses do not even have a double

strand DNA phase. How did those viruses successfully integrate into animal genome and subsequently become domesticated and co-adapted? 4) In total, 66 copies of complete or nearly full length EVEs were recorded in FabriEVEs. Are those EVEs functional, either working as regulatory elements or actively expressed? To what extends do they differ from their extant counterparts? How could they remain intact after millions of years since the initial integration event. 5) As the progress of genomic, transcriptomics and proteomics studies, there will be plenty of transcription and translation data available for those EVEs. It would be interesting to screen the expression of those EVEs to explore their functions. Overall, those questions illustrate that fascinating biological questions could be inspired by comparing EVEs data sets from various sources integrated into a systematic database such as FabriEVEs.

Though thousands of EVEs in fishes, amphibians, birds, reptiles and invertebrates have been reported, a specialized and dedicated database for them is still missing. FabriEVEs is the first database specially designed to collect, classify, annotate and present non-mammalian EVEs, which will lay the foundation for systematic analysis of EVEs. More importantly, it provides an opportunity to compare EVEs from species with various evolutionary distances, which might speed up our understanding regarding the evolutionary patterns and phylogenic relationships among those species. Currently, there are 1.5 million records in FabriEVEs. We are planning to update our database regularly to include the latest EVEs data reported by researchers in this field. Hopefully, FabriEVEs will help us to get a deeper understanding of the roles EVEs play during the evolution history of non-mammalian animals and throw light upon the co-adaptation between the endogenous viruses and their hosts.

REFERENCES

1. Feschotte, C. and Gilbert, C. (2012) Endogenous viruses: insights into viral evolution and impact on host biology., *Nat. Rev. Genet.*, **13**, 283–296.
2. Katzourakis, A. and Gifford, R. J. (2010) Endogenous Viral Elements in Animal Genomes, *PLoS Genet.*, **6**, e1001191.
3. Holmes, E. C. (2011) The Evolution of Endogenous Viral Elements, *Cell Host Microbe*, **10**, 368–377.
4. Cui, J., Zhao, W., Huang, Z., et al. (2014) Low frequency of paleoviral infiltration across the avian phylogeny., *Genome Biol.*, **15**, 539.
5. Emerman, M. and Malik, H. S. (2010) Paleovirology—Modern Consequences of Ancient Viruses, *PLoS Biol.*, **8**, e1000301.
6. Chen, D.-S., Wu, Y.-Q., Zhang, W., et al. (2016) Horizontal gene transfer events reshape the global landscape of arm race between viruses and homo sapiens, *Sci. Rep.*, **6**, 26934.
7. Feschotte, C. (2008) Transposable elements and the evolution of regulatory networks, *Nat. Rev. Genet.*, **9**, 397–405.
8. Conley, A. B., Piriyaopongsa, J. and Jordan, I. K. (2008) Retroviral promoters in the human genome, *Bioinformatics*, **24**, 1563–1567.
9. Cohen, C. J., Lock, W. M. and Mager, D. L. (2009) Endogenous retroviral LTRs as promoters for human genes: a critical assessment., *Gene*, **448**, 105–114.
10. Aswad, A. and Katzourakis, A. (2012) Paleovirology and virally derived immunity., *Trends Ecol. Evol.*, **27**, 627–636.

11. Dupressoir, A., Vernochet, C., Bawa, O., et al. (2009) Syncytin-A knockout mice demonstrate the critical role in placentation of a fusogenic, endogenous retrovirus-derived, envelope gene., *Proc. Natl. Acad. Sci. U. S. A.*, **106**, 12127–12132.
12. Dupressoir, A., Vernochet, C., Harper, F., et al. (2011) A pair of co-opted retroviral envelope syncytin genes is required for formation of the two-layered murine placental syncytiotrophoblast., *Proc. Natl. Acad. Sci. U. S. A.*, **108**, E1164--73.
13. Chuong, E. B., Elde, N. C. and Feschotte, C. (2016) Regulatory evolution of innate immunity through co-option of endogenous retroviruses, *Science (80-.)*, **351**, 1083–1087.
14. Arnaud, F., Caporale, M., Varela, M., et al. (2007) A paradigm for virus-host coevolution: sequential counter-adaptations between endogenous and exogenous retroviruses., *PLoS Pathog.*, **3**, e170.
15. Mura, M., Murcia, P., Caporale, M., et al. (2004) Late viral interference induced by transdominant Gag of an endogenous retrovirus, *Proc. Natl. Acad. Sci.*, **101**, 11117–11122.
16. Stoye, J. P. (1998) Fv1, the mouse retrovirus resistance gene., *Rev. Sci. Tech.*, **17**, 269–277.
17. Yan, Y., Buckler-White, A., Wollenberg, K., et al. (2009) Origin, antiviral function and evidence for positive selection of the gammaretrovirus restriction gene Fv1 in the genus *Mus*., *Proc. Natl. Acad. Sci. U. S. A.*, **106**, 3259–3263.
18. McDougall, A. S., Terry, A., Tzavaras, T., et al. (1994) Defective endogenous proviruses are expressed in feline lymphoid cells: evidence for a role in natural resistance to subgroup B feline leukemia viruses., *J. Virol.*, **68**, 2151–2160.

19. modENCODE Consortium, S., Roy, S., Ernst, J., et al. (2010) Identification of functional elements and regulatory circuits by Drosophila modENCODE., *Science*, **330**, 1787–1797.
20. (2009) Genome 10K: A Proposal to Obtain Whole-Genome Sequence for 10 000 Vertebrate Species, *J. Hered.*, **100**, 659–674.
21. Zhang, G., Li, C., Li, Q., et al. (2014) Comparative genomics reveals insights into avian genome evolution and adaptation., *Science*, **346**, 1311–20.
22. Zhang, G. (2015) Genomics: Bird sequencing project takes off, *Nature*, **522**, 34–34.
23. Liu, H., Fu, Y., Xie, J., et al. (2011) Widespread endogenization of densoviruses and parvoviruses in animal and human genomes., *J. Virol.*, **85**, 9863–9876.
24. Han, G.-Z., Hart, D., Frerichs, G., et al. (2015) Extensive retroviral diversity in shark, *Retrovirology*, **12**, 34.
25. Gilbert, C., Meik, J. M., Dashevsky, D., et al. (2014) Endogenous hepadnaviruses, bornaviruses and circoviruses in snakes., *Proceedings. Biol. Sci.*, **281**, 20141122.
26. Horie, M., Kobayashi, Y., Suzuki, Y., et al. (2013) Comprehensive analysis of endogenous bornavirus-like elements in eukaryote genomes., *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, **368**, 20120499.
27. Sinzelle, L., Carradec, Q., Paillard, E., et al. (2011) Characterization of a *Xenopus tropicalis* endogenous retrovirus with developmental and stress-dependent expression., *J. Virol.*, **85**, 2167–2179.
28. Crisp, A., Boschetti, C., Perry, M., et al. (2015) Expression of multiple horizontally acquired genes is a hallmark of both vertebrate and invertebrate genomes., *Genome Biol.*, **16**, 50.

29. Thézé, J., Leclercq, S., Moumen, B., et al. (2014) Remarkable diversity of endogenous viruses in a crustacean genome., *Genome Biol. Evol.*, **6**, 2129–2140.
30. Paces, J., Pavlícek, A., Zika, R., et al. (2004) HERVd: the Human Endogenous RetroViruses Database: update, *Nucleic Acids Res.*, **32**, 50D--50.
31. Paces, J., Pavlícek, A. and Paces, V. (2002) HERVd: database of human endogenous retroviruses., *Nucleic Acids Res.*, **30**, 205–206.
32. gEVE version 1.1 – Genome-based Endogenous Viral Element Database, version 1.
33. Database.
34. Jarvis, E. D., Mirarab, S., Aberer, A. J., et al. (2014) Whole-genome analyses resolve early branches in the tree of life of modern birds., *Science*, **346**, 1320–31.
35. Zhang, G., Li, B., Li, C., et al. (2014) Comparative genomic data of the Avian Phylogenomics Project, *Gigascience*, **3**, 26.
36. Zhang, G., Li, C., Li, Q., et al. (2014) Comparative genomics reveals insights into avian genome evolution and adaptation., *Science*, **346**, 1311–1320.
37. Hillier, L. W., Miller, W., Birney, E., et al. (2004) Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution., *Nature*, **432**, 695–716.
38. Yu, G., Wang, L. G. and He, Q. Y. (2015) ChIP seeker: An R/Bioconductor package for ChIP peak annotation, comparison and visualization, *Bioinformatics*.

Table 1. Number of EVEs in fishes, amphibians, birds, reptiles and invertebrates deriving from several viruses families(*hepadnaviridae*, *bornaviridae*, *circoviridae*, *parvoviridae*, *filoviridae* and *retroviridae*).

Class	Hepadnaviridae	Bornaviridae	Circoviridae	Parvoviridae	Filoviridae	Retroviridae	Others	Total
Fishes	0	0	0	2	0	19	0	21
Amphibians	0	0	0	0	0	1	0	1
Birds	225	6	5	13	0	1,478,857	0	1,479,106
Reptiles	4	3	6	0	0	0	2	15
Invertebrates	0	1	20	12	28	4183	269	4513

Figure 1

bioRxiv preprint doi: <https://doi.org/10.1101/529354>; this version posted January 24, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

Figure 2

bioRxiv preprint doi: <https://doi.org/10.1101/529354>; this version posted January 24, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.

a

1. Search Endogenous non-retroviruses (ENRVs)

Class:

Species:

EVEs type: ☐ Bornaviridae ☐ Bunyaviridae ☐ Circoviridae ☐ Densoviridae ☐ Flaviviridae ☐ Hepadnaviridae ☐ Mononegavirales ☐ Orthomyxoviridae ☐ Parvoviridae ☐ Reoviridae ☐ Retroviridae ☐ Rhabdoviridae ☐ Totiviridae ☐ Unassigned

Partial or entire: ☒ Entire ☐ Partial ☐ Unknown

2. Search Endogenous retroviruses (ERVs)

Species:

b

Fish ERVs

Callorhinchus milii, elephant shark

2 ERVs

Amphibians ERVs

Xenopus tropicalis, Western clawed frog

1 ERVs

Bird ERVs

Acanthisitta chloris Rifleman
38,983 ERVs

Apaloderma vittatum Bar-tailed trogon
28,039 ERVs

Aptenodytes forsteri Emperor penguin
45,107 ERVs

Buceros rhinoceros Rhinoceros hornbill
26,366 ERVs

Calypte anna Anna's hummingbird
27,693 ERVs

Cariama cristata Red-legged seriema
26,829 ERVs

Cathartes aura Turkey vulture
26,404 ERVs

Chaetura pelagica Chimney swift
22,562 ERVs

Charadrius vociferus Killdeer
31,218 ERVs

c

Home Search Browse **Download** Contact

FISHES EVES

AMPHIBIANS EVES

BIRDS EVES

REPTILES EVES

INVERTEBRATES EVES

ALL EVES FOR 5 CLASSES

d

Home Search Browse Download **Contact**

Contact Us

If you have any questions regarding FabriEVEs, please do not hesitate to contact us

Dong-Sheng Chen,
Email: chendongsheng@genomics.cn
Maternal and Child Health Institute, BGI Research

e

Home **Search** Browse Download Contact

Search Results

EVE_ID	Species_name	EVEs_type	Locus	Class
Acanthisitta_chloris_Retroviridae_1	Acanthisitta_chloris	Retroviridae	Scaffold18431:9120-9662	Birds
Acanthisitta_chloris_Retroviridae_2	Acanthisitta_chloris	Retroviridae	Scaffold32211:3-890	Birds
Acanthisitta_chloris_Retroviridae_3	Acanthisitta_chloris	Retroviridae	Scaffold45681:7205-7277	Birds
Acanthisitta_chloris_Retroviridae_4	Acanthisitta_chloris	Retroviridae	Scaffold46351:1-139	Birds
Acanthisitta_chloris_Retroviridae_5	Acanthisitta_chloris	Retroviridae	scaffold10031:10177-10267	Birds
Acanthisitta_chloris_Retroviridae_6	Acanthisitta_chloris	Retroviridae	scaffold10091:8-159	Birds
Acanthisitta_chloris_Retroviridae_7	Acanthisitta_chloris	Retroviridae	scaffold10281:3-128	Birds
Acanthisitta_chloris_Retroviridae_8	Acanthisitta_chloris	Retroviridae	scaffold10281:214-407	Birds
Acanthisitta_chloris_Retroviridae_9	Acanthisitta_chloris	Retroviridae	scaffold10281:20820-21130	Birds
Acanthisitta_chloris_Retroviridae_10	Acanthisitta_chloris	Retroviridae	scaffold10281:21131-22126	Birds
Acanthisitta_chloris_Retroviridae_11	Acanthisitta_chloris	Retroviridae	scaffold10281:22131-22193	Birds
Acanthisitta_chloris_Retroviridae_12	Acanthisitta_chloris	Retroviridae	scaffold10301:2908-3268	Birds
Acanthisitta_chloris_Retroviridae_13	Acanthisitta_chloris	Retroviridae	scaffold10301:3868-4015	Birds
Acanthisitta_chloris_Retroviridae_14	Acanthisitta_chloris	Retroviridae	scaffold10331:1219-1498	Birds
Acanthisitta_chloris_Retroviridae_15	Acanthisitta_chloris	Retroviridae	scaffold10331:1700-1752	Birds

f

Detail information

EVE ID	Acanthisitta_chloris_Retroviridae_1
Class	Birds
Species name	Acanthisitta_chloris
EVEs type	Retroviridae
Locus	Scaffold18431:9120-9662
Entire or partial	Unknown
Reference	Zhang, G. et al. Comparative genomics reveals insights into avian genome evolution and adaptation. Science 346, 1311-20 (2014).

View the location in genome browser of Avianbase

NATIONAL AVIAN RESEARCH FACILITY

BLAST/BLAT | Tools | Downloads | Search Rifleman...

Rifleman Location: Scaffold18431:9,120-9,662 Jobs

Location-based displays

- Whole genome
- Chromosome summary
- Region overview
- Region in detail**
- Comparative Genomics
- Alignments (image)
- Alignments (text)
- Region Comparison
- Synteny

Scaffold18431: 9,120-9,662

scaffold Scaffold18431

Region in detail

Figure 3

bioRxiv preprint doi: <https://doi.org/10.1101/529354>; this version posted January 24, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC 4.0 International license.