Targeted Cleavage and Polyadenylation of RNA by CRISPR-Cas13

Kelly M. Anderson^{1,2}, Pornthida Poosala^{1,2}, Sean R. Lindley^{1,2} and Douglas M. Anderson^{1,2,*}

¹Center for RNA Biology, ²Aab Cardiovascular Research Institute, University of Rochester School of Medicine and Dentistry, Rochester, New York, U.S.A., 14642

*Corresponding Author:

Douglas M. Anderson: doug_anderson@urmc.rochester.edu

Keywords: NUDT21, PspCas13b, PAS, CPSF, CFIm, Poly(A), SREBP1

Post-transcriptional cleavage and polyadenylation of messenger and long noncoding RNAs is coordinated by a supercomplex of ~20 individual proteins within the eukaryotic nucleus^{1,2}. Polyadenylation plays an essential role in controlling RNA transcript stability, nuclear export, and translation efficiency³⁻⁶. More than half of all human RNA transcripts contain multiple polyadenylation signal sequences that can undergo alternative cleavage and polyadenylation during development and cellular differentiation^{7,8}. Alternative cleavage and polyadenylation is an important mechanism for the control of gene expression and defects in 3' end processing can give rise to myriad human diseases^{9,10}. Here we show that fusion of catalytically dead Cas13 to a single mammalian polyadenylation factor, Nudix hydrolase 21 (NUDT21), allows for site-specific CRISPR-Cas13-guided cleavage and polyadenylation of RNA in mammalian cells. This approach, which we named *Postscriptr*, can be utilized for the non-genomic manipulation of gene expression and may have potential future therapeutic applications for treating human RNA processing diseases.

The 3' site of RNA cleavage and addition of a poly(A) tail is precisely determined by intrinsic polyadenylation signal (PAS) sequences typically composed of a canonical hexamer motif AAUAAA and upstream and downstream sequence elements (USE and DSE, respectively)¹¹. The AAUAAA PAS motif is found ~25 nucleotides (nts) upstream of the RNA cleavage site and is directly bound by two components of the cleavage and polyadenylation specific factor (CPSF) complex: cleavage and polyadenylation factor 30 (CPSF30) and WD repeat-containing protein 33 (WDR33)¹². Components of the cleavage factor Im (CFIm) complex bind directly to the USE motif UGUA which occurs

~50 nts upstream of the RNA cleavage site⁸. In transcripts that lack the canonical AAUAAA motif, the CFIm complex functions as the primary determinant of poly(A) signal recognition¹³. Nudix Hydrolase 21 (NUDT21/CPSF5), the RNA binding component of the CFIm complex, functions as an activator of 3' end processing and regulator of alternative polyadenylation site choice¹⁴. Binding of the CFIm complex is among the first steps in 3' end processing and functions to recruit additional co-factors to the 3' end processing machinery^{15,16}. Direct interactions between components of the polyadenylation supercomplex and the C-terminal domain of RNA polymerase II trigger the disassembly of the elongation complex and coordinate 3' end processing with transcription termination¹⁷.

Bacterial-derived Type VI CRISPR-Cas systems encode a family of RNA-guided endoribonucleases, named Cas13¹⁸⁻²⁰. Similar to Cas9, mutation of residues within the nuclease domains of Cas13 generate a catalytically dead enzyme that retains RNA binding affinity (dCas13). Recently, fusion of dCas13 to mammalian RNA modifying enzymes has been shown to be useful for manipulating RNA in mammalian cells²¹⁻²³. To direct polyadenylation complex formation using dCas13, we designed three fusion proteins combining the catalytically dead Type VI-B Cas13 enzyme from *Prevotella sp. P5-125* (dPspCas13b) with RNA binding components of the human 3' end processing machinery, including CPSF30, WDR33, and NUDT21 (Fig. 1a). We included an N-terminal 3x FLAG epitope tag for protein detection and a long flexible peptide linker [GGGGSGGGGS] between dPspCas13b and the polyadenylation components to reduce the likelihood of steric hindrance.

While previous dCas13-mediated RNA editing applications occurred within the cytosol, post-transcriptional cleavage and polyadenylation of RNA occurs within the nucleus and therefore requires efficient nuclear localization of dCas13 fusion proteins. The large size of dCas13 proteins and their lack of intrinsic nuclear localization signals (NLSs) could prevent their efficient nuclear localization in mammalian cells. Consistent with this, dPspCas13b fusion proteins lacking a mammalian NLS were retained in the cytoplasm when expressed in mammalian COS7 cells, as detected by immunocytochemistry using an anti-FLAG antibody (Supplementary Fig. 1). Surprisingly, the addition of a classical SV40 NLS or bipartite NLS from nucleoplasmin (NPM) did not promote efficient nuclear localization of dPspCas13b fusion proteins (Supplementary Fig. 1). However, the addition of a single copy of the non-classical bipartite NLS derived from the yeast Ty1 retrotransposon, which utilizes normal cellular protein import machinery but contains a linker sequence nearly three times as long as a typical bipartite NLS^{24,25}, resulted in robust nuclear localization of all three dPspCas13b fusion proteins (Fig. 1b).

We designed a reporter gene to detect alternative cleavage and polyadenylation (APA) in mammalian cells capable of switching between fluorescent and non-fluorescent open reading frames of superfolder GFP (sfGFP) (Fig. 2). sfGFP forms a beta barrel comprised of 11 antiparallel beta strands, which can tolerate sequence insertions between the 10th and 11th beta strands but loses fluorescence if the 11th beta strand is removed²⁶⁻²⁸. We generated an sfGFP reporter construct with the coding sequence for the 11th beta strand embedded within a prototypical mammalian intron (second intron of the rabbit beta globin gene) (sfGFP_{Papa}) (Fig. 2a). The coding

sequence of the 11th beta strand was designed in-frame with the upstream sfGFP coding sequence so that translation of the proximal open reading frame would encode a complete sfGFP sequence, albeit with a 14 amino acid linker sequence between the 10th and 11th beta strand resulting from translation of the intervening intronic sequence [sfGFP(1-10)-L-11]. Expression of the sfGFP_{Papa} reporter in mammalian COS7 cells resulted in almost no detectable fluorescence, suggesting efficient removal of the intron containing the 11th beta strand and translation of the open reading frame encoding the non-fluorescent sfGFP(1-10) protein (Fig. 2b-d). However, cells expressing the sfGFP_{Papa} reporter treated with the splicing inhibitor isoginkgetin²⁹ resulted in detectable green fluorescence after 24 hours, demonstrating that the modified sfGFP open reading frame containing the linker between the 10th and 11th beta strands is functional (Supplementary Fig. 2a,b). Insertion of an SV40 polyadenylation cassette downstream of coding sequence for the 11th beta strand of sfGFP (sfGFP_{Papa}-pA) resulted in robust green fluorescence, demonstrating that intronic cleavage and polyadenylation of the sfGFP_{Papa} reporter promotes utilization of the functional sfGFP(1-10)-L-11 open reading frame (Fig. 2e-h). Together, these data demonstrate that the sfGFP_{Papa} reporter gene encodes both functional and non-functional open reading frames which can be switched in response to changes in splicing or intronic cleavage and polyadenylation.

To determine whether the dPspCas13b fusion proteins could promote CRISPR-Cas13-mediated cleavage and polyadenylation of the sfGFP_{Papa} reporter mRNA, we designed a crRNA targeting an intronic sequence downstream of the coding sequence of the 11th beta strand of sfGFP in the sfGFP_{Papa} reporter (Fig. 3a). In live cell assays, no fluorescence signal was detected in cells expressing any of the dPspCas13b fusion

proteins using a non-targeting crRNA, relative to the sfGFP_{Papa} reporter alone (Fig. 3b, Supplementary Fig. 3). However, expression of the intron targeting crRNA with the dPspCas13b-NUDT21 fusion protein, but not the CPSF30 or WDR33 fusions proteins, resulted in detectable green fluorescent cells after 24 hours (Fig. 3b, Supplementary Fig. 3).

We performed 3' rapid amplification of cDNA ends (3' RACE) to determine whether the fluorescence signal was due to CRISPR-Cas13-mediated targeted cleavage and polyadenylation of the sfGFP_{Papa} reporter mRNA. RACE compatible cDNA was generated from total RNA using a poly(T) oligonucleotide containing two 5' nested primer sequences. Using two nested reporter specific primers upstream of the crRNA target sequence, we detected a high molecular weight band corresponding to the predicted size of the sfGFP_{Papa} pre-mRNA transcript for both samples (Fig. 3c). Interestingly, in cells targeted with the intronic crRNA, we detected an additional broad band of smaller RACE products, suggesting that cleavage and polyadenylation may have occurred near the targeted crRNA sequence (Fig. 3c,d). To determine the sequence of the 3' RACE products, we gel purified the 200-500 base pair region and subcloned them using TOPO TA cloning. We isolated and sequenced 11 unique clones which revealed that sfGFP_{Papa} mRNAs were cleaved and polyadenylated at sites ranging from -7 to +110 nts relative to the 3' side of the crRNA target sequence (Fig. 3e). In contrast to mammalian cleavage which occurs primarily after a CA dinucleotide, *Postscriptr*-induced cleavage most often occurred at a C or T nucleotide (Fig. 3f). Interestingly, four of the 11 clones showed a 3' addition of non-templated nucleotides prior to poly (A) tail elongation, which is thought to be rare in mammalian species but

observed in plants³⁰. These data demonstrate that the dPspCas13b-NUDT21 fusion protein was sufficient to induce cleavage and polyadenylation of a reporter mRNA at an intronic sequence targeted by a crRNA.

To determine whether *Postscriptr* could promote alternative cleavage and polyadenylation of an endogenously expressed human mRNA, we targeted transcripts encoding the human sterol regulatory element binding protein 1 (SREBP1). SREBP1 is a ubiquitously expressed transcription factor which transactivates genes that contain sterol regulatory elements (SREs) and encode proteins controlling lipid synthesis and uptake, such as the low density lipoprotein receptor (*LDLR*) (Fig. 4)³¹. SREBP1 is first synthesized as an inactive precursor protein anchored to the membrane of the endoplasmic reticulum (ER)³² (Fig. 4a). In sterol depleted cells, proteolytic cleavage of SREBP1 by S2P liberates an N-terminal DNA-binding fragment that allows SREBP1 to enter the nucleus and activate gene transcription³³ (Fig. 4a). We note that *SREBP1* transcripts in the liver can utilize an endogenous intronic PAS and generate a protein fragment terminated at a site adjacent to the transcriptionally active S2P cleavage product [SREBP1aΔ, AB373958; SREBP1cΔ, AB373959] (Fig. 4b,c). We were interested in whether *Postscriptr* could promote the utilization of the SREBP1 intronic PAS in a non-liver cell line and upregulate *SREBP1* target gene expression. We co-expressed the dPspCas13b-NUDT21 fusion protein with either an *SREBP1*-targeting or non-targeting crRNA in human HEK293T cells (Fig. 4d). Performing 3' RACE, we detected a strong band corresponding to the approximate size of the predicted 3'RACE product in cells transfected with the *SREBP1*-targeting crRNA, and no bands were detected in cells transfected with the control crRNA (Supplementary Fig. 4). Direct

sequencing of this 3' RACE product revealed that *SREBP1* transcripts were cleaved and polyadenylated 71 nts downstream of the crRNA target sequence and that splicing of the terminal exon was inhibited (Fig. 4d,e). Cleavage occurred at a CA dinucleotide 36 nts downstream of the reported cleavage site in humans and generated an open reading frame which terminated at a stop codon within the retained intron sequence, adding an additional 16 amino acids to the C-terminus of SREBP1 (Fig. 4e,f). Gene expression analyses in a mixed population of *SREBP1*-targeted and non-targeted HEK293T cells showed that the overall *SREBP1* transcript levels were unchanged when using qRT-PCR and PCR primers upstream of the crRNA target sequence (Fig. 4g). However, primers spanning the targeted exon showed a significant decrease in *SREBP1* transcript levels in targeted versus non-targeted cells, suggesting that transcription termination may be coupled with *Postscriptr*-mediated cleavage and polyadenylation (Fig. 4h). Interestingly, in *SREBP1*-targeted cells, we further detected a significant increase in transcript levels of the well-characterized SREBP-responsive gene, *LDLR* (Fig. 4i).

In conclusion, we have developed a unique CRISPR-Cas13 RNA modifying technology to induce site-specific cleavage and polyadenylation of RNA in mammalian cells. This approach utilizes a dPspCas13b-NUDT21 fusion protein, which we have shown requires a non-classical bipartite NLS for nuclear localization in mammalian cells. We demonstrated that the dPspCas13b-NUDT21 fusion protein was sufficient to promote cleavage and polyadenylation at an intronic sequence within a novel gain-of-function fluorescent reporter, sfGFP_{papa}. The ability of *Postscriptr* to promote cleavage and polyadenylation of the sfGFP_{papa} reporter was striking, given that the second intron

of the rabbit beta globin gene does not contain any known endogenous PAS sequences. These findings show that NUDT21 can facilitate the recruitment of a functional 3' end processing complex at a crRNA targeted sequence. While only 25 kDa in size, NUDT21 forms a number of direct protein interactions with key RNA processing components, including Poly(A) Polymerase³⁴. These findings support the role of NUDT21 as an emerging master regulator of alternative cleavage and polyadenylation site choice in mammals^{35,36}.

We further showed that *Postscriptr* could direct cleavage and polyadenylation at an endogenous intronic PAS of the human *SREBP1* gene. For both the sfGFPapa reporter and *SREBP1*, the dPspCas13b-NUDT21 fusion protein induced cleavage and polyadenylation 3' to the crRNA target sequence. However, it is currently unclear why editing of the intronic *SREBP1* PAS resulted in a single cleavage site, whereas cleavage of the sfGFPapa reporter occurred across a range of ~100 nts. We speculate that intrinsic PAS flanking sequences may be required to help guide cleavage site specificity. Future studies are necessary to delineate the minimal upstream and downstream sequence elements which may enhance the precision of *Postscriptr* targeting at ectopic target sites.

It is currently unclear why fusion of dPspCas13b to either CPSF30 or WDR33 were not sufficient to promote cleavage and polyadenylation of the sfGFPapa reporter in our assays. Since both CPSF30 and WDR33 have been shown to make direct contacts with the AAUAAA PAS motif, one possibility may be a requirement of the PAS RNA to promote a ternary complex formation among these components. Alternatively, tethering of CPSF30 and WDR33 may alone be insufficient to induce 3' end processing, a

possibility consistent with data showing that insertion of a minimal synthetic PAS (including an AAUAAA motif and downstream sequence elements) within the second intron of a rabbit beta globin gene reporter was insufficient to promote intronic cleavage and polyadenylation³⁷.

While the endonuclease activity of Cas13 is highly effective for gene knockdown in mammalian cells²¹, *Postscriptr* technology allows for a unique approach to manipulate RNA which may be useful for the study of 3' end RNA processing and for the manipulation of endogenous gene expression. Here we have utilized this technology to manipulate the alternative cleavage and polyadenylation at intronic pre-mRNA transcript sequences, however this approach could be applied to other mRNA transcript regions or different RNA species. One interesting example would be the interrogation of lncRNA function, many of which control gene expression through recruitment of chromatin modifying complexes or by establishing permissive local chromatin environments³⁸⁻⁴¹. *Postscriptr*-guided cleavage, polyadenylation, and transcriptional termination of lncRNAs could allow for the identification of functional lncRNA domains or functional genomic regions. Lastly, numerous loss-of-function mutations identified within poly(A) signal sequences have been shown to result in abnormal RNA processing and can cause human disease⁴²⁻⁴⁴. Here, future therapeutic applications of *Postscriptr* technology may be a useful approach for the rescue of normal RNA cleavage and polyadenylation site choice.

REFERENCES

- 1 Mandel, C. R., Bai, Y. & Tong, L. Protein factors in pre-mRNA 3'-end processing. *Cell Mol Life Sci* **65**, 1099-1122, doi:10.1007/s00018-007-7474-3 (2008).
- 2 Xiang, K., Tong, L. & Manley, J. L. Delineating the structural blueprint of the pre-mRNA 3'-end processing machinery. *Mol Cell Biol* **34**, 1894-1910, doi:10.1128/MCB.00084-14 (2014).
- 3 Colgan, D. F. & Manley, J. L. Mechanism and regulation of mRNA polyadenylation. *Genes Dev* **11**, 2755-2766 (1997).
- 4 Guhaniyogi, J. & Brewer, G. Regulation of mRNA stability in mammalian cells. *Gene* **265**, 11-23 (2001).
- 5 Wu, X. & Brewer, G. The regulation of mRNA stability in mammalian cells: 2.0. *Gene* **500**, 10-21, doi:10.1016/j.gene.2012.03.021 (2012).
- 6 Carmody, S. R. & Wente, S. R. mRNA nuclear export at a glance. *J Cell Sci* **122**, 1933-1937, doi:10.1242/jcs.041236 (2009).
- 7 Tian, B. & Manley, J. L. Alternative polyadenylation of mRNA precursors. *Nat Rev Mol Cell Biol* **18**, 18-30, doi:10.1038/nrm.2016.116 (2017).
- 8 Elkon, R., Ugalde, A. P. & Agami, R. Alternative cleavage and polyadenylation: extent, regulation and function. *Nat Rev Genet* **14**, 496-506, doi:10.1038/nrg3482 (2013).
- 9 Chang, J. W., Yeh, H. S. & Yong, J. Alternative Polyadenylation in Human Diseases. *Endocrinol Metab (Seoul)* **32**, 413-421, doi:10.3803/EnM.2017.32.4.413 (2017).
- 10 Curinha, A., Oliveira Braz, S., Pereira-Castro, I., Cruz, A. & Moreira, A. Implications of polyadenylation in health and disease. *Nucleus* **5**, 508-519, doi:10.4161/nucl.36360 (2014).
- 11 Tian, B. & Graber, J. H. Signals for pre-mRNA cleavage and polyadenylation. *Wiley Interdiscip Rev RNA* **3**, 385-396, doi:10.1002/wrna.116 (2012).

- 12 Chan, S. L. *et al.* CPSF30 and Wdr33 directly bind to AAUAAA in mammalian mRNA 3' processing. *Genes Dev* **28**, 2370-2380, doi:10.1101/gad.250993.114 (2014).
- 13 Venkataraman, K., Brown, K. M. & Gilmartin, G. M. Analysis of a noncanonical poly(A) site reveals a tripartite mechanism for vertebrate poly(A) site recognition. *Genes Dev* **19**, 1315-1327, doi:10.1101/gad.1298605 (2005).
- 14 Zhu, Y. *et al.* Molecular Mechanisms for CFIm-Mediated Regulation of mRNA Alternative Polyadenylation. *Mol Cell* **69**, 62-74 e64, doi:10.1016/j.molcel.2017.11.031 (2018).
- 15 Ruegsegger, U., Blank, D. & Keller, W. Human pre-mRNA cleavage factor Im is related to spliceosomal SR proteins and can be reconstituted in vitro from recombinant subunits. *Mol Cell* **1**, 243-253 (1998).
- 16 Brown, K. M. & Gilmartin, G. M. A mechanism for the regulation of pre-mRNA 3' processing by human cleavage factor Im. *Mol Cell* **12**, 1467-1476 (2003).
- 17 Zhao, J., Hyman, L. & Moore, C. Formation of mRNA 3' ends in eukaryotes: mechanism, regulation, and interrelationships with other steps in mRNA synthesis. *Microbiol Mol Biol Rev* **63**, 405-445 (1999).
- 18 Abudayyeh, O. O. *et al.* C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* **353**, aaf5573, doi:10.1126/science.aaf5573 (2016).
- 19 Smargon, A. A. *et al.* Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory Proteins Csx27 and Csx28. *Mol Cell* **65**, 618-630 e617, doi:10.1016/j.molcel.2016.12.023 (2017).
- 20 East-Seletsky, A., O'Connell, M. R., Burstein, D., Knott, G. J. & Doudna, J. A. RNA Targeting by Functionally Orthogonal Type VI-A CRISPR-Cas Enzymes. *Mol Cell* **66**, 373-383 e373, doi:10.1016/j.molcel.2017.04.008 (2017).
- 21 Abudayyeh, O. O. *et al.* RNA targeting with CRISPR-Cas13. *Nature* **550**, 280-284, doi:10.1038/nature24049 (2017).

- 22 Cox, D. B. T. *et al.* RNA editing with CRISPR-Cas13. *Science* **358**, 1019-1027, doi:10.1126/science.aag0180 (2017).
- 23 O'Connell, M. R. Molecular Mechanisms of RNA Targeting by Cas13-containing Type VI CRISPR-Cas Systems. *J Mol Biol* **431**, 66-87, doi:10.1016/j.jmb.2018.06.029 (2019).
- 24 McLane, L. M., Pulliam, K. F., Devine, S. E. & Corbett, A. H. The Ty1 integrase protein can exploit the classical nuclear protein import machinery for entry into the nucleus. *Nucleic Acids Res* **36**, 4317-4326, doi:10.1093/nar/gkn383 (2008).
- 25 Kenna, M. A., Brachmann, C. B., Devine, S. E. & Boeke, J. D. Invading the yeast nucleus: a nuclear localization signal at the C terminus of Ty1 integrase is required for transposition in vivo. *Mol Cell Biol* **18**, 1115-1124 (1998).
- 26 Kamiyama, D. *et al.* Versatile protein tagging in cells with split fluorescent protein. *Nat Commun* **7**, 11046, doi:10.1038/ncomms11046 (2016).
- 27 Feng, S. *et al.* Improved split fluorescent proteins for endogenous protein labeling. *Nat Commun* **8**, 370, doi:10.1038/s41467-017-00494-8 (2017).
- 28 Pedelacq, J. D., Cabantous, S., Tran, T., Terwilliger, T. C. & Waldo, G. S. Engineering and characterization of a superfolder green fluorescent protein. *Nat Biotechnol* **24**, 79-88, doi:10.1038/nbt1172 (2006).
- 29 O'Brien, K., Matlin, A. J., Lowell, A. M. & Moore, M. J. The biflavonoid isoginkgetin is a general inhibitor of Pre-mRNA splicing. *J Biol Chem* **283**, 33147-33154, doi:10.1074/jbc.M805556200 (2008).
- 30 Jin, Y. & Bian, T. Nontemplated nucleotide addition prior to polyadenylation: a comparison of Arabidopsis cDNA and genomic sequences. *RNA* **10**, 1695-1697, doi:10.1261/rna.7610404 (2004).
- 31 Horton, J. D., Goldstein, J. L. & Brown, M. S. SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* **109**, 1125-1131, doi:10.1172/JCI15593 (2002).

- 32 Brown, M. S. & Goldstein, J. L. The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell* **89**, 331-340 (1997).
- 33 Brown, M. S., Ye, J., Rawson, R. B. & Goldstein, J. L. Regulated intramembrane proteolysis: a control mechanism conserved from bacteria to humans. *Cell* **100**, 391-398 (2000).
- 34 Kim, H. & Lee, Y. Interaction of poly(A) polymerase with the 25-kDa subunit of cleavage factor I. *Biochem Biophys Res Commun* **289**, 513-518, doi:10.1006/bbrc.2001.5992 (2001).
- 35 Brumbaugh, J. *et al.* Nudt21 Controls Cell Fate by Connecting Alternative Polyadenylation to Chromatin Signaling. *Cell* **172**, 106-120 e121, doi:10.1016/j.cell.2017.11.023 (2018).
- 36 Sartini, B. L., Wang, H., Wang, W., Millette, C. F. & Kilpatrick, D. L. Pre-messenger RNA cleavage factor I (CFIm): potential role in alternative polyadenylation during spermatogenesis. *Biol Reprod* **78**, 472-482, doi:10.1095/biolreprod.107.064774 (2008).
- 37 Levitt, N., Briggs, D., Gil, A. & Proudfoot, N. J. Definition of an efficient synthetic poly(A) site. *Genes Dev* **3**, 1019-1025 (1989).
- 38 Anderson, K. M. *et al.* Transcription of the non-coding RNA upperhand controls Hand2 expression and heart development. *Nature* **539**, 433-436, doi:10.1038/nature20128 (2016).
- 39 Joung, J. *et al.* Genome-scale activation screen identifies a lncRNA locus regulating a gene neighbourhood. *Nature* **548**, 343-346, doi:10.1038/nature23451 (2017).
- 40 Engreitz, J. M. *et al.* Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature* **539**, 452-455, doi:10.1038/nature20149 (2016).
- 41 Rinn, J. L. & Chang, H. Y. Genome regulation by long noncoding RNAs. *Annu Rev Biochem* **81**, 145-166, doi:10.1146/annurev-biochem-051410-092902 (2012).

- 42 Higgs, D. R. *et al.* Alpha-thalassaemia caused by a polyadenylation signal mutation. *Nature* **306**, 398-400 (1983).
- 43 Orkin, S. H., Cheng, T. C., Antonarakis, S. E. & Kazazian, H. H., Jr. Thalassemia due to a mutation in the cleavage-polyadenylation signal of the human beta-globin gene. *EMBO J* **4**, 453-456 (1985).
- 44 Rund, D. *et al.* Two mutations in the beta-globin polyadenylation signal reveal extended transcripts and new RNA polyadenylation sites. *Proc Natl Acad Sci U S A* **89**, 4324-4328 (1992).

ACKNOWLEDGEMENTS

We thank Dr. Lynne Maquat for comments on the manuscript and Leisha Manchin for technical assistance. pC0043-PspCas13b crRNA backbone was a gift from Feng Zhang (Addgene plasmid # 103854; <http://n2t.net/addgene:103854> ; RRID:Addgene_103854). pC0049-EF1a-dPSPCas13b-NES-HIV, H133A/H1058A was a gift from Feng Zhang (Addgene plasmid # 103865; <http://n2t.net/addgene:103865> ; RRID:Addgene_103865).

AUTHOR CONTRIBUTIONS

K.M.A. and D.M.A. designed the experiments. K.M.A., P.P., S.R.L and D.M.A performed the experiments and analyzed data. K.M.A. and D.M.A generated the figures and wrote the manuscript.

AUTHOR INFORMATION

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to D.M.A. (doug_anderson@urmc.rochester.edu).

FIGURE LEGENDS

Figure 1 Design and expression of fusion proteins encoding catalytically dead Cas13 and human polyadenylation complex factors. **(a)** Diagram of the vectors encoding fusions between catalytically dead PspCas13b and components of the mammalian polyadenylation supercomplex, CPSF30, WDR33, and NUDT21. F – 3xFLAG epitope; NLS – Ty1 nuclear localization sequence; pA – SV40 polyadenylation sequence. **(b)** Immunohistochemistry using a primary anti-FLAG antibody and an Alexa488 conjugated secondary antibody detecting the nuclear localization of the dPspCas13b fusion proteins expressed in mammalian COS7 cells. Scale bars, 10 μ m. Results are representative of 3 independently performed experiments.

Figure 2 Design and Generation of a Fluorescent Reporter for RNA Cleavage and Polyadenylation in Mammalian Cells. **(a)** Diagram of the sfGFPapa reporter gene. **(b)** The green fluorescent protein superfolder GFP (sfGFP) forms a beta barrel composed of 11 beta strands. Removal of the 11th beta strand abolishes fluorescence. **(c)** Transcription of the sfGFPapa reporter and processing of the resulting transcript removes the coding sequence of the 11th sfGFP beta strand. **(d)** Removal results in a lack of fluorescent signal in mammalian cells. **(e)** Diagram of the sfGFPapa-pA reporter plasmid, which contains an upstream polyadenylation sequence. **(f)** sfGFP can tolerate linker sequences between the 10th and 11th beta strands without abolishing fluorescence. **(g)** Transcription of the sfGFPapa-pA reporter and processing of the resulting transcript. **(h)** Processed transcripts result in robust green fluorescence due to

translation of the sfGFP(1-10)-L-(11) functional open reading frame. Scale bars in (d) and (h), 100 μ m. Results are representative of 3 independently performed experiments.

Figure 3 CRISPR-Cas13-mediated Cleavage and Polyadenylation of a Reporter mRNA in Mammalian Cells. (a) Diagram showing the *Postscriptr* targeting strategy to induce alternative cleavage and polyadenylation of the sfGFP_{Papa} reporter construct. A crRNA was designed to target an intronic sequence downstream of the coding sequence of the 11th beta strand of sfGFP. (b) Expression of the dPspCas13b-NUDT21 fusion protein and intronic targeting crRNA resulted in green fluorescent cells after 24 hours. (c) 3'RACE products amplified from cells expressing dPspCas13b-NUDT21 fusion protein and non-targeting control or intronic targeting crRNA. (d,e) Sequences of the five most proximal 3'RACE products relative to the crRNA target site. Please see Table 1 for complete list of 3'RACE sequences. Green underlined and lowercase nucleotides highlight 3' non-templated nucleotide addition. (f) Nucleotide frequencies at *Postscriptr*-mediated cleavage and polyadenylation sites.

Figure 4 *Postscriptr*-mediated Alternative Cleavage and Polyadenylation of human SREBP1 in HEK293T cells. (a) Diagram of the SREBP1 maturation pathway. (b) The human SREBP1 locus contains an intronic PAS between exon 7 and exon 8, which results in translation of an SREBP1 Δ isoform which terminates translation adjacent to the normal S2P cleavage site (c). (d) Diagram of the *Postscriptr*-mediated targeting strategy and sequence of the recovered *Postscriptr*-induced cleaved and polyadenylated transcript (e) and predicted translational stop (f). Quantitative realtime-

PCR (qRT-PCR) gene expression analysis of SREBP1 transcript levels upstream of the crRNA target sequence (**g**), spanning the crRNA target sequence (**h**) and transcript levels of the LDLR gene (**i**) in HEK293T cells targeted with a non-targeted or SREBP1-targeted crRNA.

FIGURES

Figure 1

Figure 2

Figure 4

METHODS

Synthetic DNA and Cloning

The coding sequences of human CPSF30, WDR33 and NUDT21 were designed and synthesized for assembly as gBlocks (IDT, Integrated DNA Technologies) and cloned as C-terminal fusions to dPspCas13b into a modified version of the CS2 mammalian expression vector containing a 5' T7 promoter (CSX). The sfGFP_{Papa} reporter was designed and synthesized as two separate gBlock gene fragments and subcloned into the CS2 mammalian expression vector. Guide RNAs were designed antisense to pre-mRNA sequences of the sfGFP_{Papa} reporter or human *SREPB1* transcripts and were cloned into pC00043²². All guide RNAs were designed to be 30 nts in length and start with a 5' G. All plasmid insert sequences were fully verified by Sanger sequencing. Please see Supplementary Table 1 for fusion protein and reporter sequences.

Cell Lines, Transient Transfections, and Drug Treatments

The COS7 cell line was maintained in DMEM supplemented with 10% Fetal Bovine Serum (FBS) with penicillin/streptomycin at 37°C in an atmosphere of 5% CO₂. Cells were seeded and transiently transfected using Fugene6 (Promega) according to manufacturer's protocol. Twenty-four hours after transfection, cells were either imaged, or treated with DMSO or Isoginkgetin and cultured for an additional 24 hours prior to imaging. For live cell imaging, cells were washed twice with DPBS, counterstained with 10 µg/ml Hoechst 33342 (Invitrogen) in DPBS for 10 minutes at room temperature and imaged on a ZOE Fluorescent Cell Imager (Bio-Rad).

Antibodies and Immunohistochemistry

Transiently transfected COS7 cells were fixed in 4% formaldehyde in PBS for 20 minutes, blocked in 3% Bovine Serum Albumin (BSA) and incubated with primary anti-FLAG (Sigma, F1864) at 1:1000 in 1% BSA for 1 hour at room temperature. Cells were subsequently incubated with an Alexa488 conjugated secondary antibody (Thermofisher) in 1% BSA for 30 minutes at room temperature. Coverslips were mounted using anti-fade fluorescent mounting medium containing DAPI (Vector Biolabs, H-1200) and imaged using confocal microscopy.

3' Rapid Amplification of cDNA Ends (3' RACE)

Transfected cells were washed twice with DPBS and harvested in 1 ml of Trizol to isolate total RNA. RACE compatible cDNA was generated using SuperScript III (Invitrogen) according to manufacturer's protocol, with the exception of the addition of a custom oligo d(T) oligonucleotide containing two 5' nested primer sequences. RACE PCR was performed using two nested target gene specific primers and touchdown PCR protocol as follows: 94°C for 1 minutes (1 cycle); 94°C for 30 seconds, 72°C for 1 minute (5 cycles); 94°C for 30 seconds, 70°C for 4 minutes (5 cycles); 94°C for 20 seconds, 68°C for 1 minute (25 cycles); 72°C for 5 minutes. Please see Supplementary Table 1 for a list of qRT-PCR primers and 3'RACE PCR primers.

SUPPLEMENTARY DATA

SUPPLEMENTARY FIGURES

Supplementary Figure 1 Classic mammalian nuclear localization signals were insufficient to promote nuclear localization of dPspcas13b fusion proteins. Immunohistochemistry using a primary anti-FLAG antibody and secondary Alexa488 conjugated secondary antibody were used to detect the localization of dPspCas13b fusion proteins expressed in mammalian COS7 cells. Fusion proteins contained either no NLS, a classic SV40 NLS or the bipartite NLS from Nucleoplasmin (NPM). Scale bars, 10 μ m.

Supplementary Figure 2 Inhibiting splicing promotes expression of the sfGFP(1-10)-L-(11) open reading frame. **(a)** Diagram of the sfGFPapa reporter and predicted transcription and processing steps resulting from treatment with the splicing inhibitor isoginkgetin. **(b)** COS7 cells transiently transfected with the sfGFPapa reporter for 24 hours were treated with either DMSO or isoginkgetin. Cells treated with isoginkgetin resulted in detectable green fluorescence after 24 hours. Scale bars in **(b)**, 100 μ m.

Supplementary Figure 3 The dPspCas13b fusions to CPSF30 or WDR33 were not sufficient to promote cleavage and polyadenylation of the sfGFP_{Papa} reporter mRNA. (a) *Postscriptr* targeting of the sfGFP_{Papa} reporter using the dPspCas13b-CPSF30 or (b) – WDR33 fusion proteins using the sfGFP_{Papa} intronic-targeting crRNA did not result in detectable green fluorescence relative to a control non-targeting crRNA. Scale bars in (a) and (b), 100 μm.

Supplementary Figure 4 *Postscriptr*-mediated alternative cleavage and polyadenylation of human SREBP1 transcripts. PCR amplified 3' RACE products from cells expressing dPspCas13b-NUDT21 with non-targeting control or SREBP1-targeting crRNAs.

qRT-PCR Oligonucleotides

SREBP1 (Upstream)

F: 5' -AGCTCAAGGATCTGGTGGTG-3'

R: 5' -GACAGCAGTGCAGACTTA-3'

SREBP1 (Downstream)

F: 5' -GACAGCCCAGTCTTTGAGGA-3'

R: 5' -CAGGACAGGCAGAGGAAGAC-3'

LDLR

F: 5' -AAGGACACAGCACACAACCA-3'

R: 5' -CTGGGCTTCTTCTCATTTCC-3'

sfGFP_{Papa} reporter

ATGAGCAAAGGAGAAGAACTTTTCACTGGAGTTGTCCCAATTCTTGTTGAATTAGATGGTGATG
TTAATGGGCACAAATTTTCTGTCCGTGGAGAGGGTGAAGGTGATGCTACAAACGGAAACTCAC
CCTTAAATTTATTTGCACTACTGGAAAACCTGTTCCATGGCCAACACTTGTCACTACTCTG
ACCTATGGTGTTCATGCTTTTCCCGTTATCCGGATCACATGAAACGGCATGACTTTTTCAAGA
GTGCCATGCCGAAGGTTATGTACAGGAACGCACTATATCTTTCAAAGATGACGGGACCTACAA
GACGCGTGCTGAAGTCAAGTTTGAAGGTGATACCCTTGTTAATCGTATCGAGTTAAAAGGTATT
GATTTTAAAGAAGATGGAAACATTCTCGGACACAAACTgGAGTACAACCTTTAACTCACACAATG
TATACATCACGGCAGACAAACAAAAGAATGGAATCAAAGCTAACTTCAAATTCGCCACAACGT
TGAAGATGGTTCCGTTCAACTAGCAGACCATTATCAACAAAATACTCCAATTGGCGATGGCCCT
GTCCTTTTACCAGACAACCATTACCTGTCGACACAATCTGTCCTTTCGAAAGATCCCAACGAAA
AGAGGCCTGATCCTGAGAACTTCAGGgtgagtttggggacccttCGTGACCACATGGTCCTTCT
TGAGTTTGTAAGTCTGCTGGGATTACATGAttgttctttctttttcgctattgtaaaattcat
gttatatggagggggcaagttttcaggggtgttgtttagaatggggtcttcctcgaggaagacc
tggaagatgtcccttgatcaccatggaccctcatgataattttgtttctttcactttctactc
tgttgacaaccattgtctcctcttattttcttttcattttctgttaacttttcggttaaacttta
gcttgcatttgtaacgaatttttaaaattcacttttggtttatttgtcagattgtaagtactttct
ctaatacactttttttcaaggcaatcagggatatattatattgtacttcagcacagttttagaga
acaattgttataattaaatgataaggtagaatatttctgcatataaattctggctggcgtggaa
atattcttatttggtagaacaactacatcctggatcatcctgcctttctctttatggttaca
atgatatacactgtttgagatgaggataaaaatactctgagtccaaaccgggcccctctgctaac
catgttcatgccttctttctttttcctacagCTCCTGGGCAACGTGCTGGTTATTGTGCTGTCTC
ATCATTTTGGCAAAGAATAATAA

sfGFP(1-10)

MSKGEELFTGVVPILVELDGDVNGHKFSVRGEGEGDATNGKLTLKFICTTGKLPVPWPTLVTTL
TYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGTYKTRAEVKFEGDTLVNRIELKGI
DFKEDGNILGHKLEYNFSNVYITADKQKNGIKANFKIRHNVEDGSVQLADHYQNTPIGDGP
VLLPDNHVLTQSVLSKDPNEKRPDPENFRLLGNVLVIVLSHHFGKE**

sfGFP(1-10)-L-(11)

MSKGEELFTGVVPIVLVDGDVNGHKFSVRGEGGDATNGKLTLLKFICTTGKLPVPWPTLVTTL
TYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGTYKTRAEVKFEGDTLVNRIELKGI
DFKEDGNILGHKLEYNFNSHNVYITADKQKNGIKANFKIRHNVEDGSVQLADHYQNTPIGDGP
VLLPDNHYLSTQSVLSKDPNEKRPDPENFRVSLGTLRDHMLLEFVTAAGIT*

dPspCas13b fusion protein sequences

ps-CPSF30

3xFLAG-Ty1NLS-dPspCas13b-linker-CPSF30

ATGGACTACAAAGACCATGACGGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATG
ACAAGGGCGGAAACTCAAAGAAAAGGTCACTGGAGGACAACGAAACGGAAATCAAAGTCTCCAG
AGACACCTGGAACACTAAGAACATGCGGTCCCTGGAACCGCCACGAAGCAAGAAACGGATACAT
GGAGGCAACATCCCCGCTCTGGTGGAAAACAGAAAGTACTTTGGCACCTACAGCGTGATGG
CCATGCTGAACGCTCAGACCGTGCTGGACCACATCCAGAAGGTGGCCGATATTGAGGGCGAGCA
GAACGAGAACAACGAGAATCTGTGGTTTCACCCCGTGATGAGCCACCTGTACAACGCCAAGAAC
GGCTACGACAAGCAGCCGAGAAAACCATGTTTCATCATCGAGCGGCTGCAGAGCTACTTCCCAT
TCCTGAAGATCATGGCCGAGAACCAGAGAGAGTACAGCAACGGCAAGTACAAGCAGAACCGCGT
GGAAGTGAACAGCAACGACATCTTCGAGGTGCTGAAGCGCGCCTTCGGCGTGCTGAAGATGTAC
AGGGACCTGACCAACGCATACAAGACCTACGAGGAAAAGCTGAACGACGGCTGCGAGTTCCTGA
CCAGCACAGAGCAACCTCTGAGCGGCATGATCAACAATACTACACAGTGGCCCTGCGGAACAT
GAACGAGAGATACGGCTACAAGACAGAGGACCTGGCCTTCATCCAGGACAAGCGGTTCAAGTTC
GTGAAGGACGCTACGGCAAGAAAAGTCCCAAGTGAATACCGGATTCTTCCTGAGCCTGCAGG
ACTACAACGGCGACACACAGAAGAAGCTGCACCTGAGCGGAGTGGGAATCGCCCTGCTGATCTG
CCTGTTCTGAGACAAGCAGTACATCAACATCTTTCTGAGCAGGCTGCCCATCTTCTCCAGCTAC
AATGCCCAGAGCGAGGAACGGCGGATCATCATCAGATCCTTCGGCATCAACAGCATCAAGCTGC
CCAAGGACCGGATCCACAGCGAGAAGTCCAACAAGAGCGTGGCCATGGATATGCTCAACGAAGT
GAAGCGGTGCCCCGACGAGCTGTTCAACAACCTGTCTGCCGAGAAGCAGTCCCGGTTCAAGATC
ATCAGCGACGACCACAATGAAGTGCTGATGAAGCGGAGCAGCGACAGATTCGTGCCTCTGCTGC
TGCAGTATATCGATTACGGCAAGCTGTTTCGACCACATCAGGTTCCACGTGAACATGGGCAAGCT
GAGATACCTGCTGAAGGCCGACAAGACCTGCATCGACGGCCAGACCAGAGTCAGAGTGATCGAG
CAGCCCCCTGAACGGCTTCGGCAGACTGGAAGAGGCCGAGACAATGCGGAAGCAAGAGAACGGCA
CCTTCGGCAACAGCGGCATCCGGATCAGAGACTTCGAGAACATGAAGCGGGACGACGCCAATCC
TGCCAACTATCCCTACATCGTGGACACCTACACACACTACATCCTGGAAAACAACAAGGTCGAG
ATGTTTATCAACGACAAAGAGGACAGCGCCCCACTGCTGCCCCTGATCGAGGATGATAGATACG
TGGTCAAGACAATCCCCAGCTGCCGGATGAGCACCTTGAAATTCCAGCCATGGCCTTCCACAT
GTTTCTGTTTCGGCAGCAAGAAAACCGAGAAGCTGATCGTGGACGTGCACAACCGGTACAAGAGA
CTGTTCCAGGCCATGCAGAAAGAAGAAGTGACCGCCGAGAATATCGCCAGCTTCGGAATCGCCG
AGAGCGACCTGCCTCAGAAGATCCTGGATCTGATCAGCGGCAATGCCACGGCAAGGATGTGGA
CGCCTTCATCAGACTGACCGTGGACGACATGCTGACCGACACCGAGCGGAGAATCAAGAGATTC
AAGGACGACCGGAAGTCCATTTCGAGCGCCGACAACAAGATGGGAAAGAGAGGCTTCAAGCAGA
TCTCCACAGGCAAGCTGGCCGACTTCCTGGCCAAGGACATCGTGCTGTTTCAGCCAGCGTGAA
CGATGGCGAGAACAAGATCACCGGCCTGAACTACCGGATCATGCAGAGCGCCATTGCCGTGTAC
GATAGCGGCGACGATTACGAGGCCAAGCAGCAGTTCAAGCTGATGTTTCGAGAAGGCCCGGCTGA

TCGGCAAGGGCACAAACAGAGCCTCATCCATTTCTGTACAAGGTGTTGCCCCGAGCATCCCCGC
CAATGCCGTCGAGTTCTACGAGCGCTACCTGATCGAGCGGAAGTTCTACCTGACCGGCCTGTCC
AACGAGATCAAGAAAGGCAACAGAGTGGATGTGCCCTTCATCCGGCGGGACCAGAACAAGTGGA
AAACACCCGCCATGAAGACCCTGGGCAGAATCTACAGCGAGGATCTGCCCGTGGAAGTGGCCAG
ACAGATGTTTCGACAATGAGATCAAGTCCCACCTGAAGTCCCTGCCACAGATGGAAGGCATCGAC
TTCAACAATGCCAACGTGACCTATCTGATCGCCGAGTACATGAAGAGAGTGCTGGACGACGACT
TCCAGACCTTCTACCAGTGAACCGCAACTACCGGTACATGGACATGCTTAAGGGCGAGTACGA
CAGAAAGGGCTCCCTGCAGCACTGCTTCACCAGCGTGGAAGAGAGAGAAGGCCTCTGGAAAGAG
CGGGCCTCCAGAACAGAGCGGTACAGAAAGCAGGCCAGCAACAAGATCCGCAGCAACCGGCAGA
TGAGAAACGCCAGCAGCGAAGAGATCGAGACAATCCTGGATAAGCGGCTGAGCAACAGCCGGAA
CGAGTACCAGAAAAGCGAGAAAGTGATCCGGCGCTACAGAGTGCAGGATGCCCTGCTGTTTCTG
CTGGCCAAAAAGACCCTGACCGAACTGGCCGATTTGACGGCGAGAGGTTCAAAGTAAAGAAA
TCATGCCCCGACGCCGAGAAGGGAATCCTGAGCGAGATCATGCCATGAGCTTCACCTTCGAGAA
AGGCGGCAAGAAGTACACCATCACCAGCGAGGGCATGAAGCTGAAGAACTACGGCGACTTCTTT
GTGCTGGCTAGCGACAAGAGGATCGGCAACCTGCTGGAAGTCTGTTGGCAGCGACATCGTGTCCA
AAGAGGATATCATGGAAGAGTTCAACAAATACGACCAGTGCAGGCCCCGAGATCAGCTCCATCGT
GTTCAACCTGGAAGAGTGGGCCTTCGACACATACCCCGAGCTGTCTGCCAGAGTGGACCGGGAA
GAGAAGGTGGACTTCAAGAGCATCCTGAAAATCCTGCTGAACAACAAGAACATCAACAAAGAGC
AGAGCGACATCCTGCGGAAGATCCGGAACGCCCTTCGATGCAAACAATTACCCCGACAAAGGCGT
GGTGGAATCAAGGCCCTGCCTGAGATCGCCATGAGCATCAAGAAGGCCTTTGGGGAGTACGCC
ATCATGAAGTCGCGAGGCGGTGGAGGTAGTGGAGGGGGAGGATCAACGCGTCAGGAAATCATCG
CCAGCGTGGACCACATCAAGTTTGACTTGGAGATCGCGGTGGAGCAGCAGCTGGGGGCGCAGCC
GCTGCCCTTCCCCGGCATGGACAAGTCGGGCGCTGCTGTCTGTGAATTCTTTTGAAGCTGCC
TGCGGCAAAGGGGGCATGTGTCCGTTTTGCCACATCAGTGGTGAGAAAACAGTTGTGTGCAAAC
ACTGGCTGCGTGGCCTATGCAAGAAAGGGGACCAGTGTGAGTTCTGTCATGAGTATGACATGAC
CAAGATGCCCCGAGTGCTACTTCTACTCCAAGTTCGGGGAGTGCAGCAACAAGGAATGTCCCTTC
CTGCACATCGACCCCGAGTCCAAGATCAAGGACTGTCCTTGGTATGACCGTGGCTTCTGCAAGC
ACGGTCCCCTCTGCAGGCACCGGCACACAGGAGATCATCTGTGTGAATTACCTCGTGGGATT
CTGCCCCGAGGGGCCCTCGTGTAATTCATGCACCCTCGATTTGAACTGCCCATGGGAACCACC
GAGCAGCCCCCACTGCCGAGCAGACACAGCCTCCAGCAAAGCAGAGAACCCCGCAGGTCATCG
GGGTATGAGAGTCAAAACAGCAGCGCGGGCAACCGGGGACCCCGGCCACTGGAGCAGGTCAC
CTGTTACAAGTGTGGCGAGAAAGGACACTACGCCAACAGATGCACCAAAGGGCACTTGGCCTTT
CTCAGTGGACAGTAGTAA

3xFLAG-Ty1NLS-dPspCas13b-linker-CPSF30

MDYKDHDGDYKDHDIDYKDDDDKGGNSKKRSLEDNETEIKVSRDTWNTKNMRSLEPPRSKKRIH
GGNIPALVENQKKYFGTYSVMAMLNAQTVLDHIQKVADIEGEQNENNENLWFHPVMSHLYNAKN
GYDKQPEKTMFIIRLQSYFPFLKIMAENQREYSNGKYKQNRVEVNSNDIFEVLKRAFGVLKMY
RDLTNAYKTYEEKLNDGCEFLTSTEQPLSGMINNYTVALRNMNERYGYKTEDLAFIQDKRFKF
VKDAYGKKKSQVNTGFFLSLQDYNGDTQKKLHLSGVGIALLICFLDKQYINIFLSRLPIFFSY
NAQSEERRIIIRSFINSIKLPKDRIHSEKSNKSVAMDMLNEVKRCPDELFTTLSAEKQSRFRI
ISDDHNEVLMKRSSDRFVPLLLQYIDYGKLFDHIRFHVNMGLRYLLKADKTCIDGQTRVRVIE
QPLNGFGRLEEAEATMRKQENGTFGNSGIRIRDFENMKRDDANPANYPYIVDTYTHYLENNKVE
MFINDKEDSAPLLPVIEDDRYVVKTIPTSCRMSTLEIPAMAFHMFLLFGSKKTEKLIVDVHNRYKR
LFQAMQKEEVTAENIASFGIAESDLPQKILDLISGNAHGKDVDAFIRLTVDDMLTDTERRIKRF
KDDRKSIRSADNMKGKRGFKQISTGKLADFLAKDIVLFQPSVNDGENKITGLNYRIMQSAIAVY

DSGDDYEAKQQFKLMFEKARLIGKGTTEPHPLYKVFARSIPANAVEFYERYLIERKFYLTGLS
 NEIKKGNRVDVPFIRRDQNKWKTPAMKTLGRIYSEDLVELPRQMFNEIKSHLKS LPQMEGID
 FNNANVTYLIAEYMKRVLDDDFQTFYQWNRNYRYMDMLKGEYDRKGS LQHCFTSVEEREGLWKE
 RASRTERYRKQASNKIRSNRQMRNASSEIEIETILDKRLSNSRNEYQKSEKVI RRYRVQDALLFL
 LAKKTLTELADFDGERFKLKEIMPDAEKILSEIMPMSFTFEKGGK KYTITSEG MKLKNYGDF
 VLASDKRIGNLLELVGSDIVSKEDIMEEFNKYDQCRPEISSIVFNLEK WAFDTYPELSARVDRE
 EKVD FKSILKILLNNKNINKEQSDILRKIRNAFDANNYPDKGVVEIKALPEIAMS IKKAFGEYA
 IMKSRGGGGSGGGGSTRQEIIASVDHIKFDLEIAVEQQ LGAQPLPFP GMDKSGAAVCEFFLKAA
 CGKGGMCPFRHISGEKTVVCKHWLRGLCKKGDQCEFLHEYDMTKMPECYFY SKFGECSNKECPF
 LHIDPESKIKDCPWYDRGFCKHGPLCRHRHTRR VICVNYLVGFCPEGPSCKFMHPRFELPMGTT
 EQPPLPQQTQPPAKQRT PQVIGVMQSQNSSAGNRGPRPLEQVTCYKCGEKGHYANRCKGHLAF
 LSGQ**

ps-WDR33

3xFLAG-Ty1NLS-dPspCas13b-linker-WDR33

ATGGACTACAAAGACCATGACGGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATG
 ACAAGGGCGGAAACTCAAAGAAAAGGTCACTGGAGGACAACGAAACGGAAATCAAAGTCTCCAG
 AGACACCTGGAACACTAAGAACATGCGGTCCCTGGAACCGCCACGAAGCAAGAAACGGATACAT
 GGAGGCAACATCCCCGCTCTGGTGGAAGAACAGAAAGTACTTTGGCACCTACAGCGTGATGG
 CCATGCTGAACGCTCAGACCGTGCTGGACCACATCCAGAAGGTGGCCGATATTGAGGGCGAGCA
 GAACGAGAACACGAGAATCTGTGGTTTCACCCCGTGATGAGCCACCTGTACAACGCCAAGAAC
 GGCTACGACAAGCAGCCCGAGAAAACCATGTTTCATCATCGAGCGGCTGCAGAGCTACTTCCCAT
 TCCTGAAGATCATGGCCGAGAACCAGAGAGAGTACAGCAACGGCAAGTACAAGCAGAACC GCGT
 GGAAGTGAACAGCAACGACATCTTCGAGGTGCTGAAGCGCGCCTTCGGCGTGCTGAAGATGTAC
 AGGGACCTGACCAACGCATACAAGACCTACGAGGAAAAGCTGAACGACGGCTGCGAGTTCCTGA
 CCAGCACAGAGCAACCTCTGAGCGGCATGATCAACAATACTACACAGTGGCCCTGCGGAACAT
 GAACGAGAGATACGGCTACAAGACAGAGGACCTGGCCTTCATCCAGGACAAGCGGTTCAAGTTC
 GTGAAGGACGCTACGGCAAGAAAAAGTCCCAAGTGAATACCGGATTCTTCTGAGCCTGCAGG
 ACTACAACGGCGACACACAGAAGAAGCTGCACCTGAGCGGAGTGGGAATCGCCCTGCTGATCTG
 CCTGTTCTTGACAAGCAGTACATCAACATCTTTCTGAGCAGGCTGCCCCTCTTCTCCAGCTAC
 AATGCCCAGAGCGAGGAACGGCGGATCATCATCAGATCCTTCGGCATCAACAGCATCAAGCTGC
 CCAAGGACCGGATCCACAGCGAGAAGTCCAACAAGAGCGTGGCCATGGATATGCTCAACGAAGT
 GAAGCGGTGCCCCGACGAGCTGTTCAACAACCTGTCTGCCGAGAAGCAGTCCCGGTT CAGAATC
 ATCAGCGACGACCACAATGAAGTGCTGATGAAGCGGAGCAGCGACAGATTCGTGCCTCTGCTGC
 TGCAGTATATCGATTACGGCAAGCTGTTTCGACCACATCAGGTTCCACGTGAACATGGGCAAGCT
 GAGATACCTGCTGAAGGCCGACAAGACCTGCATCGACGGCCAGACCAGAGTCAGAGTGATCGAG
 CAGCCCCCTGAACGGCTTCGGCAGACTGGAAGAGGCCGAGACAATGCGGAAGCAAGAGAACGGCA
 CCTTCGGCAACAGCGGCATCCGGATCAGAGACTTCGAGAACATGAAGCGGGACGACGCCAATCC
 TGCCAACTATCCCTACATCGTGGACACCTACACACACTACATCCTGGAAAACAACAAGGTCGAG
 ATGTTTATCAACGACAAAGAGGACAGCGCCCCACTGCTGCCCCTGATCGAGGATGATAGATACG
 TGGTCAAGACAATCCCAGCTGCCGGATGAGCACCTGGAAATTCCAGCCATGGCCTTCCACAT
 GTTTCTGTTTCGGCAGCAAGAAAACCGAGAAGCTGATCGTGGACGTGCACAACCGGTACAAGAGA
 CTGTTCCAGGCCATGCAGAAAGAAGAAGTGACCGCCGAGAATATCGCCAGCTTCGGAATCGCCG
 AGAGCGACCTGCCTCAGAAGATCCTGGATCTGATCAGCGGCAATGCCACGGCAAGGATGTGGA

CGCCTTCATCAGACTGACCGTGGACGACATGCTGACCGACACCGAGCGGAGAATCAAGAGATTC
AAGGACGACCGGAAGTCCATTTCGGAGCGCCGACAACAAGATGGGAAAGAGAGGCTTCAAGCAGA
TCTCCACAGGCAAGCTGGCCGACTTCTGGCCAAGGACATCGTGCTGTTTCAGCCCAGCGTGAA
CGATGGCGAGAACAAGATCACCGGCCTGAACTACCGGATCATGCAGAGCGCCATTGCCGTGTAC
GATAGCGGCGACGATTACGAGGCCAAGCAGCAGTTCAAGCTGATGTTTCGAGAAGGCCCGGCTGA
TCGGCAAGGGCACAACAGAGCCTCATCCATTTCTGTACAAGGTGTTTCGCCCGCAGCATCCCCGC
CAATGCCGTGAGTTCTACGAGCGCTACCTGATCGAGCGGAAGTTCTACCTGACCGGCCTGTCC
AACGAGATCAAGAAAGGCAACAGAGTGGATGTGCCCTTCATCCGGCGGGACCAGAACAAGTGGA
AAACACCCGCCATGAAGACCCTGGGCAGAATCTACAGCGAGGATCTGCCCGTGGAAGTGGCCAG
ACAGATGTTTCGACAATGAGATCAAGTCCCACCTGAAGTCCCTGCCACAGATGGAAGGCATCGAC
TTCAACAATGCCAACGTGACCTATCTGATCGCCGAGTACATGAAGAGAGTGCTGGACGACGACT
TCCAGACCTTCTACCAGTGGAAACCGCAACTACCGGTACATGGACATGCTTAAGGGCGAGTACGA
CAGAAAGGGCTCCCTGCAGCACTGCTTCACCAGCGTGGAAGAGAGAGAAGGCCTCTGGAAAGAG
CGGGCCTCCAGAACAGAGCGGTACAGAAAGCAGGCCAGCAACAAGATCCGCAGCAACCGGCAGA
TGAGAAACGCCAGCAGCGAAGAGATCGAGACAATCCTGGATAAGCGGCTGAGCAACAGCCGGAA
CGAGTACCAGAAAAGCGAGAAAGTGATCCGGCGCTACAGAGTGCAGGATGCCCTGCTGTTTCTG
CTGGCCAAAAAGACCCTGACCGAACTGGCCGATTTTCGACGGCGAGAGGTTCAAAGTAAAGAAA
TCATGCCCCGACGCCGAGAAGGGAATCCTGAGCGAGATCATGCCCATGAGCTTCACCTTCGAGAA
AGGCGGCAAGAAGTACACCATCACCAGCGAGGGCATGAAGCTGAAGAACTACGGCGACTTCTTT
GTGCTGGCTAGCGACAAGAGGATCGGCAACCTGCTGGAAGTCTGTTGGCAGCGACATCGTGTCCA
AAGAGGATATCATGGAAGAGTTCAACAAATACGACCAGTGCAGGCCCGAGATCAGCTCCATCGT
GTTCAACCTGGAAAAGTGGGCCTTCGACACATACCCCGAGCTGTCTGCCAGAGTGGACCGGGAA
GAGAAGGTGGACTTCAAGAGCATCCTGAAAATCCTGCTGAACAACAAGAACATCAACAAAGAGC
AGAGCGACATCCTGCGGAAGATCCGGAACGCCTTCGATGCAAACAATTACCCCGACAAAGGCGT
GGTGGAAATCAAGGCCCTGCCTGAGATCGCCATGAGCATCAAGAAGGCCTTTGGGGAGTACGCC
ATCATGAAGT**CGCGAGGCGGTGGAGGTAGTGGAGGGGGAGGATCAACGCGTGGCTCACCCCTC**
GATTCTTCCACATGCCGCGCTTTCAACACCAAGCCCCAGGCAACTCTTTTACAAACGCCCAGA
CTTTGCGCAGCAGCAGGCGATGCAGCAGCTTACTTTTCGACGGAAAGCGCATGCGAAAAGCGGTC
AACCGCAAGACAATAGATTACAATCCCAGTGTCAATTAATACCTCGAGAATCGCATATGGCAAA
GAGATCAACGAGACATGAGAGCCATTCAACCGGATGCAGGTTATTACAATGATCTGGTCCCTCC
GATCGGCATGTTGAATAATCCAATGAACGCTGTCACGACCAAATTTGTCCGCACAAGTACTAAT
AAGTTAAATGCCCAGTATTTGTAGTGCAGTGGACACCCGAAGGAAGGAGGCTCGTCACAGGAG
CATCTTCCGGTGAATTTACTCTGTGGAATGGATTGACATTTAATTTTGAACAATACTTCAGGC
CCATGACTCTCCGGTGAAGGCCATGACCTGGTCCACAACGACATGTGGATGCTGACCGCAGAC
CATGGTGGGTATGTAAAATACTGGCAGTCAAACATGAATAATGTGAAGATGTTTCAGGCCATA
AAGAAGCTATACGGGAGGCTTCTTTTTCACCAACTGATAATAAGTTTCGCCACCTGTTTCAGACGA
TGGTACGGTCCGCATTTGGGATTTTCTCCGATGTATGAGGAGAGAATACTGAGGGGTACGGA
GCCGATGTCAAATGTGTGGAAGTGGCATCCAACCAAGGGTCTTGTCGTTTCAGGCTCCAAAGATT
CCCAACAGCCGATTAAGTTTTGGGACCCTAAAACCGGACAATCCCTGGCGACACTGCATGCCCA
TAAGAACACGGTCATGGAAGTCAAGCTCAATCTTAATGGGAATTGGCTGCTCACGGCTTCTCGG
GACCACCTTTGCAAGCTGTTTCGATATTAGAAATTTGAAAGAAGAACTGCAGGTATTTTCAGGGC
ATAAGAAGGAAGCCACTGCGGTAGCCTGGCACCCAGTCCATGAAGGTCTGTTTCGCGAGTGGGGG
GTCTGATGGTTCCCTGCTCTTTTGGCACGTCGGGGTAGAGAAGGAAGTTGGTGGGATGGAGATG
GCGCATGAGGGCATGATATGGAGCTTGGCCTGGCACCCCTCTCGACATATACTCTGTAGTGGCA
GCAATGATCACACATCTAAGTTTTGGACACGGAACAGACCAGGCGATAAAATGCGGGATCGATA
CAACCTTAATCTGCTTCCCGGGATGAGCGAGGACGGAGTAGAGTACGATGATCTCGAGCCAAAC

AGTCTCGCAGTGATACCCGGCATGGGCATCCCTGAGCAACTCAAATTGGCCATGGAGCAAGAGC
AGATGGGCAAAGATGAGTCTAACGAAATAGAAATGACCATTCTGGCCTTGATTGGGGGATGGA
GGAAGTAATGCAGAAAGACCAGAAGAAAGTGCCACAGAAAAAGGTCCCATACGCGAAACCGATT
CCCGCGCAATTCCAGCAGGCATGGATGCAAAACAAGGTGCCTATCCCGCTCCAAATGAAGTCC
TGAACGATCGAAAAGAGGATATTAAGTGGAGGAGAAAAAAGACGCAAGCAGAAATTGAGCA
GGAGATGGCCACGTTGCAGTACACGAACCCACAACCTGCTGGAGCAATTGAAAATCGAGCGACTT
GCGCAGAAACAGGTTGAGCAGATCTAGTAA

3xFLAG-Ty1NLS-dPspCas13b-linker-WDR33

MDYKDHGDKDHDIDYKDDDDKGGNSKKRSLDNETEIKVSRDTWNTKNMRSLEPPRSKKRIH
GGNIPALVENQKKYFGTYSVMAMLNAQTVLDHIQKVADIEGEQNENNENLWFHPVMSHLYNAKN
GYDKQPEKTMFIIRLQSYFPFLKIMAENQREYSNGKYKQNRVEVNSNDIFEVLKRAFGVLKMY
RDLTNAYKTYEEKLNDGCEFLTSTEQPLSGMINNYTVALRNMNERYGYKTEDLAFIQDKRFKF
VKDAYGKKKSQVNTGFFLSLQDYNGDTQKKLHLSGVGIALLICFLDKQYINIFLSRLPIFSSY
NAQSEERRIIIRSGINSIKLPKDRIHSEKSNKSVAMDMLNEVKRCPDELFTTLSAEKQSRFRI
ISDDHNEVLMKRSSDRFVPLLLQYIDYGKLFDHIRFHVNMGKLYLLKADKTCIDGQTRVRVIE
QPLNGFGRLEEAEATMRKQENGTFGNSGIRIRDFENMKRDDANPANYPIVDYTHYLENNKVE
MFINDKEDSAPLLPVIEDDRYVVKTIPTSCRMSTLEIPAMAFHMFLLFGSKKTEKLIVDVHNRKYR
LFQAMQKEEVTAENIASFGIAESDLQKILDLISGNAHGKDVDAFIRLTVDDMLTDTERRIKRF
KDDRKSIRSADNMKGKRGFKQISTGKLADFLAKDIVLFQPSVNDGENKITGLNYRIMQSAIAVY
DSGDDYEAKQQFKLMFEKARLIGKGTTEPHPLYKVFARSIPANAVEFYERYLIERKFYLTGLS
NEIKKGNRVDVPFIRRDQNKWKTTPAMKTLGRIYSEDLVELPRQMFNDNEIKSHLKSLLPQMEGID
FNNANVTYLIAEYMKRVLDDDFQTFYQWNRNYRYMDMLKGEYDRKGSLLQHCFTSVEEREGLWKE
RASRTERYRKQASNKIRSQRNASSEIEIETILDKRLSNSRNEYQKSEKVIIRRYRVQDALLFL
LAKKTLTELADFDGERFKLKEIMPDAEKILSEIMPMSFTFEKGGKKTITSEGMKLKNYGDF
VLASDKRIGNLLELVGSDIVSKEDIMEEFNKYDQCRPEISSIVFNLEKWAFTYPELSARVDRE
EKVDFKSILKILLNNKNINKEQSDILRKIRNAFDANNYPDKGVVEIKALPEIAMSIKKAFFEY
IMKSRGGGSGGGGSTRGSPPRFFHMPRFQHQAPRQLFYKRPDFAQQQAMQQLTFDGKMRKAV
NRKTIDYNPSVIKYLENRIWQRDQDMRAIQPDAGYYNDLVPPIGMLNPNMNAVTTKFVRTSTN
KVKCPVFVVRWTPEGRRLVTGASSGEFTLWNGLTFFNFETILQAHDSPVRAMTWSHNDMWMLTAD
HGGYVKYWQSNMNNVVMFQAHKEAIREASFSPDKNKFATCSDDGTVRIWDFLRCEERILRGHG
ADVCKVDWHPTKGLVVSQKDSQQPIKFWDPKTGQSLATLHAHKNTVMEVKLNLLNGNWLTLASR
DHLCKLFDIRNLKEELQVFRGHKKEATAVAWHPVHEGLFASGGSDGSLLFWHVGEVEVGGMEM
AHEGMIWSLAWHPLGHILCSGSNDHTSKFWTRNRPDGMKRDYRNLNLLPGMSDGEVYDDLEPN
SLAVIPGMGIPEQLKLAMEQEQMGKDESNEIEMTIPGLDWGMEEVMQKDQKKVPQKKVPYAKPI
PAQFQQAWMQNKVPIAPNEVLNDRKEDIKLEKKKTQAEIEQEMATLQYTNPQLLEQLKIERL
AQKQVEQI**

ps-NUDT21

3xFLAG-Ty1NLS-dPspCas13b-linker-NUDT21

ATGGACTACAAAGACCATGACGGTGATTATAAAGATCATGACATCGATTACAAGGATGACGATG
ACAAGGGCGGAAACTCAAAGAAAAGGTCACTGGAGGACAACGAAACGGAAATCAAAGTCTCCAG
AGACACCTGGAACACTAAGAACATGCGGTCCCTGGAACCGCCACGAAGCAAGAAACGGATACAT
GGAGGCAACATCCCCGCTCTGGTGGAAAACCAGAAGAAGTACTTTGGCACCTACAGCGTGATGG

CCATGCTGAACGCTCAGACCGTGCTGGACCACATCCAGAAGGTGGCCGATATTGAGGGCGAGCA
GAACGAGAACAACGAGAATCTGTGGTTTCACCCCGTGATGAGCCACCTGTACAACGCCAAGAAC
GGCTACGACAAGCAGCCCGAGAAAACCATGTTTCATCATCGAGCGGCTGCAGAGCTACTTCCCAT
TCCTGAAGATCATGGCCGAGAACAGAGAGAGTACAGCAACGGCAAGTACAAGCAGAACC GCGT
GGAAGTGAACAGCAACGACATCTTCGAGGTGCTGAAGCGCGCCTTCGGCGTGCTGAAGATGTAC
AGGGACCTGACCAACGCATACAAGACCTACGAGGAAAAGCTGAACGACGGCTGCGAGTTCCTGA
CCAGCACAGAGCAACCTCTGAGCGGCATGATCAACAACCTACTACACAGTGGCCCTGCGGAACAT
GAACGAGAGATACGGCTACAAGACAGAGGACCTGGCCTTCATCCAGGACAAGCGGTTCAAGTTC
GTGAAGGACGCCTACGGCAAGAAAAAGTCCCAAGTGAATACCGGATTCTTCCTGAGCCTGCAGG
ACTACAACGGCGACACACAGAAGAAGCTGCACCTGAGCGGAGTGGGAATCGCCCTGCTGATCTG
CCTGTTCTCTGGACAAGCAGTACATCAACATCTTTCTGAGCAGGCTGCCCATCTTCTCCAGCTAC
AATGCCCAGAGCGAGGAACGGCGGATCATCATCAGATCCTTCGGCATCAACAGCATCAAGCTGC
CCAAGGACCGGATCCACAGCGAGAAGTCCAACAAGAGCGTGGCCATGGATATGCTCAACGAAGT
GAAGCGGTGCCCCGACGAGCTGTTCAACAACCTGTCTGCCGAGAAGCAGTCCCGGTTTCAAGTTC
ATCAGCGACGACCACAATGAAGTGCTGATGAAGCGGAGCAGCGACAGATTCTGTCCTCTGCTGC
TGCAGTATATCGATTACGGCAAGCTGTTTCGACCACATCAGGTTCCACGTGAACATGGGCAAGCT
GAGATACCTGCTGAAGGCCGACAAGACCTGCATCGACGGCCAGACCAGAGTCAGAGTGATCGAG
CAGCCCCCTGAACGGCTTCGGCAGACTGGAAGAGGCCGAGACAATGCGGAAGCAAGAGAACGGCA
CCTTCGGCAACAGCGGCATCCGGATCAGAGACTTCGAGAACATGAAGCGGGACGACGCCAATCC
TGCCAACTATCCCTACATCGTGGACACCTACACACACTACATCCTGGAAAACAACAAGGTCGAG
ATGTTTATCAACGACAAAGAGGACAGCGCCCCACTGCTGCCCCTGATCGAGGATGATAGATACG
TGGTCAAGACAATCCCCAGCTGCCGGATGAGCACCTTGAAATTCCAGCCATGGCCTTCCACAT
GTTTCTGTTTCGGCAGCAAGAAAACCGAGAAGCTGATCGTGGACGTGCACAACCGGTACAAGAGA
CTGTTCCAGGCCATGCAGAAAGAAGAAGTGACCGCCGAGAATATCGCCAGCTTCGGAATCGCCG
AGAGCGACCTGCCTCAGAAGATCCTGGATCTGATCAGCGGCAATGCCACGGCAAGGATGTGGA
CGCCTTCATCAGACTGACCGTGGACGACATGCTGACCGACACCGAGCGGAGAATCAAGAGATTC
AAGGACGACCGGAAGTCCATTTCGAGCGCCGACAACAAGATGGGAAAGAGAGGCTTCAAGCAGA
TCTCCACAGGCAAGCTGGCCGACTTCTGGCCAAGGACATCGTGCTGTTTCAGCCCAGCGTGAA
CGATGGCGAGAACAAGATCACCGGCCTGAACTACCGGATCATGCAGAGCGCCATTGCCGTGTAC
GATAGCGGCGACGATTACGAGGCCAAGCAGCAGTTCAAGCTGATGTTTCGAGAAGGCCCGGCTGA
TCGGCAAGGGCACAACAGAGCCTCATCCATTTCTGTACAAGGTGTTTCGCCCGCAGCATCCCCGC
CAATGCCGTGAGTTCTACGAGCGCTACCTGATCGAGCGGAAGTTCTACCTGACCGGCCTGTCC
AACGAGATCAAGAAAGGCAACAGAGTGGATGTGCCCTTCATCCGGCGGGACCAGAACAAGTGGA
AAACACCCGCCATGAAGACCCTGGGCAGAATCTACAGCGAGGATCTGCCCGTGGAAGTGGCCAG
ACAGATGTTTCGACAATGAGATCAAGTCCCACCTGAAGTCCCTGCCACAGATGGAAGGCATCGAC
TTCAACAATGCCAACGTGACCTATCTGATCGCCGAGTACATGAAGAGAGTGCTGGACGACGACT
TCCAGACCTTCTACCAGTGGAACCGCAACTACCGGTACATGGACATGCTTAAGGGCGAGTACGA
CAGAAAGGGCTCCCTGCAGCACTGCTTCACCAGCGTGGAAGAGAGAGAAGGCCTCTGGAAAGAG
CGGGCCTCCAGAACAGAGCGGTACAGAAAGCAGGCCAGCAACAAGATCCGCAGCAACCGGCAGA
TGAGAAACGCCAGCAGCGAAGAGATCGAGACAATCCTGGATAAGCGGCTGAGCAACAGCCGGAA
CGAGTACCAGAAAAGCGAGAAAGTATCCGGCGCTACAGAGTGCAGGATGCCCTGCTGTTTCTG
CTGGCCAAAAAGACCCTGACCGAACTGGCCGATTTTCGACGGCGAGAGGTTCAAAGTAAAGAAA
TCATGCCCGACGCCGAGAAGGGAATCCTGAGCGAGATCATGCCCATGAGCTTACCTTCGAGAA
AGGCGGCAAGAAGTACACCATCACCAGCGAGGGCATGAAGCTGAAGAACTACGGCGACTTCTTT
GTGCTGGCTAGCGACAAGAGGATCGGCAACCTGCTGGAAGTCTGTTGGCAGCGACATCGTGTCCA
AAGAGGATATCATGGAAGAGTTCAACAATACGACCAGTGCAGGCCCCGAGATCAGCTCCATCGT

GTTCAACCTGGAAAAGTGGGCCTTCGACACATACCCCGAGCTGTCTGCCAGAGTGGACCGGGAA
GAGAAGGTGGACTTCAAGAGCATCCTGAAAATCCTGCTGAACAACAAGAACATCAACAAAGAGC
AGAGCGACATCCTGCGGAAGATCCGGAACGCCTTCGATGCAAACAATTACCCCGACAAAGGCGT
GGTGGAAATCAAGGCCCTGCCTGAGATCGCCATGAGCATCAAGAAGGCCCTTTGGGGAGTACGCC
ATCATGAAGTCGCGAGGCGGTGGAGGTAGTGGAGGGGGAGGATCAACGCGTCTCTGTGGTACCGC
CCAATCGCTCGCAGACCGGCTGGCCCCGGGGGGTCACTCAGTTCGGCAACAAGTACATCCAGCA
GACGAAGCCCCTCACCCTGGAGCGCACCATCAACCTGTACCCTCTTACCAATTATACTTTTGGT
ACAAAAGAGCCCCTCTACGAGAAGGACAGCTCTGTTGCAGCCAGATTTTCAGCGCATGAGGGGAAG
AATTTGATAAAATTGGAATGAGGAGGACTGTAGAAGGGGTTCTGATTGTACATGAGCACCGGCT
ACCCCATGTGTTACTGCTGCAGCTGGGAACAACCTTTCTTCAAACCTACCTGGTGGTGAACCTAAC
CCAGGAGAAGATGAAGTTGAAGGACTAAAACGCTTAATGACAGAGATACTGGGTCGTCAGGATG
GAGTTTTGCAAGACTGGGTCATTGACGATTGCATTGGTAACTGGTGGAGACCAAATTTTGAACC
TCCTCAGTATCCATATATTCTGCACATATTACAAAGCCTAAGGAACATAAGAAGTTGTTTCTG
GTTTCAGCTTCAAGAAAAAGCCTTGTTCAGTCCCTAAAAATTACAAGCTGGTAGCTGCACCAT
TGTTTGAATTGTATGACAATGCACCAGGATATGGACCCATCATTTCTAGTCTCCCTCAGCTGTT
GAGCAGGTTCAATTTTATTTACAAC TAGTAA

3xFLAG-Ty1NLS-dPspCas13b-linker-NUDT21

MDYKDHDGDYKDHDIDYKDDDDKGGNSKKRSLDNETEIKVSRDTWNTKNMRSLEPPRSKKRIH
GGNIPALVENQKKYFGTYSVMAMLNAQTVLDHIQKVADIEGEQNNENNENLWFHPVMSHLYNAKN
GYDKQPEKTMFIIERLQSYFPFLKIMAENQREYSNGKYQNRVEVNSNDIFEVLKRAFGVLKMY
RDLTNAYKTYEEKLNDGCEFLTSTEQPLSGMINNYTVALRNMNERYGYKTEDLAFIQDKRFKF
VKDAYGKKKSQVNTGFFLSLQDYNGDTQKKLHLSGVGIALLICLFLDKQYINIFLSRLPIFFSY
NAQSEERRIIIRSFINSIKLPKDRIHSEKSNKSVAMDMLEVKRCPDELFTTLSAEKQSRFRI
ISDDHNEVLMKRSSDRFVPLLLQYIDYGKLFDFHIFHVMGKLRLLKADKTCIDGQTRVRVIE
QPLNGFGRLEEAEATMRKQENGTFGNSGIRIRDFENMKRDDANPANYPIVDYTHYLENNKVE
MFINDKEDSAPLLPVIEDDRYVVKTIPTSCRMSTLEIPAMAFHMFLLFGSKKTEKLIVDVHNRYKR
LFQAMQKEEVTAENIASFGIAESDLPQKILDLISGNAHGKDVDAFIRLTVDMLTDTERRIKRF
KDDRKSIRSADNKMGRGFKQISTGKLADFLAKDIVLFQPSVNDGENKITGLNYRIMQSAIAVY
DSGDDYEAKQQFKLMFEKARLIGKGTTEPHPFYKVFARSIPANAVEFYERYLIERKFYLTGLS
NEIKKGNRVDVPFIRRDQNKWKTPAMKTLGRIYSEDLPVELPRQMFNDNEIKSHLKSLLPQMEGID
FNNANVTYLIAEYMKRVLDDDFQTFYQWNRNYRYMDMLKGEYDRKGSLLQHCFTSVEEREGLWKE
RASRTERYRKQASNKIRSQRMRNASSEIETILDKRLSNSRNEYQKSEKVIRRYRVQDALLFL
LAKKTLTELADFDGERFKLKEIMPDAEKGILSEIMPMSFTFEKGGKKTITSEGMLKNYGDFF
VLASDKRIGNLLELVGSDIVSKEDIMEEFNKYDQCRPEISSIVFNLEKWAFTYPELSARVDRE
EKVDFKSILKILLNNKNINKEQSDILRKIRNAFDANNYPDKGVVEIKALPEIAMSIKKAFFEYA
IMKSRGGGGSGGGGSTRSVPPNRSQTGWPRGVTQFGNKYIQQTKPLTLERTINLYPLTNYTTFG
TKEPLYEKDSSVAARFQRMREEFDKIGMRRTVEGVLI VHEHRLPHVLLLQLGTTFFKLPGGELN
PGEDEVEGLKRLMTEILGRQDGVLDWVIDDCIGNWWRPNFEPQYPYIPAHITKPKHEHKKLFL
VQLQEKALFAVPKNYKLVAAPL FELYDNAPGYGPIISSLPQLLSRFNFIYN**