

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Letter

Experimentally reduced insulin/IGF-1 signalling in adulthood extends lifespan of parents and improves Darwinian fitness of their offspring

Running title: Lifespan extension improves offspring fitness

**Martin I. Lind^{1,†,*}, Sanjana Ravindran^{1,†}, Zuzana Sekajova¹, Hanne Carlsson^{1,2},
Andrea Hinas³ and Alexei A. Maklakov^{1,2,*}**

¹ Animal Ecology, Department of Ecology and Genetics, Uppsala University, Norbyvägen 18D, Uppsala 752 36, Sweden

² School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

³ Department of Cell and Molecular Biology, Uppsala University, PO Box 596, Uppsala 751 24, Sweden

† these authors contributed equally to the study

* Authors for correspondence:

Martin I. Lind - martin.i.lind@gmail.com

Alexei A. Maklakov - a.maklakov@uea.ac.uk

20 **Total word count: 5714**

21 Keywords: ageing, antagonistic pleiotropy, functional trade-offs, hyperfunction, IIS

22 signalling, parental effects, senescence

23 **Abstract**

24 Classical theory maintains that ageing evolves via energy trade-offs between reproduction
25 and survival leading to accumulation of unrepaired cellular damage with age. In contrast, the
26 emerging new theory postulates that ageing evolves because of deleterious late-life hyper-
27 function of reproduction-promoting genes leading to excessive biosynthesis in late-life. The
28 hyper-function theory uniquely predicts that optimizing nutrient-sensing molecular signalling
29 in adulthood can simultaneously postpone ageing and increase Darwinian fitness. Here we
30 show that reducing evolutionarily conserved insulin/IGF-1 nutrient-sensing signalling via
31 *daf-2* RNA interference (RNAi) fulfils this prediction in *Caenorhabditis elegans* nematodes.
32 Long-lived *daf-2* RNAi parents showed normal fecundity as self-fertilizing hermaphrodites
33 and improved late-life reproduction when mated to males. Remarkably, the offspring of *daf-2*
34 RNAi parents had higher Darwinian fitness across three different genotypes. Thus, reduced
35 nutrient-sensing signalling in adulthood improves both parental longevity and offspring
36 fitness supporting the emerging view that sub-optimal gene expression in late-life lies at the
37 heart of ageing.

38

39

40 **Impact Statement**

41 Understanding mechanisms underpinning ageing is fundamental to improving quality of life
42 in an increasingly long-lived society. Recent breakthroughs have challenged the long-
43 standing paradigm that the energy trade-off between reproduction and somatic maintenance
44 causes organismal senescence via slow accumulation of unrepaired cellular damage with age.
45 The emerging new theory of ageing provides a conceptually novel framework by proposing
46 that ageing is a direct consequence of physiological processes optimized for early-life
47 function, such as growth and early-life reproduction, that are running ‘too high’ (i.e. at
48 hyperfunction) in late adulthood. Contrary to the classic view based on damage accumulation,
49 the hyperfunction theory proposes that suboptimal gene expression in late-life causes ageing
50 via excessive biosynthesis. Thus, the hyperfunction theory uniquely predicts that longevity
51 and Darwinian fitness can be simultaneously increased by reducing unnecessarily high levels
52 of nutrient-sensing signalling in adulthood. Here we show that reducing evolutionarily
53 conserved nutrient-sensing signalling pathway fulfils this prediction in *Caenorhabditis*
54 *elegans* nematodes. We found that downregulation of the insulin/IGF-1 signalling in adult *C.*
55 *elegans* nematodes not only improves longevity but, most intriguingly, increases fitness of
56 the resulting offspring in the next generation. We found support for increase in offspring
57 fitness across different genetic backgrounds. Our findings contradict the theoretical
58 conjecture that energy trade-offs between growth, reproduction and longevity is the universal
59 cause of senescence and provide strong experimental support for the emerging hyperfunction
60 theory of ageing.

61

62 **Introduction**

63 Understanding mechanisms underpinning healthy ageing is fundamental to improving quality
64 of life in an increasingly long-lived society. The long-standing paradigm postulates that
65 energy trade-offs between reproduction and somatic maintenance underlie organismal ageing
66 (Kirkwood 1977, Kirkwood and Austad 2000, Kirkwood 2017). This theory is supported by a
67 large number of studies in different taxa that reported a negative correlation between
68 reproduction and survival (reviewed in Hughes and Reynolds 2005, Partridge et al. 2005,
69 Edward and Chapman 2011, Flatt 2011, Nussey et al. 2013). However, the discoveries of
70 environmental interventions that dramatically increase healthy lifespan in model organisms
71 without the cost of reduced reproduction have challenged the current paradigm and suggested
72 that our understanding of the evolution of ageing is incomplete (Dillin et al. 2002, Kenyon
73 2005, Antebi 2013, Gems and Partridge 2013, Maklakov and Immler 2016, Flatt and
74 Partridge 2018). Specifically, experimental downregulation of nutrient-sensing insulin/IGF-
75 like (IIS) signalling pathway that governs biosynthesis in response to nutrient availability can
76 achieve increased longevity without a concomitant decrease in reproduction in model
77 organisms (Dillin et al. 2002, Kenyon 2010, Kenyon 2011).

78

79 Since cost-free lifespan extension contradicts the traditional view of how ageing evolves,
80 several studies investigated the fitness consequences of reduced IIS signalling (Gems et al.
81 1998, Walker et al. 2000, Jenkins et al. 2004, Savory et al. 2014, Maklakov et al. 2017).
82 Indeed, mutations that reduce nutrient-sensing signalling during the whole life, as well as
83 environmental interventions aimed at mimicking the mutational effect, often have detrimental
84 pleiotropic effects on key life-history traits, such as development, growth rate, body size and
85 early-life reproduction resulting in reduced Darwinian fitness even if total reproduction is

86 unaffected (Gems et al. 1998, Briga and Verhulst 2015). The first longevity mutant
87 discovered in *C. elegans*, *age-1*, is a good example because increased longevity, stress
88 resistance and late-life reproduction come at a cost of reduced early-life reproduction and
89 total individual fitness (Maklakov et al. 2017). Moreover, a recent literature survey suggests
90 that all classic longevity-extending mutations across taxa from worms to flies to mice
91 detrimentally affect life-history traits resulting in reduced fitness (Briga and Verhulst 2015).
92 Similarly, experimental evolution studies showed that when longevity and fecundity are
93 increased simultaneously through selection, the organisms pay the price in slow development
94 and delayed sexual maturation, again resulting in reduced individual fitness (Lind et al.
95 2017). These results support the theoretical conjecture that genes with antagonistically
96 pleiotropic effects between early-life and late-life fitness play an important role in the
97 evolution of ageing (Williams 1957). However, the mechanisms of antagonistic pleiotropy
98 (AP) remain elusive. The leading hypothesis, the “disposable soma” theory of ageing (DS)
99 suggests that ageing results from competitive energy allocation between somatic maintenance
100 and reproduction (Kirkwood 1977, Kirkwood and Holliday 1979, Kirkwood and Austad
101 2000). Indeed, increased reproductive performance in early-life correlates with reduced
102 survival and/or reduced performance in late-life in natural populations (Gustafsson and Part
103 1990, Boonekamp et al. 2014, Lemaitre et al. 2015) and in laboratory experiments (Rose
104 1984, Charlesworth 1993, Partridge et al. 1999, but see Chen and Maklakov 2012, Kimber
105 and Chippindale 2013, Chen and Maklakov 2014, Curtsinger 2019).

106

107 However, this hypothesis suffered several setbacks in recent years, with many empirical
108 studies challenging the importance of energy trade-offs in organismal senescence (reviewed
109 in Flatt 2011, Kenyon 2011, Antebi 2013, Gems and Partridge 2013, Maklakov and Immler
110 2016, Flatt and Partridge 2018). Instead, several authors proposed that ageing can result from

111 molecular signalling networks being optimized for development, growth and early-life
112 reproduction rather than for late-life reproduction and longevity (Blagosklonny 2010, Kenyon
113 2010, Antebi 2013, Gems and Partridge 2013, Ezcurra et al. 2018). For example, the
114 hyperfunction theory maintains that ageing is driven by excessive nutrient-sensing molecular
115 signalling in adulthood, which results in cellular hypertrophy leading to age-related
116 pathologies (Blagosklonny 2006, 2010, Ezcurra et al. 2018). These ideas can be traced back
117 to the original AP theory by George Williams, who suggested that the same physiological
118 processes that are beneficial for fitness early in life can become detrimental for organismal
119 fitness with age because of the reduced strength of natural selection on late-life function
120 (Williams 1957).

121

122 Williams's AP theory (Williams 1957) provides the population genetic framework for the
123 evolution of ageing via two different physiological routes: energy trade-offs (the “disposable
124 soma” theory of ageing) or functional trade-offs (e.g. “hyperfunction” theory of ageing) (Fig.
125 1). While both of these physiological theories rely on the same underlying principle, they
126 make uniquely distinct predictions with respect to reproduction costs of longevity. The
127 “disposable soma” theory maintains that organismal senescence is caused by slow
128 accumulation of unrepaired cellular damage with age because insufficient energy resources
129 are allocated to repair as organisms are maximising fitness rather than longevity (Kirkwood
130 1977, Kirkwood 2017). Therefore, the “disposable soma” theory predicts that increased
131 allocation of resources to somatic maintenance will increase longevity at the cost of reduced
132 resources available to current growth and reproduction. On the contrary, the functional trade-
133 off theory suggests that longevity is compromised by suboptimal physiology in adulthood
134 because, as discussed first by Williams (1957), selection is not strong enough to fully
135 optimize the age-specific expression of an allele whose effects are strongly beneficial in

136 early-life (e.g. during development) and slightly detrimental in late-life (e.g. during
137 adulthood). Consequently, the functional trade-off theory predicts that experimental
138 optimization of physiology in adulthood can increase longevity without any cost to
139 reproduction, or even simultaneously increase longevity and reproduction.

140

141 Because the main cost of longevity appears to be associated with reduced early-life function,
142 it seems plausible that age-specific modification of gene expression can potentially
143 circumvent this problem. In their landmark study, Dillin et al. (2002) used age-specific RNA
144 interference (RNAi) approach to knock down *daf-2* gene expression in *C. elegans* nematodes
145 across the life cycle of the worms. While early-life feeding with bacteria expressing *daf-2*
146 double-stranded (ds) RNA resulted in reduced early-life reproduction, there was no
147 detrimental effect of *daf-2* RNAi in adult worms, which enjoyed two-fold lifespan extension
148 without any cost to reproduction (Dillin et al. 2002). This study provides the strongest
149 support to date for the hypothesis that ageing results from molecular nutrient-sensing
150 signalling that is optimized for early-life function but is suboptimal for late-life function.

151

152 Nevertheless, while this study provided a powerful example for the cost-free lifespan
153 extension, it is possible that certain fitness costs have been overlooked. One possibility is that
154 fecundity costs become apparent only in mated hermaphrodites. In nature, *C. elegans* live in
155 populations with small (~0.3%) yet appreciable number of males living among self-fertilising
156 hermaphrodites with sometimes high levels of outcrossing (Sivasundar and Hey 2005), and
157 mating, as well as mere presence of male-derived pheromones, has pronounced effects of the
158 life-history of hermaphrodites (Maures et al. 2013, Shi and Murphy 2014, Aprison and
159 Ruvinsky 2016). While it is certainly interesting to consider how male-derived effects affect

160 resource allocation in hermaphrodites, it is unlikely that putative trade-off that is only visible
161 to selection in such rare circumstances can shape the evolution of ageing in this species.
162 Perhaps more importantly, it is possible that the fitness of the offspring and, therefore,
163 Darwinian fitness of the parents are compromised. The trade-off between offspring number
164 and quality is well known from a number of study systems (Stearns 1992), and is a potential
165 explanation for the apparent lack of fitness costs in the previous studies (Maklakov and
166 Immler 2016). Alternatively, longevity and Darwinian fitness can be simultaneously
167 increased by reducing unnecessarily high levels of nutrient-sensing signalling in adulthood.
168 To distinguish between these possibilities, we need to understand how reduction in nutrient-
169 sensing signalling in adulthood affects longevity, offspring number and offspring quality.
170 Here we show that *daf-2* RNAi in adult *C. elegans* results in increased offspring fitness
171 across three genetic backgrounds. We discuss these findings in the light of the emerging new
172 theories of ageing and suggest that they support the hypothesis that functional trade-offs
173 between early-life fitness and late-life fitness shape the evolution of ageing.

174

175 **Materials and Methods**

176

177 ***Strains***

178 We used the *Caenorhabditis elegans* strains Bristol N2 wild-type (Brenner, Genetics 1974),
179 as well as the mutants *ppw-1(pk2505)* and *rrf-1(pk1417)*, obtained from Caenorhabditis
180 Genetics Center (CGC, Missouri, USA).

181

182 ***Maintenance***

183 Before each assay, worms were recovered from freezing and synchronised by bleaching for

184 two generations to remove any freezing effects. The nematode populations were maintained
185 at 20°C and 60% relative humidity in an environmental test chamber. For regular
186 maintenance, the worms were kept on NGM agar supplemented with the antibiotics
187 streptomycin, kanamycin and nystatin (following Lionaki and Tavernarakis 2013), seeded
188 with the antibiotic-resistant *E. coli* strain OP50-1 (pUC4K).

189

190 ***Outline of the study***

191 The study was run in three separate experiments. In the first experiment, we investigated
192 lifespan and reproduction of mated and unmated N2 hermaphrodites reared from sexual
193 maturity onwards on *daf-2* RNAi or empty vector (EV, control) plates. For logistic reasons,
194 this experiment was conducted in two blocks for mated worms and one block for unmated
195 worms. In the second experiment, we investigated the lifespan and egg size of unmated N2,
196 *rrf-1(pk1417)* and *ppw-1(pk2505)* hermaphrodites on raised from sexual maturity onwards on
197 *daf-2* RNAi or EV plates. In the third experiment, we collected one egg from each parent at
198 their second day of adulthood (from *daf-2* RNAi and EV treatments) and investigated the
199 lifespan and reproduction of these offspring on control plates. Because different experiments
200 differed in setup time, daily reproduction values (and calculations based upon these, such as
201 λ_{ind}) are only meaningful for comparison between treatments within each experiment.

202

203 ***RNAi***

204 RNase-III deficient, IPTG-inducible HT115 *Escherichia coli* bacteria with empty plasmid
205 vector (L4440) was used as control (Timmons et al. 2001) and the same HT115 bacteria with
206 *daf-2* RNAi construct from the Vidal library was used as RNAi treatment. RNAi treatment
207 started from sexual maturity, and continued until the death of the individual. During the
208 experiments, worms were maintained on 35 mm NGM agar plates (supplemented with 1 mM

209 IPTG and 50 µg/ml ampicillin) seeded with 0.1 ml L4440 empty vector control or *daf-2*
210 bacteria grown in LB supplemented with 50 µg/ml ampicillin for 16-20 hours and seeded
211 (incubated) on the NGM agar plates again for 24 hours (following. Hinas et al. 2012).

212

213 ***Lifespan Assays***

214 Lifespan assays were set up for all treatment combinations described above. In the lifespan
215 assays, the individual age-synchronised L4 worms were placed on separate 35 mm plates and
216 the plates were checked daily to record any instances of death. The surviving worms were
217 moved to new plates daily until their death. Fertile worms, which showed odd developmental
218 characteristics and low offspring numbers (<36 offspring), were excluded from the final
219 analysis (3 mated control worms and 7 mated *daf-2* worms).

220

221 ***Reproduction assays***

222 Offspring production was scored in the reproduction assays using the same worms as those
223 scored for lifespan, except for the parental N2, *ppw-1* and *rrf-1* worms in the second
224 experiment, where only lifespan was recorded. Unmated individual hermaphrodites were
225 moved to new plates daily and scored for offspring produced 2.5 days later. In the “mated”
226 treatment, two male *C. elegans* (from the initial sample population of N2 strain) were placed
227 on a plate with a single hermaphrodite for two hours every day to allow time for mating.
228 Offspring production was scored 2.5 days later, as in the “unmated” treatment.

229

230 ***Egg size assays***

231 Egg size was measured in N2, *ppw-1* and *rrf-1* strains (unmated hermaphrodites) growing on
232 either *daf-2* RNAi or empty vector (EV) plates. Two days after maturation, worms were
233 placed individually on new plates and observed continually during five hours for the presence
234 of newly laid eggs, of which the first two eggs were collected. Eggs were picked immediately
235 after laying and placed under a Leica M165C microscope set on 12x magnification; photos
236 were taken using a Lumenera Infinity 2-6C digital microscope camera. Egg size was analysed
237 from photos using *ImageJ* (<https://imagej.nih.gov/ij/>). Only eggs laid during gastrulation
238 stage (the normal developmental stage at egg laying) were included in the analyses.

239

240 ***Statistical analyses***

241 Survival was analysed for each experiment in Cox proportional hazard models in *R* 3.3.3.
242 Mated (EV: n=72, *daf-2* n=68) and unmated (n=25 per treatment) individuals were analysed
243 separately, as they were run in different blocks. Unmated individuals were analysed using the
244 *coxph* function in the package *survival*, with *daf-2* RNAi treatment as a fixed factor. For
245 mated individuals, we used the *coxme* package in order to fit block as a random effect, in
246 addition to the fixed effect of RNAi treatment. In the second experiment (n=25 per
247 treatment), in addition to RNAi treatment, we also fitted the fixed factor strain (N2, *ppw-1*,
248 *rrf-1*) and its interaction with treatment using the *coxph* function in the *survival* package.

249 Reproduction was analysed as total reproduction as well as rate-sensitive individual fitness
250 λ_{ind} , which encompasses the timing and number of offspring (Brommer et al. 2002, Lind et al.
251 2016). λ_{ind} is estimated by solving the Euler-Lotka equation for each individual using the
252 *lambda* function in the *popbio* package and is analogous to the intrinsic rate of population
253 growth (Stearns 1992). For all unmated worms (n=25 per treatment), we estimated the fixed
254 effect of treatment (*daf-2* RNAi or empty vector). For offspring of the three mutants (n=25

255 per treatment), we also estimated the fixed effect of strain, using linear models. For the mated
256 worms (EV: n=72, *daf-2* n=68), we also estimated the random effect of block, in addition to
257 RNAi treatment. These models were implemented as mixed effect models using the *lme4*
258 package in R 3.3.3, and chi-square tests of fixed effects were performed using the *car*
259 package. Egg size was analysed in a mixed effect model in *lme4*, treating strain and RNAi
260 treatment as crossed fixed effects, and parent ID as well as block as random effects. We
261 obtained the following n: N2 on EV: 56, N2 on *daf-2*: 54, *ppw-1* on EV: 44, *ppw-1* on *daf-2*:
262 42, *rrf-1* on EV: 59, *rrf-1* on *daf-2*: 42.

263

264 **Results**

265 First off, we confirmed that *daf-2* RNAi significantly extended the lifespan of unmated N2
266 wild-type hermaphrodite worms (censoring matricide: $z = -4.94$, $df = 1$, $p < 0.001$, Fig. 2A;
267 including matricide as dead: $z = -4.97$, $df = 1$, $p < 0.001$), as expected from previous studies
268 (Dillin et al. 2002). In addition, for mated N2, *daf-2* RNAi extended lifespan when matricide
269 was censored ($z = -2.42$, $df = 1$, $p = 0.016$, Fig. 2B) but not if matricidal worms were
270 included as dead ($z = 0.16$, $df = 1$, $p = 0.87$) because of an increase in matricide in the late
271 reproducing mated *daf-2* RNAi N2.

272

273 We did not find any effect of *daf-2* RNAi on total reproduction (unmated: $F = 0.32$, $df = 1$, p
274 $= 0.58$; mated: $\chi^2 = 1.11$, $df = 1$, $p = 0.29$) or individual fitness λ_{ind} (unmated: $F = 0.30$, $df = 1$,
275 $p = 0.59$; mated: $\chi^2 = 0.43$, $df = 1$, $p = 0.51$) for neither unmated nor mated N2 (Table 1, Fig.
276 3). However, *daf-2* RNAi had a positive effect on late (day 5+) reproduction for mated
277 hermaphrodites ($\chi^2 = 24.76$, $df = 1$, $p < 0.001$, Fig. 3B).

278

279 In a second experiment, using unmated hermaphrodites only, we investigated the effect of
280 *daf-2* RNAi on parent lifespan and offspring lifespan and reproduction across three genetic
281 backgrounds (N2 wild-type and the mutants *ppw-1* and *rrf-1*, that are deficient for germline
282 and somatic RNAi, respectively). Parental treatment with *daf-2* RNAi increased lifespan
283 across all genetic backgrounds, both when matricide was censored (treatment: $\chi^2 = 90.39$, df
284 = 1, $p < 0.001$; strain: $\chi^2 = 21.8$, df = 2, $p < 0.001$; treatment \times strain: $\chi^2 = 10.46$, df = 2, $p =$
285 0.005, Fig. 2A) and included as dead (treatment: $\chi^2 = 85.25$, df = 1, $p < 0.001$; strain: $\chi^2 =$
286 20.45, df = 2, $p < 0.001$; treatment \times strain: $\chi^2 = 9.43$, df = 2, $p = 0.009$). In accordance with
287 previously published research (Hibshman et al. 2016), parental *daf-2* RNAi increased egg
288 size (treatment: $\chi^2 = 5.11$, df = 1, $p = 0.024$; strain: $\chi^2 = 13.89$, df = 2, $p < 0.001$; treatment \times
289 strain: $\chi^2 = 2.68$, df = 2, $p = 0.262$, Fig. 4). However, we found that the effect was most
290 pronounced in N2 wildtype worms, and relatively weak in both somatic and germline *daf-2*
291 knockdown (see Fig. 4), suggesting that *daf-2* knockdown in both somatic and reproductive
292 tissues is required to maximize the effect on egg size.

293

294 Parental *daf-2* RNAi treatment did not, however, influence the lifespan of their offspring,
295 neither when matricidal worms were censored (treatment: $\chi^2 = 0.04$, df = 1, $p = 0.85$; strain:
296 $\chi^2 = 24.2$, df = 2, $p < 0.001$; treatment \times strain: $\chi^2 = 0.61$, df = 2, $p = 0.74$, Fig. 5A) nor when
297 included as dead (treatment: $\chi^2 = 0.01$, df = 1, $p = 0.92$; strain: $\chi^2 = 21.8$, df = 2, $p < 0.001$;
298 treatment \times strain: $\chi^2 = 0.48$, df = 2, $p = 0.79$).

299

300 In contrast, parental *daf-2* RNAi treatment significantly increased offspring total reproduction
301 (treatment: $F = 15.9$, df = 1, $p < 0.001$; strain: $F = 33.7$, df = 2, $p < 0.001$; treatment \times strain: F
302 = 0.09, df = 2, $p = 0.91$, Fig. 5B-C) and individual fitness λ_{ind} (treatment: $F = 11.8$, df = 1, p

303 <0.001; strain: $F = 13.1$, $df = 2$, $p < 0.001$; treatment \times strain: $F = 0.18$, $df = 2$, $p = 0.84$, Fig.
304 5D) across all genetic backgrounds. Importantly, there was no correspondence between the
305 effect of parental *daf-2* RNAi on egg size (see above) and offspring total reproduction /
306 individual fitness, suggesting that factors beyond the amount of resources in the egg
307 contribute to increased fitness of offspring of *daf-2* RNAi parents.

308

309 **Discussion**

310 The “disposable soma” theory of ageing postulates that senescence results from competitive
311 energy allocation between key life-history traits, such as growth, reproduction and somatic
312 maintenance (Kirkwood 1977, Kirkwood 2017). This theory predicts that genetic and
313 environmental manipulations that increase energy allocation to growth and somatic
314 maintenance will result in detrimental effects to reproduction. This is why the findings by
315 Dillin et al. (2002), which suggested that adult-only downregulation of insulin/IGF-1 by *daf-2*
316 RNAi can substantially increase lifespan without any detrimental effect to reproduction, were
317 subsequently scrutinized in an attempt to find the hidden costs of longevity (Jenkins et al.
318 2004, Partridge et al. 2005). Nonetheless, both the original findings (Dillin et al. 2002) and
319 our results here, suggest that adult-only *daf-2* RNAi can more than double longevity without
320 any negative effect on reproduction. Moreover, when supplied with sperm from males, *daf-2*
321 RNAi-treated parents have improved fecundity in late-life. However, the key question that we
322 asked in this study was whether treatments that improve parental performance have positive
323 or negative effects on their offspring. The trade-off between offspring number and offspring
324 quality is a well known concept in life-history evolution (Stearns 1992) but is rarely
325 considered in biogerontological research (reviewed in Maklakov and Immler 2016). Germline
326 maintenance is costly (Sniegowski et al. 2000, Agrawal and Wang 2008, Maklakov and

327 Immler 2016, Berger et al. 2017), and increased investment into somatic maintenance can, in
328 theory, result in increased mutation rate and reduced fitness of progeny.

329

330 Alternatively, it is possible that instead of energy trade-offs, the evolution of senescence is
331 governed by functional trade-offs. Functional trade-offs can occur because the physiological
332 requirements of a young organism can differ substantially from those of a mature one
333 (Williams 1957). In his classic 1957 paper, George Williams (Williams 1957) described a
334 hypothetical example of a mutation that positively affects bone calcification in a developing
335 young organism but increases calcification of the connective tissues of arteries in a mature
336 one with detrimental consequences. More recently, it has been suggested that nutrient sensing
337 IIS/TOR molecular signalling pathways that govern growth and development result in
338 excessive biosynthesis in late-life leading to different pathologies and increased mortality
339 (Blagosklonny 2006, 2010, Gems and Partridge 2013, Ezcurra et al. 2018). These proximate
340 explanations rest on the fundamental assumption that the strength of natural selection
341 declines with age because of environmental mortality from a range of biotic and abiotic
342 hazards (e.g. predation, pathogens, competition, starvation) (Williams 1957). Because of such
343 environmental mortality, optimizing development, growth and instantaneous reproduction
344 may be more important for organismal fitness than optimizing future survival and
345 reproduction (Williams 1957, Hamilton 1966). Thus, progressively weakening selection in
346 adulthood may result in suboptimal levels of IIS/TOR signalling leading to pathology and
347 senescence (Ezcurra et al. 2018). However, unlike the classic energy trade-off theory, the
348 functional trade-off hypothesis predicts that it should be possible to modify adult physiology
349 to improve both longevity and fitness.

350

351 Here we found that reduced insulin/IGF-1 signalling in adult worms not only improved
352 longevity, but also increased reproduction and Darwinian fitness of the resulting offspring in
353 three different genetic backgrounds. This result contradicts the hypothesis that improved
354 longevity and postponed ageing of *daf-2* RNAi parents comes at the cost of offspring fitness.
355 Instead, our findings are in line with the hypothesis that suboptimal levels of nutrient-sensing
356 signalling in adult life accelerate ageing, curtail lifespan and reduce individual fitness. This
357 result was not caused by direct inheritance of *daf-2* RNAi, since we did not recover the
358 lifespan extension effect of *daf-2* knockdown in these offspring. Because *daf-2* is essential for
359 successful development and growth of a young worm (Dillin et al. 2002), these results
360 suggest that wildtype *C. elegans* nematodes trade-off improved pre-adult performance for
361 reduced offspring quality. Such trade-offs are at heart of AP theory, but are usually
362 interpreted as evidence for energy-based trade-offs. Our results clearly demonstrate that this
363 is not the case, and that adulthood-only *daf-2* RNAi increases offspring fitness. In summary,
364 our findings suggest that selection on expression of *daf-2* in adulthood is not sufficiently
365 strong in nature. We predict that such effects may be very common, and suggest that future
366 studies should aim to quantify the fitness consequences of experimental manipulation of age-
367 specific gene expression across a broad range of taxa. Such approach will allow us to
368 estimate the relative importance of energy trade-offs versus functional trade-offs in the
369 evolution of ageing across the tree of life.

370

371 Because previous research found that both dietary restriction and reduction in insulin-like
372 signalling by *daf-2* RNAi knockdown increased embryo size in *C. elegans* nematodes
373 (Hibshman et al. 2016), we replicated these results to test whether increased fitness of adult
374 progeny results from increased resource allocation to eggs by *daf-2* RNAi mothers. While
375 *daf-2* knockdown increased egg size to a different degree in N2, *ppw-1* and *rif-1* strains, there

376 was no correlation between the effect of parental *daf-2* RNAi on egg size and offspring
377 reproductive performance. We provisionally conclude that increased egg size under reduced
378 maternal insulin-like signalling can contribute to increased offspring fitness, but it is likely
379 not the sole source of variation in this trait. Recent work has identified oocyte-specific IIS
380 targets that are different from soma-specific IIS targets suggesting that IIS signalling
381 regulates reproduction and longevity through different mechanisms (Templeman et al. 2018).
382 In the future, it will be interesting to test for individual fitness of offspring produced via
383 genetic manipulation of oocyte-specific targets of IIS signalling pathway. In recent years,
384 there has been a vigorous debate regarding whether mechanistic understanding of life-history
385 trade-offs is necessary to advance life-history theory (Flatt and Heyland 2011, Stearns 2011a,
386 b). Here we used the mechanistic approach to separate between two conceptually different
387 evolutionary theories of ageing – energy trade-offs and functional trade-offs – in an empirical
388 study. We argue that unification between the conceptual approach and the mechanistic
389 understanding may often prove fruitful in this regard.

390

391 **Author contributions**

392 MIL and AAM designed the study, with the aid of AH. SR, ZS, MIL and HC collected the
393 data, MIL analysed the data, MIL and AAM drafted the manuscript. All authors contributed
394 to the revision of the manuscript.

395

396 **Acknowledgements**

397 This work was supported by ERC Consolidator Grant 724909 GermlineAgeingSoma to AAM
398 and Swedish Research Council VR Grant 2016-05195 to MIL. The authors declare no
399 conflicts of interest.

400

401 **Data accessibility**

402 Upon acceptance, the data will be archived at Dryad.

403

404

405 **References**

- 406 Agrawal, A. F., and A. D. Wang. 2008. Increased transmission of mutations by low-condition
407 females: Evidence for condition-dependent DNA repair. *Plos Biology* **6**:389-395.
- 408 Antebi, A. 2013. Regulation of longevity by the reproductive system. *Experimental*
409 *Gerontology* **48**:596-602.
- 410 Aprison, E. Z., and I. Ruvinsky. 2016. Sexually antagonistic male signals manipulate
411 germline and soma of *C. elegans* hermaphrodites. *Current Biology* **00**:XXX-XXX.
- 412 Berger, D., J. Stangberg, K. Grieshop, I. Martinossi-Allibert, and G. Arnqvist. 2017.
413 Temperature effects on life-history trade-offs, germline maintenance and mutation
414 rate under simulated climate warming. *Proceedings of the Royal Society B-Biological*
415 *Sciences* **284**.
- 416 Blagosklonny, M. V. 2006. Aging and immortality - Quasi-programmed senescence and its
417 pharmacologic inhibition. *Cell Cycle* **5**:2087-2102.
- 418 Blagosklonny, M. V. 2010. Revisiting the antagonistic pleiotropy theory of aging TOR-
419 driven program and quasi-program. *Cell Cycle* **9**:3151-3156.
- 420 Boonekamp, J. J., M. Salomons, S. Bouwhuis, C. Dijkstra, and S. Verhulst. 2014.
421 Reproductive effort accelerates actuarial senescence in wild birds: an experimental
422 study. *Ecology Letters* **17**:599-605.
- 423 Briga, M., and S. Verhulst. 2015. What can long-lived mutants tell us about mechanisms
424 causing aging and lifespan variation in natural environments? *Experimental*
425 *Gerontology* **71**:21-26.
- 426 Brommer, J. E., J. Merila, and H. Kokko. 2002. Reproductive timing and individual fitness.
427 *Ecology Letters* **5**:802-810.
- 428 Charlesworth, B. 1993. Evolutionary mechanisms of senescence. *Genetica* **91**:11-19.

- 429 Chen, H.-y., and A. A. Maklakov. 2012. Longer lifespan evolves under high rates of
430 condition dependent mortality. *Current Biology* **22**:2140-2143.
- 431 Chen, H.-y., and A. A. Maklakov. 2014. Condition dependence of male mortality drives the
432 evolution of sex differences in longevity. *Current Biology* **24**:2423-2427.
- 433 Curtsinger, J. W. 2019. Fecundity for free? Enhanced oviposition in longevous populations of
434 *Drosophila melanogaster*. *Biogerontology*.
- 435 Dillin, A., D. K. Crawford, and C. Kenyon. 2002. Timing requirements for insulin/IGF-1
436 signaling in *C-elegans*. *Science* **298**:830-834.
- 437 Edward, D. A., and T. Chapman. 2011. Mechanisms underlying reproductive trade-offs:
438 Costs of reproduction.
- 439 Ezcurra, M., A. Benedetto, T. Sornda, A. F. Gilliat, C. Au, Q. Zhang, S. van Schelt, A. L.
440 Petrache, H. Wang, Y. de la Guardia, S. Bar-Nun, E. Tuler, M. J. Wakelam, and D.
441 Gems. 2018. *C. elegans* eats its own intestine to make yolk leading to multiple
442 senescent pathologies. *Current Biology* **28**:2544-2556.
- 443 Flatt, T. 2011. Survival costs of reproduction in *Drosophila*. *Experimental Gerontology*
444 **46**:369-375.
- 445 Flatt, T., and A. Heyland. 2011. Why mechanisms are important for life-history theory: A
446 response by Flatt and Heyland. Pages 375-379 in T. Flatt and A. Heyland, editors.
447 Mechanisms of Life History Evolution: The Genetics and Physiology of Life History
448 Traits and Trade-Offs. Oxford University Press, Oxford.
- 449 Flatt, T., and L. Partridge. 2018. Horizons in the evolution of aging. *BMC Biology* **16**:93.
- 450 Gems, D., and L. Partridge. 2013. Genetics of longevity in model organisms: debates and
451 paradigm shifts. *Annual Review of Physiology* **75**:621-644.
- 452 Gems, D., A. J. Sutton, M. L. Sundermeyer, P. S. Albert, K. V. King, M. L. Edgley, P. L.
453 Larsen, and D. L. Riddle. 1998. Two pleiotropic classes of daf-2 mutation affect

- 454 larval arrest, adult behavior, reproduction and longevity in *Caenorhabditis elegans*.
455 *Genetics* **150**:129-155.
- 456 Gustafsson, L., and T. Part. 1990. Acceleration of senescence in the collared flycatcher
457 *Ficedula albicollis* by reproductive costs. *Nature* **347**:279-281.
- 458 Hamilton, W. D. 1966. The moulding of senescence by natural selection. *Journal of*
459 *Theoretical Biology* **12**:12-45.
- 460 Hibshman, J. D., A. Hung, and L. R. Baugh. 2016. Maternal diet and insulin-like signaling
461 control intergenerational plasticity of progeny size and starvation resistance. *PLoS*
462 *Genetics* DOI: [10.1371/journal.pgen.1006396](https://doi.org/10.1371/journal.pgen.1006396).
- 463 Hinas, A., A. J. Wright, and C. P. Hunter. 2012. SID-5 is an endosome-associated protein
464 required for efficient systemic RNAi in *C. elegans*. *Current Biology* **22**:1938-1943.
- 465 Hughes, K. A., and R. M. Reynolds. 2005. Evolutionary and mechanistic theories of aging.
466 *Annual Review of Entomology* **50**:421-445.
- 467 Jenkins, N. L., G. McColl, and G. J. Lithgow. 2004. Fitness cost of extended lifespan in
468 *Caenorhabditis elegans*. *Proceedings of the Royal Society of London Series B-*
469 *Biological Sciences* **271**:2523-2526.
- 470 Kenyon, C. 2005. The plasticity of aging: Insights from long-lived mutants. *Cell* **120**:449-
471 460.
- 472 Kenyon, C. 2011. The first long-lived mutants: discovery of the insulin/IGF-1 pathway for
473 ageing. *Philosophical Transactions of the Royal Society B-Biological Sciences* **366**:9-
474 16.
- 475 Kenyon, C. J. 2010. The genetics of ageing. *Nature* **464**:504-512.
- 476 Kimber, C. M., and A. K. Chippindale. 2013. Mutation, condition and the maintenance of
477 extended lifespan in *Drosophila*. *Current Biology* **23**: 2283-2287.
- 478 Kirkwood, T. B. L. 1977. Evolution of aging. *Nature* **270**:301-304.

- 479 Kirkwood, T. B. L. 2017. The Disposable Soma Theory of Ageing. Pages 23-39 *in*
480 Shefferson RP, Jones OR, and S.-G. R, editors. The Evolution of Senescence in the
481 Tree of Life. Cambridge University Press, Cambridge, United Kingdom.
- 482 Kirkwood, T. B. L., and S. N. Austad. 2000. Why do we age? *Nature* **408**:233-238.
- 483 Kirkwood, T. B. L., and R. Holliday. 1979. Evolution of aging and longevity. Proceedings of
484 the Royal Society of London Series B-Biological Sciences **205**:531-546.
- 485 Lemaitre, J.-F., V. Berger, C. Bonenfant, M. Douhard, M. Gamelon, F. Plard, and J.-M.
486 Gaillard. 2015. Early-late life trade-offs and the evolution of ageing in the wild.
487 Proceedings of the Royal Society B-Biological Sciences **282**.
- 488 Lind, M. I., H. Y. Chen, S. Meurling, A. C. G. Gil, H. Carlsson, M. K. Zwoinska, J.
489 Andersson, T. Larva, and A. A. Maklakov. 2017. Slow development as an
490 evolutionary cost of long life. *Functional Ecology* **31**:1252-1261.
- 491 Lind, M. I., M. K. Zwoinska, S. Meurling, H. Carlsson, and A. A. Maklakov. 2016. Sex-
492 specific trade-offs with growth and fitness following lifespan extension by rapamycin
493 in an outcrossing nematode, *Caenorhabditis remanei*. *Journals of Gerontology Series*
494 *a-Biological Sciences and Medical Sciences* **71**:882-890
- 495 Lionaki, E., and N. Tavernarakis. 2013. High-throughput and longitudinal analysis of aging
496 and senescent decline in *Caenorhabditis elegans*. Pages 485-500 *in* Lorenzo Galluzzi,
497 Ilio Vitale, Oliver Kepp, and G. Kroemer, editors. *Cell Senescence*.
- 498 Maklakov, A. A., H. Carlsson, P. Denbaum, M. I. Lind, B. Mautz, A. Hinas, and S. Immler.
499 2017. Antagonistically pleiotropic allele increases lifespan and late-life reproduction
500 at the cost of early-life reproduction and individual fitness. Proceedings of the Royal
501 Society B-Biological Sciences **284**.
- 502 Maklakov, A. A., and S. Immler. 2016. The expensive germline and the evolution of ageing.
503 *Current Biology* **26**:R577-R586.

- 504 Maures, T. J., L. N. Booth, B. A. Benayoun, Y. Izrayelit, F. C. Schroeder, and A. Brunet.
505 2013. Males shorten the life span of *C. elegans* hermaphrodites via secreted
506 compounds. *Science* **1244160**.
- 507 Nussey, D. H., H. Froy, J. F. Lemaitre, J. M. Gaillard, and S. N. Austad. 2013. Senescence in
508 natural populations of animals: Widespread evidence and its implications for bio-
509 gerontology. *Ageing Research Reviews* **12**:214-225.
- 510 Partridge, L., D. Gems, and D. J. Withers. 2005. Sex and death: What is the connection? *Cell*
511 **120**:461-472.
- 512 Partridge, L., N. Prowse, and P. Pignatelli. 1999. Another set of responses and correlated
513 responses to selection on age at reproduction in *Drosophila melanogaster*.
514 *Proceedings of the Royal Society of London Series B-Biological Sciences* **266**:255-
515 261.
- 516 Rose, M. R. 1984. Laboratory evolution of postponed senescence in *Drosophila*-
517 *melanogaster*. *Evolution* **38**:1004-1010.
- 518 Savory, F. R., T. G. Benton, V. Varma, I. A. Hope, and S. M. Sait. 2014. Stressful
519 environments can indirectly select for increased longevity. *Ecology and Evolution*
520 **4**:1176-1185.
- 521 Shi, C., and C. T. Murphy. 2014. Mating induces shrinking and death in *Caenorhabditis*
522 *mothers*. *Science Express*:1-7.
- 523 Sivasundar, A., and J. Hey. 2005. Sampling from natural populations with RNAi reveals high
524 outcrossing and population structure in *Caenorhabditis elegans*. *Current Biology*
525 **15**:1598-1602.
- 526 Sniegowski, P. D., P. J. Gerrish, T. Johnson, and A. Shaver. 2000. The evolution of mutation
527 rates: separating causes from consequences. *Bioessays* **22**:1057-1066.
- 528 Stearns, S. C. 1992. *The Evolution of Life Histories*. Oxford University Press, New York.


- 529 Stearns, S. C. 2011a. Does impressive progress on understanding mechanisms advance life
530 history theory? Pages 365-373 *in* T. Flatt and A. Heyland, editors. Mechanisms of
531 Life History Evolution: The Genetics and Physiology of Life History Traits and
532 Trade-Offs. Oxford University Press, Oxford.
- 533 Stearns, S. C. 2011b. Reply by Stearns: Mechanisms do not yet force the theory to change.
534 Page 379 *in* T. Flatt and A. Heyland, editors. Mechanisms of Life History Evolution:
535 The Genetics and Physiology of Life History Traits and Trade-Offs. Oxford
536 University Press, Oxford.
- 537 Templeman, N. M., S. J. Luo, R. Kaletsky, C. Shi, J. Ashraf, W. Keyes, and C. T. Murphy.
538 2018. Insulin signaling regulates oocyte quality maintenance with age via Cathepsin B
539 activity. *Current Biology* **28**:753-+.
- 540 Timmons, L., D. L. Court, and A. Fire. 2001. Ingestion of bacterially expressed dsRNAs can
541 produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene*
542 **263**:103-112.
- 543 Walker, D. W., G. McColl, N. L. Jenkins, J. Harris, and G. J. Lithgow. 2000. Natural
544 selection - evolution of lifespan in *C-elegans*. *Nature* **405**:296-297.
- 545 Williams, G. C. 1957. Pleiotropy, natural selection, and the evolution of senescence.
546 *Evolution* **11**:398-411.
- 547
- 548

549 **Table 1. The effect of *daf-2* RNAi on reproduction.** Total reproduction and individual
550 fitness (λ_{ind}) for unmated and mated *C. elegans* N2 wild-type treated with either empty vector
551 (Control) or *daf-2* RNAi from adulthood onwards. All values expressed as mean \pm SE.

RNAi treatment	Total reproduction		Fitness (λ_{ind})	
	unmated	mated	unmated	mated
Control	311.0 \pm 7.0	595.7 \pm 24.3	4.66 \pm 0.03	4.47 \pm 0.05
<i>daf-2</i> RNAi	317.4 \pm 8.9	630.5 \pm 22.2	4.63 \pm 0.05	4.50 \pm 0.05


552

553




554

555 **Fig. 1. Theoretical framework.** The relationship between population genetic theory of
 556 ageing (“antagonistic pleiotropy”, Williams 1957) and physiological theories of ageing based
 557 on either energy trade-offs (“disposable soma” (Kirkwood 1977) or functional trade-offs (e.g.
 558 “hyperfunction” (Blagosklonny 2006)). Note that Williams (1957) was the first to provide an
 559 abstract example of a functional trade-off between early-life and late-life allelic effects on
 560 organismal physiology. Recently, Blagosklonny (2006, 2010) put forward a “hyperfunction”
 561 hypothesis that specifically links suboptimal levels of nutrient-sensing signalling to excessive
 562 biosynthesis (hence, hyperfunction) leading to cellular and organismal senescence.




563

564 **Fig. 2. The effect of *daf-2* RNAi on lifespan.** Survival probability for (A) unmated or (B)
 565 mated N2 wild-type (purple), *ppw-1* (black) and *rrf-1* (orange) mutants, treated with either
 566 *daf-2* RNAi (solid lines, filled symbols) or control empty vector (broken lines, open symbols)
 567 from adulthood onwards.




568
 569 **Fig. 3. The effect of *daf-2* RNAi on reproduction.** Daily offspring number for (A) unmatd
 570 or (B) mated N2 wild-type worms, treated with either *daf-2* RNAi (solid lines, filled symbols)
 571 or control empty vector (broken lines, open symbols) from adulthood onwards. Symbols
 572 represent mean \pm SE.

573



574
 575 **Fig. 4. The effect of *daf-2* RNAi on egg size.** Egg size of unmatd, parental worms. N2 wild-
 576 type (purple), *ppw-1* (black) and *rrf-1* (orange) mutants, treated with either *daf-2* RNAi (solid

577 lines, filled symbols) or control empty vector (broken lines, open symbols) from adulthood
 578 onwards. Symbols represent mean \pm SE.



579
 580 **Fig. 5. The effect of parental *daf-2* RNAi on offspring survival and reproduction.**

581 Offspring worms, unmated, on control (empty vector) plates from parents exposed to *daf-2*
 582 RNAi or control treatment. (A) Survival probability, (B) daily offspring number, (C) Total
 583 reproduction and (D) individual fitness (λ_{ind}) of offspring (on control plates) from parents
 584 either exposed to *daf-2* RNAi (solid lines, filled symbols) or control empty vector (broken
 585 lines, open symbols). The colors reflect N2 wild-type (purple), *ppw-1* (black) and *rrf-1*
 586 (orange) mutants. Symbols represent mean \pm SE.