RUNNING TITLE: BMI polygenic score, early stress, and depression.

A Polygenic Score for Body Mass Index is Associated with Depressive Symptoms via Early Life Stress: Evidence for gene-environment correlation

Reut Avinun and Ahmad R. Hariri

Laboratory of NeuroGenetics, Department of Psychology & Neuroscience, Duke University, Durham, NC, USA

FUNDING AND DISCLOSURE

We would like to thank the participants of the Duke Neurogenetics Study and members of the Laboratory of NeuroGenetics, especially Annchen R. Knodt, Spenser R. Radtke, and Bartholomew D. Brigidi for their assistance with data collection and analysis. The DNS was supported by Duke University as well as US-National Institutes of Health grants R01DA033369 and R01DA031579. RA and ARH received further support from US-National Institutes of Health grant R01AG049789. The Brain Imaging and Analysis Center was supported by the Office of the Director, National Institutes of Health under Award Number S10OD021480. The authors declare no competing financial or other interests.

Corresponding Author: Reut Avinun, Ph.D., Laboratory of NeuroGenetics, Department of Psychology & Neuroscience, Duke University, Grey Building 2020 West Main St, Ste 0030 Durham, NC 27705. Email: reut.avinun@duke.edu

1 **ABSTRACT**

2 **Background:** Increasing childhood overweight and obesity rates are associated with
 3 not only adverse physical, but also mental health outcomes, including depression.
 4 These negative outcomes may be caused and/or exacerbated by the bullying and
 5 shaming overweight individuals experience. As body mass index (BMI) can be highly
 6 heritable, we hypothesized that a genetic risk toward higher BMI, will predict higher
 7 early life stress (ELS), which in turn will predict higher depressive symptoms in
 8 adulthood. Such a process will reflect an evocative gene-environment correlation
 9 (rGE) wherein an individual's genetically influenced phenotype evokes a reaction
 10 from the environment that subsequently shapes the individual's health.

11 **Methods:** We modeled genetic risk using a polygenic score of BMI derived from a
 12 recent large GWAS meta-analysis. Self-reports were used for the assessment of ELS
 13 and depressive symptoms in adulthood. The discovery sample consisted of 524 non-
 14 Hispanic Caucasian university students from the Duke Neurogenetics Study (DNS;
 15 278 women, mean age 19.78 ± 1.23 years) and the independent replication sample
 16 consisted of 5 930 white British individuals from the UK biobank (UKB; 3 128 women,
 17 mean age 62.66 ± 7.38 years).

18 **Results:** A significant mediation effect was found in the DNS (indirect effect=.207,
 19 bootstrapped SE=.10, 95% CI: .014 to .421), and then replicated in the UKB (indirect
 20 effect=.04, bootstrapped SE=.01, 95% CI: .018 to .066). Higher BMI polygenic scores
 21 were associated with higher depressive symptoms through the experience of higher
 22 ELS.

23 **Conclusions:** Our findings suggest that evocative rGE may contribute to weight-
24 related mental health problems and stress the need for interventions that aim to
25 reduce weight bias, specifically during childhood.

Overweight individuals suffer from stigmatization, bias, and bullying, from multiple sources including peers, health care providers, educators, and, most surprisingly perhaps, family members [1]. In a study of adolescents enrolled in weight loss camps, 37% reported being teased or bullied by a parent [2]. Another study on 2 449 women recruited from a weight loss support group organization, found that 44% experienced stigma from their mothers more than once, while 34% experienced it from their fathers [3]. As weight-related teasing has been shown to predict depression and lower self-esteem [4, 5], it may represent another form of early life stress (ELS) that is associated with various negative physical and mental health outcomes [6, 7].

Gene environment correlations [rGE; 8, 9] can represent passive, evocative, and active processes that create associations between individuals' genes and the environment. Evocative rGE, which refers to instances in which a genetically influenced phenotype of an individual evokes a certain reaction from the environment, may be relevant to weight-related teasing and bullying, so that individuals with a genetic propensity toward a higher body mass index (BMI), will be more likely to experience teasing, especially in the current Western cultural climate, which is characterized by negative and prejudicial attitudes towards overweight and obese individuals [10].

A recent meta-analysis of genome-wide association studies (GWAS; [11]), consisting of 681 275 participants on average, explained 5% of the variance in BMI with GWAS significant single nucleotide polymorphisms (SNPs). In the current study, we hypothesized that a polygenic score based on the results from this meta-analysis, will predict early life stress (ELS), consistent with evocative rGE, which in turn will

50 predict depressive symptoms in adulthood. We tested our hypothesis in two
51 independent samples: a discovery sample of 524 non-Hispanic Caucasian university
52 students from the Duke Neurogenetics Study and a replication sample of 5 930 adult
53 white British volunteers from the UK Biobank (UKB). Although the GWAS meta-
54 analysis included data from the UKB, current BMI was not a phenotype of interest in
55 our study, and therefore the overlap should not bias our analyses. Nonetheless, to
56 validate our results in the analyses that included UKB data, we also used BMI
57 polygenic scores that were based on a GWAS that did not include the UKB as a
58 discovery sample [12].

59

60 MATERIALS AND METHODS

61 *Participants*

62 Our discovery sample consisted of 524 self-reported non-Hispanic Caucasian
63 participants (278 women, mean age 19.78 ± 1.23 years) from the Duke
64 Neurogenetics Study (DNS) for whom there was complete data on genotypes, ELS,
65 depressive symptoms, and all covariates. All procedures were approved by the
66 Institutional Review Board of the Duke University Medical Center, and participants
67 provided informed consent before study initiation. All participants were free of the
68 following study exclusions: 1) medical diagnoses of cancer, stroke, diabetes requiring
69 insulin treatment, chronic kidney or liver disease, or lifetime history of psychotic
70 symptoms; 2) use of psychotropic, glucocorticoid, or hypolipidemic medication; and
71 3) conditions affecting cerebral blood flow and metabolism (e.g., hypertension).
72 Importantly, neither current nor lifetime diagnosis were an exclusion criterion, as the

DNS sought to establish broad variability in multiple behavioral phenotypes related to psychopathology.

The replication sample consisted of 5 930 white British individuals (3 128 women, mean age 62.66 ± 7.38 years), who participated in the UKB's imaging wave, completed an online mental health questionnaire [13], and had complete genotype, ELS, depressive symptoms and covariate data. The UKB [www.ukbiobank.ac.uk; 14] includes over 500,000 participants, between the ages of 40 and 69 years, who were recruited within the UK between 2006 and 2010. The UKB study has been approved by the National Health Service Research Ethics Service (reference: 11/NW/0382), and our analyses were conducted under UKB application 28174.

Race/Ethnicity

Because self-reported race and ethnicity are not always an accurate reflection of genetic ancestry, an analysis of identity by state of whole-genome SNPs in the DNS was performed in PLINK [15]. The first two multidimensional scaling components within the non-Hispanic Caucasian subgroup were used as covariates in analyses of data from the DNS. The decision to use only the first two components was based on an examination of a scree plot of the variance explained by each component. For analyses of data from the UKB, only those who were 'white British' based on both self-identification and a principal components analysis of genetic ancestry were included. Additionally, the first 10 multidimensional scaling components received from the UKB's data repository (unique data identifiers: 22009-0.1-22009-0.10) were included as covariates as previously done [e.g., 16].

97 *Body Mass Index (BMI)*

98 In both DNS and UKB samples, BMI was calculated at the time of imaging based on
99 the height and weight of the participants. In the DNS, this calculation was based on
100 imperial system values (pounds/inches²*703), while in the UKB the metric system
101 was used (kg/m²). In the DNS 1.3% of the sample was obese, compared to 18.7% in
102 the UKB.

103

104 *Depressive symptoms*

105 In the DNS, the 20-item Center for Epidemiologic Studies Depression Scale (CES-D)
106 was used to asses depressive symptoms in the past week [17]. All items were
107 summed to create a total depressive symptoms score. In the UKB, the Patient Health
108 Questionnaire 9-question version (PHQ-9) was used to asses depressive symptoms in
109 the past 2 weeks [18]. All items were summed to create a total depressive symptoms
110 score.

111

112 *Early life stress*

113 In the DNS, ELS was estimated using the Childhood Trauma Questionnaire [CTQ; 19].
114 The CTQ has 28-items and it assesses the frequency of emotional, physical, and
115 sexual abuse as well as emotional and physical neglect. The scores on the 5
116 subscales (each ranging from 5 to 25) were summed to create a total score of ELS. In
117 the UKB, the Childhood Trauma Screener – 5 item (CTS-5) was used to assess adverse
118 events during childhood [20]. CTS-5 is a short version of the CTQ consisting of 5
119 items: "Felt hated by family member as a child", "Physically abused by family as a
120 child", "Felt loved as a child" (reverse coded), "Sexually molested as a child", and

"Someone to take to doctor when needed as a child" (reverse coded). The 5 items, each ranging from 0-4, were summed to create a total score of ELS.

Genotyping

In the DNS, DNA was isolated from saliva using Oragene DNA self-collection kits (DNA Genotek) customized for 23andMe (www.23andme.com). DNA extraction and genotyping were performed through 23andMe by the National Genetics Institute (NGI), a CLIA-certified clinical laboratory and subsidiary of Laboratory Corporation of America. One of two different Illumina arrays with custom content was used to provide genome-wide SNP data, the HumanOmniExpress (N=329) or HumanOmniExpress-24 [N=195; 21, 22, 23]. In the UKB, samples were genotyped using either the UK BiLEVE (N=569) or the UKB axion (N=5,361) array. Details regarding the UKB's quality control can be found elsewhere[24].

Quality control and polygenic scoring

For genetic data from both the DNS and UK Bionbank, PLINK v1.90 [15] was used to apply quality control cutoffs and exclude SNPs or individuals based on the following criteria: missing genotype rate per individual $>.10$, missing rate per SNP $>.10$, minor allele frequency $<.01$, and Hardy-Weinberg equilibrium $p < 1e-6$. Additionally, in the UKB, quality control variables that were provided with the dataset were used to exclude participants based on a sex mismatch (genetic sex different from reported sex), a genetic relationship to another participant, outliers for heterozygosity or missingness (unique Data Identifier 22010-0.0), and UKBiLEVE genotype quality control for samples (unique Data Identifiers 22050-0.0-22052-0.0).

Polygenic scores were calculated using PLINK's [15] "--score" command based on published SNP-level summary statistics from a recent BMI GWAS meta-analysis [11]. SNPs from the GWAS of BMI meta-analysis were matched with SNPs from the DNS and the UKB. For each SNP the number of the alleles (0, 1, or 2) associated with BMI was multiplied by the effect estimated in the GWAS. The polygenic score for each individual was an average of weighted BMI-associated alleles. All SNPs matched with SNPs from the DNS and UKB were used regardless of effect size and significance in the original GWAS, as previously recommended and shown to be effective [25, 26]. A total of 442 040 SNPs from the DNS and 648 530 SNPs from the UKB were included in the polygenic scores. The approach described here for the calculation of the polygenic scores was successfully used in previous studies [e.g., 27, 28-30]. For validation of the indirect effect in the UKB, BMI polygenic scores were also calculated based on an older GWAS that did not include the UKB as a discovery sample [12].

Statistical analysis

Linear regression analyses in SPSS v25 were conducted to test for an association between the BMI polygenic score and BMI in adulthood. The PROCESS SPSS macro, version 3.1 [31], was used to conduct the mediation analyses. Participants' sex (coded as 0=males, 1=females), age, and ethnicity genomic components were entered as covariates in all analyses. In the mediation analyses, bias-corrected bootstrapping (set to 5,000) was used to allow for non-symmetric 95% confidence intervals (CIs). Specifically, indirect effects are likely to have a non-normal distribution, and consequently the use of non-symmetric CIs for the determination of significance is recommended [32]. However, bias-corrected bootstrapping also has

its faults [33] and, consequently, as supportive evidence for the indirect effect, we also present the test of joint significance, which examines whether the *a path* (BMI polygenic score to ELS) and the *b path* (ELS to depressive symptoms, while controlling for the BMI polygenic score) are significant. The BMI polygenic scores were standardized to make interpretability easier. The mediation was first analyzed in the DNS, and then a replication was tested in the UKB. As a validation of the indirect effect in the UKB, it was also tested with an older BMI polygenic score that was not based on a GWAS that included the UKB [12].

RESULTS

Descriptive statistics are presented in table 1.

Confirming an association between BMI polygenic scores and measured BMI

As a preliminary analysis we confirmed that higher BMI polygenic scores were significantly associated with higher measured BMI in both the DNS (N=522, $b=.837$, $SE=.117$, $p<.001$) and the UKB (N=5 925, $b=1.41$, $SE=.054$, $p<.001$). (notably, the UKB was included in the BMI GWAS, and consequently the significant association is expected and possibly somewhat inflated). These associations were robust to the inclusion of sex, age, and ethnicity genomic components as covariates. The sample sizes for these analyses were slightly different from the mediation analyses below because measured BMI was missing for a few participants.

BMI polygenic scores predict ELS (a path) in the DNS

The BMI polygenic scores were significantly associated with ELS ($b=.65$, $SE=.31$, $p=.038$), so that higher scores predicted higher ELS. Of the covariates, age was significantly and negatively associated with ELS ($b=-.73$, $SE=.25$, $p<.01$).

ELS predicts depressive symptoms (b path) in the DNS

With the BMI polygenic scores in the model, ELS significantly and positively predicted depressive symptoms ($b=.32$, $SE=.04$, $p<.001$).

BMI polygenic scores predict depressive symptoms in the DNS

The BMI polygenic scores did not significantly predict depressive symptoms ($b=-.34$, $SE=.31$, ns). Notably, however, the significance of a direct path from X (BMI polygenic scores) to Y (depressive symptoms) or the 'total effect' (the 'c' path), is not a prerequisite for the testing of a mediation/indirect effect [34-36].

Mediation model in the DNS

The indirect path ($a*b$), BMI polygenic scores to ELS to depressive symptoms was significant as indicated by the bias corrected bootstrapped 95% CI not including zero (Figure 1a; indirect effect $=.207$, bootstrapped $SE=.10$, 95% CI: .014 to .421).

Mediation Model in the UBK

The *a path*, from the BMI polygenic scores to ELS, and the *b path*, from ELS to depressive symptoms while controlling for BMI polygenic scores, were significant (*a path*: $b=.10$, $SE=.03$, $p<.01$; *b path*: $b=.40$, $SE=.02$, $p<.001$). The indirect path also replicated (Figure 1b; indirect effect $=.04$, bootstrapped $SE=.01$, 95% CI: .018 to .066),

supporting a mediation in which BMI polygenic scores are associated with depressive symptoms indirectly through ELS. Similar results were obtained with the BMI polygenic scores that were based on a GWAS that did not include the UKB as a discovery sample (indirect effect=.026, bootstrapped SE=.01, 95% CI: .004 to .05).

DISCUSSION

Here, in two independent samples, we provide novel evidence supporting evocative rGE as a possible mechanism in weight-related depression. We demonstrate a significant mediation in which higher GWAS-derived BMI polygenic scores are associated with higher levels of depressive symptoms in adulthood through elevated levels of ELS. These results suggest that in the current Western cultural climate, having a genetic makeup that increases the risk of a high BMI, may lead to a phenotype that evokes increased stress, which increases the experience of depressive symptoms in adulthood.

Various studies have reported links between being overweight and experiencing stigmatization, teasing, and bullying from peers, educators, co-workers, health care providers, and family members [1]. This negativity can lead to adverse mental health outcomes, including depression [5], but is not limited to mental health. Obesity, childhood trauma, and depression have all been linked to physical illness including cardiovascular disease, type 2 diabetes, and autoimmune disorders [6, 37-40].

While several strategies have been proposed to battle the growing prevalence of childhood obesity, including nutrition standards for school meals; improved early care and education; and increased access to adolescent bariatric surgery [41], our

findings further encourage weight stigma reduction efforts, specifically among family members and parents. In addition to the myriad of mental and physical health disorders that are associated with ELS and childhood trauma, one of the most prevalent coping responses to weight stigma is eating [3]. Consequently, ELS may lead to additional weight gain and is itself a risk factor for obesity [42]. Thus, interventions that aim to reduce weight stigma may have a broad positive effect on health.

Although our study has several strengths, including the use of two independent samples with markedly different characteristics (e.g., young university students versus older community volunteers) and a GWAS-derived polygenic score, it is not without limitations. First, retrospective reports were used for the estimation of ELS and childhood trauma. Ideally, prospective data should be used to model ELS in the absence of reporting bias. Second, we did not have measures of childhood BMI in either sample. Although previous research does support a link between childhood BMI, teasing, and depression, and genetic influences on BMI have been shown to be relatively stable throughout development [43, 44], genetically informed longitudinal studies across development are needed to further validate our findings. Third, the non-Hispanic Caucasian DNS sample is relatively homogeneous in terms of social background, which may have led to an underestimation of the effect in this sample. Fourth, our findings are limited to populations of European descent and to the Western culture. Additional research in diverse populations is needed to determine the extent to which the observed evocative rGE mechanism shapes weight-related mental health. Further replication is also needed to evaluate the potential of the BMI polygenic score as a risk biomarker of depression associated with ELS.

References

1. Puhl RM, Latner JD. Stigma, obesity, and the health of the nation's children. *Psychol. Bull.* 2007; **133**(4): 557.
2. Puhl RM, Peterson JL, Luedicke J. Weight-based victimization: Bullying experiences of weight loss treatment-seeking youth. *Pediatrics* 2013; **131**(1): e1-e9.
3. Puhl RM, Brownell KD. Confronting and coping with weight stigma: an investigation of overweight and obese adults. *Obesity* 2006; **14**(10): 1802-1815.
4. Greenleaf C, Petrie TA, Martin SB. Relationship of weight-based teasing and adolescents' psychological well-being and physical health. *J. Sch. Health* 2014; **84**(1): 49-55.
5. Greenleaf C, Petrie T, Martin S. Exploring weight-related teasing and depression among overweight and obese adolescents. *Revue Européenne de Psychologie Appliquée/European Review of Applied Psychology* 2017; **67**(3): 147-153.

- 286 6. Goodwin RD, Stein MB. Association between childhood trauma and physical
287 disorders among adults in the United States. *Psychol. Med.* 2004; **34**(3): 509-
288 520.
289
- 290 7. Mersky JP, Topitzes J, Reynolds AJ. Impacts of adverse childhood experiences
291 on health, mental health, and substance use in early adulthood: A cohort
292 study of an urban, minority sample in the US. *Child Abuse Negl.* 2013; **37**(11):
293 917-925.
294
- 295 8. Plomin R, DeFries JC, Loehlin JC. Genotype-environment interaction and
296 correlation in the analysis of human behavior. *Psychol. Bull.* 1977; **84**(2): 309-
297 22.
298
- 299 9. Avinun R, Knafo-Noam A. Socialization, Genetics and their Interplay in
300 Development. In: Grusec JE, Hastings PD (eds). *Handbook of Socialization:*
301 *Theory and research*, 2 edn. Guilford Press, 2014, pp 347-371.
302
- 303 10. Pearl RL. Weight bias and stigma: public health implications and structural
304 solutions. *Social Issues and Policy Review* 2018; **12**(1): 146-182.
305

- 306 11. Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN *et al.*
307 Meta-analysis of genome-wide association studies for height and body mass
308 index in ~ 700,000 individuals of European ancestry. *bioRxiv* 2018: 274654.
309
- 310 12. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR *et al.* Genetic studies
311 of body mass index yield new insights for obesity biology. *Nature* 2015;
312 **518**(7538): 197.
313
- 314 13. Davis KA, Coleman JR, Adams M, Allen N, Breen G, Cullen B *et al.* Mental
315 health in UK Biobank: development, implementation and results from an
316 online questionnaire completed by 157 366 participants. *BJPsych Open* 2018;
317 **4**(3): 83-90.
318
- 319 14. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J *et al.* UK biobank:
320 an open access resource for identifying the causes of a wide range of complex
321 diseases of middle and old age. *PLoS Med.* 2015; **12**(3): e1001779.
322
- 323 15. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D *et al.*
324 PLINK: a tool set for whole-genome association and population-based linkage
325 analyses. *The American Journal of Human Genetics* 2007; **81**(3): 559-575.
326

- 327 16. Whalley HC, Adams MJ, Hall L, Clarke T-K, Fernandez-Pujals AM, Gibson J *et*
328 *al.* Dissection of major depressive disorder using polygenic risk scores for
329 schizophrenia in two independent cohorts. *Translational Psychiatry* 2016;
330 **6**(11): e938.
331
- 332 17. Radloff LS. The CES-D scale: A self-report depression scale for research in the
333 general population. *Applied psychological measurement* 1977; **1**(3): 385-401.
334
- 335 18. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression
336 severity measure. *J. Gen. Intern. Med.* 2001; **16**(9): 606-613.
337
- 338 19. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T *et al.*
339 Development and validation of a brief screening version of the Childhood
340 Trauma Questionnaire. *Child Abuse Negl.* 2003; **27**(2): 169-190.
341
- 342 20. Glaesmer H, Schulz A, Häuser W, Freyberger HJ, Brähler E, Grabe H-J. The
343 childhood trauma screener (CTS)-development and validation of cut-off-
344 scores for classificatory diagnostics. *Psychiatr. Prax.* 2013; **40**(4): 220-226.
345
- 346 21. Do CB, Tung JY, Dorfman E, Kiefer AK, Drabant EM, Francke U *et al.* Web-
347 based genome-wide association study identifies two novel loci and a

- 348 substantial genetic component for Parkinson's disease. *PLoS Genet.* 2011;
349 **7**(6): e1002141.
- 350
- 351 22. Eriksson N, Macpherson JM, Tung JY, Hon LS, Naughton B, Saxonov S *et al.*
352 Web-based, participant-driven studies yield novel genetic associations for
353 common traits. *PLoS Genet.* 2010; **6**(6): e1000993.
- 354
- 355 23. Tung JY, Do CB, Hinds DA, Kiefer AK, Macpherson JM, Chowdry AB *et al.*
356 Efficient replication of over 180 genetic associations with self-reported
357 medical data. *PLoS ONE* 2011; **6**(8): e23473.
- 358
- 359 24. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K *et al.* Genome-
360 wide genetic data on ~ 500,000 UK Biobank participants. *bioRxiv* 2017:
361 166298.
- 362
- 363 25. Dudbridge F. Power and predictive accuracy of polygenic risk scores. *PLoS*
364 *Genet.* 2013; **9**(3): e1003348.
- 365
- 366 26. Ware EB, Schmitz LL, Faul JD, Gard A, Mitchell C, Smith JA *et al.* Heterogeneity
367 in polygenic scores for common human traits. *bioRxiv* 2017: 106062.
- 368

- 369 27. Domingue BW, Belsky DW, Conley D, Harris KM, Boardman JD. Polygenic
370 influence on educational attainment: New evidence from the National
371 Longitudinal Study of Adolescent to Adult Health. *AERA open* 2015; **1**(3):
372 2332858415599972.
373
- 374 28. Domingue BW, Liu H, Okbay A, Belsky DW. Genetic heterogeneity in
375 depressive symptoms following the death of a spouse: Polygenic score
376 analysis of the US Health and Retirement Study. *Am. J. Psychiatry* 2017;
377 **174**(10): 963-970.
378
- 379 29. Stephan Y, Sutin AR, Luchetti M, Caille P, Terracciano A. Polygenic Score for
380 Alzheimer Disease and cognition: The mediating role of personality. *J.*
381 *Psychiatr. Res.* 2018.
382
- 383 30. Marioni RE, Ritchie SJ, Joshi PK, Hagenaars SP, Okbay A, Fischer K *et al.*
384 Genetic variants linked to education predict longevity. *Proc. Natl. Acad. Sci.*
385 *U. S. A.* 2016; **113**(47): 13366-13371.
386
- 387 31. Hayes AF. *Introduction to mediation, moderation, and conditional process*
388 *analysis: A regression-based approach*, Guilford Publications, 2017.
389

- 390 32. MacKinnon DP, Lockwood CM, Williams J. Confidence Limits for the Indirect
391 Effect: Distribution of the Product and Resampling Methods. *Multivariate*
392 *Behavioral Research* 2004; **39**(1): 99.
- 393
- 394 33. Hayes AF, Scharkow M. The relative trustworthiness of inferential tests of the
395 indirect effect in statistical mediation analysis: Does method really matter?
396 *Psychol. Sci.* 2013; **24**(10): 1918-1927.
- 397
- 398 34. MacKinnon DP, Krull JL, Lockwood CM. Equivalence of the mediation,
399 confounding and suppression effect. *Prev. Sci.* 2000; **1**(4): 173-181.
- 400
- 401 35. Rucker DD, Preacher KJ, Tormala ZL, Petty RE. Mediation analysis in social
402 psychology: Current practices and new recommendations. *Social and*
403 *Personality Psychology Compass* 2011; **5**(6): 359-371.
- 404
- 405 36. Hayes AF. Beyond Baron and Kenny: Statistical mediation analysis in the new
406 millennium. *Communication monographs* 2009; **76**(4): 408-420.
- 407
- 408 37. Carney RM, Freedland KE. Depression and coronary heart disease. *Nature*
409 *Reviews Cardiology* 2017; **14**(3): 145.
- 410

- 411 38. Versini M, Jeandel P-Y, Rosenthal E, Shoenfeld Y. Obesity in autoimmune
412 diseases: not a passive bystander. *Autoimmun. Rev.* 2014; **13**(9): 981-1000.
413
- 414 39. Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin
415 resistance and type 2 diabetes. *Nature* 2006; **444**(7121): 840.
416
- 417 40. Van Gaal LF, Mertens IL, Christophe E. Mechanisms linking obesity with
418 cardiovascular disease. *Nature* 2006; **444**(7121): 875.
419
- 420 41. Gortmaker SL, Wang YC, Long MW, Giles CM, Ward ZJ, Barrett JL *et al.* Three
421 interventions that reduce childhood obesity are projected to save more than
422 they cost to implement. *Health Aff. (Millwood)* 2015; **34**(11): 1932-1939.
423
- 424 42. Danese A, Tan M. Childhood maltreatment and obesity: systematic review
425 and meta-analysis. *Mol. Psychiatry* 2014; **19**(5): 544.
426
- 427 43. Haberstick BC, Lessem JM, McQueen MB, Boardman JD, Hopfer CJ, Smolen A
428 *et al.* Stable genes and changing environments: body mass index across
429 adolescence and young adulthood. *Behav. Genet.* 2010; **40**(4): 495-504.
430

431 44. Silventoinen K, Kaprio J. Genetics of tracking of body mass index from birth to
432 late middle age: evidence from twin and family studies. *Obes. Facts* 2009;
433 2(3): 196-202.

434

435

436

437

Table 1. Descriptive statistics of study variables.

	DNS				UK Biobank			
	Min	Max	Mean	SD	Min	Max	Mean	SD
Age	18	22	19.78	1.24	45	78	62.66	7.38
BMI	16.30	39.15	22.29	2.83	14.94	58.04	26.60	4.419
Early life stress	25	74	31.29	7.16	0	20	1.68	2.32
Depressive symptoms	0	43	8.99	7.18	0	27	2.45	3.39

Figure 1. Mediation model linking genetic risk for higher BMI to higher depressive symptoms, via elevated levels of early life stress

1a. Duke Neurogenetics Study: Discovery sample

1b. UK Biobank: Replication sample

Note. * $p < .05$, ** $p < .01$, *** $p < .001$. c- the total effect of the BMI polygenic scores on depressive symptoms; c'-the effect of BMI polygenic scores on depressive symptoms, while controlling for ELS.