1 **Title: Wildebeest migration in East Africa: Status, threats and**
2 **conservation measures**

3 **Running head: Status of wildebeest migration in East Africa**

4 Fortunata U. Msoffe^{1,7,9†}, Joseph O. Ogutu^{1,2*†}, Mohammed Y. Said^{1,3,4}, Shem C.
5 Kifugo^{1,5}, Jan de Leeuw^{1,6}, Paul Van Gardingen⁷, Robin S. Reid^{1,8}, JA Stabach¹⁰,
6 Randall B. Boone¹¹

7

8 ¹International Livestock Research Institute, P.O. Box 30709-00100, Nairobi, Kenya

9

10 ²University of Hohenheim, Institute for Crop Science-340, 70599 Stuttgart, Germany

11

12 ³Kenya Market Trust, 14 Riverside, Cavendish Block 3rd Floor, Suite B, Riverside
13 Drive P.O. Box 44817 – 00100, Nairobi, Kenya

14

15 ⁴Institute for Climate Change and Adaptation, University of Nairobi, P.O. Box 30197,
16 00100, Nairobi, Kenya.

17

18 ⁵Northern Rangelands Trust, Private Bag, Isiolo 60300, Kenya.

19

20 ⁶ International Soil Reference and Information Centre (ISRIC) P.O. Box 353 6700 AJ
21 Wageningen, The Netherlands.

22

23 ⁷Centre for the Study of Environmental Change and Sustainability, School of

24 Geosciences, University of Edinburgh, Edinburgh, Scotland, UK

25

26 ⁸Centre for Collaborative Conservation and Department of Ecosystem Science and

27 Sustainability, Colorado State University, Fort Collins, CO, USA

28

29 ⁹Tanzania National Parks Authority, P.O. Box 3134, Arusha, Tanzania

30

31 ¹⁰Conservation Ecology Center, National Zoological Park, Smithsonian Conservation

32 Biology Institute, Front Royal, VA, USA

33

34 ¹¹Natural Resource Ecology Laboratory Colorado State University Fort Collins, CO

35 80523-1499

36 [†]Joint first authors

37 Fortunate Msoffe: fortu2_2@yahoo.com

38 Joseph Ogutu: jogutu2007@gmail.com

39 Mohammed Said: msaid362@gmail.com

40 Shem Kifugo: ckifugo2014@gmail.com

41 Jan de Leeuw jan.deleeuw@wur.nl

42 Paul van Gardingen p.vangardingen@gmail.com

43 Robin Reid Robin.Reid@colostate.edu

44 Jared Stabach stabachj@si.edu

45 Randall Boone Randall.Boone@colostate.edu

46

47 ***Corresponding author:**

Current Address: Institute for Crop Science-340, 70599 Stuttgart, Germany;

Tel: +49-711-459 23022; Fax: +49-711-459-24345;

E-mail: jogutu2007@gmail.com

Abstract

Migration of ungulates is under pressure worldwide from range contraction, habitat loss and degradation, anthropogenic barriers and poaching. Here, we synthesize and compare the extent of historical migrations of the white-bearded wildebeest (*Connochaetes taurinus*) to their contemporary status, in five premier East African ecosystems, namely the Serengeti-Mara, Masai Mara, Athi-Kaputiei, Amboseli and Tarangire-Manyara. The current status, threats to migration, migratory ranges and routes for wildebeest were characterized using colonial-era maps, literature reviews, GIS and aerial survey databases, GPS collared animals and interviews with long-term researchers. Interference with wildebeest migratory routes and dispersal ranges has stopped or severely threatens continuation of the historical migration patterns in all but the Serengeti-Mara ecosystem where the threat level is relatively lower. Wildebeest migration has collapsed in Athi-Kaputiei ecosystem and is facing enormous pressures from land subdivision, settlements and fences in Amboseli and Mara ecosystems and from cultivation in Tarangire-Manyara ecosystem. Land use change, primarily expansion in agriculture, roads, settlements and fencing, increasingly restrict migratory wildebeest from accessing traditional grazing resources in unprotected lands. Privatization of land tenure in group ranches in Kenya and settlement policy (villagization) in Tanzania have accelerated land subdivision, fencing and growth in permanent settlements, leading to loss of key wildebeest habitats including their migratory routes and wet season calving and feeding grounds.

These processes, coupled with increasing human population pressures and climatic variability, are exerting tremendous pressures on wildebeest migrations. Urgent conservation interventions are necessary to conserve and protect the critical wildebeest habitats and migration routes in East Africa.

Keywords: Wildebeest; population declines; migration; migratory routes, migratory corridors, land use change; land tenure change; wildlife conservancies; agriculture; settlements; fences; human population growth; poaching; Kenya; Tanzania; Serengeti-Mara Ecosystem; Masai Mara Ecosystem; Loita Plains; Athi-Kaputiei Ecosystem; Athi-Kaputiei Plains; Amboseli Ecosystem, Western Kajiado Ecosystem, Ngorongoro Conservation Area; Tarangire-Manyara Ecosystem

Introduction

Large mammal migrations are among the most awe inspiring of all migrations [1]. These migrations, the seasonal and round-trip movement of large herbivores between discrete areas, are under increasing pressures worldwide. Globally, migrations of 6 out of 24 species of ungulates are either already extinct or their status is unknown [2]. Of the remaining ungulate mass migrations, most occur in six locations in Africa, including the white-bearded wildebeest (*Connochaetes taurinus* Burchell, 1823) migration in the Serengeti-Mara ecosystem of Kenya and Tanzania [2]. Range restriction and alteration, degradation and loss of habitat due to agriculture, poaching and barriers that block migration, such as fences, roads, railroads, pipelines and settlements have progressively disrupted historical migratory routes and decimated or driven rapid population declines of many of the once spectacular migratory herds over the 20th century [1,3-5] .

97

98 Because migration enables populations to grow to large abundances, its disruption
 99 leads to restricted ranges and consequent population declines [1,6,7]. The preservation
 100 of the phenomenon of migration requires conservation of both the migratory species
 101 and the habitats along their routes. It also requires a sound understanding of the
 102 factors and processes underlying the degradation and loss of migratory routes and
 103 declines of populations to devise effective strategies for protecting migratory routes,
 104 habitats and populations [1]. Although causes of ungulate migrations are not yet fully
 105 understood [8], the temporal regularity of migrations suggests that they are a response
 106 to seasonal fluctuations in spatial patterns of resource availability and quality [9,10].
 107 Thus, in the Serengeti-Mara ecosystem, rainfall through its effect on food supply and
 108 salinity of drinking surface water has been suggested as a trigger for the northward
 109 migration [11] whereas high nutrient availability on the short grass plains is thought
 110 to attract lactating female wildebeest southwards [12]. This migration results in the
 111 movement of wildebeest from the open, highly nutritive grasslands with low biomass
 112 in the wet season, to wooded grasslands with high biomass of lesser nutritive quality
 113 during the dry season [13].

114

115 We focus on populations of wildebeest sub-species in East Africa because they (1) are
 116 taxonomically closely related, (2) represent some of the most important remaining
 117 large mammal migrations on earth, (3) share similar conservation problems, (4) all
 118 have ranges within and outside protected areas, and (5) have a range of current and
 119 potential pathways to protection. The threats facing wildebeest migrations involve the
 120 interplay of multiple factors and processes [1-3,14]. In the Masailand of Kenya and
 121 Tanzania, ungulate population declines, particularly of wildebeest, are linked to

habitat loss due to land use change or habitat degradation caused mainly by expansion of cultivation [9,15,16]. Illegal hunting might, however, have contributed more to dwindling populations of migratory ungulates in some areas, including the Tarangire-Manyara ecosystem of Tanzania [17-19]. Wildebeest also cause problems for livestock, including competition for forage and transfer of the deadly malignant catarrhal fever virus from wildebeest calves to cattle [20,21]. The type and intensity of these factors and processes vary among migratory species and across their meta-populations or ecosystems. Effective wildlife conservation and protection thus requires clear prioritization of the factors leading to population declines both in the short-and long- term. Integral to this process is reviewing the history, status, trends and threats facing populations of particular migratory species across a range of ecosystems along the entirety of their migratory routes to extract general insights into the threats they face as a basis for developing approaches likely to succeed in conserving their populations and migrations.

We describe and compare the extent of historical migrations of the western (*C.t. mearnsi*) and eastern (*C.t. albojubatus*) white-bearded wildebeest with the current status of these migrations and migratory routes in five ecosystems of East Africa. We evaluate long-term wildebeest population trends, putative drivers of change and their impacts on the critical habitat and migratory ranges of wildebeest in each of the five ecosystems. We suggest potential strategies for conserving these migrations, some of which rank among the Earth's most spectacular remaining terrestrial migrations (Fig 1). Lastly, we evaluate causes of wildebeest population declines and range contraction, including human population expansion, land-use change, poaching, land uses incompatible with wildlife conservation, deficiencies in existing wildlife

policies, institutions and markets in Kenya and Tanzania and suggest conservation strategies to alleviate the population declines.

-----Fig 1 about here -----.

Materials and methods

Study Area

This study covers the five ecosystems in East Africa with migratory wildebeest populations (Fig. 2). These include the Serengeti-Mara, Loita Plains, Athi-Kaputiei Plains, Amboseli Basin, and Tarangire-Manyara ecosystems. Across these five ecosystems, we focus on eight populations of either the western (Serengeti-Mara, Ngorongoro, Loita Plains, Narok County) or the eastern (Athi-Kaputiei, Machakos County, Amboseli, West Kajiado, Tarangire-Manyara) subspecies of the white bearded wildebeest [22]. We consider three (Ngorongoro, Narok County and Machakos) of the eight populations only superficially because they are part of at least one of the other populations considered in detail. We do not consider small, resident wildebeest populations occupying the western corridor in Serengeti and the Loliondo Game Controlled Area (LGCA) in north-eastern Tanzania [14].

The Serengeti-Mara Ecosystem covers about 40,000 km² in Tanzania and Kenya [14,23]. The ecosystem encompasses the Serengeti National Park, Ngorongoro Conservation Area, Maswa, Grumeti, Ikorongo and Kijereshi Game Reserves, Loliondo Game Controlled Area, Ikona and Makao Wildlife Management Areas in

170 Tanzania and the Masai Mara National Reserve and adjoining wildlife conservancies
171 and pastoral ranches in Kenya.

172 The Ngorongoro Conservation Area (NCA, 8,292 km²) is part of the Greater
173 Serengeti-Mara ecosystem. It includes the Ngorongoro Crater (310 km²) and is
174 bordered to the north by the Loliondo Game Controlled Area (4000 km²). Lake
175 Natron Game Controlled Area (LNGCA, 3000 km²) borders the LGCA to the
176 southeast and the NCA to the northeast (Fig. 2).

177

178 The Narok County (17,814 km²) encompasses the Loita Plains and the Masai Mara
179 Ecosystem in Kenya. The Athi-Kaputiei ecosystem (2,200 km²) covers the Nairobi
180 National Park (117 km²) and the adjacent Athi-Kaputiei Plains in Kenya. Machakos
181 County (14,225 km²) is contiguous with the Athi-Kaputiei ecosystem. The Greater
182 Amboseli ecosystem of Kenya (7730.32 km²) covers the Amboseli National Park (392
183 km²) and surrounding dispersal areas on pastoral rangelands, covering some 3,000
184 km² [24-26] . Western Kajiado (11388.54 km²) is bounded by the Greater Amboseli
185 Ecosystem to the East. Both ecosystems are found in Kajiado County of Kenya.

186

187 The Tarangire-Manyara ecosystem of Tanzania covers the Tarangire (2,850 km²) and
188 Lake Manyara (649 km²) National Parks and Manyara Ranch (177 km²), a private
189 conservancy that supports livestock rearing, wildlife conservation and tourism. This
190 ecosystem is adjoined by rangelands managed primarily for cultivation, livestock
191 grazing, legal game hunting, and tourism on community land designated as Open
192 Areas, Game Controlled Areas or Wildlife Management Areas [15]. These include the
193 Simanjiro Plains, the Mkungunero Game Reserve (800 km²) and Lolkisale Game

Controlled Area (1500 km²). Altogether, the range for the migratory wildebeest covers about 35,000 km² [17,27-30].

Human population growth drives sedentarization, expansion of settlements, fences and other land use developments in the study ecosystems [4,5]. These changes promote land use intensification and illegal livestock incursions into protected areas to the detriment of migratory wildebeest [31,32]. In Kenya, human population size increased in Narok County by 673% from 110,100 in 1962 to 850,920 in 2009; in Kajiado County by 905% from 68,400 in 1962 to 687,312 in 2009 and in Machakos County by 247% from 571,600 in 1962 to 1,983,111 in 2009 [33]. Similarly, in Tanzania human population size increased in the Serengeti District by 11.6% from 249,420 in 2012 to 282,080 in 2017 and in Monduli and Simanjiro Districts, containing the Tarangire-Manyara Ecosystem, by 13.5% from 460,775 people in 2012 to 532,939 in 2017 (www.nbs.go.tz).

Across the Serengeti-Mara ecosystem, Narok County, Masai Mara ecosystem and the Loita Plains, rainfall is markedly bimodal and increases steeply along a southeast–northwest gradient, from east to west, south to north and over time [34]. Notably, rainfall increases from 500 mm on the Serengeti Plains to the Southeast to 1400 mm to the north-west of Masai Mara National Reserve. Across the Kajiado County in which the Amboseli, Athi-Kaputiei and Western Kajiado Ecosystems are found, rainfall is low, bimodal and highly variable, and total annual rainfall averages 685 mm (range 327-1576 mm). The short rains fall from November to December ($30.97 \pm 27.85\%$ of the annual total) and the long rains from March to May ($47.5 \pm 15.06\%$ of the annual total). The dry season rains fall during June- September. Rainfall is

markedly variable in space and increases with elevation such that it averages 300 mm/yr in the low-lying Amboseli basin and rises to 1250 mm/yr on the slopes of Mt. Kilimanjaro and Chyulu Hills in the southeast of the County to 800 mm in Nairobi National Park and 971 mm at Ngong hills in the northwest of the County [26]. Rainfall increases from under 500 mm in the extreme southeast of the Athi-Kaputiei Plains to over 800 mm in northern Nairobi Park [35]. In the Tarangire-Manyara ecosystem, rainfall is bimodal and averages 650 mm per annum. The short rains span from October to December and the long rains from March to May. The rains are unreliable and frequently fail, especially the short rains [15]. Land use patterns in the study ecosystems are described comprehensively elsewhere [15,26,34-36].

-----Fig. 2 about here -----.

Historical wildebeest migrations in East Africa

Information on the migratory wildebeest range, routes and status was compiled from literature reviews, colonial-era records, maps, GIS databases, Global Positioning System (GPS) collared wildebeest and interviews with local residents and researchers knowledgeable about the study ecosystems. We reviewed historical records to provide a context for assessing changes in wildebeest migrations in East Africa.

Mapping contemporary wildebeest migratory routes and ranges

To obtain information on contemporary wildebeest movements, we placed GPS collars on 15 wildebeest in the Loita Plains in the Mara Ecosystem in May 2010, 12 in the Athi-Kaputiei Plains and 9 in the Amboseli Basin in October 2010. The collars

were programmed to collect the position of each wildebeest 16 times each day (every hour from 6:00 AM to 6:00 PM and every three hours from 6:00 PM to 6:00 AM) for a 2-year study period. Data are available on Movebank (www.movebank.org)

In the Tarangire-Manyara ecosystem, OIKOS and Tanzania National Parks [37] tracked movements of radio collared wildebeest and zebra and GPS collared elephants (*Loxodonta africana*) during 1995-2002 to establish if they still used the main migratory routes identified earlier [38]. OIKOS also established the presence or absence of migratory routes and assessed wildlife species abundance in the ecosystem during 1995-2002 by interviewing local communities, hunting operators, employees and residents and conducting multiple aerial reconnaissance and systematic reconnaissance flights. Several studies later mapped and analysed land use changes along the migratory corridors [36,39]. We did additional unstructured interviews on the status of the migration routes in the ecosystem during 2006-2007. Our interviews targeted long-term local residents and researchers and were carried out during ground truthing work for imagery analysis on historical land use and cover changes in the ecosystem from 1984 through 2000 to 2006-2007. Local Masai elders who knew the history of the ecosystem well helped with the ground truthing and interviewing local residents about land use and cover changes. The field data form used for our interviews is provided in Table S1.

Wildebeest population trends

Wildebeest population estimates were compiled from aerial surveys conducted in Kenya by the Directorate of Resource Surveys and Remote Sensing (DRSRS) and in Tanzania by the Tanzania Wildlife Research Institute (TAWIRI), Tanzanian Wildlife

Conservation Monitoring Unit (TWCM) and Frankfurt Zoological Society (FZS). The methods used in the aerial surveys and for estimating population size are described in detail elsewhere [33,40-42]. Aerial surveys began in the Athi-Kaputiei ecosystem in 1949 [43], in the Serengeti-Mara ecosystem in 1957 [44-46]), in the Tarangire-Manyara ecosystem in 1964 [27] and in Amboseli in 1973 [47].

Distribution of cultivation and fences

Data on the distribution of agriculture were obtained from the FAO Africover project 2000 [48]. The project mapped land cover for the year 2000 for the whole of East Africa from Landsat images (30 m resolution) and updated the Kenya map in 2008. The map category 'agriculture' was extracted from the Africover data set and clipped according to the study area boundary. In the Athi-Kaputiei ecosystem fences were mapped in 2004 and 2009 by the International Livestock Research Institute (ILRI) and African Wildlife Foundation (AWF) in collaboration with the local communities and local NGO's using hand-held (GPS, with scientific, technical and logistical support provided by ILRI [4,5,35]. Fences, settlements, roads and other infrastructures were similarly mapped with hand held GPS in Amboseli in 2004-2006 [49,50] and in Masai Mara in 1999, 2002 and 2015 [51-54]. A few fences also exist in the ecosystem in Tanzania.

Wildlife conservation initiatives and gaps in policies, institutions and markets in Kenya and Tanzania

We reviewed official records on contemporary wildlife conservation initiatives and identified important gaps in wildlife policies, institutions and markets in Kenya and Tanzania.

Statistical Analysis

Estimates of wildebeest population size for each ecosystem were obtained using Jolly's method II for transects of unequal lengths [55] and related to the year of survey using negative binomial regression models with linear and quadratic polynomial terms and serial autocorrelation in the counts accounted for using the first-order autoregressive model. Selection between the linear and quadratic models was based on the Akaike Information Criterion [56]. The models were fitted using the SAS GLIMMIX procedure [57]. Temporal trends in wildebeest population size were modeled using a semiparametric generalized linear mixed model with a negative binomial error distribution and a log link function in SAS GLIMMIX procedure [33]. The percentage change in population size between the start and end dates of the surveys was estimated for each ecosystem. For some ecosystems predicted population size for two to three consecutive surveys were averaged and used to compute the averages to minimize the effect of stochastic noise due to small sample size or areal coverage.

Results

Historic and contemporary migratory routes

The seasonal migration of the white-bearded wildebeest and zebra (*Equus quagga burchelli*) from the Serengeti Plains in Tanzania to Masai Mara in Kenya [14,46] is called the southern migration. It differs from the northern migration that involves seasonal movements of wildebeest, zebra and Thomson's gazelle (*Gazella thomsoni*) between the Loita Plains and Masai Mara National Reserve within the Narok County of Kenya [9,10,16,58]. The wildebeest involved in the northern migration form the bulk of the Narok County population.

The migration ranges, routes and population trends of the Serengeti-Mara wildebeest have been extensively studied (e.g., [59,60], Table S2). After rinderpest killed about 95% of this population between 1890 and 1892, it remained low till 1962. The population increased from 263,362 in 1961 to 483,292 in 1967 following veterinary removal of rinderpest in wildebeest in 1962 and again from 1967 to 1.4 million in 1977, coincident with an increase in the dry season rainfall [61,62].

The migration pattern of the Serengeti-Mara wildebeest in the 1940s and 1950s was different from what it is today. Then, wildebeest migrated periodically between the Kenya's Loita Plains and Tanzania [63,64]. Heavy harvesting of wildebeest (plus zebra and other species) in Narok County during World War II to provide meat for labour and prisoners of war and reduce competition with Masai livestock until 1947 [64 reduced their population. Even so, wildebeest population on the Loita Plains numbered about 50,000-100,000 individuals prior to 1947 [45]. Thereafter, the

333 population further declined drastically because the Loita Plains were opened up to
 334 uncontrolled commercial meat-hunters for a short period after World War II [67].
 335 Subsequently, the Mara population (including the Loita Plains) numbered about
 336 15000 by 1958 [45] and 17,817 by 1961 [65].
 337
 338 The Mara wildebeest [45,46,58,66] were formerly more numerous and their
 339 distribution extended far beyond their contemporary range in the Mara region of
 340 Narok County. But major land use and cover changes have progressively degraded
 341 and reduced the historical wildebeest habitats [9,68].
 342
 343 Wildebeest occupied parts of the Rift Valley extending from the Tanzania border
 344 north to Kedong Valley in Kenya and in the early 1900s, to the eastern and southern
 345 shores of Lake Naivasha [46, 69-75]. Thus, in 1902, Meinertzhagen [74] recorded
 346 wildebeest on the flats east of Lake Naivasha. Moreover, based on annual hunting
 347 returns, wildebeest were shot in Naivasha in 1906-07 [70] and in the Rift Valley in
 348 1909-10 [71]. The Mosiro Plateau in Narok County was the probable link between the
 349 Rift and Serengeti-Mara wildebeests when the Plateau landscape was still open and
 350 had tall grass, few gullies and bushes [46]. Livestock overgrazing degraded the
 351 plateau making it impassable for wildebeest by early 1960s [46]. Part of this
 352 wildebeest population also occupied the centre and west of the Rift Valley, near Mt.
 353 Suswa, south of Naivasha in 1909 and 1910 [73]. Their distribution extended to the
 354 East Rift Wall, near Kijabe in central Kenya [76]. But, all the wildebeest populations
 355 in the northern area of the Rift Valley in Kenya became extinct before 1962 because
 356 of hunting and fencing of ranches around Lake Naivasha and in the Rift neighbouring
 357 Suswa area [46]. Wildebeest were later re-introduced on the Crescent Island in Lake

Naivasha and slowly spread, or were physically moved, to other surrounding areas. Their population increased steeply in Nakuru Conservancy in the Nakuru-Naivasha region of Kenya during 1996-2015 [77]. Wildebeest were also found in several other parts of Kenya in earlier years where they have since been exterminated. In particular, returns of game animals shot on license in 1909-10, show wildebeest were found on the Mau Plateau in Narok County, the Kisii region in Southwestern Kenya, Laikipia and North Uaso Nyiro in Central Kenya, Makindu and Voi in southeastern Kenya [71].

A small wildebeest population occurs in the Ngorongoro Conservation Area in Tanzania [78,79]. It concentrates in the Crater in the dry season but disperses to areas outside the Crater in the wet season, including the Lake Natron Game Controlled Area. During the early dry season wildebeest sometimes move east from the Serengeti short grass plains into Ngorongoro Crater/Conservation Area and leave at the onset of the rains but a smaller residual population remains in the Crater during the rainy season [46,63]. Wildebeest from the NCA and the Serengeti short grass plains also migrate to the south-eastern part of the LGCA.

The Athi-Kaputiei, Amboseli, Western Kajiado and Machakos County wildebeest populations were historically part of a single large migratory population that used to range over most of the present day Kajiado County in Kenya until the 1960s before it split up into three rather distinct populations [22,26,35,76]. The western Kajiado population is currently non-migratory. The Athi-Kaputiei wildebeest population uses the Nairobi National Park during the dry season due to its reliable water supply and abundant grass and move to calve on the pastoral lands to the southeast of the park

during the wet season [4,5,35,80-83]. The Athi-Kaputiei wildebeest population was centred on the Athi-Kaputiei Plains in the wet season prior to the 1920s. Historically, some Athi-Kaputiei wildebeest may have migrated south to the Amboseli ecosystem [46,67,76]. They intermingled with the then larger Amboseli population centred on the Amboseli Plains north of Kilimanjaro in the wet seasons. Both populations were migratory and moved to water in the hills and woodlands in the dry seasons and returned to the plains in the wet seasons. In very dry periods many wildebeest from both populations moved northeast and south, including into Tanzania [46].

The Athi-Kaputiei wildebeest moved north, east, and south in the wet season but spent the dry season on the Athi-Kaputiei Plains until at least 1927 [46,67,76]. The Athi-Kaputiei wildebeest population migrated as far north as the Thika River in the dry season but only few went beyond this point [46,67,76]. A resident wildebeest population north of the Thika River and another near Juja, both northeast of Nairobi in Kenya [46,75,76] went extinct. The Athi-Kaputiei wildebeest population also migrated as far north as Muranga (Fort Hall) and the Yatta Plateau, south of the Tana River [63]. De Beaton [84] recorded animal movements from the Nairobi National Park westwards towards the Ngong Hills and to the south. The Nairobi-Mombasa Road and later the park fence bordering this road interrupted the northward migration of wildlife to Nairobi, Ruiru-Thika and Ol Donyo Sapuk in the dry season. When a fence was first erected around Nairobi in the early 1900s, it killed many animals, including wildebeest [85]. A fence constructed in 1967 along the eastern side of the Ngong Hills and South of the Kiserian River and that joined the south-western corner of the Nairobi Park fence, further disrupted the dry-season migration of wildlife to the Ngong Hills [81,86].

408

409 Large areas of the Athi-Kaputiei Plains were ploughed and planted with wheat to
 410 contribute to war time food production during World War II. The wheat attracted
 411 large wildlife herds, including wildebeest, which were shot as part of crop protection
 412 [87]. This population dispersed periodically to the adjoining Machakos County,
 413 especially during droughts, but more recently, due to displacement by extreme land
 414 use changes and developments in the Athi-Kaputiei Ecosystem [5,33,35,83,88].

415

416 The migration and ranges of the Amboseli and West Kajiado wildebeest populations
 417 are described by several authors [22,25,46,63,67,75,76]. Occasional old bulls moved
 418 from the Rift near Lake Magadi area in western Kajiado to Nairobi area in the 1920s
 419 [76]. Wildlife, especially wildebeest and zebra, were also harvested in large numbers
 420 in Kajiado County (Amboseli Ecosystem) during World War II to provide meat for
 421 labour and prisoners of war; free meat for the Kamba people because of famine
 422 caused by severe drought; and after the end of the war to reduce competition with
 423 Masai livestock for forage [64,67].

424

425 In the Tarangire-Manyara Ecosystem, the migratory wildebeest occupy the Tarangire
 426 National Park in the dry season but disperse to their wet season ranges and calving
 427 grounds on the Simanjiro Plains, the Mkungunero Game Reserve, Lolkisale Game
 428 Controlled Area, Manyara Ranch, Lake Manyara National Park and adjacent game
 429 controlled areas (used mainly as hunting areas) in the wet season. OIKOS and
 430 TANAPA [37] confirmed that nine main migratory routes that Lamprey [38] had
 431 identified earlier in the Tarangire-Manyara ecosystem, were still being used during
 432 1995-2002.

433

434 Wildebeest migrated from within protected areas in the dry season to dispersal areas
435 outside conservation areas in the wet season in all the ecosystems but the Serengeti-
436 Mara ecosystem where the migration occurred mostly within the protected areas (Fig.
437 2). Wildebeest migration has discontinued altogether in parts of the ecosystems, and
438 reduced along a number of historical routes (Fig. 3).

439

440 -----Fig. 3 about here -----.

441

442 Notably, the decline and discontinuation of migration happened in four out of the five
443 ecosystems where wildebeest migrated outside protected areas. No discontinuation of
444 migration is reported from the Serengeti-Mara ecosystem where wildebeest migrates
445 almost entirely within protected areas. Discontinued or currently less intensively used
446 migration routes overlapped with agricultural and settlement expansion in the Mara
447 and Tarangire-Manyara ecosystems and fences, settlements and roads in the Athi-
448 Kaputiei Plains (Fig. 3). Fig. 3 does not include settlements, which is another main
449 cause of change to the migratory routes in the study ecosystems.

450

451 -----Fig 4 about here-----.

452

453 Movement data ($n = 279,718$ fixes) from the GPS collared wildebeest showed the
454 migration routes during 2010-2013 (Fig. 4). Several features of the wildebeest
455 movements and space use are noteworthy. Wildebeest primarily used habitats outside
456 of the protected areas in the Mara, Athi-Kaputiei and Amboseli ecosystems ($> 87\%$ of

the 279,718 fixes). This emphasizes the importance of pastoral lands and community-based conservation to the protection of the three wildebeest populations. In particular, the Loita Plains wildebeest heavily used the wildlife conservancies adjoining the Masai Mara National Reserve to the north. Hence, when both the reserve and conservancies are considered, 73.4% (85,194 of 116,061 fixes) of the Loita Plains wildebeest locations fell within the conservation area boundaries. Further, one wildebeest collared in Loita Plains moved south through the LGCA to the NCA in Tanzania, covering a total of 205.4 km from its initial collaring location (Fig. 4b). This route approximates the historical migration route of the Loita wildebeest up to the 1950s [63]. This reinforces the critical importance of LGCA to Serengeti-Mara, Loita and Ngorongoro wildebeest migrations and to the ecological integrity of the Greater Serengeti-Mara ecosystem.

The Nairobi-Namanga tarmac road, bisecting the wet season range of the Athi-Kaputiei wildebeest, has split the population into two distinct sub-populations, concentrated on the eastern and western sides of the road (Fig. 4b). Collaring locations and direct field observations showed that no collared wildebeest crossed the tarmac road during the 2010-2013 study period. Lastly, the Amboseli wildebeest population also moved widely, including into the adjoining Longido District in Tanzania, reflecting the historical migration routes for this population [46,76]. One wildebeest collared in the Amboseli Basin travelled 6,197.8 km over 728 days during the study period. Further details on the collared wildebeest movements can be found in Stabach [88].

Wildebeest population trends

The Serengeti-Mara wildebeest population grew steadily from 190,000 in 1957 following the veterinary eradication of rinderpest in cattle in 1962 [90], until 1977 when it stabilized with one noticeable decline during 1993, when a severe drought reduced the population from around 1.2 million to less than 900,000 animals [91] (Fig. 5a). The population has since then recovered and stabilized at around 1.3 million animals [14] though the more recent population size estimates suggest some slight upward trend (Fig. 5a). The estimated population size and standard errors and other details of the aerial surveys for all the eight study ecosystems are provided in S1-S5 Datas.

The Loita Plains wildebeest population declined steadily from about 123,930 animals in 1977-1978 to around 19,650 animals by January 2016 (Fig. 4b), a decrease of 80.9%. This decline was highly significant (Table 1). The population of the Serengeti migrants coming to the Mara ecosystem in the dry season (July-October) similarly decreased by 73.4% from 587,500 in July-August 1979 to 157,124 animals in November 2016. The dramatic decline was also evident for the Narok County wildebeest population (Table 1, Fig. 5c, S2 Data).

The Athi-Kaputiei wildebeest population suffered a 95% decline in numbers from over 26,800 animals in 1977-1978 to less than 10,000 by the mid-1990s and under 3,000 animals in 2007-2014. The decline of this population has been much more dramatic in recent decades, leading to a virtual collapse of the migration (Fig. 5d). The catastrophic decline is highly significant (Table 1, S3 Data). A recent 1298% increase in Machakos County population, coincident with the decrease in the Athi-

Kaputiei population (Fig. 5e), is not statistically significant likely because of a large variance in the population estimates (Table 1, S4 Data). This strongly suggests that some wildebeest migrated from the Athi-Kaputiei rather than died.

The migratory wildebeest population in the Amboseli ecosystem also declined by 84.5% from about 16,290 animals in 1977-1979 to 2,375 by 2010-2014 (Fig. 5f). The population fluctuated between 16,290 and 20,000 individuals and increased to 33,000-37,000 individuals during 1978-1986 and fell to 16,779 animals by 2007. The population declined to under 5,000 animals in 2010 following a severe drought in 2008-2009 (Fig. 5f) and has not recovered ever since. This decline is highly statistically significant (Table 1). The non-migratory wildebeest population in West Kajiado decreased by 44% from 5,700 animals in 1977-1979 to 3200 animals in 2010-2014 but this decrease is not statistically significant likely due to large variances in population size estimates (Table 1, Fig. 5g, S5 Data).

The Tarangire-Manyara population first increased from an estimated 24,399 animals in 1987 to 48,783 animals in 1990. Thereafter the population fell precipitously to 13,603 animals by 2016 without signs of recovery (Fig. 5h). This extreme population decline is statistically significant (Table 1) despite the large variances in the population estimates (S5 Data).

-----Fig. 5 about here-----

529 **Table 1.** Results of the regression of wildebeest population size on year of survey.

530 NDF and DDF are the numerator and denominator degrees of freedom, respectively.

Region	Number of surveys	Intercept	Linear Slope	Quadratic slope	NDF	DDF	F	P > F
Serengeti-Mara	22	-4785.20	4.8075	-0.00120	^a 1	19	39.84	<0.0001
					^b 1	19	39.41	<0.0001
Mara	21	13272	-13.2237	0.0033	^a 1	18	11.40	0.0034
					^b 1	18	11.26	0.0035
Narok County	17	62.966	-0.0261		1	15	7.52	0.0151
Athi-Kaputiei	25	11.68	-0.00024		1	23	18.74	0.0002
Machakos	5	-94.57	0.0511		1	3	1.53	0.3046
Amboseli	21	-15810	15.8951	-0.0023	^a 1	18	13.04	0.0020
				-0.00399	^b 1	18	13.11	0.0020
West Kajiado	18	42.56	-0.01715		1	16	2.39	0.1417
Tarangire	8	156.64	-0.07808		1	5	11.16	0.0205

531 ^aLinear slope, ^bQuadratic slope

532

533 Discussion

534 Wildebeest movements and migratory routes

535 Animal movement depends on individual fitness and is essential for accessing
536 favoured resources, finding potential mates and escaping deteriorating habitat
537 conditions [92]. As expected, the GPS collared wildebeest moved more, in virtually
538 all measured aspects, in Amboseli, the least productive and least anthropogenically
539 disturbed of the three Kenyan ecosystems, than in the Loita Plains and Athi-Kaputiei.
540 The productivity of Amboseli grasslands has reduced even further in recent years

[93,94], apparently forcing wildebeest to move over larger areas in search of food in the dry season [95]. Wildebeest surprisingly moved less in Athi-Kaputiei than in either the Amboseli or Loita Plains even though the Loita Plains had the greatest availability of resources of the three landscapes. This is unexpected even if the wildebeest decline in the Athi-Kaputiei has reduced intraspecific competition and the need to move to locate resources. High livestock density likely heightens interspecific competition with wildebeest for resources and thus could force wildebeest to move more in Athi-Kaputiei. The reduced wildebeest movements in the Athi-Kaputiei landscape therefore reflect its high degree of anthropogenic disturbance and truncation [5], preventing needed further movement [88]. It follows that resource availability and anthropogenic disturbance determine wildebeest movements. Consequently, because wildebeest occur primarily outside protected areas, except in the Serengeti-Mara, controlling the rate and type of anthropogenic change in these areas is crucial to maintaining the long-term viability of their populations and migrations.

Wildebeest population declines

Migratory wildebeest population size and their routes declined in all the five ecosystems except the Serengeti-Mara. The declines are related to expansion of agriculture, settlements, fences and roads that progressively occlude wildebeest grazing resources and migratory routes (Table 2). Even though it was not possible to formally test if these processes caused the declines, literature review, interviews and collared wildebeest movements, suggest that they are all important. In all the four ecosystems where they are declining, agricultural encroachment excludes wildebeest from part of their seasonal ranges. Notably, irrigated agriculture encroached the

swamps that ring the base of Mt. Kilimanjaro, denying wildebeest access to their critical dry season dispersal areas in Amboseli [24,25,96]. Settlements also interfere with wildebeest movements in the Mara [31,34,52], Tarangire-Manyara [15,18] and Athi-Kaputiei [4,5,35] by blocking their migratory routes and access to resources. Further, although wildebeest avoid anthropogenic disturbances [97], they are attracted to short grass created by livestock grazing outside protected areas on pastoral lands with moderate densities of pastoral settlement and livestock [21,98].

Land fragmentation through fencing, roads and settlements primarily exclude wildebeest from their grazing ranges in the Athi-Kaputiei ecosystem [5] (Table 2). In Kitengela, a major part of the Athi-Kaputiei Plains adjoining Nairobi National Park, fenced land parcels have spread throughout the range of wildlife and movements of people, livestock, dogs and vehicles harass wildlife [4,5,35]. Fences impede wildebeest movements between the Nairobi Park and the Athi-Kaputiei Plains [4,5,35]. Similarly, the Nairobi-Namanga road has effectively truncated the ecosystem, splitting the Athi-Kaputiei population into two separate sub-populations [89]. The Athi-Kaputiei landscape is also fragmented and degraded by large, un-rehabilitated mines, mining waste, unregulated development, commercial charcoal burning and sand harvesting, all of which restrict wildebeest habitats and obstruct their migratory routes. Invasive weeds are also spreading in the rangelands and at abandoned settlement sites in Athi-Kaputiei, degrading wildebeest habitats [5,35,51,99]. Fences [100,101] are also increasing rapidly in the Mara, including in the Loita Plains, following land subdivision and privatization of land ownership.

The loss of connectivity restricts the mobility and flexibility of migratory wildebeest, especially during droughts when heavy mortality can result where wildebeest access to water and food is blocked [102-104]. The risk of outbreaks of zoonotic diseases and population declines can also increase if ungulate migrations are curtailed by degraded habitats yet climate change increases the frequency and severity of droughts [105]. Climate change may amplify the frequency of outbreaks of zoonotic diseases by modifying host and vector population characteristics that control pathogen transmission, including concentration in key resource areas, population density, prevalence of infection by zoonotic pathogens, and the pathogen load in individual hosts and vectors [106,107]. Also, calving wildebeest transmit bovine malignant catarrhal fever (BMCF) virus to livestock where the two species co-occur, causing livestock losses [20]. The risk of transmitting the BMCF virus is elevated where habitat loss and degradation force livestock and wildebeest to use the same areas.

Another leading cause of wildebeest decline is poaching, which removes 6-10% of the Serengeti-Mara wildebeest annually [108,109]. Poaching is also common in the other ecosystems, including the Mara [34] and Athi-Kaputiei [35]. The status and threats facing the five ecosystems with migratory wildebeest populations in East Africa are summarized in Table S2.

Table 2. Summary of the processes likely associated with the declining migratory wildebeest populations and patterns in the East African rangelands.

Processes	*Serengeti- Mara	Masai Mara	Athi- Kaputiei	Greater Amboseli	Tarangire- Manyara
-----------	---------------------	---------------	-------------------	---------------------	-----------------------

Direct interferences/causes					
Agricultural encroachment	-	+++	+	+++	+++
Fencing	-	++	+++	++	-
Settlements	+	++	+++	++	++
Urbanization	-	+	+++	-	-
Roads & Infrastructural developments	+	++	+++	+	++
Poaching by increasing human populations	++	+	++	+	++
Competition with livestock for forage, water and space	+	+++	+++	+++	+++
Declining drinking water supply and quality	+	++	+++	++	+
Drivers					
Human population increase	+	++	+++	++	+++
Land tenure change	+	++	+++	+++	+
Land subdivision	-	+++	+++	++	++
Settlement policies	++		+++		+++
Wildlife conservation and management policies	+++	+++	+++	+++	+++
Wildlife management institutions	-	+++	+++	++	++
Wildlife markets or benefits to landowners	-	+++	+++	+++	+

‡+++ High importance; ++ Important; + less importance; - not important. Source: Interviews with resident researchers [4,5,15,18,111].

Additional factors that adversely affect access of migratory wildebeest to critical habitats, food and water include human population expansion, land subdivision and privatization of land tenure, development of urban centres and intensification of land use following sedentarization of formerly semi-nomadic pastoralists [4,26,31,33-35].

Why is the Serengeti-Mara wildebeest population stable while the other populations are declining? Moreover, given that the Serengeti-Mara ecosystem protects nearly 1.5 million wildebeest, why should we worry about conserving the other smaller wildebeest populations? First, the Serengeti-Mara populations are not declining because over 80% of the wildebeest live in the large and relatively well-protected ecosystem. Second, it is important to conserve the smaller populations in other areas for at least three reasons. a) Some of the other areas support populations of wildebeest belonging to a different subspecies from that found in Serengeti-Mara. b) Migratory wildebeest provide important ecosystem services, even at low densities, such as promoting calf survival among other ungulate species by reducing predation pressure when present in an area [112]. c) Wildebeest migrations are a magnificent spectacle and thus can provide significant tourism revenue opportunities in specific areas.

Land-use change and poaching as causes of wildebeest population declines

Land use change, particularly expansion of agriculture, settlements and fences and commercial charcoal production linked to human population growth degrade and reduce wildlife habitats [5,26]. In the Athi-Kaputiei ecosystem, expansion of the neighbouring Nairobi Metropolis, urbanization in the ecosystem plus relatively lower land prices compared with Nairobi, strongly drive land use change [114]. Development of new industries, businesses and infrastructure attract more people from Nairobi and elsewhere to the Athi-Kaputiei [4,83]. In Amboseli and Western Kajiado, commercial charcoal production is causing widespread deforestation of wildlife habitats [115].

Agriculture, particularly large-scale commercial cultivation, is a leading cause of habitat loss for the migratory wildebeest. Previously, mainly outsiders practiced cropped agriculture in the study ecosystems, but the Masai have recently started cultivating next to their settlements [116,117]. Widespread adoption of subsistence agriculture in small plots right around a household's compound can threaten wildebeest populations migrating outside protected areas [118]. Remarkably, nearly 500 km² of natural vegetation in the Loita Plains were converted to wheat farms and other uses between 1975 and 1995 [67] and even more has been converted in recent years [119,120]. In Tanzania, people moved into, and cultivated for several years, parts of Game Controlled Areas or Open Areas, such as the LGCA, which had functioned much like game reserves in the past, interfering with wildlife migrations.

In the Tarangire-Manyara ecosystem, about 710 km² of land was converted from rangelands to farms between 1984 and 2000 [15], cutting-off large portions of forage and dispersal areas and blocking routes traditionally used by migratory wildebeest. Villagization promoted by government settlement policies in Tanzania is another key driver of land conversion to agriculture in the Tarangire-Manyara ecosystem [15] and western Serengeti [14].

Wildebeest prefer land suitable for agriculture [121] and therefore face a high risk of displacement by agriculture and competition with livestock for space, forage and water in pastoral lands. Such high potential lands tend to generate higher economic returns from cropping than from livestock or conservation [114,119]. Land users are thus likely to opt for cultivation rather than conservation thereby accentuating

encroachment of agriculture and wildebeest population declines. But once land is cultivated, it is difficult to restore to its former rangeland status where returns from agriculture overwhelm those from either livestock or wildlife [119]. Even where wildlife tourism benefits are competitive with those from agriculture, the benefits often accrue to the rich so that the poor land owners, who bear the burden of supporting wildlife on their lands, typically receive meagre benefits [122]. This calls for schemes for more equitable sharing of wildlife benefits [116].

Land tenure change from group ranches to private ownership is another important driver of land use change in Masailand in Kenya [116]. The land sub-divisions and individualization of tenure associated with fencing in Masai Mara, Athi-Kaputiei and Amboseli ecosystems amplify habitat fragmentation and interfere with the migratory wildebeest [4,5,123].

Poaching is associated with increasing human population size and resource use intensity [108-110]. On commercial wheat farms in the Mara, poaching is very common (R. Lamprey, pers comm), especially far from pastoral settlements, because pastoralists often discourage poaching. Poaching is also common inside the protected areas in the Mara and Serengeti [34,109,110]. In the Athi-Kaputiei, poachers killed many wildebeest by running them up against fences [35].

Ways to make land use compatible with wildlife conservation

Human population explosion, unplanned urbanization, settlements, cultivation and other developments pose unprecedented challenges to conservation and maintenance of migrations as the spaces available for wildlife and their habitats shrink, leading to

population declines. It is thus important to conserve spatially extensive migratory systems while balancing human and wildlife needs. In Kenya, wildlife conservancies are expanding conservation areas for wildlife beyond the state-owned parks and reserves onto land owned privately by local communities or individuals who benefit by receiving land rents and job opportunities [34,113].

It is primarily tourism income that pays for conservancy land leases and management in Kenya. Thus, the success of the common conservancy model in Kenya is contingent upon sustainable wildlife tourism making it worthwhile for landowners to allow conservancies to be set up on their lands. This conservancy model can thus only be viable in areas with low tourism potential if tourism revenue is supplemented with other revenue streams.

Nevertheless, wildebeest can and do benefit from community-based wildlife conservation endeavours where wildlife conservancies have been established on private and communal rangelands, including in areas of high rainfall [14,25,124,125]. By 2015, 178 wildlife or mixed livestock-wildlife conservancies had been established across Kenya [51] and new ones continue to be established on private and communal lands in Masai Mara, Amboseli, Athi-Kaputiei and Machakos (Tables S3 and S4), Naivasha-Nakuru and other parts of Kenya [26,34,77,125,126]. The total area of wildlife conservancies and ranches in Kenya's rangeland counties by 2017 was 54,265 km² of which Narok, Kajiado and Machakos counties that support wildebeest populations had set aside 2,219, 2,837 and 463 km², respectively (KWCA, Unpublished data, <https://kwcakenya.com/>).

What makes the conservancies so popular with local communities is that they also protect land rights; create jobs; provide income to communities through tourism; and provide increased security for people, livestock and wildlife [51]. Conservancies are crucial for wildlife conservation because all state protected areas cover only about 10% of Kenya's land surface (an additional 10.7% is in conservancies, benefiting close to 700,000 people nationally) and 70% of these areas are found in the rangelands of Kenya. Moreover, about 65% of Kenya's wildlife are found outside the protected areas [127]. As limited public land constrains expansion of public protected areas, the private and communal conservancies are crucial for expanding the space for wildlife in Kenya. The conservancies are promoting positive attitudes towards wildlife and restoration of degraded rangelands by regulating livestock grazing, restricting settlements and other developments. They act as buffers for parks and reserves, besides offering increased protection to wildlife, enabling many wildlife species to increase within conservancies [77,126,128].

Effective wildlife conservation would require permanent conservancies, land purchases or conservation easements on land used by wildlife. In the Kenya wildlife conservancies, landowners typically amalgamate adjacent individual plots to create large, viable game viewing areas. They then broker land lease agreements with a coalition of commercial tourism operators under institutional arrangements modelled in the form of payments for ecosystem services [125]. There is a strong interest in this wildlife conservancy model in Kenya. Thus, starting with only two conservancies in 2005-2006 covering 145.76 km², there were eight conservancies covering about 1000 km² by 2010 [125] and 10 conservancies by 2016 (Table S3). The Mara conservancies covered 1355 km² by 2018 and are expanding rapidly. The development of the Mara

conservancies has helped partially unblock the movements of migratory wildebeest between the Mara Reserve and the Loita Plains.

In certain areas, such as the Athi-Kaputiei, land owners are paid conservation land lease fees since 2000 to keep land open for use by wildlife and livestock, not building fences and for collecting poachers' snares [35,129-131]. The cost of financing such land leases over large areas year after year would, however, require creating conservancies able to maintain viable conservation enterprises, such as a vibrant tourism industry, to ensure their long-term sustainability. The benefits derived from such enterprises would be an important incentive for the landowners to continue keeping their land open for use by wildlife and desisting from other uses incongruent with conservation. The changes taking place in Athi-Kaputiei are, however, so dramatic and fast that unless these conservation efforts are undertaken immediately, the opportunity to save even the very few remaining and most critical portions of this once magnificent ecosystem, is highly likely to be lost for good.

In Tanzania, various conservation initiatives have been launched to protect the remaining migratory routes and dispersal ranges beyond the borders of protected areas. These include reducing illegal hunting and livestock grazing in Manyara Ranch, recently converted to a private Conservancy. Provision of artificial water holes in the Manyara Ranch Conservancy keeps migratory wildebeest and zebra in the vicinity of the Conservancy until late in the dry season. On the communal grazing lands, initiatives have been launched to enhance the wildlife benefits going to the local communities. In Simanjiro Plains, hunting companies, tour operators and

conservation organizations have teamed up together to pay for conservation land lease fees to community members to refrain from farming or expanding settlements into critical areas of communal grazing lands [132]. Certificate of Customary Right of Occupancy (CCRO) agreements are also being used to protect grazing ranges for wildlife and pastoral livestock, including in areas neighbouring migratory corridors. One such CCRO was established in Selela Village situated north of Manyara and includes an important but narrow corridor [133]. Other initiatives include establishment of Wildlife Management Areas (WMAs) two of which were recently established to the north and west of Tarangire National Park [18]. The WMAs not only protect communal land but also reduce the incentive for poaching by distributing tourism revenue to the local communities. The Tarangire National Park and the Tanzania Wildlife Authority are also supporting community game rangers to intensify anti-poaching patrols in the WMAs and Manyara Ranch Conservancy and among villages in the Simanjiro Plains.

Wildlife conservation initiatives and gaps in wildlife policies, institutions and markets in Kenya and Tanzania

What else can be done to stop the declines and allow migratory wildebeest access to at least the few remaining critical portions of their former habitats? A significant challenge to wildlife conservation in East Africa remains incoherent government development policies that promote incompatible land uses, such as promoting cultivation in pastoral rangelands occupied by wildlife to combat food insecurity while also promoting wildlife-based tourism in the same areas. Such policies should be harmonised to minimize the adverse impacts on wildlife conservation of incongruent land uses in pastoral rangelands. Another weakness of the wildlife policy

in Kenya is that the state owns all wildlife whereas land owners in the rangelands do not have access to or user rights over wildlife. The land owners do not get any compensation for the opportunity cost of supporting wildlife on their private lands nor for wildlife damage to their private property, thus fuelling indifference or hostility towards wildlife. There is also no public institution specifically charged with conserving and managing wildlife on the private lands. Although these shortcomings are well documented [114,134] and have partly been addressed by the Wildlife Conservation and Management Act 2013 [135] and the National Wildlife Strategy [136], the Act should be fully implemented to address these glaring policy, institutional and market deficiencies.

In Tanzania, several national initiatives are being undertaken to restructure the institutions that manage the wildlife sector in order to contain a spiralling poaching crisis. Key among these is the dissolution of the former Wildlife Division (WD) that used to manage all the Game Reserves and Game Controlled Areas, including overseeing all wildlife in village lands (i.e., WMAs), and its reconstitution as the Tanzania Wildlife Authority (TAWA) in October 2015. TAWA is empowered and better funded compared with its predecessor, the WD, to improve the management of the wildlife areas under its jurisdiction.

The second is the re-organization of the entire wildlife sector in the country into paramilitary style organizations to intensify the fight against run-away poaching in protected and unprotected areas, most especially in game reserves. Because many game reserves and game controlled areas share open borders with national parks, wildlife population declines due to poaching are occurring even inside the national parks. But, to be successful in curbing poaching, these efforts should be accompanied with

enhanced economic incentives to communities neighbouring wildlife areas or sharing land with wildlife to discourage poaching and destruction of wildlife habitats. Tanzania is also working on expanding the Serengeti National Park by adding to it about 1500 km² from the Loliondo Game Controlled Area to the east and extending the western side of the park to reach the shores of Lake Victoria.

A major initiative in both Tanzania and Kenya is the development of national policies on wildlife corridors, dispersal areas, buffer zones and migratory routes to promote habitat connectivity [137,138]. Regional initiatives linking the two countries are, however, needed to foster close cooperation between Kenya and Tanzania in conserving the trans-boundary wildebeest migrations and implementing regional and international conservation conventions and treaties. Such initiatives should include harmonization of policies, legal and regulatory frameworks for the conservation of wildlife and other species involved in trans-boundary migrations.

Conclusions

Migratory wildebeest populations in four out of five key ecosystems in East Africa are under severe threats and two populations are on their way to total collapse if the trends are left to continue unabated. Such collapse in migratory wildlife population in East Africa has been documented for zebra and Thomson's gazelle populations that used to migrate between Lakes Nakuru and Elementaita and Baringo regions of Kenya [76,77,139] that went extinct because of fences and uncontrolled shooting [85]. The migration of the Athi-Kaputiei wildebeest to Nairobi National Park had also virtually collapsed by 2011 [35]. Recent surveys in the park show that the wildebeest

involved in this migration remained under 350 animals from 2012 to 2015 [5].

Agricultural encroachment, settlements, poaching, roads and fencing are the major proximate threats responsible for the extreme wildebeest losses and degradation of their habitats as they directly kill, displace, or reduce wildebeest access to forage, water and calving areas. The fundamental causes of wildebeest population declines seem to be expanding unplanned land use developments driven by human population growth; poaching, policy, institutional and market deficiencies. Consequently, the Kenyan and Tanzanian governments need to strongly promote and lead the conservation of the remaining key wildebeest habitats, migration corridors and populations to ensure their continued access to grazing resources in these rangelands. More wildlife conservancies or management areas should be established to protect migratory routes or corridors, buffer zones, dispersal areas and calving grounds for the species. Land use and development planning should be enhanced and gaps in wildlife policies, institutions and markets addressed. Where migration occurs across international boundaries, such as in the Serengeti-Mara, Loita Plains and Amboseli ecosystems, wildlife policies, land use plans, conservation and management goals should be harmonized to ensure the long-term survival of migratory species and the sustainability of the rangelands upon which they depend. All areas currently under protection should ideally have binding legal restrictions on future developments to minimize their vulnerability to future changes. The various conservation initiatives should be coordinated spatially and across bureaucratic lines to enhance their effectiveness.

Acknowledgements

We thank the Tanzania Wildlife Research Institute (TAWIRI) and the Directorate of Resource Surveys and Remote Sensing of Kenya (DRSRS) for permission to use their aerial survey data. JOO was supported by a grant from the German National Research Foundation (DFG; Grant # OG 83/1-1). MYS was supported by the Pathways to Resilience in Semi-Arid Economies (PRISE, Project # 107643-001). This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 641918 through the AfricanBioservices Project. National Science Foundation (NSF) DEB Grant #0919383) supported this work through the project: Wildebeest Forage Acquisition in Fragmented Landscapes under Variable Climates.

References

1. Bolger DT, Newmark WD, Morrison TA, Doak DF (2008) The need for integrative approaches to understand and conserve migratory ungulates. *Ecology Letters* 11: 63-77.
2. Harris G, Thirgood S, Hopcraft JGC, Crooms J, Berger J (2009) Global decline in aggregated migrations of large terrestrial mammals. *Endangered Species Research* 7: 55-76.
3. Berger J (2004) The Last Mile: How to sustain long-distance migrations in mammals. *Conservation Biology* 18: 320-331.
4. Reid RS, Gichohi H, Said MY, Nkedianye D, Ogutu JO, Kshatriya M, Kristjanson P, Kifugo SC, Agatsiva JL, Andanje SA, Bagine R (2008) Fragmentation of a peri-

888 urban savanna, Athi-Kaputiei plains, Kenya. In: Fragmentation in Semi-arid and Arid
889 Landscapes; Consequences for Human and Natural Systems (eds KA Galvin, RS Reid,
890 RH Behnke, NT Hobbs), pp. 195-224. Springer.

891

892 **5.** Said MY, Ogutu JO, Kifugo SC, Makui O, Reid RS, de Leeuw J (2016) Effects of
893 extreme land fragmentation on wildlife and livestock population abundance and
894 distribution. *Journal of Nature Conservation* 34: 151-164.

895

896 **6.** Fryxell JM, Greever J, Sinclair ARE (1988) Why are migratory ungulates so
897 abundant? *American Naturalist* 131: 781-98.

898

899 **7.** Hopcraft JGC, Holdo RM, Mwangomo E, Mduma SAR, Thirgood SJ, Borner M,
900 Sinclair ARE (2015) Why are wildebeest the most abundant herbivore in the
901 Serengeti ecosystem? *Serengeti IV: sustaining biodiversity in a coupled human-*
902 *natural system*, 125.

903

904 **8.** Sinclair ARE (1995) Serengeti past and present. Pp. 3-30 In: *Serengeti II*. Sinclair,
905 ARE Arcese P (Eds). Chicago University Press, Chicago.

906

907 **9.** Ottichilo WK, de Leeuw J, Prins HHT (2001) Population trends of resident
908 wildebeest *Connochaetes taurinus hecki* (Neumann) and factors influencing them in
909 the Masai Mara ecosystem, Kenya. *Biological Conservation* 97: 271-282.

910

911 **10.** Serneels S, Lambin EF (2001). Proximate causes of land-use change in Narok

912 District, Kenya: a spatial statistical model. *Agriculture, Ecosystems Environment* 85:
913 65-81.
914

915 **11.** Wolanski E, Gereta E, Borner M, Mduma SAR (1999) Water, migration and the
916 Serengeti ecosystem. *American Scientist* 87: 523-526.
917

918 **12.** McNaughton SJ (1990) Mineral nutrition and seasonal movements of African
919 migratory ungulates. *Nature* 345: 613-615.
920

921 **13.** Fryxell JM, Sinclair ARE. (1988) Causes and consequences of migration by
922 large herbivores. *Trends in Ecology and Evolution* 9: 237-241.
923

924 **14.** Thirgood S, Mosser A, Tham S, Hopcraft JGC, Mwangomo E, Mlengeya T,
925 Kilewo M, Fryxell J, Sinclair ARE, Borner M (2004) Can Parks protect migratory
926 ungulates? The case of the Serengeti wildebeest. *Animal Conservation* 7: 113-120.
927

928 **15.** Msoffe FU, Kifugo SC, Said MY, Neselle M, van Gardingen P, Reid RS, Ogutu
929 JO, Herrero M and de Leeuw J (2011) Drivers and impacts of land-use change in the
930 Masai-Steppe of Northern Tanzania; a ecology-socio-political analysis. *Land Use*
931 *Science* 6: 261-281.
932

933 **16.** Homewood K, Lambin EF, Kariuki A, Kikula I, Kivulia J, Said MY, Serneels S,
934 Thompson M (2001) Long-term changes in Serengeti-Mara wildebeest and land
935 cover: pastoralism, population or policies? *Proceedings of the National Academy of*
936 *Science* 98: 12544-12549.

937

938 **17.** Morrison TA, Bolger DT (2012) Wet season range fidelity in a tropical migratory
939 ungulate. *Journal of Animal Ecology* 81: 543-552.

940

941 **18.** Morrison TA, Link WA, Newmark WD, Foley CA, Bolger DT (2016) Tarangire
942 revisited: Consequences of declining connectivity in a tropical ungulate population.
943 *Biological Conservation* 197: 53-60.

944

945 **19.** Foley C, Foley L (2015) Wildlife trends and status of migratory corridors in the
946 Tarangire Ecosystem, Ed. TP Wildlife Conservation Society, Arusha.

947

948 **20.** Bedelian C, Nkedianye, D, Herrero M (2007) Masai perception of the impact and
949 incidence of malignant catarrhal fever (MCF) in southern Kenya. *Preventive*
950 *Veterinary Medicine* 78: 296-316.

951

952 **21.** Reid RS (2012) *Savannas of our birth: People, wildlife, and change in East Africa.*
953 Univ of California Press, Berkeley, CA.

954

955 **22.** Estes RD, East R (2009) Status of the wildebeest (*Connochaetes taurinus*) in
956 the wild 1967-2005. Wildlife Conservation Society.

957

958 **23.** Pennycuik L (1975) Movements of the migratory wildebeest population in the
959 Serengeti area between 1960 and 1973. *African Journal of Ecology* 13: 65-87.

960

- 961 **24.** Western D (1975) Water availability and its influence on the structure and
962 dynamics of a savannah large mammal community. *African Journal of Ecology* 13:
963 265-286.
964
- 965 **25.** Western D (1982) Amboseli National Park: Enlisting Landowners to Conserve
966 Migratory Wildlife. *Ambio* 5: 302-305.
967
- 968 **26.** Ogutu JO, Piepho H-P, Said MY, Kifugo SC (2014) Herbivore Dynamics and
969 Range Contraction in Kajiado County Kenya: Climate and Land Use Changes,
970 Population Pressures, Governance, Policy and Human-wildlife Conflicts. *Open*
971 *Ecology Journal* 7: 9-31.
972
- 973 **27.** Lamprey HF (1964) Estimation of large mammal densities, biomass and energy
974 exchange in the Tarangire Game Reserve and the Masai Steppe inTanganyika. *East*
975 *African Journal of Ecology* 2: 1-46.
976
- 977 **28.** Borner M (1985) The Increasing Isolation of Tarangire National Park. *Oryx* 19:
978 91-96.
979
- 980 **29.** Kahurananga J, Silkilwasha, F (1997) The migration of zebra and wildebeest
981 between Tarangire National Park and Simanjiro Plains, northern Tanzania, in 1972
982 and recent trends. *African Journal of Ecology* 35: 179-185.
983
- 984 **30.** OIKOS (2002) Analysis of Migratory movements of large mammals and their

interactions with human activities in the Tarangire area Tanzania, as a contribution
and sustainable development strategy: Tarangire-Manyara Conservation Project
(TCMP) Final Project Report. Istituto Oikos and University of Milan, Italy in
collaboration with Tanzania National Parks.

31. Ogutu JO, Owen-Smith N, Piepho H-P, Said MY (2011) Continuing wildlife
population declines and range contraction in the Mara region of Kenya during
1977-2009. *Journal of Zoology* 285: 99-109.

32. Veldhuis MP, Ritchie ME, Ogutu JO, Beale C, Estes A, Hopcraft JGC, Morrison
TA, Mwakilema W, Ojwang GO, Parr CL, Probert J, Wargute PW, Olff H (2019) The
Serengeti squeeze: cross-boundary human impacts compromise an iconic protected
ecosystem. *Science* (In Revision).

33. Ogutu JO, Piepho H-P, Said MY, Ojwang GO, Njino LW, Wargute PW, Kifugo
SC (2016) Extreme wildlife declines and concurrent increase in livestock numbers in
Kenya: What are the causes? *PLoS ONE* 10(8): e0133744.
doi:10.1371/journal.pone.0133744.

34. Ogutu JO, Piepho H-P, Dublin HT, Bhola N, Reid RS (2009) Dynamics of Mara-
Serengeti ungulates in relation to land use changes. *Journal of Zoology* 278: 1-14.

35. Ogutu JO, Owen-Smith N, Piepho H-N, Said MY, Kifugo SC, Reid RS, Gichohi
H, Kahumbu P, Andanje S (2013) Changing Wildlife Populations in Nairobi National

1009 Park and Adjoining Athi-Kaputiei Plains: Collapse of the Migratory Wildebeest. Open
 1010 Conservation Biology Journal 7: 11-26.
 1011
 1012 **36.** Msoffe FU, Said MY, Ogutu JO, Kifugo SC, de Leeuw J, van Gardingen P, Reid
 1013 RS (2011) Spatial correlates of land-use changes in the Masai-Steppe of Tanzania:
 1014 Implications for conservation and environmental planning. International Journal of
 1015 Biodiversity Conservation 3: 280-290.
 1016
 1017 **37.** Tarangire Conservation Project (TCP) (1997) Analysis of migratory movements
 1018 of large mammals and their interactions with human activities in the Tarangire area,
 1019 Tanzania, as a contribution to a conservation and sustainable development strategy.
 1020 Final report, pp.217. University of Milan and Instituto Oikos, Italy, in collaboration
 1021 with Tanzania National Parks (TANAPA).
 1022
 1023 **38.** Lamprey HF (1963) on the ecological separation of the large mammal species in
 1024 the Tarangire Game Reserve, Tanganyika. East African Wildlife Journal 1: 63-92.
 1025
 1026 **39.** Msoffe F U, Mturi FA, Galanti V, Tosi W, Wauters LA, Tosi G (2007)
 1027 Comparing data of different survey methods for sustainable wildlife management in
 1028 hunting areas: the case of Tarangire–Manyara ecosystem, northern Tanzania.
 1029 European Journal of Wildlife Research 53: 112-124.
 1030
 1031 **40.** Norton-Griffiths M (1978) Counting Animals. A series of handbooks on
 1032 techniques currently used in African Wildlife ecology. Hand book No.1; Second
 1033 Edition, African wildlife Foundation, Nairobi, Kenya.

1034

1035 **41.** Grunblatt L, Said MY, Warugute R (1996) National Rangeland Report. Summary
1036 of population estimates of wildlife and livestock. DRSRS. Nairobi, Kenya, Ministry
1037 of Planning and National Development .

1038

1039 **42.** Woodworth B, Farm B (1996) Tanzania Wildlife Conservation Monitoring:
1040 Procedure Manual. Frankfurt Zoological Society, Arusha, Tanzania.

1041

1042 **43.** Stewart DRM, Zaphiro DRP (1963) Biomass and density of wild herbivores in
1043 different East African habitats. Mammalia 27: 483-496.

1044

1045 **44.** Pearsall MH (1957) Report on an Ecological Survey of the Serengeti National
1046 Park Tanganyika. Oryx 4: 71-136.

1047

1048 **45.** Darling (1960) An ecological reconnaissance of the Mara Plains in Kenya Colony.
1049 Wildlife Monographs 5, 41pp.

1050

1051 **46.** Talbot LM, Talbot MH (1963) The wildebeest in western Masailand, East Africa.
1052 Wildlife Monographs 12, 88p.

1053

1054 **47.** Western D, Nightingale DLM (2003) Environmental change on the vulnerability
1055 of pastoralists to drought: the Masai in Amboseli, Kenya. In: Africa Environmental
1056 Outlook: Human Vulnerability to Environmental Change. Earthprint on behalf of the
1057 United Nations Environmental Program. London.

1058 Available at: http://oceandocs.net/bitstream/1834/436/1/Amboseli_Masai.pdf

1059

1060 **48.** Africover land classification. Available at <http://www.fao.org/3/a-bd854e.pdf>

1061

1062 **49.** Okello MM, D'amour DE (2008) Agricultural expansion within Kimanaelectric
1063 fences and implications for natural resource conservation around Amboseli National
1064 Park, Kenya. Journal of Arid Environments 72: 2179-2192.

1065

1066 **50.** Okello MM (2009) Contraction of wildlife dispersal area and displacement by
1067 human activities in Kimana Group Ranch near Amboseli National Park, Kenya. Open
1068 Conservation Biology Journal 3: 49-56.

1069

1070 **51.** Reid RS, Kaelo D, Galvin KA, Harmon R (2016) Pastoral Wildlife Conservancies
1071 in Kenya: A Bottom-up Revolution in Conservation, Balancing Livelihoods and
1072 Conservation? Proceedings of the International Rangelands Congress 18-22 July
1073 2016, Saskatoon, Canada.

1074

1075 **52.** Lamprey RH, Reid RS (2004) Expansion of human settlement in Kenya's Masai
1076 Mara: what future for pastoralism and wildlife?. Journal of Biogeography 31:997-
1077 1032.

1078

1079 **53.** Ogutu JO, Piepho HP, Reid RS, Rainy ME, Kruska RL, Worden JS, Hobbs NT
1080 (2010) Large herbivore responses to water and settlements in savannas. Ecological
1081 Monographs 80: 241-266.

1082

1083 **54.** Ogutu JO, Reid RS, Piepho H-P, Hobbs NT, Rainy ME, Kruska RL, Nyabenge M
1084 (2014). Large herbivore responses to surface water and land use in an East African
1085 savanna: implications for conservation and human- wildlife conflicts. *Biodiversity*
1086 and Conservation 23: 573-596.
1087
1088 **55.** Jolly GM (1969) Sampling methods for aerial censuses of wildlife populations.
1089 East African Agricultural and Forestry Journal 34(sup1): 46-49.
1090
1091 **56.** Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a
1092 practical information-theoretic approach. Springer Science and Business Media.
1093
1094 **57.** SAS Institute Inc (2018) SAS system for windows. Cary, North Carolina, USA.
1095
1096 **58.** Stelfox JG, Peden DG, Epp H, Hudson RJ, Mbugua SW, Agastiva JL, Amuyunzu
1097 C L (1986) Herbivore Dynamics in Southern Narok, Kenya. *Journal of Wildlife*
1098 Management 50: 339-347.
1099
1100 **59.** Maddock L (1979) The “migration” and grazing succession. Pp. 104-129 In: ARE
1101 Sinclair, M Norton-Griffiths (1979), University of Chicago Press, Chicago, USA.
1102
1103 **60.** Hopcraft JGC, Morales J, Beyer H, Borner M, Mwangomo E, Sinclair ARE, Olf
1104 H, Haydon D (2014) Competition, predation, and migration: individual choice
1105 patterns of Serengeti migrants captured by hierarchical models. *Ecological*
1106 Monographs 84: 355–372.
1107

- 1108 **61.** Sinclair ARE (1973) Population increases of buffalo and wildebeest in the
1109 Serengeti. *African Journal of Ecology* 11: 93-107.
1110
- 1111 **62.** Sinclair ARE, Norton-Griffiths M (1982) Does competition or facilitation regulate
1112 ungulate populations in the Serengeti? A test of hypotheses. *Oecologia*. 53: 364-369.
1113
- 1114 **63.** Sidney J (1965) The past and present distribution of some African ungulates.
1115 Transactions of the Zoological Society of London 30: 1-396.
1116
- 1117 **64.** Jenkins P (2001) Wildlife use in World War II. In: An impossible dream, some of
1118 Kenya's last colonial wardens recall the game department in the closing years of the
1119 British Empire, Pages 39-41, eds, I Parker and S Bleazard. Librario Publishing
1120 Limited.
1121
- 1122 **65.** Talbot LM, Stewart DRM (1964) First wildlife census of the entire Serengeti-
1123 Mara Region, East Africa. *Journal of Wildlife Management* 28: 815-827.
1124
- 1125 **66.** Talbot LM, Talbot MH (1961) Preliminary observations on the population
1126 dynamics of wildebeest in Narok District, Kenya. *East African Agricultural and*
1127 *Forestry Journal* 27: 108-116.
1128
- 1129 **67.** Simon N (1962) Between the sunlight and the Thunder: The Wild Life of Kenya.
1130 London: Collins.
1131
- 1132 **68.** Serneels S, Lambin EF (2001) Impacts of Land-use changes on the wildebeest

1133 migration in the northern part of the Serengeti-Mara ecosystem. Journal of
1134 Biogeography 28: 391-407.
1135
1136 **69.** Thomson J (1885) Through Masailand. Samson Low, Marston, Searle and
1137 Rivington, London.
1138
1139 **70.** Game Report (GAR) (1907) Game Annual Report 1906-07. Kenya National
1140 Archives.
1141
1142 **71.** Game Report (GAR) (1910) Game Report and lists of game killed 1909-1910.
1143 Kenya National Archives.
1144
1145 **72.** Heller E (1913) The white rhinoceros. Smithsonian Miscellaneous Collections 61:
1146 1-77.
1147
1148 **73.** Roosevelt T, Heller E (1914) Life Histories of African Game Animals. Charles
1149 Scribner's Sons, New York.
1150
1151 **74.** Meinertzhagen R (1957) Kenya Diary (1902-1906). Edinburgh: Oliver and Boyd.
1152
1153 **75.** Stewart DRM, Stewart J (1963) The distribution of some large mammals in
1154 Kenya. Journal of East African Natural History Society 24:1-52.
1155
1156 **76.** Percival AB (1928) A game Ranger on Safari. Nesbit Co Ltd, London.
1157

- 1158 **77.** Ogutu JO, Kuloba B, Piepho H-P, Kanga E (2017) Wildlife population dynamics
1159 in human-dominated landscapes under community-based conservation: Example of
1160 Nakuru Wildlife Conservancy, Kenya. PLoS One 12(1), e0169730.
1161
- 1162 **78.** Estes RD, Atwood JL, Estes AB (2006) Downward trends in Ngorongoro Crater
1163 ungulate populations 1986–2005: Conservation concerns and the need for ecological
1164 research. Biological Conservation 131: 106-120.
1165
- 1166 **79.** Oates L, Rees PA (2013) The historical ecology of the large mammal populations
1167 of Ngorongoro Crater, Tanzania, east Africa. Mammal Review 43: 124-141.
1168
- 1169 **80.** Foster JB, Kearney D (1967) Nairobi National Park game census, 1966. East
1170 African Wildlife Journal 5: 112-120.
1171
- 1172 **81.** Foster JB, Coe MJ (1968) The biomass of game animals in Nairobi National Park,
1173 1960-66. Journal of Zoology 155: 413-25.
1174
- 1175 **82.** Gichohi HW (1996) The Ecology of a truncated ecosystem, The Athi-Kapiti
1176 Plains. PhD Thesis, University of Leicester, Leicester.
1177
- 1178 **83.** Gichohi H (2000) Functional relationships between parks and agricultural areas in
1179 East Africa: the case of Nairobi National Park. In: Prins HHT, GrootenhuisJG,
1180 Thomas TD (Eds.) Wildlife Conservation by Sustainable Use.
1181
- 1182 **84.** De Beaton KP (1949) A warden’s diary. East African Standard.

- 1183
- 1184 **85.** Percival AB (1924) A Game Ranger's Note Book. Nisbet, London.
- 1185
- 1186 **86.** Hillman JC, Hillman AK (1977) Mortality of wildlife in Nairobi National Park,
- 1187 during the drought of 1973–1974. African Journal of Ecology 15: 1-18.
- 1188
- 1189 **87.** Simon N (2001) New Directions in the 1950s. In: An impossible dream, some of
- 1190 Kenya's last colonial wardens recall the game department in the closing years of the
- 1191 British Empire, Pages 83-92, eds, I Parker and S Bleazard. Librario Publishing
- 1192 Limited. Kinloss, Scotland.
- 1193
- 1194 **88.** Stanley J (2000) The Machakos Wildlife Forum: The story from a woman on the
- 1195 land. In Wildlife Conservation by Sustainable Use (pp. 13-20). Springer Netherlands.
- 1196
- 1197 **89.** Stabach JA (2015) Movement, resource selection, and the physiological stress
- 1198 response of white-bearded wildebeest. PhD Thesis, Colorado State University,
- 1199 Fort Collins, USA.
- 1200
- 1201 **90.** Sinclair ARE, Dublin H, Borner M (1985) Population regulation of Serengeti
- 1202 Wildebeest: a test of the food hypothesis. Oecologia 65: 266-268.
- 1203
- 1204 **91.** Mduma SAR, Sinclair ARE, Hilborn R (1999) Food regulates the Serengeti
- 1205 wildebeest: a 40-year record. Journal of Animal Ecology 68: 1101-1122.
- 1206

- 1207 **92.** Hobbs NT, Galvin KA, Stokes CJ, Lockett JM, Ash AJ, Boone RB, Reid RS,
1208 Thornton PK (2008) Fragmentation of rangelands: Implications for humans, animals,
1209 and landscapes. *Global Environmental Change* 18:776-785.
1210
- 1211 **93.** Western D (2007) A half a century of habitat change in Amboseli National Park,
1212 Kenya. *African Journal of Ecology* 45: 302-310.
1213
- 1214 **94.** Western D (2007) The ecology and changes of the Amboseli ecosystem.
1215 Recommendations for planning and conservation. Amboseli Conservation Program
1216 Report, 53pp. ACC, Nairobi, Kenya.
1217
- 1218 **95.** Mose VN, Nguyen-Huu T, Western D, Auger P, Nyandwi C (2013) Modelling the
1219 dynamics of migrations for large herbivore populations in the Amboseli National
1220 Park, Kenya. *Ecological Modelling* 254: 43-49.
1221
- 1222 **96.** Andere DK (1981) Wildebeest *Connochaetes taurinus* (Burchell) and its food
1223 supply in Amboseli Basin. *African Journal of Ecology* 19: 239-250.
1224
- 1225 **97.** Stabach JA, Wittemyer G, Boone RB, Reid RS, Worden JS (2016) Variation in
1226 habitat selection by white-bearded wildebeest across different degrees of human
1227 disturbance. *Ecosphere* 7(8):e01428. 10.1002/ecs2.1428.
1228
- 1229 **98.** Bhola N, Ogutu JO, Said MY, Olff H (2012) Herbivore hotspots in the Mara
1230 Region of Kenya in relation to land use. *Journal of Animal Ecology* 81: 1268-1287.
1231

99. Morara MK, MacOpiyo L, Kogi-Makau W (2014) Land use, land cover change in urban pastoral interface. A case of Kajiado County, Kenya. *Journal of Geography and Regional Planning* 7: 192-202.

100. Reid RS, Rainy M, Ogutu JO, Kruska RL, Kimani K, Nyabenge M, McCartney M, Kshatriya M, Worden J, Ng'ang'a L, Owuor J, Kinoti J, Njuguna E, Wilson CJ, Lamprey R (2003) People, wildlife and livestock in the Mara ecosystem: The Mara Count 2002. International Livestock Research Institute, Nairobi, Kenya. Available online at:

https://www.researchgate.net/profile/Joseph_Ogutu2/publication/266852551_People_wildlife_and_livestock_in_the_Mara_Ecosystem_the_Mara_Count_2002/links/54508a6d0cf249aa53da977c.pdf

101. Løvschal M, Bøcher PK, Pilgaard J, Amoke I, Odingo A, Thuo A, Svenning JC (2017) Fencing bodes a rapid collapse of the unique Greater Mara ecosystem. *Nature Scientific Reports* 7:41450. DOI: 10.1038/srep41450.

102. Williamson D, Williamson J (1985) Botswana's fences and the depletion of Kalahari wildlife. *Oryx* 18: 218-222.

103. Tambling CJ, Du Toit JT (2005) Modelling wildebeest population dynamics: implications of predation and harvesting in a closed system. *Journal of Applied Ecology* 42: 431-441.

- 1256 **104.** Western D (2010) The Worst Drought: Tipping point or Turning point. Swara
1257 2:16-20.
1258
- 1259 **105.** Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P
1260 (2008) Global trends in emerging infectious diseases. Nature 451: 990–993.
1261
- 1262 **106.** Mills JN, Gage KL, Khan AS (2010) Potential influence of climate change on
1263 vector-borne and zoonotic diseases: a review and proposed research plan.
1264 Environmental Health Perspectives 118:1507-1514.
1265
- 1266 **107.** Bryony AJ, Grace D, Kock R, Alonso S, Rushton J, Said MY, McKeever D,
1267 Mutua F, Young J, McDermott J, Pfeiffer DO (2013) Zoonosis emergence linked to
1268 agricultural intensification and environmental change. Proceedings of the National
1269 Academy of Science 110(21): 8399-8404. <http://dx.doi.org/10.1073/pnas.1208059110>
1270
- 1271 **108.** Mduma SAR, Hilborn R, Sinclair ARE (1998) Limits to exploitation of
1272 Serengeti wildebeest and implications for its management. Pp 243–265 in Dynamics
1273 of tropical communities. Newbury DM, Prins, HHT Brown N (Eds). Oxford:
1274 Blackwell Science.
1275
- 1276 **109.** Rentsch D, Packer C (2015) The effect of bushmeat consumption on migratory
1277 wildlife in the Serengeti ecosystem, Tanzania. Oryx 49: 287-294.
1278

- 1279 **110.** Knapp EJ (2012) Why poaching pays: a summary of risks and benefits illegal
 1280 hunters face in Western Serengeti, Tanzania. *Tropical Conservation Science* 5: 434-
 1281 445.
 1282
- 1283 **111.** Msoffe FU, Ogutu JO, Kaaya J, Bedelian C, Said MY, Kifugo SC, Reid RS,
 1284 Neselle M, van Gardingen P, Thirgood S (2010) Participatory wildlife surveys in
 1285 communal lands: a case study from Simanjiro, Tanzania. *African Journal of Ecology*
 1286 48: 727-735.
 1287
- 1288 **112.** Lee DE, Kissui BM, Kiwango YA, Bond ML (2016) Migratory herds of
 1289 wildebeests and zebras indirectly affect calf survival of giraffes. *Ecology and*
 1290 *Evolution* 6: 8402-8411.
 1291
- 1292 **113.** Bedelian C, Ogutu JO (2017) Trade-offs for climate-resilient pastoral livelihoods
 1293 in wildlife conservancies in the Mara ecosystem, Kenya. *Pastoralism* 7: 10.
 1294
- 1295 **114.** Norton-Griffiths M, Said MY (2010) The future for wildlife on Kenya's
 1296 rangelands: an economic perspective. *Wild Rangelands: Conserving wildlife while*
 1297 *maintaining livestock in semi-arid ecosystems*, p. 367-392.
 1298
- 1299 **115.** KWS 2010 Aerial total count: Amboseli – West Kilimanjaro Natron cross border
 1300 landscape, Wet season, March 2010. Available at
 1301 <https://www.kws.go.ke/kws/sites/default/files/Amboseli%20West%20Kilimanjaro%20and%20Magadi%20->
 1302

1303 %20Natron%20Cross%20Border%20Landscape%20March%202010%20Wet%20Sea
1304 son.pdf.
1305
1306 **116.** Thompson DM, Homewood K (2002) Entrepreneurs, elites and exclusion in
1307 Masailand: trends in wildlife conservation and pastoral development. Human Ecology
1308 30: 107-138.
1309
1310 **117.** McCabe JT, Leslie PW, DeLuca L (2010) Adopting cultivation to remain
1311 pastoralists: The diversification of Masai livelihoods in northern Tanzania. Human
1312 Ecology 38: 321-334.
1313
1314 **118.** Boone RB, Galvin KA, Thornton PK, Swift DM, Coughenour MB (2006)
1315 Cultivation and conservation in Ngorongoro conservation area, Tanzania.
1316 Human Ecology 34: 809-828.
1317
1318 **119.** Norton-Griffiths M, Said MY, Serneels S, Kaelo DS, Coughenour M, Lamprey
1319 RH, Thompson DM, Reid RS (2008) Land use economics in the Mara Area of the
1320 Serengeti Ecosystem. Pp. 379-416In: Serengeti III: The future of an ecosystem, Eds
1321 ARE Sinclair, C Packer, SAR Mduma, JM Fryxell. University of Chicago Press.
1322
1323 **120.** Mundia NC, Murayama Y (2009) Analysis of land use/cover changes and animal
1324 population dynamics in a wildlife sanctuary in East Africa. Remote Sensing 1: 952-
1325 970.
1326

- 1327 **121.** Norton-Griffiths M (1996) Property rights and the marginal wildebeest: An
1328 economic analysis of wildlife conservation options in Kenya. Biodiversity and
1329 Conservation 5: 1557-1577.
1330
- 1331 **122.** Homewood K (2009) Policy and practice in Kenya rangelands: Impacts on
1332 livelihoods and wildlife. Pages 335-367 in K Homewood, P Kristjanson, PC Trench,
1333 editors. Staying Masai? Livelihoods, conservation and development in East African
1334 rangelands. Springer, New York.
1335
- 1336 **123.** Western D, Groom R, Worden J (2009) The impact of subdivision and
1337 sedentarization of pastoral lands on wildlife in an African savanna ecosystem.
1338 Biological Conservation 142: 2538-2546.
1339
- 1340 **124.** Kideghesho JR (2002) Trends in areas adjacent to Tarangire National
1341 Park, Tanzania: What Community-Based land use planning can offer?
1342 Kakakuona[Jan-March]. Ministry of Natural Resources and Tourism, Tanzania.
1343
- 1344 **125.** Osano PM, Said MY, Leeuw J, Ndiwa N, Kaelo D, Schomers S, Ogutu JO
1345 (2013) Why keep lions instead of livestock? Assessing wildlife tourism-based
1346 payment for ecosystem services involving herders in the Masai Mara, Kenya. Natural
1347 Resources Forum 37: 242-256.
1348
- 1349 **126.** Blackburn S, Hopcraft JGC, Ogutu JO, Matthiopoulos J, Frank L (2016)
1350 Human–wildlife conflict, benefit sharing and the survival of lions in pastoralist
1351 community-based conservancies. Journal of Applied Ecology 53: 1195–1205.

1352

1353 **127.** Western D, Russell S, Cuthill I (2009) The status of wildlife in protected areas
1354 compared to non-protected areas of Kenya. PloS One 4: e6140.

1355

1356 **128.** Dougherty LS (2014) The ecological viability of wildlife conservancies in the
1357 Mara ecosystem. Transfer report to spatial ecology and land use unit, faculty of health
1358 and life sciences, Oxford Brookes University. Unpublished Report.

1359

1360 **129.** Nkedianye D, Radeny M, Kristjanson P, Herrero M (2009) Assessing returns to
1361 land and changing livelihood strategies in Kitengela Pages 115-150 in K Homewood,
1362 P Trench, and P Kristjanson, editors. Staying Masai? Livelihoods, Conservation and
1363 Development in East African Rangelands. Springer-Verlag, London.

1364

1365 **130.** de Leeuw JM, Said MY, Kifugo S, Ogutu JO, Osano P, de Leeuw J (2014)
1366 Spatial variation in the willingness to accept payments for conservation of a migratory
1367 wildlife corridor in the Athi-Kaputiei Plains, Kenya. Ecosystem Services 8: 16-24.

1368

1369 **131.** Matiko D (2014) Wildlife conservation leases are considerable conservation
1370 options outside Protected Areas: The Kitengela-Nairobi National Park Wildlife
1371 Conservation Lease Program. Journal of Ecosystem and Ecography. 4:2.
1372 <http://dx.doi.org/10.4172/2157-7625.1000146>.

1373

1374 **132.** Nelson F, Foley C, Foley LS, Leposo A, Loure E, Peterson D, Peterson
1375 MPeterson T,Sachedina H,Williams A (2010) Payments for ecosystemservices as a

1376 framework for community-based conservation in Northern Tanzania. Conservation
1377 Biology 24: 78–85.
1378
1379 **133.** Morrison TA, Bolger DT (2014) Connectivity and bottlenecks in a migratory
1380 wildebeest *Connochaetes taurinus* population. Oryx 48: 613-621.
1381
1382 **134.** Norton-Griffiths, M (2000) Wildlife losses in Kenya: An analysis of
1383 conservation policy. Natural Resources Modelling 13: 13-34.
1384
1385 **135.** Republic of Kenya (2013) The Wildlife Conservation and Management Act,
1386 2013. Kenya Gazette Supplement No. 181, Acts No. 47, Sixth Schedule. Nairobi.
1387 Available at: [http://kenyalaw.org/kl/fileadmin/pdfdownloads/Acts/](http://kenyalaw.org/kl/fileadmin/pdfdownloads/Acts/WildlifeConservationandManagement%20Act2013.pdf)
1388 [WildlifeConservationandManagement%20Act2013.pdf](http://kenyalaw.org/kl/fileadmin/pdfdownloads/Acts/WildlifeConservationandManagement%20Act2013.pdf). Kenya Gazette
1389 Supplement No. 18/ (Acts No. 47). Republic of Kenya.
1390
1391 **136.** Ministry of Tourism and Wildlife (2018) National Wildlife Strategy 2030.
1392 Government of Kenya Publication, Nairobi.
1393
1394 **137.** Ojwang' GO, Wargute PW, Said MY, Worden JS, Davidson Z, Muruthi P, Kanga
1395 E, Ihwagi F, Okita-Ouma B (2017) Wildlife migratory corridors and dispersal areas:
1396 Kenya rangelands and coastal terrestrial ecosystems. Government of the Republic of
1397 Kenya, Nairobi.
1398

- 1399 **138.** United Republic of Tanzania (2018) Wildlife Conservation (Wildlife corridors,
1400 dispersal areas, buffer zones and migratory routes) Regulations 2018. Government
1401 Printer, Dar es Saalam.
1402
- 1403 **139.** Ogutu J O, Owen-Smith N, Piepho H P, Kuloba B, Edebe J (2012) Dynamics of
1404 ungulates in relation to climatic and land use changes in an insularized African
1405 savanna ecosystem. *Biodiversity Conservation* 21:1033-1053.
1406
- 1407 **140.** Swynnerton GH (1958) Fauna of the Serengeti National Park. *Mammalia* 22:
1408 435-450.
1409
- 1410 **141.** Grizmek B, Grizmek M (1960) Serengeti shall not die. Hamish Hamilton, Ltd.
1411 London. 344pp.
1412
- 1413 **142.** Stewart D RM, Talbot LM (1962) Census of wildlife in the Serengeti, Mara and
1414 Loita Plains. *East African Agricultural and Forestry Journal* 28: 58-60.
1415
- 1416 **143.** Anderson GD, Talbot, LM (1965) Soil factors affecting the distribution of the
1417 grassland types and their utilisation by wild animals on the Serengeti plains,
1418 Tanganyika. *Journal of Ecology* 53: 33-56.
1419
- 1420 **144.** Watson RM (1967) The population ecology of the wildebeest (*Connochaetes*
1421 taurus albojubatus Thomas) in the Serengeti. PhD Thesis, Cambridge University.
1422

1423 **145.** Bell R V H (1971) A grazing ecosystem in the Serengeti. Scientific American
1424 224: 86-93.
1425
1426 **146.** Kreulen D (1975) Wildebeest habitat selection on the Serengeti Plains, Tanzania,
1427 in relation to Calcium and lactation. African Journal of Ecology 13: 297-304.
1428
1429 **147.** McNaughton S J (1976) Serengeti migratory wildebeest: Facilitation of energy
1430 flow by grazing. Science 191 (4222): 92-94.
1431
1432 **148.** Hilborn R, ARE Sinclair (1979) A simulation of the wildebeest population, other
1433 ungulates and their predators. Pages 287–309 in ARE Sinclair and MNorton-Griffith,
1434 editors. Serengeti: dynamics of an ecosystem. University of Chicago Press, Chicago.
1435
1436 **149.** Sinclair ARE (1979) The eruption of the ruminants. Pp 82–103 In: A R E
1437 Sinclair and M Norton-Griffith, editors. Serengeti: dynamics of an ecosystem.
1438 University of Chicago Press, Chicago.
1439
1440 **150.** Sinclair ARE, Norton-Griffiths M (1979) Serengeti: Dynamics of an ecosystem.
1441 University of Chicago Press, Chicago, USA.
1442
1443 **151.** Bell RVH (1982) The effect of soil nutrient availability on community structure
1444 in African ecosystems. In: Ecology of tropical savannahs, p. 193-216. ed. BJHuntley,
1445 Walker BH. Springer, New York.
1446

- 1447 **152.** Broten MD, Said MY (1995) Population trends of ungulates in and around
1448 Kenya's Masai Mara Reserve. In: Serengeti II: Dynamics, Management,
1449 and Conservation of an Ecosystem, p. 169-193. ed. ARE Sinclair. P Arcese. Univ. of
1450 Chicago Press, Chicago.
- 1451
- 1452 **153.** Fryxell JM (1995) Aggregation and migration by grazing ungulates in relation
1453 to resources and predators. In: Serengeti II. Dynamics, Management, and Conservation
1454 of an Ecosystem, p. 257-273. ed. ARE Sinclair, P Arcese. Univ. of Chicago Press,
1455 Chicago.
- 1456
- 1457 **154.** Murray MG (1995) Specific nutrients requirements and migration of wildebeest.
1458 Pp 231-56 In: Serengeti II; dynamics, management and conservation of an ecosystem.
1459 University of Chicago Press, Chicago, USA.
- 1460
- 1461 **155.** Wilmshurst JF, Fryxell JM, Farm BP, Sinclair ARE, Henschel CP (1999) Spatial
1462 distribution of Serengeti wildebeest in relation to resources. Canadian Journal of
1463 Zoology 77: 1223-1232.
- 1464
- 1465 **156.** Gereta E, Wolanski E, Chiombola EAT (2003) Assessment of the environmental,
1466 social and economic impacts on the Serengeti ecosystem of the developments in the
1467 Mara. River catchment in Kenya. Amala Project Report, 59pp. TANAPA, FZS,
1468 Arusha, Tanzania.
- 1469
- 1470

157. Musiega DE, Kazadi SN (2004) Simulating the East African wildebeest migration patterns using GIS and remote sensing. *African Journal of Ecology* 42: 355-62.

158. Boone RB, Thirgood SJ, Hopcraft JGC (2006) Serengeti wildebeest migratory patterns modeled from rainfall and new vegetation growth. *Ecology* 87: 1987-94.

159. Sinclair ARE, Mduma SA, Hopcraft JGC, Fryxell J M, Hilborn R A Y, Thirgood S (2007) Long-Term Ecosystem Dynamics in the Serengeti: Lessons for Conservation. *Conservation Biology* 21: 580-590.

160. Holdo R M, Holt R D, Fryxell J M (2009) Grazers, browsers, and fire influence the extent and spatial pattern of tree cover in the Serengeti. *Ecological Applications* 19: 95-109.

161. Bhola N, Ogutu J O, Piepho H-P, Said MY, Reid RS, Hobbs NT, Olff H (2012b) Comparative changes in density and demography of large herbivores in the Masai Mara Reserve and its surrounding human-pastoral ranches in Kenya. *Biodiversity Conservation* 21: 1509-1530.

162. Bedelian C (2014) Saving the Great Migrations: Declining wildebeest in East Africa? *Environmental Development* 9: 101-109.

- 1494 **163.** Stabach JA, Boone RB, Worden JS, Florant G (2015) Habitat disturbance effects
1495 on the physiological stress response in resident Kenyan white-bearded wildebeest
1496 (*Connochaetes taurinus*). Biological Conservation 182:177–186.
1497
1498 **164.** Ottichilo WK (2000) Wildlife Dynamics: An Analysis of Change in the Masai
1499 Mara Ecosystem of Kenya. PhD Dissertation, ITC, The Netherlands.
1500
1501 **165.** Sheehan MM (2016) Determining drivers for wildebeest (*Connochaetes*
1502 *taurinus*) distribution in the Masai Mara National Reserve and surrounding Group
1503 Ranches (Doctoral dissertation, Miami University).
1504
1505 **166.** McCutcheon JT (1910) In Africa. The Bobbs-Merrill Company. Indianapolis,
1506 USA.
1507
1508 **167.** Foster JB, McLaughlin R (1968) Nairobi National Park game census, 1967.
1509 East African Wildlife Journal 6: 152-54.
1510
1511 **168.** Casebeer RL, Koss GG (1970) Food habits of wildebeest, zebra, hartebeest and
1512 cattle in Kenya Masailand African Journal of Ecology 8: 25-36.
1513
1514 **169.** Petersen JCB, Casebeer RL (1972) Distribution, population status and group
1515 composition of wildebeest (*Connochaetes taurinus* Burchell) and zebra (*Equus*
1516 *burchelli* Gray) on the Athi-Kapiti plains, Kenya. Wildlife Management Project.
1517 UNDP/FAO KEN/71 /526, Project Working Document No. 1.
1518

- 1519 **170.** Casebeer RL, Mbai HJ (1974) Animai mortality 1973/74. Kajiado District,
1520 FAOProject DP/KEN/71/526. Working Document. Report No. 5.
1521
- 1522 **171.** Owaga M L (1975) The feeding ecology of wildebeest and zebra in Athi-Kaputei
1523 Plains. African Journal of Ecology 13: 375-83.
1524
- 1525 **172.** Hillman JC (1979) The biology of the eland (*Taurotraaus oryx* Pallas) in the
1526 wild. PhD. University of Nairobi.
1527
- 1528 **173.** Trzebinski E (1985) The Kenya Pioneers. Cox and Wyman Ltd, Great Britain.
1529
- 1530 **174.** Gichohi HW (2003) Direct payments as a mechanism for conserving important
1531 wildlife corridor links between Nairobi National Park and its wider ecosystem: The
1532 Wildlife Conservation Lease Program. In Vth World Parks Congress.
1533
- 1534 **175.** Ego WK, Mbuvi D M, Kibet PFK (2003) Dietary composition ofwildebeest
1535 (*Connochaetes taurinus*), kongoni (*Alcephalus buselaphus*) andcattle (*Bos indicus*),
1536 grazing on a common ranch in south-central Kenya. AfricanJournal of Ecology 41:
1537 83-92.
1538
- 1539 **176.** Imbahale SS, Githaiga J M, Chira RM, Said MY (2008) Resource utilization by
1540 large migratory herbivores of the Athi-Kapiti ecosystem. African Journal of Ecology
1541 46: 43-51.
1542

177. Croze H (1978) Aerial surveys undertaken by the Kenya Wildlife Management Project: methodologies and results. FAO Project DP/KEN/71/526. Working document. Report No. 16.
178. Campbell DJ, Gichohi H, Mwangi A, Chege L (2000) Land Use Conflicts in Kajiado District, Kenya. Land Use Policy 17: 337-48.
179. Worden J, Reid RS, Gichohi H (2003) Land-use impacts on large wildlife and livestock in the swamps of the Greater Amboseli Ecosystem, Kajiado District, Kenya Lucid Project. International Livestock Research Institute, Nairobi, Kenya. https://cgspace.cgiar.org/bitstream/handle/10568/1901/Lucid_wp27_part1.pdf?sequence=1.
180. Okello M M (2005) Land use changes and human–wildlife conflicts in the Amboseli Area, Kenya. Human Dimensions of Wildlife 10: 19-28.
181. Sitati N, Lekishon K, Bakari S, Warinwa F, Mwiu S N, Gichohi N, Mukeka J (2014). Wildebeest (*Connochaetes taurinus*) Population densities and distribution in dry and wet season in the Kilimanjaro landscape. Natural Resources 5: 810.
182. Galanti V, Tosi G, Rossi R, Foley C (2000) The Use of GPS radio-collars to track elephants (*Loxodonta africana*) in the Tarangire National Park (Tanzania). Hystrix 11: 27-37.

183. TAWIRI (2001) Tarangire Ecosystem: Wet Season Systemmatic

ReconnaissanceFlight Count, May 2001. TAWIRI, Arusha, Tanzania.

184. Gereta E, Meing'ataki GEO, Mduma SAR, Wolanski E (2004) The role

ofwetlands in wildlife migration in the Tarangire ecosystem, Tanzania. Wetlands

Ecology and Management 12: 285-99.

185. Newmark WD (2008) Isolation of African protected areas. Frontiers in Ecology

and the Environment 6:321-8.

Figure legends

Fig 1. Wildebeest migration in the greater Serengeti-Mara ecosystem in East

Africa. (Photo credit: ARE Sinclair).

Fig 2. Map showing the general extent of Masailand in Kenya and Tanzania and

the five study ecosystems with eight populations: 1 = Serengeti Ecosystem, 2 =

Masai Mara Ecosystem, 3 = Narok County, 4 = Athi-Kaputiei Ecosystem, 5 =

Machakos County, 6 = Greater Amboseli Ecosystem, 7 = West Kajiado and 8 =

Tarangire-Manyara Ecosystem populations. Notes: NCA = Ngorongoro Conservation

Area, MGR = Maswa Game Reserve, SNP =Serengeti National Park, IGR = Ikorongo

Game Reserve, GGR = Grumeti Game Reserve, MMNR = Masai Mara National

Reserve, NNP = Nairobi National Park, ANP = Amboseli National Park, LMNP =

Lake Manyara National Park, LGCA = Lokisale Game Controlled Area, TNP =

Tarangire National Park and MGR* = Mkungunero Game Reserve. Use of each

seasonal area by the study populations is described in the text.

Fig 3. Map showing the general area occupied by the (a) Greater Serengeti-Mara, b) Athi-Kaputiei, (c) Greater Amboseli and (d) Tarangire-Manyara ecosystems. For each ecosystem, the status of routes of migratory wildebeest post-2000 in relation to the distribution of agriculture and fences is highlighted. The current wildlife conservancies (and wildlife-livestock ranches) are provided for the Masai Mara, Athi-Kaputiei and Greater Amboseli ecosystems of Kenya. Also shown are extreme land fragmentation through fences in Athi-Kaputiei and recent emergence of fences along the eastern and south eastern borders of the Mara Conservancies.

Fig 4. Movement tracks of GPS collared wildebeest during 2010-2013 (colored lines) in Kenya (A = Loita Plains in Masai Mara Ecosystem, B = Athi-Kaputiei Ecosystem, C = Greater Amboseli Ecosystem). Protected areas (1 = Masai Mara National Reserve, 2 = Serengeti National Park in Tanzania, 3 = Nairobi National Park, 4 Amboseli National Park) are partially obscured.

Fig 5. Trends in population size of migratory wildebeest populations in a) Serengeti-Mara ecosystem, b) Masai Mara ecosystem, c) Narok County in which Masai Mara is located, d) Athi-Kaputiei ecosystem, e) Machakos County, f) Greater Amboseli ecosystem, g) West Kajiado and h) Tarangire-Manyara ecosystem.

Supporting information

Table S1. The field data form used to collect information for ground truthing historical information on land use and cover changes in the Tarangire-Manyara ecosystem of Tanzania. We uploaded the coordinates and their Ids into a Global

Positioning System (GPS) and used them to locate sampling points. We collected information on the following attributes for each sampling point. (1) Was there agriculture at the location in 1984, 2000 or 2006-2007? (2) If yes, was agriculture practiced on small or large scale? (3) Was agriculture irrigated or not? (4) When did agriculture start? (5) Photographs of the sampling point. (6) Crop types cultivated at each sampling location. (7) We empirically assessed the change type we had identified during initial image interpretation in the office and assigned change codes A, B, C or D described in the table to the observed changes. (8) General comments.

Supporting information

Table S1.

Table S2. The five ecosystems with migratory wildebeest populations in East Africa, their current status and earlier studies.

Table S3. Wildlife conservancies in Masai Mara, their names, size, number of landowners that pooled land to form the conservancy, tourist camps, tourist beds, rangers and scouts and jobs created by each conservancy and year of establishment.

Table S4. Wildlife conservancies or ranches, their names and sizes in Machakos Plains (adjoining the Athi-Kaputiei), Athi-Kaputiei and Greater Amboseli ecosystem. The total area covered in each ecosystem is 347.0, 40.6 and 1046.5 km² in the Machakos Plains, Athi-Kaputiei and Greater Amboseli ecosystems, respectively.

1641 **S1 Data. The estimated population size and standard errors and other details of**
 1642 **the aerial surveys of wildebeest for the Serengeti-Mara (1957-2012) and**
 1643 **Tarangire-Manyara (1987-2016) ecosystems.**

1644

1645 **S2 Data. The estimated population size and standard errors and other details of**
 1646 **the aerial surveys of wildebeest for the Masai Mara Ecosystem (1977-2016)**
 1647 **ecosystems.**

1648

1649 **S3 Data. The estimated population size and standard errors and other details of**
 1650 **the aerial surveys of wildebeest for the Athi-Kaputiei Ecosystem (1977-2014).**

1651

1652 **S4 Data. The estimated population size and standard errors and other details of**
 1653 **the aerial surveys of wildebeest for the Machakos County (1977-2015).**

1654

1655 **S5 Data. The estimated population size and standard errors and other details of**
 1656 **the aerial surveys of wildebeest for the Greater Amboseli (1977-2014) and**
 1657 **Western Kajiado (1977-2014) ecosystems.**

bioRxiv preprint doi: <https://doi.org/10.1101/548747>; this version posted February 11, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Fig1

Fig2

Fig3

Fig4

Fig5