The Process Pathway Model of bacterial growth

Daniel Biro¹, Ximo Pechuan¹, Maryl Lambros¹, Aviv Bergman^{1,2,3,4,*}

¹Department of Systems and Computational Biology,

²Dominick P. Purpura Department of Neuroscience,

³Department of Pathology, Albert Einstein College of Medicine,

1301 Morris Park Ave, Bronx, NY 10461, USA

⁴Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

*To whom correspondence should be addressed; E-mail: aviv@einstein.yu.edu.

Abstract

The growth profile of microorganisms in an enclosed environment, such as a bioreactor or flask, is a well studied and characterized system. Despite a long history of examination, there are still many competing mathematical models used to describe an output of the microorganisms, namely the number of bacteria as a function of time. However, these descriptions are either purely phenomenological and give no intuition as to the biological mechanisms underlying the growth curves, or extremely complex and become computationally unfeasible at the population level. In this paper, we develop the Process Pathway Model by modifying a model of sequential processes, which was first used to model robustness in metabolic pathways, and demonstrate that the Process Pathway Model encapsulates many features and temperature dependence of bacterial growth. We verify the predictions of the model against growth data for multiple species of microorganisms, and confirm that the model generates accurate predictions on temperature dependence of bacterial growth. The model has five free parameters, and the simplifying assumptions used to build the model are built upon biologically realistic notions. The Process Pathway Model accurately models a microorganism's growth profile at an intermediate level of complexity that is computationally feasible. This model can be used as both an conceptual model for thinking about systems of bacterial growth, as well as a computational model that operates at level of complexity that is amenable to large scale simulation. This balance in accuracy and intuitiveness was accomplished by using realistic biological assumptions to simplify the underlying biology, which may point the way forward for future models of this type.

1 Introduction

1 The growth profile of microorganisms in an enclosed environment, such as a bioreactor or flask, known
2 as batch culture, is a commonly used and well studied and characterized system [8, 20]. It has ap-
3 plications to many fields, including food science, microbiology, experimental evolution, and bioreactor
4 engineering [4, 5, 15, 24]. However, despite a long history of examination, there are still many compet-
5 ing mathematical models used to describe the output of the system, namely the number of bacteria as
6 a function of time. Furthermore, these descriptions are either purely phenomenological, which give no
7 biological intuition into the mechanisms underlying the growth curves, or extremely complex, becoming
8 computationally unfeasible at the population level. The most common empirical models are the Logistic,
9 Gompertz, van Impe, and Baranyi-Roberts, which describe growth profiles as concentrations over time
10 for a population [2, 7, 23]. At the higher end of complexity, the full cell computational model of Karr
11 *et al.* [13] predicts division time as a function of fundamental metabolic processes for a single low com-
12 plexity species, *Mycoplasma genitalium*. Here we build off of a model of sequential processes developed
13 by Kacser and Burns, originally used to model enzymatic robustness in metabolic pathways [11, 12], in

order to develop a model of intermediate complexity between an empirical model and full cell metabolic integration. By adding temperature dependence to the simple model of Kacser and Burns [11], our model, termed the *Process Pathway Model*, encapsulates many features and temperature dependence of bacterial growth. Our results demonstrate that a relatively simple mechanistic model can be used to accurately describe and predict the dynamics of a complex biological system while maintaining biological relevance, computational tractability, and broad applicability.

Previous models of bacterial growth have been developed to predict bacterial growth rates as functions of time and temperature [2, 3, 9, 21, 22, 25]. Additional features that models are designed to predict are lag time, which is the time the bacteria spend in a stationary state before growing; carrying capacity, which is the maximal concentration that the bacteria grow to; and maximal growth rate, all as functions of variables such as temperature and pH. In addition to closed-form equations, differential equation models have also been applied to the problem of bacterial growth rates. The most common of these models is the Baryani-Roberts model [2, 3]. The equations of the model are integrated to obtain growth curves, and the predictions are typically better than those of closed-form equations, though parameters such as lag time, maximal growth rate, and carrying capacity have to be explicitly added to the differential equation models (see Supplementary Data: Fig. S4 for full comparison of models). Refinements and additions to the Baranyi-Roberts and van Impe models have had some success in predicting population level phenomena by approximating underlying processes, but these models remain highly phenomenological with little ability to extract biological insight into key parameters, such as the lag time [1, 14, 17–19, 25]. Other empirical models such as the Ratkowsky model, asymptote model, and hyperbola model describe a single aspect of bacterial growth, such as growth rate, carrying capacity, or lag, respectively, as functions of temperature, and are adaptable to numerous species of bacteria [21, 22, 26–28]. However, each empirical model is designed to explain only a single aspect of bacterial growth, such as density as a function of time, carrying capacity, or lag time, but no model has successfully integrated all of these features into a greater framework.

To further the understanding of the underlying phenomenon of bacterial growth and predict many key features of growth in a way that is computationally feasible, we introduce the Process Pathway Model. While abstract in nature, the model is rooted in the fundamental processes of biology, without becoming overly complex. The model is derived from a model of enzymatic robustness first put forward by Kacser and Burns [11]. However, rather than individual enzymes, gene expression and other cellular and physiological processes are modelled as being the underlying phenomena in the model. The model predicts key features of growth, specifically the lag time, maximal growth rate, and carrying capacity, as functions of temperature with similar accuracy to existing models of bacterial growth, but with more biological meaning and computational tractability. Additionally, the model can predict growth under fluctuating temperatures. Thus the model is useful on two levels; firstly as a model of simplifying biological assumptions to distill the most important abstractions involved in bacterial growth. Secondly the Process Pathway Model can be used as an accurate model to make predictions of the quantitative parameters involved in bacterial growth.

2 Material and Methods

2.1 Mathematical Model

The Process Pathway Model consists of a chain of N processes, each processing an input and generating an output. Each process can be thought of as a series of physiological and gene regulatory activities. For example, one process could model the up regulation of gene expression in a particular metabolic pathway as a result of exposure to a novel resource rich environment. The totality of these processes, each representing a different cellular activity, comprise the network of cellular metabolism.

Each process in the chain of N processes is governed independently by Michaelis-Menten kinetics,

60 resulting in a flux, ϕ_i between processes given by:

$$\phi_i = \frac{S_{i-1} * V_{max,i}}{K_{M,i} + S_{i-1}} \quad (1)$$

61 for $i = 1, 2, 3, \dots, N$, where S_N is the concentration of the output of process N . For this model we
62 assume no flux into S_0 and the flux out of S_N is given by a linear death rate. The dynamics for the
63 variables S_i is then given by:

$$\frac{dS_i}{dt} = \frac{S_{i-1} * V_{max,i}}{K_{M,i} + S_{i-1}} - \frac{S_i * V_{max,i+1}}{K_{M,i+1} + S_i} \quad (2)$$

64 for $i = 1, 2, 3, \dots, N - 1$,

$$\frac{dS_0}{dt} = -\frac{S_0 * V_{max,1}}{K_{M,1} + S_0} \quad (3)$$

65 for $i = 0$ and

$$\frac{dS_N}{dt} = \frac{S_{N-1} * V_{max,N}}{K_{M,N} + S_{N-1}} - D * S_N \quad (4)$$

66 for $i = N$ (Fig. 1 Panel A). S_0 represents the concentration of the limiting resource, and is subsequently
67 reduced as it is consumed by the process generating S_1 . The initial value of S_0 is a free parameter, while
68 the initial value of all other S_i are set to 0 to model an initial state before growth has begun. The
69 concentration of bacteria at time t is taken to be the value of the final process, $S_N(t)$, while the value
70 of all other $S_i(t)$ represent the activity of the intermediate processes (Supplemental Figures: Fig. S1).
71 Here, the final process is taken to represent the progress of the final metabolic pathway in the chain, in
72 this case that of reproduction. The parameter D represents the natural death rate of the population.
73 $V_{max,i}$ and $K_{M,i}$ here follow the same intuition as traditional Michaelis-Menten kinetics, where $V_{max,i}$
74 represents the maximal activity of each process, and $K_{M,i}$ is the concentration of substrate at which the
75 process is at half of its maximal activity. In the original model of Kacser and Burns [11], $K_{M,i}$ were
76 free to vary between enzymes; however here we take $K_{M,i}$ to be equal for all processes to reduce the
77 number of free parameters without sacrificing significant accuracy. Other instantiations of this model
78 may benefit from the relaxation of this constraint.

79 Temperature dependence of each process is incorporated by modelling a temperature dependence of
80 the parameter $V_{max,i}$. The functional form is a modified version of the function described by Daniels
81 *et al.* [6], although the exact form of the temperature dependence did not significantly alter the results.
82 The primary features of the temperature dependence that was salient for the model were the peak and
83 the minima of the temperature dependence.

84 It is important to note here that while the equation for temperature dependence contain multiple
85 parameters, they are not all free parameters, as the biological constraints effectively reduce the set of
86 free parameters for temperature dependence of growth to a single value, visualized as the temperature
87 of maximal growth. Thus, in all, this model has effectively three free parameters for defining bacterial
88 growth; the maximal growth temperature defining the growth curve, S_0 , and $K_{M,i}$. With the use of these
89 parameters and the above assumptions which were used to construct the model, we can then generate
90 predictions throughout the entire biokinetic range for the pertinent characteristics of bacterial growth.

91 In order to determine the optimal value for N , the number of processes in the chain, we compared
92 experimental data from growth of *Escherichia coli* to the predictions of the model, yielding a optimal
93 prediction of $N = 8$ (Supplemental Figures: Fig. S2). This is in line with predictions of the diameter of
94 process networks in bacteria, as demonstrated by the whole cell computation model of Karr [13] utilizing
95 6 independent metabolic "compartments". Furthermore, this result is robust for a wide range of values

of N (see **Supplemental Figures: Fig. S3**), and these values are likely to be widely applicable, as network diameter scales with the logarithm of network size [10].

Applying the Pathway Process Model to data obtained from *E. coli* at a single temperature resulted in an accurate prediction of growth (**Fig. 2 Panel A**). The prediction was comparable to existing models of bacterial growth (**Supplemental Figures: Fig. S4**). Furthermore, the application of the Pathway Process model to data obtained of growth under a fluctuating temperature profile (**Fig. 2 Panel B**) fit the data as well as existing phenomenological models (**Supplemental Figures: Fig. S5**)

In order to test the ability of the model to predict the effects of temperature on the properties of bacterial growth, we determined a single set of parameters that best matched the data for growth rate, μ , for *Lactobacillus plantarum* data from Zwietering *et al.* [27] (**Fig. 3 Panel A**). The plot of the function for $V_{max,i}(T)$ using the parameters in **Table S1** is shown in (**Supplemental Figures: Fig. S5**). It should be noted that the curve in this figure is derived experimentally, and is one of the few free parameters fed into the model. The results of the growth rate across the full dynamic range of the model are shown in **Fig. 1 Panel C** alongside the data raw data and the prediction from the empirical fit model. Zwietering *et al.* obtained values of μ , lag time, and carrying capacity by fitting bacterial count data to a Gompertz curve, so for consistency this is the same method used to extract parameters from the growth curves obtained from the Pathway Process Model. The algorithm used to estimate parameters and derive results is summarized in **Fig. 1 Panel B**.

2.2 Computational Modelling

All computations were performed in MATLAB 2017a (Mathworks, Natick, MA). Solving of differential equations were performed using the function ode23s. Gompertz curves were fit using the Marquardt algorithm implemented in MATLAB.

Parameters for the temperature modelling were optimized by applying a gradient descent algorithm across all free parameters to minimise the least squares error between the predicted maximal growth rate, μ , and the data from Zwietering *et al.* Figure 2, except for D , which was chosen to be small. The least squares values were calculated by fitting a Gompertz curve to the output of the model integration, and extracting the parameter μ from this fit at each temperature. Once the parameters for temperature dependence were established using this procedure, the lag time and carrying capacity were similarly extracted from the Gompertz fit using the same parameters as for the maximal growth rate, μ . Parameters for all values are given in **Table S1**.

2.3 Data for Bacterial Growth Measurements

A derivative strain of *Escherichia coli* REL606 was used to seed the experiment. Bacteria were grown in M9 media with glucose supplemented to 4g/L. Tubes containing media were placed in BioSan LV Personal Bioreactor RTS-1C (Riga, Latvia), which controlled temperature to within 0.1°C. OD850nm measurements were taken by the RTS-1C at 2 minute intervals. To seed the culture, 100 μ L of old culture was transferred to a new tube containing 20mL of fresh media.

Temperature switching and control was performed by the built in function of the Bioreactor RTS-1C. Temperature switching was programmed to occur after specified measurements of OD850nm occurred for the first time in a growth cycle. After temperature switching occurred, new temperature equilibrium was reached in approximately 30 minutes, which is less than the expected doubling time for the bacteria.

Each of the experiments started by inoculating the medium with 0.1 mL of an overnight culture grown at 37 ° in minimal media M9 with 10 % glucose. For the evaluation of the temperature dependence of the growth parameters, an additional day of acclimation was allowed [16].

2.4 Data for Growth Rate, Carrying Capacity, and Lag

Data for Figure 3 Panels A-C was used with permission from Figures 2, 5, & 6 of Zwietering *et al.* 1991. Data was originally collected from growth measurements of *Lactobacillus plantarum* in MRS media, with growth rates calculated by measuring CFU by titers, and parameters were calculated by fitting the data to a Gompertz curve.

3 Results

The two primary predictions of the model concerning the growth of bacteria over the entire dynamic temperature range, namely carrying capacity and lag time, are shown in Fig. 3 Panels A and B alongside the data from Zwietering *et al* [27]. In fact, aspects of the lag time and carrying capacity as functions of temperature are predicted by the model to a higher degree of precision than is seen in the empirical predictions. These results were obtained with minimal assumptions about the biology of the system and few free parameters.

The predictions for growth rate as a function of temperature generated by the process pathway model are equivalent to the empirical model in predicting the actual growth data (Fig. 3 Panel a). Similarly, the carrying capacity data derived from the process pathway model are shown to be similar to the empirical model in the middle of the growth range (Fig. 3 Panel C). Additionally, for carrying capacity, a slight decrease of the carrying capacity at low temperatures is predicted by the Process Pathway Model but not by the Ratkowsky asymptote model [27]. This results is shown both qualitatively and quantitatively in the data.

Additionally, for the lag time, the slight increase in lag time at higher temperatures was not accurately captured by the hyperbola model of lag time [27] but is predicted by the Process Pathway Model (Fig. 3 Panel B). Again, this result is shown both quantitatively and qualitatively as a prediction of the process pathway model and in the empirical data, but is not expected or shown in the traditional empirical models of bacterial growth.

The results here are derived from a model that takes in effectively five free parameters, and uses simplifying assumptions about the dynamics of bacterial growth in order to predict the biokinetic proprieties of growth throughout the viable temperature range.

4 Discussion

Bacterial growth is a highly complex phenomenon with many influences and complex behaviors. While there are numerous models describing growth, all are either empirical or detailed to the extent of being computationally burdensome, and none to our knowledge incorporate a framework based on simple abstractions of fundamental metabolic processes. Additionally, each empirical model is designed to explain only a single aspect of bacterial growth, such as density as a function of time, carrying capacity, or lag time, but no model has successfully integrated all of these phenomenon into a greater framework. Here we have put forward a model that encapsulates many of the pertinent features of bacterial growth, particularly in regards to temperature sensitivity, while being computationally tractable enough to be used for population level modelling. While biologically, the different phases of growth involve numerous different mechanisms and pathways, we were able to successfully abstract them away into their fundamental contributions in this model. This was surprisingly able to be successfully done with a single set of parameters for all phases. Many studies, including experimental evolution and protein structure and stability studies take interest in the effects of temperature on metabolic processes. Our model can successfully predict the above phenomenon with minimal assumptions and few free parameters. We reiterate that this model effectively has three free parameters, and the simplifying assumptions used to build the model are built upon biologically realistic assumptions.

183 One of the interesting and counter-intuitive insights gleaned from this model is the relationship be-
 184 tween lag time and maximum growth rate. Previous models have treated these two variables as inde-
 185 pendent entities, though perhaps correlated, can be independently modelled. However a direct result of
 186 the process pathway model is that the lag time is simply a result of the rate of process activity, V_{max} of
 187 each individual process. While this intuitively expected to be the case for the maximum growth rate, the
 188 more interesting result is that it is also the case for the lag time, which has been traditionally modelled
 189 as a phenomenon in its own right.

190 While this model is not a comprehensive model of bacterial metabolism or reproduction, we believe
 191 that it represents a form of a "minimal model", where the pertinent features of the metabolic processes in-
 192 volved in reproduction are included, but the extraneous features are abstracted away. We have attempted
 193 to keep the number of free parameters to a minimum as to not overfit, while still being able to fit the
 194 data as accurately as models that do not have a basis in the fundamental biology. As such, this model
 195 represents an example what the authors feel is the appropriate abstraction of biological systems that
 196 renders them conducive to mathematical modelling and quantification while still retaining fundamental
 197 biological intuition.

References and Notes

1. Antonio A Alonso, Ignacio Molina, and Constantinos Theodoropoulos. Modeling bacterial population growth from stochastic single-cell dynamics. *Applied and environmental microbiology*, 80(17):5241–53, sep 2014.
2. J Baranyi, J Baranyi, T A Roberts, T A Roberts, P McClure, P McClure, Earley Gate, and Earley Gate. A non-autonomous differential equation to model bacterial growth. *Food Microbiology*, pages 43–59, 1993.
3. József Baranyi. Mathematics of predictive food microbiology. *International Journal of Food Microbiology*, 26(2):199–218, 1995.
4. József Baranyi and Terry A. Roberts. A dynamic approach to predicting bacterial growth in food. *International Journal of Food Microbiology*, 23(3-4):277–294, 1994.
5. Zachary D Blount, Christina Z Borland, and Richard E Lenski. Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *PNAS*, 105(23):7899–906, jun 2008.
6. Bryan C Daniels, Yan-Jiun Chen, James P Sethna, Ryan N Gutenkunst, and Christopher R Myers. Sloppiness, robustness, and evolvability in systems biology. *Current opinion in biotechnology*, 19(4):389–95, aug 2008.
7. Benjamin Gompertz. On the Nature of the Function Expressive of the Law of Human Mortality, and on a New Mode of Determining the Value of Life Contingencies. *Source: Philosophical Transactions of the Royal Society of London*, 115:513–583, 1825.
8. Joseph Horowitz, Mark D. Normand, Maria G. Corradini, and Micha Peleg. Probabilistic model of microbial cell growth, division, and mortality. *Applied and Environmental Microbiology*, 76(1):230–242, 2010.
9. J F Van Impe, T Martens, and J De Baerdemaeker. Dynamic mathematical model to predict microbial growth and inactivation during food Dynamic Mathematical Model To Predict Microbial Growth and Inactivation during Food Processing. 58(9):2901–2909, 1992.
10. H Jeong, B Tombor, R Albert, Z N Oltvai, and a L Barabási. The large-scale organization of metabolic networks. *Nature*, 407(6804):651–4, oct 2000.
11. H. Kacser and J. A. Burns. The molecular basis of dominance. *Genetics*, 97(3-4):639–666, 1981.
12. H Kacser and J a Burns. The control of flux. *Biochemical Society transactions*, 23(2):341–366, 1995.
13. Jonathan R. Karr, Jayodita C. Sanghvi, Derek N. Macklin, Miriam V. Gutschow, Jared M. Jacobs, Benjamin Bolival, Nacyra Assad-Garcia, John I. Glass, and Markus W. Covert. A Whole-Cell Computational Model Predicts Phenotype from Genotype. *Cell*, 150(2):389–401, jul 2012.
14. Y LEMARC, C PIN, and J BARANYI. Methods to determine the growth domain in a multi-dimensional environmental space. *International Journal of Food Microbiology*, 100(1-3):3–12, apr 2005.
15. R E Lenski, M R Rose, S C Simpson, and S C Tadler. Long-term experimental evolution in *Escherichia coli*. *The American Naturalist*, 138(6):1315–1341, 1991.

- 261 16. Armand M. Leroi, Richard E. Lenski, and Albert F. Bennett. Evolutionary Adaptation to Tem-
262 perature. III. Adaptation of *Escherichia coli* to a Temporally Varying Environment. *Evolution*,
263 48(4):1222, aug 1994.
- 264 17. A METRIS, Y LEMARC, A ELFWING, A BALLAGI, and J BARANYI. Modelling the vari-
265 ability of lag times and the first generation times of single cells of. *International Journal of Food*
266 *Microbiology*, 100(1-3):13–19, apr 2005.
- 267 18. Micha Peleg and Maria G. Corradini. Microbial Growth Curves: What the Models Tell Us and
268 What They Cannot. *Critical Reviews in Food Science and Nutrition*, 51(10):917–945, 2011.
- 269 19. F POSCHET, K VEREECKEN, A GEERAERD, B NICOLAI, and J VANIMPE. Analysis of a
270 novel class of predictive microbial growth models and application to coculture growth. *International*
271 *Journal of Food Microbiology*, 100(1-3):107–124, apr 2005.
- 272 20. E. O. POWELL. Growth Rate and Generation Time of Bacteria, with Special Reference to Con-
273 tinuous Culture. *Journal of General Microbiology*, 15(3):492–511, dec 1956.
- 274 21. D A Ratkowsky, R K Lowry, T A McMeekin, A N Stokes, and R E Chandler. Model for bacterial
275 culture growth rate throughout the entire biokinetic temperature Model for Bacterial Culture
276 Growth Rate Throughout the Entire Biokinetic Temperature Range. *Journal of Bacteriology*,
277 154(3):1222–1226, 1983.
- 278 22. DA Ratkowsky, J Olley, TA McMeekin, and A Ball. Relationship between temperature and growth
279 rate of bacterial cultures. *Journal of bacteriology*, 149(1):1–5, 1982.
- 280 23. Jon Schnute. A Versatile Growth Model with Statistically Stable Parameters. *Canadian Journal*
281 *of Fisheries and Aquatic Sciences*, 38(9):1128–1140, sep 1981.
- 282 24. O. Tenaillon, A. Rodriguez-Verdugo, R. L. Gaut, P. McDonald, a. F. Bennett, a. D. Long, and
283 B. S. Gaut. The Molecular Diversity of Adaptive Convergence. *Science*, 335(6067):457–461, jan
284 2012.
- 285 25. J F Van Impe, B M Nicolaï, M Schellekens, T Martens, and J De Baerdemaeker. Predictive
286 microbiology in a dynamic environment: a system theory approach. *International journal of food*
287 *microbiology*, 25(3):227–49, may 1995.
- 288 26. J VANIMPE, F POSCHET, A GEERAERD, and K VEREECKEN. Towards a novel class of
289 predictive microbial growth models. *International Journal of Food Microbiology*, 100(1-3):97–105,
290 apr 2005.
- 291 27. M H Zwietering, J T de Koos, B E Hasenack, J C de Witt, and K van’t Riet. Modeling of bacterial
292 growth as a function of temperature. *Applied and environmental microbiology*, 57(4):1094–101, apr
293 1991.
- 294 28. M H Zwietering, I Jongenburger, F M Rombouts, and K van ’t Riet. Modeling of the bacterial
295 growth curve. *Applied and environmental microbiology*, 56(6):1875–81, jun 1990.

$$\phi_1 = \frac{S_0 * V_{max,1}}{K_{M,1} + S_0}$$

$$\phi_2 = \frac{S_1 * V_{max,2}}{K_{M,2} + S_1}$$

$$\phi_{N-1} = \frac{S_{N-2} * V_{max,N-1}}{K_{M,N-1} + S_{N-2}}$$

$$\phi_N = \frac{S_{N-1} * V_{max,N}}{K_{M,N} + S_{N-1}}$$

<https://doi.org/10.1101/553982>

CC-BY 4.0 International license

Figure 1

a

b

Figure 2

Figure 3