GUT MICROBIOME ANALYSIS IN ADULT TROPICAL GARS (*Atractosteus tropicus*)

Roberto Méndez-Pérez^{1†}, Rodrigo García-López^{3,4†}, J. Santiago Bautista-López¹, Jorge F. Vázquez-Castellanos^{3,4}, Emyr S. Peña-Marín⁵, Rafael Martínez-García⁵, Verónica I. Domínguez-Rodríguez², Randy H. Adams-Schroeder², Eduardo Baltierra-Trejo^{2,6}, Carolina Melgar Valdés⁷, Andrés Moya^{3,4,✉}, Carlos A. Alvarez-González⁵, [Rodolfo Gómez-Cruz](#)^{1,✉}

[†] Equal contribution

¹ Biotech & Environmental Genomics and ² Biorremediation Laboratories. Academic Body Environmental Sciences (CACIAM). Research Center for Conservation and Sustainable Use of Tropical Resources ([CICART](#)). Biological Sciences Academic Division ([DACBioI](#)). Juárez Autonomous University of Tabasco State ([UJAT](#)), Villahermosa, Tabasco, Mexico. ³ [Department of Genomics and Health](#). Valencian Region Foundation for the Promotion of Health and Biomedical Research ([FISABIO](#)), Valencia, Spain. ⁴ Integrative Systems Biology Institute (I2Sysbio), University of Valencia, Valencia, Spain. ⁵ Tropical Aquaculture Laboratory. [CICART-DACBioI-UJAT](#), Villahermosa, Tabasco, Mexico. ⁶ [CONACyT-UJAT](#). ⁷ General Aquaculture Laboratory. Multidisciplinary Academic Division of the Rivers ([DAMRios](#))-[UJAT](#). Tenosique, Tabasco, Mexico.

✉ Corresponding authors: [Rodolfo Gómez – rodolfo.gomez@ujat.mx](#), Andrés Moya – [moya_and@gva.es](#)

Abstract. Tropical gar (*Atractosteus tropicus*), is freshwater and estuarine fish that has inhabited the Earth since the Mesozoic era, undergoing limited physiological variation ever since. This omnivorous fish is endemic to southern Mexico and part of Central America. Besides its recognized cultural and scientific relevance, the species has seen remarkable growth in its economic impact due to pisciculture. Previous studies have highlighted the role of microbial communities in fish, particularly those in the gut microbiome, in maintaining their host homeostasis or disease. In this study, we present the first report of the whole taxonomic composition of microbial communities in gut contents of adults' *A. tropicus*, by sex (female/male) and origin (wild/cultivated). Using culture-independent techniques, we extracted metagenomic DNA that was used for high throughput 16S rDNA profiling by amplifying the V4 – V5 hypervariable regions of the bacterial gene. A total of 364,735 total paired-end reads were obtained on an Illumina MiSeq sequencing platform, belonging to 508 identified genera, with the most and least abundant are *Cetobacterium*, *Edwardsiella*, *Serratia*, *Clostridium sensu stricto*, *Paludibacter* and *Campylobacter*, *Snodgrassella*, *Albirehodobacter*, *Lentilitoribacter*, respectively. We detected that, by sex and origin, Proteobacteria, Fusobacteria, Firmicutes and Bacteroidetes phyla are the core gut microbiome of the adults' *A. tropicus*. We discover the Deinococcus-Thermus phylum sequence, wildtype males only, with extremophile capacity in another freshwater fish. We also identified the species *Lactococcus lactis* strains CAU929 and CAU6600, Cp6 and CAU9951, *Cetobacterium* strain H69, *Aeromonas hydrophila* strains P5 and WR-5-3- 2, *Aeromonas sobria* strain CP DC28 and *Aeromonas hydrophila* with probiotic potential in aquaculture within the three dominant phyla, especially in wild-type organisms.

Keywords: *Atractosteus tropicus*, gut microbiome, metagenomics, 16S rRNA profiling

1. Introduction. Aquaculture produces 76.6 million tons of fish for human consumption and economically ensures the livelihood of 10 to 12 % of the world population (FAO 2014, 2017). It has become the fastest growing food sector in the world, with an average annual growth rate of 8.9 % since 1970 (Subasinghe 2005) and there is a growing global trend towards diversifying the spectrum of cultured aquatic species. Biological diversity in Latin America is one of the richest on

the planet, including an important variety of its freshwater ichthyofauna (Flores-Nava y Brown 2010). Some of the greatest challenges for aquaculture during this millennium have been the creation of integral studies on pisciculture-relevant endemic species and the development of technologies that may allow for controlled production of these fish in a profitable, innocuous and environmentally-conscious approach (Márquez-Couturier y Vázquez-Navarrete 2015). Studies in other fish have explored the bacterial populations in varying habitats such as the skin, gills, eggs and gut microbiome (GMB) and the way they influence the host's general health and physiology (MacFarlane *et al.* 1986, Cahill 1990, Ringø & Tabachek 1995, Givens 2012, Austin & Austin 2016). They have reported a large variation of microbiota among the different niches and between species with the GMB as one of the most studied due to its high microorganism concentration. Of interest for aquaculture-relevant species are outbreaks of viral, bacterial and fungal infections as they may cause devastating economic losses worldwide due to poor environmental conditions in the farms, unbalanced feeding, generation of toxins and genetic factors (Martínez-Cruz *et al.* 2012).

Tropical gar (*Atractosteus tropicus*, also known as pejelagarto) is freshwater and brackish fish species of the Lepisosteidae family, which has an ample fossil record since the Cretaceous period of the Mesozoic Era (Wiley 1976, Reséndez-Medina & Salvadores-Baledón 1983). The morphology of these species has remained mostly unaltered, with current specimens having lengths between 1.0 and 1.2 m and weighing between 1000 and 3000 g in the wild. In their natural habitat, this species exists in coastal wetlands of the tropical rainy areas of southeastern Mexico, Belize, Guatemala, El Salvador, Honduras, Nicaragua and Costa Rica (Wiley 1976, Bussing 1998, Miller *et al.* 2005, Nelson 2006). It is an omnivore, feeding on other fish, decomposing organic matter, crustaceans, plants, etc., depending on the availability, preferring carnivorous habits. This specially secluded species has seen a drastic decrease in wild populations, caused by anthropogenic activities that have led to the loss of habitats and severe ecological alterations (Méndez-Marín *et al.* 2012). Currently, *A. tropicus* is cultivated in fish farms for human consumption in Mexico (Márquez *et al.* 2015).

The gastrointestinal tract of *A. tropicus*, is formed by the buccopharyngeal, esophagus, stomach, gut, pyloric blind, rectum and anus, which is rapidly developing during the larval period (Frias-Quintana *et al.* 2015). Having absorbed and secretory function, its intestine is a long tube, narrower than the stomach with an inner epithelial mucosa that forms long, and numerous folds comprised of high columnar cells with clearly defined brush border (Márquez-Couturier *et al.* 2006). Having absorbed and secretory function, its intestine is a long tube, narrower than the stomach with an inner epithelial mucosa that forms long, and numerous folds comprised of high columnar cells with clearly defined brush border (Márquez-Couturier *et al.* 2006). In the anterior and middle regions, long folds divide these regions into a series of compartments called a spiral valve. The shorter posterior region forms the rectum, which opens in the anus. The external muscle in the intestine is thinner than in the stomach. Early juveniles 20 days after hatching show a GMB like that of adults. Before colonization, microorganisms may access the GMB through food and water intake during the larval phase (Núñez de la Rosa 2011). The high abundance of certain groups of gastrointestinal bacteria in fish, when compared to the microbial composition of the surrounding water suggests that GMB poses as a unique niche for a specific, but diverse group of bacteria (Cahill 1990, Givens 2012). It has been reported that GMB fish has between 10^7 and 10^{11} bacteria per gram of feces (Nayak 2010). GMB fish plays an important role and directly influences the host's nutrition and general homeostasis. Normal GMB fish may contain both beneficial and potentially pathogenic bacteria. The loss of the microbiota equilibrium (dysbiosis) has been reported to impact the host's physiological state, potentially compromising immunity, growth, general development as well as

the overall quality of the aquaculture production due to an increase in fish morbidity and mortality (Núñez de la Rosa 2011, Al-Harbi and Udding, 2005).

Prior to 2005, microbial studies on fish relied exclusively on culture techniques for enumerating and identifying bacteria (Newman *et al.* 1972, MacFarlane *et al.* 1986, Spanggaard *et al.* 2000, Aschfalk & Müller 2002, Verner-Jeffreys *et al.* 2003, Al-Harbi & Uddin 2004, Martin-Antonio *et al.* 2007, Skrodenyté 2007), providing valuable information as isolates provide a suitable framework for studying individual culturable strains on a finer level but limited appreciation of the whole spectrum in microbial communities. In 2007, Izvekova and collaborators reviewed GMB fish studies published between 1929 and 2006 reporting 73 different bacteria. This is because culturable bacteria typically account for less than 1 % of the cells that are present by direct microscopic enumeration (Ferguson *et al.* 1984, Head *et al.* 1998).

The advent of culture-independent metagenomic studies during the mid-2000s enabled the simultaneous analysis of complex genomic information contained in a hundreds of microbial species in a single niche (Nielsen *et al.* 2014). These techniques circumvent most culturing requirements of microorganisms, avoiding collection and sampling biases, effectively representing the actual diversity of a microbial community. Under intensive production conditions for sustainable aquaculture, aquatic species are subjected to high-stress conditions, leading to an increased incidence of diseases that decrease productivity (Bondad *et al.* 2005). It has been proposed that microbiota dysbiosis may be avoided through the regulation of their microbiota (Verschuere *et al.* 2000).

Microbial studies in aquaculture focus on the understanding of the symbiotic or antagonistic interactions between microbes and their eukaryotic hosts such as fish, crustaceans and molluscs. In this sense, metagenomics can provide a deeper understanding of these relationships through information revealed by sequencing microbial DNA extracted from specific niches within host organisms and, in the case of 16S profiling, taxa are representative of the medium (Suttle 2007, Gianoulis *et al.* 2009). This latter consists of surveying the 16S rRNA gene of all present microorganisms as this marker is found in all prokaryotes with enough mutations to discern each taxon. Formerly, some bacteria had been difficult to isolate because some of them are obligate intracellular microorganisms that could only be cultured in semi-aqueous and/or cellular culture media (Avila-Villa *et al.* 2011). Current sequencing platforms and bioinformatic tools enable the research on the diversity of intracellular bacteria, but also to ignore culture other culture requirements altogether as the DNA from the whole community are sequenced as is, with majority and accessory species, in order to elucidate the relevant community configurations for the improvement of aquaculture techniques. Consequently, the objective of this research was to explore the bacterial composition in the gut of adults' *A. tropicus*, by analyzing the 16S rRNA gene profiles from adult male and female organisms, cultivated and wild, for biotechnological-relevant applications future.

2. Materials and methods

Specimen collection. Fifteen totals live adults' *A. tropicus* were collected for the study, 7 of them were cultivated in the Tropical Aquaculture Laboratory, [Research Center for Conservation and Sustainable Use of Tropical Resources](#) (CICART) at [Biological Sciences Academic Division](#) (DACBiol), [Juárez Autonomous University of Tabasco State](#) (UJAT), Mexico, and 8 wildtype, with an average weight and length of 5 kg and 1 m, respectively. This specimens were provided by fishermen of the municipalities of Nacajuca ([18°14'50"N 92°49'58"O](#); n female, m male) and Centla ([18°20'00"N 92°30'00"O](#); n female, m male), Tabasco, Mexico (Figure 1). All specimens were cut

lengthwise under sterile conditions to remove the intestine for extracting its contents with scissors disinfected in absolute ethyl alcohol, for storage at -20 °C in a 2 mL Eppendorf tubes.

Samples collection. All organisms were sacrificed according to the protocol was approved by Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación (NOM-062-ZOO-1999) on 18 June 2001 by percussion stunning method, consisting of striking a quick blow on the head of the fish manually with a bat, to obtain fresh samples, squeezing the GMB and obtaining the feces separately, considering the geographical origin and sex by sterile conditions.

DNA extraction, pyrosequencing and sequence for metagenomic analysis. 200 mg of fresh feces were used and 2 mL were enough to ensure the target DNA with a capacity of the Eppendorf tubes selected. A whole genomic DNA (gDNA) extraction was carried out for each sample with a QIAamp DNA Stool Mini Kit (Qiagen, Valencia, CA, USA), following the manufacturer's instructions. DNA integrity and concentration was evaluated by electrophoresis submerged with 1.2 % agarose gel and by spectrophotometry with a GenovaNano spectrophotometer (Jenway, Stone, Staffs, UK), respectively. Universal primers 533F (5'-GTGCCAGCAGCCGCGGTAA-3') (Weisburg 1991) and 909R (5'-CCCCGYCAATTCMTTTRAGT-3') (Tamaki *et al.* 2011) were selected for the amplification a region including the V4 and V5 hypervariable region in the 16S rRNA gene. PCR amplifications were carried out using Phusion High-Fidelity DNA Polymerase (Finnzymes OY, Espoo, Finland; Klindworth *et al.* 2013) and conditions were as follows: Library preparation and high throughput sequencing was performed through Research & Testing Labs (Lubbock, Texas, USA) services, with an Illumina MiSeq platform, using 16S rDNA profiling by amplifying the V4 – V5 hypervariable regions of the bacterial gene, reagents sequencing for 2x 300 bp paired-end reads.

Bioinformatic analysis. The sequences were subjected to the standard quality protocols, which included the sequencing adapters removal with the Cutadapt tool (1.18), the too short and low-quality sequences filtering based on the Phred quality score implemented in the PRINSEQ tool.lite (0.20.4). The paired-end were concatenated with -fastq_join command of the USEARCH program (V.11) and the chimeras were removed with -uchime2_ref command of the same program, using as reference the Gold database (microbiomeutil-r20110519). The sequences were clustered in OTUs with 97 % identity by usearch61_ref algorithm, using the SILVA132 database as a reference through the pick_otus.py command in QIIME (1.9) and OTUs were filtered with 3 or fewer sequences.

Alpha diversity was measured through the Chao1, Shannon and P. D. (Phylogenetic Diversity) diversity indexes, which were calculated by alpha_rarefaction.py command in QIIME (1.9). The Alpha diversity among samples and origin of the samples (wild and farmed) were compared using Kruskal-Wallis statistic (Multiple range test, Bonferroni post-hoc) and Mann-Whitney W statistic (Wilcoxon), respectively, both implemented in the ggpubr library of R software (V.3.4.4).

Beta Diversity was analyzed using a PCoA constructed from the Unweighted UniFrac distance, calculated with the beta_diversity_through_plots.py command from QIIME (1.9), the difference among groups was evaluated by ANOSIM statistical test, and the main coordinates of the PCoA were graphed in the factoextra library in R (V.3.4.4) and Differential Analysis of Abundance was compared the abundance patterns of the taxa between the groups of samples through LefSe (Linear discriminant analysis effect size).

Phylogenetic analysis. A sequences base was constructed according to certain genera with probiotic potential in fish that we find in the literature sought (Table 1), 16S rRNA gene sequences were obtained from the NCBI GenBank database. The 6,266 sequences obtained were clustered to eliminate identical sequences and generate a table of OTUs with 97 % similarity, using the

usearch61 algorithm of QIIME (1.9). The database was indexed (Query Sequence) to be used in a local blast with the BLAST + tool (NCBI). The local blast was performed using the sequences of the samples (Subject sequences) clustered at 97 % as mentioned in the section on the processing of the sequences. The identity percentage was adjusted to 99 % (-perc_identity) and a coverage of 70 % (-qcov_hsp_perc 70). For phylogenetic analysis, the blast resulting Query and Subject sequences were subjected to a multiple alignment with the ClustalW algorithm, in MEGAX (Molecular Evolutionary Genetics Analysis), and the phylogenetic relationships were inferred by the Neighbor-joining method in MEGAX, with the predefined settings.

3. Results. The pyrosequencing method of the Illumina MiSeq platform was evaluated that uses the Research & Testing Labs (Lubbock, Texas, USA) services, applying 16S rRNA in V4 – V5 hypervariable regions. A total of 364,735 sequencing reads were generated, organized by sex and origin, according to Table 2. We can visualize in Table 2 that readings obtained of wildtype male organisms are far greater than of wildtype and cultivated females, and cultivated males. Sequences were clustered into 16,503 different OTUs (97 % identity), which were classified into 11 phyla, 22 class, 37 order, 86 families and 179 genera. Table 2 shows that about twice as many reads correspond to wild male.

Microbiota composition. The cultivated organisms, wildtype females and males' samples are dominated by the *Fusobacteria* (37.32 %), *Proteobacteria* (30.49 %) and *Firmicutes* (18.55 %) phyla, respectively (Figure 2a). In wildtype females the most abundant phyla were *Proteobacteria* (0.70 ± 0.28 SD), *Firmicutes* (0.38 ± 0.32 SD) and *Fusobacteria* (0.24 ± 0.29 SD) and in cultivated females are *Proteobacteria* (0.64 ± 0.20 SD), *Fusobacteria* (0.54 ± 0.25 SD) and *Bacteroidetes* (0.45 ± 0.006 SD). In wildtype and cultivated males, the most abundant phyla are *Fusobacteria* (0.63 ± 0.39 SD), *Proteobacteria* (0.40 ± 0.26 SD) and *Fusobacteria* (0.76 ± 0.36 SD), *Proteobacteria* (0.19 ± 0.21 SD), respectively. *Actinobacteria* (0.16 ± 0.17 SD) and *Deinococcus-thermus* (0.003 ± 0.006 SD) were identified only in wildtype females and wildtype males, respectively. The same tendency is observed in male and female specimens, since the dominant phyla in both wild type and cultivated ones are *Fusobacteria* (0.63 ± 0.39 SD), *Proteobacteria* (0.70 ± 0.28 SD) and *Fusobacteria* (0.76 ± 0.36 SD) and *Proteobacteria* (0.64 ± 0.20 SD), respectively (Figure 2a). At the genus level, the most abundant in wildtype females are *Edwardsiella* (0.32 ± 0.53 SD) and *Cetobacterium* (0.24 ± 0.29 SD), and in cultivated females are *Serratia* (0.57 ± 0.17 SD) and *Cetobacterium* (0.54 ± 0.25 SD). In wildtype males, the most abundant genera are *Cetobacterium* (0.63 ± 0.39 SD) and *Clostridium sensu stricto* (0.32 ± 0.49 SD) in cultivated males *Cetobacterium* (0.75 ± 0.29 SD) and *Edwardsiella* (0.22 ± 0.30 SD) (Figure 2b). In the case *Cetobacterium* (37.34 %) and *Edwardsiella* (10.55 %) genera are the most abundant in cultivated organisms and wildtype females' samples, respectively (Figure 2b); whereas *Clostridium sensu stricto* (13.53 %) was more abundant in wildtype males' samples. The cultivated and wildtype organisms' samples are dominated by the *Fusobacteria* (38.95 %) and *Firmicutes* (19.57 %) phyla, respectively (Figure 2a). For the *Cetobacterium* (38.95 %) and *Clostridium sensu stricto* (15.09 %) genera they are more abundant in the cultured organisms and samples of wildtype organisms' samples, respectively (Figure 2b).

Likewise, the core microbiome composition at phylum and genus taxonomic levels, on a per-sample basis of adults' *A. tropicus* gut (figure 3), shows that the most abundant phyla per sample were *Fusobacteria* (47.01 %), *Proteobacteria* (29.02 %) and *Firmicutes* (12.90 %) (Figure 3a), while the most abundant genera were *Cetobacterium* (47.01 %), *Edwardsiella* (12.00 %) and *Serratia* (10.55 %) (Figure 3b). Although the primers are designed only to 16S rRNA gene amplify of bacteria, OTUs belonging to the archaea domain were also identified, represented by the genera *Salinirubrum*, *Salinigranum* and *Methanoculleus*, which together represent about 0.01 % of the total abundance.

Alpha diversity. In this work, diversity, dominance and richness were evaluated using the Shannon index about the total number of OTUs. Microbiota composition is influenced by source and sex

factors. The Shannon index indicates that the microbial types in adults' *A. tropicus* are more evenly distributed and thus may be more diverse in wild (5.32 ± 0.13 SD) and cultivated (3.75 ± 0.09 SD) female individuals than in the wild (3.14 ± 0.07 SD) and cultivated (3.38 ± 0.08 SD) male individuals (Table 3). We found significant differences in diversity indices and only some samples were not significant. The wild and cultivated females turned out to be the most diverse, but by origin, only the wild samples were the most diverse. In a total of 50 taxa, significant differences can be observed between cultivated and wild conditions.

Beta diversity. Bacterial communities beta diversity associated to TGI of adults' *A. tropicus* in origin conditions were measured through the ordination analysis from Principal Coordinates Analysis (PCoA), using Unweighted unifrac distances (UniFrac). The analysis produced an ordination of the dissimilarities, where similar individuals are close to one another and dissimilar ones are more distant (Figure 4). All ordination analysis showed a clear separation between wild and cultivated individuals samples. ANOSIM statistic (R: 0.60933; p-value <0.002) found significant statistical differences by origin, between the cultivated and wild groups (Figure 4).

Significant microbial components associated with the origin and sex. 8 genera were identified in wild organisms and these are *Methanoculleus*, *Flavobacterium*, *Psychrobacter*, *Acinetobacter*, *Pseudomonas*, *Paracoccus*, *Massilia* and *Shewanella*. Likewise, 13 genera were identified in cultivated organisms, which are *Paludibacter*, *Intestinibacter*, *Cellulosilyticum*, *Odoribacter*, *Turicibacter*, *Deffluviitalea*, *Vallitalea*, *Acetivibrio*, *Terrisporobacter*, *Bacteroides*, *Acidaminobacter*, *Sporacetigenium* and *Macellibacteroides*.

Specific taxa distributed differentially between wild and cultivated organisms were identified by LEfSe tool. This allows to obtain statistical differences per each taxon, where linear discriminant analysis (LDA) score barplot is shown for both conditions, cultivated and wild organisms (female & male; figure 5a), cultivated and wild organisms (female; figure 5b), male organisms, cultivated & wild (Figure 6a), cultivated female organisms (Figure 6b) and wild female organisms (Figure 6c), representing each phylum, class, order, family and genus by a histogram.

Phylogenetic reconstruction analysis. The sequences-base of rRNA 16S gene constructed of some organisms found in the literature with probiotic potential in fish, according to table 1, allowed us to find and identify possible organisms with probiotic potential in microbiome gut of adults' *A. tropicus*, adjusted to 99% identity and 70% coverage. In figure 7 we obtained a phylogenetic tree based on the identified sequences of the most abundant phyla per sample, such as Fusobacteria, Proteobacteria and Firmicutes. Besides, we *Methanosarcina thermophila* and *Archaeoglobus profundus* sequences data used thermophilic methanogens and sulfate-reducing archaea, respectively, as outgroups for the root identification, considering that they may be related only distantly with identified sequences in our work.

In phylogenetic tree reconstruction (Figure 7) we determined that the evolutionary conclusion of these relationships is that the two species of archaea or outgroups selected are ancestors of the nine species or ingroups identified with blue squares. Likewise, they belong to the three most dominant of phyla groups that integrate the core microbiome gut in adults' *A. tropicus*. On the other hand, these nine species, identified by blue squares, are those that have probiotic potential.

4. Discussion. We obtained the 16S rRNA Gene Amplicons preparation with high-throughput sequencing using Illumina MiSeq platform by Research & Testing Labs (Lubbock, Texas, USA) services, facilitating discovery and analysis of rare taxa and detection of previously unrecognized eukaryotic and prokaryotic microbiome (Paul *et al.* 2018). In this sense, we identified and compared, by indexes attachment of multiplexing samples and readings of sequencing products for 2x300 bp

paired-end, the core microbiome and microbial diversity, respectively, in the gut of adults' *A. tropicus* by origin and sex.

Previous studies have shown that the fish gut hosts an estimated 10^7 to 10^{11} bacteria per gram intestinal content (Nayak 2010) and the bacterial colonizers in fish gut include Proteobacteria, Fusobacteria, Firmicutes, Bacteroidetes, Actinobacteria, Clostridia, Bacilli and Verrucomicrobia (Ringø *et al.* 2006, Desai *et al.* 2012, Li *et al.* 2013a,b, Carda-Diéguez *et al.* 2014, Ingerslev *et al.* 2014a,b) with the first 4 being the most abundant, depending on environmental conditions and the host's diet (Wang *et al.* 2017). Almost all Proteobacteria, Fusobacteria, Firmicutes, Bacteroidetes and Deinococcus-Thermus 16S rRNA sequences that we detect in CGMB of adults' *A. tropicus* belonged to the *Cetobacterium*, *Edwardsiella*, *Serraria* and *Clostridium sensu stricto* genres. The observed bacterial profiles in the gut of adults' *A. tropicus*, by sex and origin, may reflect specific central microbiota, beyond the differences most likely attributable to feeding behavior. At the phylum level, almost 90 % of the total bacterial abundance only we were classified into total four phyla. Among these phyla, Proteobacteria and Fusobacteria were dominant in the fifteen fish species samples, female/male and cultivated/wild (Figs. 2, 3). The inferred physiological roles of the dominant prokaryotes are related to the metabolism of carbohydrates and nitrogenous compounds (Kormas *et al.* 2014). In contrast, dominant microbiota of marine fish is facultative anaerobes, including *Vibrio*, *Pseudomonas*, *Acinetobacter*, *Corynebacterium*, *Alteromonas*, *Flavobacterium* and *Micrococcus* (Onarheim *et al.* 1994, Blanch *et al.* 1997, Verner-Jeffreys *et al.* 2003).

Considering the above-mentioned, fish core gut microbiota (CGMB) can influence nutrition, growth, reproduction, general population dynamics and the host's vulnerability to diseases thus supporting a crucial role in aquaculture practice (Ghanbari *et al.* 2015). Current DNA sequencing technologies and bioinformatic analysis have contributed towards a deeper understanding of the complex microbial communities associated to diverse habitats, including CGMB of fish in response to a variety of factors affecting the host, including temperature variations, salinity, growth stage, digestive physiology and feeding strategy (Cahill 1990, Jammal *et al.* 2017). The concept 'core microbiota' referred to a set of abundant microbial lineages that are shared by all individuals from the same species (Wong *et al.* 2013). The concept of CGMB has been explored both in mammalian host's context and in freshwater fish (Turnbaugh *et al.* 2009, Roeselers *et al.* 2011, Nam *et al.* 2011, Wu *et al.* 2012, Wong *et al.* 2013).

On the other hand, Alpha diversity results indicate that the greatest gut microbiota abundance and richness is found in adults' *A. tropicus* wildtype female, rather than adult's wildtype males and the least gut microbiota abundance and richness is found in adults *A. tropicus*, female or male cultivated (Table 3). Ley *et al.* (2008) concluded that gut microflora of herbivorous mammals have the greatest richness and phylogenetic diversity and that both richness and phylogenetic diversity decreased among omnivores and decreased further among carnivores. We found the lowest richness and phylogenetic diversity (Table 3) in gut microbiomes from adults *A. tropicus* defined as top piscivores (carnivores; e.g., cultivated female and male *A. tropicus*). Indeed, MacFarlane *et al.* (1986) observed that farmraised fish had a simpler gut flora than their wild counterparts.

Likewise, our coefficients of similarity or dissimilarity of unweighted unifracs distances (UniFrac) by PCoA ordination analysis, indicate that gut microbiota of wild and cultivated adults *A. tropicus* are only similar by two wild samples and one cultivated and dissimilarity by most of the wild and cultivated samples (Figure 4). Principal coordinate analysis (PCoA) revealed that gut bacterial communities from adults' *A. tropicus* by origin formed different clusters. Cultivated and wild organisms formed distinctly clusters in PCoA space (Figure 4), suggesting that the enrichment and diversity of gut microbiota are affected by the origin. This result is similar to the research of Ni *et*

al. (2012) that the origin and host phylogeny they are quite related to the composition of adults' *A. tropicus* gut bacteria. Moreover, previous studies have shown that the microbiotic diversity content in all intestinal sections depends on the fish size (Moran *et al.* 2005, Bolnick *et al.* 2014, Clements *et al.* 2014).

Numerous works have built on this, demonstrating that many species of herbivorous and omnivorous fishes contain diverse intestinal communities (Rimmer & Wiebe 1987, Clements *et al.* 1989, Clements 1991, 1997, Martínez-Díaz & Pérez-España 1999, Ray *et al.* 2012) and that herbivorous and detritivorous fish species harbour distinctive microbial populations (Clements *et al.* 2014). Feeding habits are also an important factor that generally influences the microbial diversity in the fish CGMB, displaying a higher diversity in the following order: carnivores> omnivores> herbivores (Ward *et al.* 2009, Larsen *et al.* 2014, Li *et al.* 2014a,b, Miyake *et al.* 2015, Liu *et al.* 2016). He *et al.* (2013) study revealed that the herbivorous carp (*Ctenopharyngodon idellus*) reported a wider variety of bacterial species than the dark carnivorous carps and Gibel (*Carassius gibelio*) which are exclusively omnivorous, and also the sea bream, under the same culture conditions. This same tendency we identify in gut core microbiome of adults *A. tropicus* by sex and origin, because although these are omnivores, they prefer carnivorous habits. (Méndez-Marín *et al.* 2012, Figure 5a). At the sex level, we observed that core microbiome is more diverse in female organisms, and particularly in wild type organisms (5b, 6a,b)

More recent sequence-based approaches show that fish hindgut microbial communities much more closely resemble those of mammals than environmental microbial communities (Fidopiastis *et al.* 2006; Sullam *et al.* 2012), especially in the prevalence of Proteobacteria, Firmicutes and Bacteroidetes (Clements *et al.* 2007, Smriga *et al.* 2010, Sullam *et al.* 2012, Ye *et al.* 2014). These findings indicate that fish, like other vertebrates, harbour specialized gastrointestinal communities (Clements *et al.* 2014). We identified Proteobacteria as the most dominant phylum in gut core microbiome of the adults' *A. tropicus*. According to Rudi *et al.* (2018), Proteobacteria phylum is very characteristic in wildtype fish and denotes of a diet high in fat presence.

We identified Fusobacteria as the second dominant phylum in the gut of the male adults *A. tropicus*, wild and cultivated (Figure 2a). A few studies have shown Fusobacteria as dominant members of the gut microbiota of freshwater fishes (van Kessel *et al.* 2011, Di Maiuta *et al.* 2013). Fusobacteria is anaerobic, Gram-negative bacilli that produce butyrate (Bennett and Eley 1993), a short-chain fatty acid that is often the end product of the fermentation of carbohydrates including those found in mucins (Titus and Ahearn 1988, von Engelhardt *et al.* 1998). In mammals, butyrate provides many benefits to the host, including providing a majority of the energy supply to gastrointestinal cells (von Engelhardt *et al.* 1998, Collinder *et al.* 2003), enhancing mucus production, acting as an anti-carcinogen and anti-inflammatory, as well as playing a role in satiation (McBain *et al.* 1997, von Engelhardt *et al.* 1998, Andoh *et al.* 1999, Hamer *et al.* 2007). This fatty acid has been found in the gut of herbivorous and omnivorous fishes only (Clements *et al.* 1994, Clements and Choat 1995). Nuez-Ortín *et al.* (2012) demonstrated that the ability of butyric acid to inhibit potential freshwater fish pathogens, and sodium butyrate is currently sold as a food additive to promote fish health and growth. However, trials using blends of sodium butyrate and other additives have not proven beneficial (Owen *et al.* 2006, Gao *et al.* 2011).

OTUs sequences of the *Cetobacterium* genus were identified as dominant mainly in gut microbiome of the adults' *A. tropicus*, wild and cultivated male organisms. *Cetobacterium* genus is widely identified in freshwater and warm water fish species (Tsuchiya *et al.* 2007, Larsen *et al.* 2014, Li *et al.* 2017). *Cetobacterium* genus members' presence can perform fermentative metabolism of peptides and carbohydrates and produce vitamin B12 (cobalamin) (Larsen *et al.* 2014). *Cetobacterium* and

Bacteroides were reported as major producers of the vitamin B12 in the intestine (Tsuchiya *et al.* 2008, Vogiatzoglou *et al.* 2009) and they were the dominant genera in grass carp's intestine, with the abundance of more than 50 % (Li *et al.* 2015a). Animals, plants and fungi are incapable of cobalamin production and it is the only vitamin that is exclusively produced by microorganisms, particularly by anaerobes (Roth *et al.* 1996, Martens *et al.* 2002, Smith *et al.* 2007). Qi *et al.* (2017) showed that different concentrations of ammonia would affect the abundance of *Bacteroides* and *Cetobacterium* in gut fish, and the higher the concentration, the lower the abundance.

Despite we detected OTUs of Deinococcus-Thermus bacterial phyla non-dominant (0.003 ± 0.006 SD) in gut microbiome of the adults' *A. tropicus*, wildtype males only. However, Deinococcus-Thermus species are known for their resistance to extreme stresses, such as radiation, oxidation, desiccation and high temperature (Li *et al.* 2015b). The deeply branching Deinococcus-Thermus lineage is recognized as one of the most extremophilic phylum of bacteria (Theodorakopoulos *et al.* 2013). Sequence information from Deinococcus-Thermus phylum is presently available for only a limited number of species. However, the sequenced genomes include species from both the main families (i.e., *Deinococcaceae* and *Thermaceae*) within this phylum (Griffiths & Gupta 2007). In recent years, researchers have begun using Deinococcus spp in biotechnologies and bioremediation due to their specific ability to grow and express novel engineered functions. More recently, the sequencing of several Deinococcus spp and comparative genomic analysis have provided new insight into the potential of this genus. Features such as the accumulation of genes encoding cell cleaning systems that eliminate organic and inorganic cell toxic components are widespread among Deinococcus spp. Other features such as the ability to degrade and metabolize sugars and polymeric sugars make Deinococcus spp. an attractive alternative for use in industrial biotechnology (Gerber *et al.* 2015). That is why their functional role in the gut of the adults' *A. tropicus* deserves further research.

Bacteroidetes is a phylum composed of three large classes of Gram-negative, nonsporeforming, anaerobic or aerobic, and rod-shaped bacteria that are widely distributed in the environment, including in soil, sediments, and seawater, as well as in the guts and on the skin of animals (Ley *et al.* 2008), this was also present as the dominant phylum in adults *A. tropicus* female, wild and cultivated, usually; only this type of bacteria was detected as dominant in sample 4, cultivated male organism (Figure 2a). Likewise, a large part of the proteins synthesized by the genome of *Bacteroides*, a genus of Bacteroidetes, can break down polysaccharides and metabolize their sugars, playing a fundamental role in the degradation of complex molecules in the gut of the host. Their ability to harvest alternative energy sources from food could allow *Bacteroides* to be more competitive than other bacteria in CGMB of fish during starvation stage (Xu *et al.* 2003, Xia *et al.* 2014).

We identify *Clostridium sensu stricto* genus sequences in GMB of the adults' *A. tropicus*, wildtype males only. This genus has also been identified in GMB of carp fish (Li *et al.* 2015a). The members of the genera *Clostridium sensu stricto* that are dominant in the intestinal microbiota of grass carp (*Ctenopharyngodon idellus*), they are also versatile in their ability to utilise various polysaccharides, such as cellulose, xylan and hemicelluloses, which constitute the major part of vegetal fibres (Uffen 1997, Uz & Ogram 2006, Li *et al.* 2015a). Others also include not only species with saccharolytic and fiber-fermenting activities but also proteolytic species (Lubbs *et al.* 2009, Pikuta *et al.* 2009, Li *et al.* 2015a).

Serratia, *Edwardsiella*, *Plesiomonas* and *Reyranella* are genera belonging to Enterobacteriaceae family and Proteobacteria phylum. This isolated bacterial species are facultative pathogens for fish and humans and may be isolated from fish without apparent symptoms of the disease (Walczak *et al.*

2017). *Serratia* produces serrawettin that acts as a wetting agent to reduce the surface tension of the environment (Chan *et al.* 2013). *Edwardsiella* has been isolated from tortoises (Iveson 1971), crocodiles (Iveson 1971), aquarium water (Bartlett *et al.* 1977) and from seagull roosting areas (Berg & Anderson 1972). *Edwardsiella* was isolated on several occasions during the examination of dressed catfish for *Salmonella* (Wyatt *et al.* 1979). This report provides information on the isolation, identification, and incidence of *Edwardsiella* in freshwater catfish and their environment. There are many unanswered questions regarding their importance in freshwater fish. Liu *et al.* (2015) are reports of pathogenicity of *Plesiomonas shigelloides* to fish. *P. shigelloides* can occur as natural intestinal flora of fish, but in case of stress conditions, the following symptoms are observed: darkening of body, hemorrhaging, fin rotting, ascitic fluid in the abdominal cavity, and lesions in internal organs (Liu *et al.* 2015). Phylogenetically, *Reyranella* genus has an evolutionary lineage within the family Rhodospirillaceae in the class Alphaproteobacteria. The type species *Reyranella massiliensis* was originally identified by Pagnier *et al.* (2011). Subsequently, Kim *et al.* (2013) emended several characteristics (e.g., nitrate reduction, respiratory quinone information) into the genus description, and more recently, Cui *et al.* 2017 found that reduction of nitrate to nitrite is variable, the predominant isoprenoid quinone is ubiquinone-10 (Q-10), major polar lipids are PME, DPG, PG, PE and one unknown aminolipid. We detect these genera mainly in gut microbiome of the adults' *A. tropicus*, wildtype females and cultivated females and males organisms (Figures 2b, 3b). We assume that, although commonly known for their ability to cause deadly infectious diseases, this is populations of bacteria (identified as normal flora of the adults *A. tropicus*), which despite symbiotically living on and between wildtype females and cultivated females and males organisms of the adults' *A. tropicus*, really have a positive impact on host survival (Figures 2b, 3b).

Paludibacter genus was found exclusively in African microbiota. This is probably due to their increased fitness to grow on polysaccharides abundant in xylan or cellulose diets (De Filippo *et al.* 2010, Thomas *et al.* 2011). We identified OTUs of *Paludibacter* genus in core gut microbiome of the adults' *A. tropicus*, females and males cultivated organisms, only (Figure 2b).

In phylogenetic trees reconstruction (Figure 7) and according to Table 1, we identified the species *Lactococcus lactis* strains CAU929 and CAU6600, Cp6 and CAU9951, *Cetobacterium* strain H69, *Aeromonas hydrophila* strains P5 and WR-5-3- 2, *Aeromonas sobria* strain CP DC28 and *Aeromonas hydrophila* with probiotic potential within the three dominant phyla in core gut microbiome of the adults *A. tropicus*.

5. Conclusions. We conclude that adults' *A. tropicus* core gut microbiome is constituted by Proteobacteria, Fusobacteria, Firmicutes and Bacteroidetes phyla. Definitely, diversity and richness in core gut microbiome are higher of the female than males' organisms and wild than cultivated organisms of the adults' *A. tropicus*. We can also suppose that *Serratia*, *Edwardsiella*, *Plesiomonas* and *Reyranella* genera lives in symbiosis and has a positive impact on the survival of the adults' *A. tropicus*. There is great potential in the biotechnology industry with *Deinococcus-Thermus* phylum, due to its extremophile capacity. Further, adults' *A. tropicus* by sex and origin have been adapting to the environment and the diet due to their great capacity to saccharolytic, fiber fermentation and starvation activities. Finally, we can confirm that nine species can be found with the probiotic potential of 3 phyla that shape up the core gut microbiome of the adults *A. tropicus*. The CGMB of *A. tropicus* is increasingly regarded as an integral component of the host, due to important roles in the modulation of the immune system, the proliferation of the intestinal epithelium and the regulation of the dietary energy intake. Understanding the factors that influence the composition of these microbial communities is essential to health management, and the application to aquatic animals still requires basic investigation.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgments

The authors thanks the National Council of Science and Technology of Mexico for the thesis scholarship for postgraduate studies and the mixed scholarship granted to carry out the research stay at the University of Valencia (Spain). This research received external funding from the National Council of Science and Technology of Mexico “Strengthening of the Master’s Degree in Environmental Sciences for its Permanence in the National Register of Quality Graduates of CONACYT TAB-2014-C29-245836 key” and the project “Estudio de la fisiología digestiva en larvas y juveniles de pejelagarto (*Atractosteus tropicus*) con base en técnicas histológicas, bioquímicas y moleculares” CB-2016-01-282765 key.

References

- Al-Harbi AH, Uddin N (2004) Seasonal variation in the intestinal bacterial flora of hybrid tilapia (*Oreochromis niloticus*, *Oreochromis aureus*) cultured in earthen ponds in Saudi Arabia. *Aquaculture* 229: 37 – 44.
- Al-Harbi AH, Uddin N (2005). Bacterial diversity of tilapia (*Oreochromis niloticus*) cultured in brackish water in Saudi Arabia. *Aquaculture*, 250: 566 – 572.
- Andoh A, Bamba T, Sasaki M (1999). Physiological and anti-inflammatory roles of dietary fiber and butyrate in intestinal functions. *JPEN J Parenter Enteral Nutr* 23: S70 – S73.
- Aschfalk A, Müller W (2002) Clostridium perfringens toxin types from wild-caught Atlantic cod (*Gadus morhua* L.), determined by PCR and ELISA. *Can J Microbiol* 48:365–368.
- Austin B, Austin DA (2016) Bacterial Fish Pathogens: Disease of Farmed and Wild Fish. Springer Sixth Edition. 259 pp. Doi: 10.1007/978-3-319-32674-0
- Avila-Villa LA, Martínez-Porchas M, Gollas-Galván T, López-Elías JA, Mercado L, Murguía-López A, et al. (2011). Evaluation of different microalgae species and Artemia (*Artemia franciscana*) as possible vectors of necrotizing hepatopancreatitis bacteria. *Aquaculture* 318: 273 – 276.
- Bartlett KH, Trust TJ, Lior H (1977). Small pet aquarium frogs as a source of Salmonella. *Appl. Environ. Microbiol.* 33: 1026 – 1029.
- Bennett KW, Eley A (1993). Fusobacteria – new taxonomy and related diseases. *J Med Microbiol* 39: 246 – 254.
- Berg RW, Anderson AW (1972). Salmonellae and *Edwardsiella tarda* in gull feces: a source of contamination in fish processing plants. *Appl. Microbiol.* 24: 501 – 503.
- Blanch A, Alsina M, Simon M, Jofre J (1997). Determination of bacteria associated with reared turbot (*Scophthalmus maximus*) larvae. *Journal of Applied Microbiology* 82: 729 – 734.
- Bolnick BI, Snowberg LK, Hirsch PE, Lauber CHL, Knight R, Caporaso JG, Svanbäck R (2014). Individuals’ diet diversity influences gut microbial diversity in two freshwater fish

494 (threespine stickleback and Eurasian perch). Ecology Letters, 17: 979 – 987.
 495 <https://doi.org/10.1111/ele.12301>

496 Bondad MG, Subasinghe RP, Arthur JR (2005). Disease and health management in Asian
 497 aquaculture. *Veterinary parasitology*, vol. 132, no. 3-4, pp. 249 – 272.

498 Boutin S, Bernatchez L, Audet C, Derôme N (2012). Antagonistic effect of indigenous skin bacteria
 499 of brook charr (*Salvelinus fontinalis*) against *Flavobacterium columnare* and *F.*
 500 *psychrophilum*. *Vet. Microbiol.* 155: 355 – 361.
 501 <https://doi.org/10.1016/j.vetmic.2011.09.002>

502 Brunt J, Austin B (2005). Use of a probiotic to control lactococcosis and streptococcosis in rainbow
 503 trout, *Oncorhynchus mykiss* (Walbaum). *J Fish Dis* 28: 693–701.

504 Bussing WA (1998). Peces de las aguas continentales de Costa Rica. Ed. Univ. Costa Rica. 271 p.

505 Cahill M (1990). Bacterial flora of fishes: A review. *Microbial Ecology*, 19: 21 – 41.
 506 <https://doi.org/10.1007/BF02015051>

507 Carda-Diéguez M, Mira A, Fouz B (2014). Pyrosequencing survey of intestinal microbiota diversity
 508 in cultured sea bass (*Dicentrarchus labrax*) fed functional diets. *FEMS Microbiology Ecology*
 509 87: 451 – 459.

510 Chan XY, Chang CY, Hong KW, Tee KK, Yin WF, Chan KG (2013). Insights of biosurfactant
 511 producing *Serratia marcescens* strain W2.3 isolated from diseased tilapia fish: a draft genome
 512 analysis *Gut Pathogens*, 5: 29. <https://doi.org/10.1186/1757-4749-5-29>

513 Clements KD, Sutton DC, Choat JH (1989). The occurrence and characteristics of unusual protistan
 514 symbionts from surgeonfishes (F. Acanthuridae) of the Great Barrier Reef, Australia.
 515 *Marine Biology*, 102: 403 – 412.

516 Clements KD (1991). Endosymbiotic communities of two herbivorous labroid fishes, *Odax*
 517 *cyanomelas* and *O. pullus*. *Marine Biology*, 106: 223 – 229.

518 Clements KD, Gleeson VP, Slaytor M (1994). Shortchain fatty-acid metabolism in temperate marine
 519 herbivorous fish. *J Comp Physiol B*. 164: 372 – 377.

520 Clements KD Choat JH (1995). Fermentation in tropical marine herbivorous fishes. *Physiol Zool*.
 521 68: 355 – 378.

522 Clements KD (1997). Fermentation and gastrointestinal microorganisms in fishes. In:
 523 *Gastrointestinal Microbiology. Vol. 1: Gastrointestinal Ecosystems and Fermentations* (eds
 524 Mackie RI, White BA), pp. 156–198. Chapman and Hall, New York.

525 Clements KD, Pasch IBY, Moran D, Turner SJ (2007). Clostridia dominate 16S rRNA gene libraries
 526 prepared from the hindgut of temperate marine herbivorous Fishes. *Mar. Biol.* 150: 1431 –
 527 1440. <https://doi.org/10.1007/s00227-006-0443-9>

528 Clements KD, Angert ER, Montgomery WL, Choat JH (2014). Intestinal microbiota in fishes: what's
 529 known and what's not. *Molecular Ecology*, 23: 1891 – 1898.
 530 <https://doi.org/10.1111/mec.12699>

531 Collinder E, Bjornhag G, Cardona M, Norin E, Rehbinder C, Midtvedt T (2003). Gastrointestinal
532 host-microbial interactions in mammals and fish: comparative studies in man, mice, rats,
533 pigs, horses, cows, elk, reindeer, salmon and cod. *Microb Ecol Health Dis* 15: 66 – 78.

534 Cui Y, Chun SJ, Ko SR, Lee HG, Srivastava A, Oh HM, Ahn CY (2017). *Reyranella aquatilis* sp. nov.,
535 an alphaproteobacterium isolated from a eutrophic lake. *Int J Syst Evol Microbiol*,
536 67: 3496 – 350. <https://doi.org/10.1099/ijsem.0.00215>

537 De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S, Pieraccini G,
538 Lionetti P (2010). Impact of diet in shaping gut microbiota revealed by a comparative study
539 in children from Europe and rural Africa. *Proc. Natl. Acad. Sci. U.S.A.* 107: 14691 – 14696.

540 Desai AR, Links MG, Collins SA, Mansfield GS, Drew MD, Van Kessel AG et al. (2012). Effects of
541 plant-based diets on the distal gut microbiome of rainbow trout (*Oncorhynchus mykiss*).
542 *Aquaculture* 350: 134 – 142.

543 Di Maiuta N, Schwarzentruher P, Schenker M, Schoelkopf J (2013). Microbial population dynamics
544 in the faeces of wood-eating loricariid catfishes. *Lett Appl Microbiol* 56, 401 – 407.

545 von Engelhardt W, Bartels J, Kirschberger S, Duttingdorf HDMZ, Busche R (1998). Role of short-
546 chain fatty acids in the hind gut. *Vet Q* 20: S52 – S59.

547 FAO (2014). *The State of World Fisheries and Aquaculture*. Roma.

548 FAO (2017). *The State of World Fisheries and Aquaculture*. Roma.

549 Ferguson RL, Buckley E, Palumbo A (1984). Response of marine bacterioplankton to differential
550 filtration and confinement. *Appl Environ Microbiol* 47: 49–55.

551 Fidopiastis P, Bezdek D, Horn M, Kandel J (2006). Characterizing the resident, fermentative
552 microbial consortium in the hindgut of the temperate-zone herbivorous fish, *Hermosilla*
553 *azurea* (Teleostei: Kyphosidae). *Mar Biol*, 148: 631 – 642. [https://doi.org/10.1007/s00227-005-](https://doi.org/10.1007/s00227-005-0106-2)
554 [0106-2](https://doi.org/10.1007/s00227-005-0106-2)

555 Flores-Nava A, Brown A (2010). Peces nativos de agua dulce de América del Sur de interés para la
556 acuicultura: Una síntesis del estado de desarrollo tecnológico de su cultivo. *Serie*
557 *Acuicultura en Latinoamérica*, FAO, 1: 3.

558 Frías-Quinatana CA, Marquez-Couturier G, Alvarez-González CA, Tovar-Ramírez D, Nolasco-
559 Soria H, Galaviz-Espinosa M, Martínez-García R, Camarillo-Coop S, Martínez-Yáñez R,
560 Gisbert E (2015). Development of digestive tract and enzyme activities during the early
561 ontogeny of the tropical gar *Atractosteus tropicus*. *Fish Physiol Biochem* 41:1075–1091.

562 Gao YL, Storebakken T, Shearer KD, Penn M, Overland M (2011). Supplementation of fishmeal and
563 plant protein-based diets for rainbow trout with a mixture of sodium formate and butyrate.
564 *Aquaculture*, 311: 233–240.

565 Gerber E, Bernard R, Castang S, Chabot N, Coze F, Dreux-Zigha A, Hauser E, Hivin P, Joseph P,
566 Lazarelli C, Letellier G, Olive J, Leonetti JP (2015). *Deinococcus* as new chassis for industrial
567 biotechnology: biology, physiology and tools. *Journal of Applied Microbiology*, 119: 1 – 10.
568 <https://doi.org/10.1111/jam.12808>

569 Ghanbari M, Kneifel W, Domig KJ (2015). A new view of the fish gut microbiome: Advances from
570 next-generation sequencing. *Aquaculture* 448: 464 – 475.
571 <https://doi.org/10.1016/j.aquaculture.2015.06.033>

572 Gianoulis TA, Raes J, Patel PV, Bjornson R, Korbel JO, Letunic I, et al. (2009). Quantifying
573 environmental adaptation of metabolic pathways in metagenomics. *Proceedings of the*
574 *National Academy of Sciences*, 106: 1374 – 1379.

575 Givens CE (2012). A fish tale: comparison of the gut microbiome of 15 fish species and the influence
576 of diet and temperature on its composition. Thesis of Philosophy. University of Georgia.
577 Pp 10 – 229.

578 Griffiths E, Gupta RS (2007). Identification of signature proteins that are distinctive of the
579 *Deinococcus-Thermus* phylum. *International Microbiology*, 10: 201 – 208.
580 <https://doi.org/10.2436/20.1501.01.28>

581 Hamer H, Jonkers D, Troost FJ, Bast A, Vanhoutvin S, Venema K, Brummer RJ (2007). Butyrate
582 modulates oxidative stress in the colonic mucosa of healthy humans. *Gastroenterology* 132,
583 A79.

584 Head IM, Saunders JR, Pickup RW (1998) Microbial evolution, diversity, and ecology: a decade of
585 ribosomal RNA analysis of uncultivated microorganisms. *Microb Ecol* 35: 1–21.

586 Ingerslev HC, Strube ML, von Gersdorff Jørgensen L, Dalsgaard I, Boye M, Madsen L (2014a). Diet
587 type dictates the gut microbiota and the immune response against *Yersinia ruckeri* in
588 rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology* 40: 624–633.

589 Ingerslev HC, von Gersdorff Jørgensen L, Strube ML, Larsen N, Dalsgaard I, Boye M et al. (2014b).
590 The development of the gut microbiota in rainbow trout (*Oncorhynchus mykiss*) is affected
591 by first feeding and diet type. *Aquaculture* 424: 24 – 34.

592 Irianto A, Austin B (2002a). Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus*
593 *mykiss* (Walbaum). *Journal of Fish Diseases*, 25: 333 – 342.

594 Irianto A, Austin B (2002b). Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus*
595 *mykiss* (Walbaum). *J. Fish Dis.* 25: 1 – 10. <https://doi.org/10.1046/j.1365-2761.2002.00375.x>

596 Iveson JB (1971). Strontium chloride B and E. E. broth media for the isolation of *Edwardsiella*,
597 *Salmonella* and Arizona species from tiger snakes. *J. Hyg.* 69: 323 – 330.

598 Izvekova GI, Izvekov E, Plotnikov A (2007). Symbiotic microflora in fishes of different ecological
599 groups. *Biol Bull* 34: 610 – 618.

600 Jammal A, Bariche M, zu Dohna H, Kambris Z (2017). Characterization of the Cultivable Gut
601 Microflora in Wild-Caught Mediterranean Fish Species. *Current Nutrition & Food Science*,
602 13:147-154. <https://doi.org/10.2174/1573401313666170216165332>

603 Joborn A, Dorsch M, Olsson JC, Westerdahl A, Kjelleberg S (1999). *Carnobacterium inhibens* sp nov.,
604 isolated from the intestine of Atlantic salmon (*Salmo salar*). *Int. J. Syst. Bacteriol.* 49: 1891 –
605 1898.

606 van Kessel MA, Dutilh BE, Neveling K, Kwint MP, Veltman MP, Flik G, Jetten MS, Klaren PH *et al.*
607 (2011). Pyrosequencing of 16S rRNA gene amplicons to study the microbiota in the
608 gastrointestinal tract of carp (*Cyprinus carpio* L.). *AMB Express* 1, 41

609 Kim SJ, Ahn JH, Lee TH, Weon HY, Hong SB *et al.* (2013). *Reyranella soli* sp. nov., isolated from
610 forest soil, and emended description of the genus *Reyranella* Pagnier *et al.* 2011. *Int J Syst*
611 *Evol Microbiol*, 63: 3164 – 3167.

612 Klindworth A, Pruesse E, Schweer T, Peplies J, Quast Ch, Horn M, Oliver Glöckner FO (2013).
613 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
614 generation sequencing-based diversity studies. *Nucleic Acids Research*, 41:1–11.
615 <https://doi.org/10.1093/nar/gks808>

616 Kormas KA, Meziti A, Mente E, Frentzos A (2014). Dietary differences are reflected on the gut
617 prokaryotic community structure of wild and commercially reared sea bream (*Sparus*
618 *aurata*). *MicrobiologyOpen*, 3(5): 718 – 728. <https://doi.org/10.1002/mbo3.202>

619 Larsen AM, Mohammed HH, Arias CR (2014). Characterization of the gut microbiota of three
620 commercially valuable warmwater fish species. *Journal of Applied Microbiology* 116:
621 1396–1404.

622 Ley RE, Hamady M, Lozupone C, Turnbaugh PJ and others (2008). Evolution of mammals and their
623 gut microbes. *Science* 320: 1647–1651. <https://doi.org/10.1126/science.1155725>

624 Li J, Ni J, Li X, Yan Q, Yu Y (2013a). Relationship between gastrointestinal bacterial structure and
625 development of *Silurus soldatovi meridodalis*. *Chen. Acta Hydrobiologica Sinica* 37: 613–619.

626 Li X, Yan Q, Xie S, Hu W, Yu Y, Hu Z (2013b). Gut microbiota contributes to the growth of fast-
627 growing transgenic common carp (*Cyprinus carpio* L.). *PLoS One* 8: e64577.

628 Li J, Ni J, Wang C, Li X, Wu S, Zhang T *et al.* (2014a). Comparative study on gastrointestinal
629 microbiota of eight fish species with different feeding habits. *Journal of Applied*
630 *Microbiology* 117: 1750–1760.

631 Li X, Zhu Y, Yan Q, Ringø E, Yang D (2014b). Do the intestinal microbiotas differ between
632 paddlefish (*Polyodon spathala*) and bighead carp (*Aristichthys nobilis*) reared in the same
633 pond? *Journal of Applied Microbiology* 117: 1245–1252.

634 Li T, Long M, Gatesoupe FJ, Zhang Q, Li A, Gong X (2015a). Comparative Analysis of the Intestinal
635 Bacterial Communities in Different Species of Carp by Pyrosequencing. *Microb Ecol.*,
636 *Microb Ecol.*, 69:25–36. DOI 10.1007/s00248-014-0480-8.

637 Li H, Zhong Q, Wirth S, Wang W, Hao Y, Wu S, Zou H, Li W, Wang G (2015b). Diversity of
638 autochthonous bacterial communities in the intestinal mucosa of grass carp
639 (*Ctenopharyngodon idellus*) (Valenciennes) determined by culture-dependent and culture-
640 independent techniques. *Aquaculture Research*, 46: 2344 – 2359.
641 <https://doi.org/10.1111/are.12391> Li T, Li H, Gatesoupe FJ, She R, Lin Q, Yan X, Li J, Li X
642 (2017). Bacterial Signatures of “Red-Operculum” Disease in the Gut of Crucian Carp
643 (*Carassius auratus*). *Microb Ecol.*, 74: 510 – 521. <https://doi.org/10.1007/s00248-017-0967-1>

644 Liu Z, Ke X, Lu M, Gao F, Cao J, Zhu H, Wang M (2015). Identification and pathological observation
645 of a pathogenic *Plesiomonas shigelloides* strain isolated from cultured tilapia (*Oreochromis*
646 *niloticus*). Wei Sheng Wu Xue Bao, 55, 96–106.

647 Liu H, Guo X, Gooneratne R, Lai R, Zeng C, Zhan F et al. (2016). The gut microbiome and
648 degradation enzyme activity of wild freshwater fishes influenced by their trophic levels.
649 Scientific Reports 6: 24340.

650 Lubbs D, Vester B, Fastinger N, Swanson K (2009). Dietary protein concentration affects intestinal
651 microbiota of adult cats: a study using DGGE and qPCR to evaluate differences in microbial
652 populations in the feline gastrointestinal tract. J Anim Physiol Anim Nutr 93:113– 121

653 MacFarlane RD, McLaughlin JJ, Bullock GL (1986). Quantitative and qualitative studies of gut flora
654 in striped bass from estuarine and coastal marine environments. J Wildl Dis 22: 344–348.

655 Márquez-Couturier G, Álvarez-González CA, Contreras-Sánchez WM, Hernández-Vidal U,
656 Hernández-Franyutti AA, Mendoza-Alfaro RE, Aguilera-González C, García-Galano T,
657 Civera-Cerecedo R & Goytortua-Bores E (2006). Avances en la alimentación y nutrición de
658 pejelagarto *Atractosteus tropicus*. p. 446-523. In: Cruz Suárez, L. E., D. Ricque-Marie, M.
659 Tapia-Salazar, M.G. Nieto-López, D.A. Villarreal-Cabazos, A.C. Puello & A. García-Ortega
660 (Eds.). Avances en Nutrición Acuícola VIII. VIII Simposium Internacional de Nutrición
661 Acuícola. 15 al 17 Noviembre. Universidad Autónoma de Nuevo León, Monterrey, Nuevo
662 León, México.

663 Márquez-Couturier G, Vázquez-Navarrete CJ (2015). Estado de arte de la biología y cultivo de
664 pejelagarto (*Atractosteus tropicus*). Agroproductividad 3:44-51. ISSN 24487546.

665 Márquez-Couturier G, Vázquez-Navarrete CJ, Contreras-Sánchez WM, Alvarez-González CA
666 (2015). Acuicultura tropical sustentable: Una estrategia para la producción y conservación
667 del pejelagarto (*Atractosteus tropicus*) en Tabasco, México. 2nd edition. Colección José
668 Narciso Roviroso. Villahermosa, Tabasco, Mexico. 87 p.

669 Martens JH, Barg H, Warren MJ, Jahn D (2002). Microbial production of vitamin B12. Appl
670 Microbiol Biotechnol, 58: 275 – 285.

671 Martin-Antonio B, Manchado M, Infante C, Zerolo R, Labella A, Alonso C, Borrego JJ (2007)
672 Intestinal microbiota variation in Senegalese sole (*Solea senegalensis*) under different feeding
673 regimes. Aquacult Res 38: 1213–1222

674 Martínez-Cruz P, Ibáñez AL, Monroy-Hermosillo O, Ramirez-Saad HC (2012). Use of Probiotics in
675 Aquaculture. ISRN Microbiology, 2012:13, ID 916845. <https://doi.org/10.5402/2012/916845>

676 Martínez-Díaz SF, Pérez-Espana H (1999) Feasible mechanisms for algal digestion in the king
677 angelfish. Journal of Fish Biology, 55: 692–703.

678 McBain JA, Eastman A, Nobel CS Mueller GC (1997). Apoptotic death in adenocarcinoma cell lines
679 induced by butyrate and other histone deacetylase inhibitors. Biochem Pharmacol, 53: 1357
680 – 1368.

681 Méndez-Marín O, Hernández-Franyutti AA, Álvarez-González CA, Contreras-Sánchez WM,
682 Uribe-Aranzábal, MC (2012). Histología del ciclo reproductor de hembras del pejelagarto

683 *Atractosteus tropicus* (Lepisosteiformes: Lepisosteidae) en Tabasco, México. Revista de
684 Biología Tropical. Universidad de Costa Rica, Costa Rica. 60:1857-1871. ISSN: 0034-7744.

685 Miller RR, Minckley WL & Norris SM (2005). Freshwater fishes of Mexico. University of Chicago
686 press, Chicago and London. 490 p.

687 Miyake S, Ngugi DK, Stingl U (2015). Diet strongly influences the gut microbiota of surgeonfishes.
688 Molecular Ecology 24: 656–672.

689 Moran D, Turner SJ, Clements KD (2005). Ontogenetic Development of the Gastrointestinal
690 Microbiota in the Marine Herbivorous Fish *Kyphosus sydneyanus*. Microbial Ecology, 49: 590
691 – 597. <https://doi.org/10.1007/s00248-004-0097-4>

692 Nam YD, Jung MJ, Roh SW, Kim MS, Bae JW (2011). Comparative analysis of Korean human gut
693 microbiota by barcoded pyrosequencing. PLoS One 6:e22109.
694 <https://doi.org/10.1371/journal.pone.0022109>

695 Nayak SK (2010). Role of gastrointestinal microbiota in fish. Aquaculture Research 41: 1553–1573.

696 Nelson SJ (2006). Fishes of the World. 4th Edition. A Wiley-Interscience publication. EEUU. 601 p.

697 Newman JT Jr, Cosenza BJ, Buck JD (1972). Aerobic microflora of the bluefish (*Pomatomus saltatrix*)
698 intestine. J Fish Res Board Can 29:333–336.

699 Ni J, Yu Y, Zhang T, Gao L (2012). Comparison of intestinal bacterial communities in grass carp,
700 *Ctenopharyngodon idellus*, from two different habitats. Chinese Journal of Oceanology and
701 Limnology, 30: 757–765. <https://doi.org/10.1007/s00343-012-1287-4>

702 Nielsen HB, Almeida M, Juncker AS, Rasmussen S, Li J, Sunagawa S, et al. (2014). Identification
703 and assembly of genomes and genetic elements in complex metagenomic samples without
704 using reference genomes. Nature Biotechnology 32: 822– 828.

705 Nuez-Ortin WG, Prado S, Toranzo AE (2012). Antimicrobial properties of butyric acid and other
706 organic acids against pathogenic bacteria affecting the main aquatic species. In AQUA 2012.
707 Prague.

708 Núñez de la Rosa MG (2011). Evaluación preliminar de las poblaciones bacterianas asociadas al
709 tracto intestinal de la tilapia (*Oreochromis niloticus*) expuestas a aceites esenciales de
710 orégano en la dieta. Tesis de Maestría. Facultad de Ciencias. Universidad Nacional de
711 Colombia. Pp 1 – 87.

712 Onarheim AM, Wiik R, Burghardt J, Stackebrandt E (1994). Characterization and identification of
713 two *Vibrio* species indigenous to the intestine of fish in cold sea water; description of *Vibrio*
714 *iliopiscarius* sp. nov. Systematic and Applied Microbiology 17: 370 – 379.

715 Owen MA, Waines P, Bradley G, Davies S (2006). The effect of dietary supplementation of sodium
716 butyrate on the growth and microflora of *Clarias gariepinus* (Burchell 1822). In Proceedings
717 of the XII International Symposium Fish Nutrition and Feeding ed. Deneri, O.E. pp. 301.
718 Biarritz, France

719 Pagnier I, Raoult D, La Scola B (2011). Isolation and characterization of *Reyranella massiliensis* gen.
720 nov., sp. nov. from freshwater samples by using an amoeba co-culture procedure. Int J Syst
721 Evol Microbiol, 61: 2151 – 2154.

722 Paul F, Otte J, Schmitt I, Dal Grande F (2018). Comparing Sanger sequencing and high-throughput
723 metabarcoding for inferring photobiont diversity in lichens. Scientific Reports, 8: 8624.
724 <https://doi.org/10.1038/s41598-018-26947-8>

725 Pikuta EV, Hoover RB, Marsic D, Whitman WB, Lupa B, Tang J, Krader P (2009). *Proteocatella*
726 *sphenisci* gen. nov., sp. nov., a psychrotolerant, spore-forming anaerobe isolated from
727 penguin guano. Int J Syst Evol Microbiol 59: 2302 – 2307.

728 Pirarat N, Kobayashi T, Katagiri T, Maita M, Endo M (2006). Protective effects and mechanisms of
729 a probiotic bacterium *Lactobacillus rhamnosus* against experimental *Edwardsiella tarda*
730 infection in tilapia (*Oreochromis niloticus*). Vet Immunol Immunopathol, 113: 339–347.

731 Qi XZ, Xue MY, Yang SB, Zha JW, Wang GX, Ling F (2017). Ammonia exposure alters the
732 expression of immune-related and antioxidant enzymes-related genes and the gut
733 microbial community of crucian carp (*Carassius auratus*). Fish & Shellfish Immunology, 70:
734 485 – 492. <https://doi.org/10.1016/j.fsi.2017.09.043>

735 Ray AK, Ghosh K, Ringø E (2012). Enzyme-producing bacteria isolated from fish gut: a review.
736 Aquaculture Nutrition, 18: 465– 492.

737 Reséndez-Medina A, Salvadores-Baledón ML (1983). Contribución al conocimiento de la biología
738 del pejelagarto *Lepisosteus tropicus* (Gill) y la tenguayaca *Petenia splendida* Günther, del
739 estado de Tabasco. Biótica 8: 413-426.

740 Rimmer DW, Wiebe RJ (1987). Fermentative microbial digestion in herbivorous fishes. Journal of
741 Fish Biology, 31:229–236.

742 Ringø E, Strøm E, Tabachek JA (1995). Intestinal microflora of salmonids: a review. Aquacult Res.
743 26: 773–789.

744 Ringø E, Sperstad S, Myklebust R, Refstie S, Krogdahl Å (2006). Characterisation of the microbiota
745 associated with intestine of Atlantic cod (*Gadus morhua* L.): the effect of fish meal, standard
746 soybean meal and a bioprocessed soybean meal. Aquaculture 261: 829–841.

747 Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, Rawls JF (2011).
748 Evidence for a core gut microbiota in the zebrafish. ISME J 5: 1595 – 1608

749 Roth JR, Lawrence JG, Bobik TA (1996). Cobalamin (coenzyme B12): synthesis and biological
750 significance. Annu Rev Microbiol, 50: 137 – 181.

751 Rudi K, Angell IL, Pope PB, Vik JO, Sandve SR, Snipen LG (2018). Stable Core Gut Microbiota across
752 the Freshwater-to-Saltwater Transition for Farmed Atlantic Salmon. Applied and
753 Environmental Microbiology, 84: e01974 – 17. <https://doi.org/10.1128/AEM.01974-17>

754 Salinas I, Cuesta A, Esteban MA, Meseguer J (2005). Dietary administration of *Lactobacillus*
755 *delbrii* and *Bacillus subtilis*, single or combined, on gilthead sea bream cellular innate
756 immune responses. Fish Shellfish Immunol 19: 67–77.

757 Skrodenytė Arbačiauskienė V, (2007). Enzymatic activity of intestinal bacteria in roach *Rutilus*
758 *rutilus* L. *Fish Sci*, 73: 964 – 966.

759 Smriga S, Sandin SA, Azam F (2010). Abundance, diversity, and activity of microbial assemblages
760 associated with coral reef fish guts and feces. *FEMS Microbiology Ecology*, 73: 31 – 42.
761 <https://doi.org/10.1111/j.1574-6941.2010.00879.x>

762 Smith AG, Croft MT, Moulin M, Webb ME (2007). Plants need their vitamins too. *Curr Opin Plant*
763 *Biol*, 10: 266 – 275.

764 Spanggaard B, Huber I, Nielsen J, Nielsen T, Appel KF, Gram L (2000). The microflora of rainbow
765 trout intestine: a comparison of traditional and molecular identification. *Aquaculture* 182:
766 1 – 15.

767 Subasinghe RP (2005). Epidemiological approach to aquatic animal health management:
768 opportunities and challenges for developing countries to increase aquatic production
769 through aquaculture. *Prev. Vet. Med.* 67: 117–124.

770 Sullam KE, Essinger SD, Lozupone CA, O'Connor MP, Rosen GL, Knight R, Kilham SS, Russell JA
771 (2012). Environmental and ecological factors that shape the gut bacterial communities of
772 fish: a meta-analysis. *Molecular Ecology*, 21: 3363 – 3378. [https://doi: 10.1111/j.1365-](https://doi.org/10.1111/j.1365-294X.2012.05552.x)
773 [294X.2012.05552.x](https://doi.org/10.1111/j.1365-294X.2012.05552.x)

774 Suttle CA (2007). Marine viruses—major players in the global ecosystem. *Nature Reviews*
775 *Microbiology*, 5: 801 – 812.

776 Tamaki H, Wright CL, Li X, Lin Q, Hwang C, Wang S, Thimmapuram J, Kamagata Y, Liu WT (2011).
777 Analysis of 16S rRNA Amplicon Sequencing Options on the Roche/454 Next-Generation
778 Titanium Sequencing Platform. *PLOS ONE* 6:
779 e25263. <https://doi.org/10.1371/journal.pone.0025263>

780 Theodorakopoulos N, Bachar D, Christen R, Alain K, Chapon V (2013). Exploration of *Deinococcus*-
781 *Thermus* molecular diversity by novel group-specific PCR primers. *Microbiologyopen*, 2: 862
782 – 72. <https://doi.org/10.1002/mbo3.119>

783 Thomas F, Hehemann JH, Rebuffet E, Czejek M, Michel G (2011). Environmental and gut
784 Bacteroidetes: the food connection. *Front Microbiol*, 2: 93.
785 <https://doi.org/10.3389/fmicb.2011.00093>

786 Titus E, Ahearn GA (1988). Short-chain fatty-acid transport in the intestine of a herbivorous teleost.
787 *J Exp Biol* 135, 77–94.

788 Tsuchiya C, Sakata T, Sugita H (2007). Novel ecological niche of *Cetobacterium somerae*, an anaerobic
789 bacterium in the intestinal tracts of freshwater fish. *Letters in Applied Microbiology* ISSN
790 0266-8254.

791 Tsuchiya C, Sakata T, Sugita H (2008). Novel ecological niche of *Cetobacterium somerae*, an anaerobic
792 bacterium in the intestinal tracts of freshwater fish. *Letters in Applied Microbiology* 46: 43–
793 48. <https://doi.org/10.1111/j.1472-765X.2007.02258.x>.

794 Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, Sogin ML, Jones WJ,
795 Roe BA, Affourtit JP (2009). A core gut microbiome in obese and lean twins. *Nature* 457:
796 480 – 484

797 Uffen R (1997). Xylan degradation: a glimpse at microbial diversity. *J Ind Microbiol Biotechnol* 19:1–
798 6.

799 Uz I, Ogram AV (2006). Cellulolytic and fermentative guilds in eutrophic soils of the Florida
800 Everglades. *FEMS Microbiol Ecol* 57:396–408

801 Verner-Jeffreys DW, Shields RJ, Bricknell IR, Birkbeck TH (2003). Changes in the gut-associated
802 microflora during the development of Atlantic halibut (*Hippoglossus hippoglossus* L.) larvae
803 in three British hatcheries. *Aquaculture* 219: 21 – 42.

804 Verschuere L, Rombaut G, Sorgeloos P, Verstraete W (2000). Probiotic Bacteria as Biological Control
805 Agents in Aquaculture Microbiology and Molecular Biology Reviews, 64: 655 – 671.

806 Villamil L, Tafalla C, Figueras A, Novoa B (2002). Evaluation of immunomodulatory effects of lactic
807 acid bacteria in turbot (*Scophthalmus maximus*). *Clin Diagn Lab Immunol* 9:1318–1323.

808 Vogiatzoglou A, Smith AD, Nurk E, Berstad P, Drevon CA, Ueland PM, Vollset SE, Tell GS, Refsum
809 H (2009). Dietary sources of vitamin B-12 and their association with plasma vitamin B-12
810 concentrations in the general population: the Hordaland Homocysteine Study. *The*
811 *American Journal of Clinical Nutrition*, 89: 1078–1087.
812 <https://doi.org/10.3945/ajcn.2008.26598>

813 Wang AR, Ran C, Ringø E, Zhou ZG (2017). Progress in fish gastrointestinal microbiota research.
814 *Aquaculture* 0:1–15. <https://doi.org/10.1111/raq.12191>

815 Walczak N, Puk K, Guz L (2017). Bacterial flora associated with diseased freshwater ornamental
816 fish. *J Vet Res*, 61: 445 – 449. <https://doi.org/10.1515/jvetres-2017-0070>

817 Ward NL, Steven B, Penn K, Methe BA, Detrich WH (2009). Characterization of the intestinal
818 microbiota of two Antarctic notothenioid fish species. *Extremophiles* 13: 679 – 685.

819 Weisburg WG, Barns SM, Pelletier DA, Lane DJ (1991). 16S ribosomal DNA amplification for
820 phylogenetic study. *J. Bacteriol.* 173: 697 – 703.

821 Wiley O (1976). The phylogeny and biogeography of fossil and recent gars (Actinopterygii:
822 Lepisosteidae). *Univ. KS. Hist. Nat. Mus. Publ.* 64: 1 – 11.

823 Wong S, Waldrop T, Summerfelt S, Davidson J, Barrows F, Kenney PB, Welch T, Wiens GD, Snekvik
824 K, Rawls JF (2013). Aquacultured rainbow trout (*Oncorhynchus mykiss*) possess a large
825 core intestinal microbiota that is resistant to variation in diet and rearing density. *Appl*
826 *Environ Microbiol* 79:4974–4984

827 Wu S, Wang G, Angert ER, Wang W, Li W, Zou H (2012). Composition, diversity, and origin of the
828 bacterial community in grass carp intestine. *PLoS One* 7:e30440. doi: 10.1371/journal.
829 pone.0030440

- 830 Wyatt LE, Nickelson R, Vanderzant C (1979). Occurrence and control of Salmonella in freshwater
831 catfish. *J. Food Sci.* 44: 1067 – 1069.
- 832 Xia JH, Lin G, Fu GH, Wan ZY, Lee M, Wang L, Liu XJ, Yue GH (2014). The intestinal microbiome
833 of fish under starvation. *BMC Genomics*, 15:266. <http://doi.org/10.1186/1471-2164-15-266>
- 834 Xu J, Bjursell MK, Himrod J, Deng S, Carmichael LK, Chiang HC, Hooper LV, Gordon JI (2003). A
835 genomic view of the human-Bacteroides thetaiotaomicron symbiosis. *Science*, 299:2074–
836 2076.
- 837 Ye L, Amberg J, Chapman D, Gaikowski M, Liu WT (2014). Fish gut microbiota analysis
838 differentiates physiology and behaviour of invasive Asian carp and indigenous American
839 fish. *ISME Journal*, 8: 541–551. <https://doi.org/10.1038/ismej.2013.181>

Figure 1. Collection sites of wild (triangle figures) and cultivated (circle figure) of the adults' tropical gars (*Atractosteus tropicus*)

Figure 2. Taxonomic summary at different levels of relative bacterial abundance. (a) Phyla and (b) Genus. First letter W: wild or F: Farming individuals. Second letter sex M: male or F: female. Only taxa with relative abundance $\geq 1\%$ are shown.

Figure 3. Core microbiome in gut of cultivated/wild and female/male adults' *A. tropicus*, at taxonomic class-level of (a) phylum and (b) genus.

Figure 4. Similarity or dissimilarity coefficients of Unweighted UniFrac distances (UniFrac) by PCoA ordination analysis of the microbiota in adults' *A. tropicus*. Blue and red circles correspond to wild and cultivated individuals, respectively

Figure 5. LefSe analysis with the LDA score histogram of microbiota in adults' *A. tropicus* composition, (a) comparison among female and male organisms, cultivated and wild, and (b) comparison among female organisms cultivated and wild. Both show statistically significant differences depending on the origin. Red and green colors indicate cultivated and wild samples, respectively.

Figure 6. LefSe analysis with the LDA score histogram of microbiota in adults' *A. tropicus* composition (a) male organisms, cultivated & wild, (b) female organisms cultivated and (c) wild female. All show statistically significant differences depending on the origin.

Figure 7. Phylogenetic tree from the rRNA 16S gene constructed of some organisms found in the literature with probiotic potential in fish (see table 1). Red circles and blue squares represent outgroup and reference sequences, respectively.