

1 **The high persister phenotype of *Pseudomonas aeruginosa* is associated with increased**  
2 **fitness and persistence in cystic fibrosis airways**

3

4 **Authors**

5 Biljana Mojsoska<sup>1\*</sup>, David R. Cameron<sup>3\*</sup>, Jennifer A. Bartell<sup>2\*</sup>, Janus Anders Juul Haagenzen<sup>2</sup>,  
6 Lea M. Sommer<sup>1</sup>, Kim Lewis<sup>3§</sup>, Søren Molin<sup>2</sup>, Helle Krogh Johansen<sup>1,4§</sup>

7

8 **Affiliations:**

9 <sup>1</sup> Department of Clinical Microbiology, Rigshospitalet, Copenhagen, Denmark

10 <sup>2</sup> The Novo Nordisk Foundation Center for Biosustainability, Technical University of  
11 Denmark, Lyngby, Denmark

12 <sup>3</sup> Antimicrobial Discovery Center, Northeastern University, Boston, USA

13 <sup>4</sup> University of Copenhagen, Department of Clinical Medicine, Copenhagen, Denmark

14

15 **\*Contributed equally to the work**

16 **§Corresponding authors**

17

18

19 BIOLOGICAL SCIENCES: Microbiology

20 Keywords: persister cells, persistent infection, *Pseudomonas aeruginosa*, cystic fibrosis

21

## 22 **Abstract**

23 Despite intensive antibiotic treatment of cystic fibrosis (CF) patients, *Pseudomonas aeruginosa*  
24 often persists in patient airways for decades, and can do so without the development of  
25 antibiotic resistance. Using a high-throughput screening assay of survival after treatment with  
26 high concentrations of ciprofloxacin, we have determined the prevalence of high-persister  
27 variants (Hip) in a large patient cohort. In a screen of 467 longitudinal clinical isolates of *P.*  
28 *aeruginosa* from 40 CF patients, we classified 25.7% as Hip. Hip were identified in 26 patients,  
29 but only a few bacterial lineages were dominated by Hip. Instead, the emergence of Hip  
30 increased over time, suggesting that CF airways treated with ciprofloxacin select for Hip with  
31 an increased fitness in this environment. We generally observed diverse genetic changes in the  
32 Hip isolate population (as many co-occurring routes to increased fitness exist), but interestingly  
33 elevated mutation counts in the RpoN gene of 18 Hip isolates suggest that this sigma factor  
34 plays a role in shaping levels of antibiotic tolerance. To probe the impact of the Hip phenotype  
35 in a CF-similar environment, we tested the fitness properties of otherwise genotypically and  
36 phenotypically similar low-persister (Lop) and Hip isolates in co-culture using a specialized  
37 flow-cell biofilm system mimicking pharmacokinetic/-dynamic antibiotic dosing. Hip survived  
38 ciprofloxacin treatment far better than Lop isolates. The results of this investigation provide  
39 novel insights into persister dynamics and fitness contributions to survival in the CF lung, and  
40 show that the Hip phenotype of antibiotic susceptible bacteria plays an important role in long-  
41 term infections.

42

## 43 **Significance**

44 Antibiotic resistance is emphasized as a rapidly increasing health threat, but antibiotic tolerance  
45 via the occurrence of persister cells in antibiotic-treated bacterial populations is clinically and  
46 publicly neglected. In 40 CF patients representing a well-established human infection model –  
47 long-term lung infections by *Pseudomonas aeruginosa* – we show the emergence and  
48 accumulation of persister variants in a clinical population heavily reliant on antibiotic therapy.  
49 We observe that the high-persister (Hip) phenotype is independent of resistance and likely the  
50 consequence of numerous genetic alterations, complicating surveillance and inhibition in the  
51 clinic. Furthermore, we find Hip are selected for over time, survive better than ‘normal’  
52 bacteria, and can outcompete them in CF-similar conditions, ultimately affecting 65% of  
53 patients in an early disease cohort.

54

55

## 56 **Introduction**

57 Antibiotic-tolerant persister cells are suspected to be a significant clinical problem that has  
58 been seriously neglected in favor of combating antibiotic-resistant bacteria, though persisters  
59 were in fact described shortly after the clinical introduction of antibiotics (1). Persisters are  
60 distinct from antibiotic-resistant mutants, as they do not grow in the presence of antibiotics.  
61 Instead, they remain dormant during antibiotic exposure but retain the capacity to resuscitate  
62 and restore the population when antibiotic concentrations drop (2–4). However, our  
63 understanding of the physiology and clinical relevance of persister cells is limited, given the  
64 difficulty in reliably isolating what is theorized to be a stochastic phenotype *in vitro*, much less  
65 monitoring this phenotype in routine clinical care. Thus, few studies have assayed persister  
66 formation in clinical scenarios (5, 6). One study of oral carriage (0-19 weeks) of *Candida*  
67 *albicans* isolates from 22 cancer patients undergoing chemotherapy found that patients with  
68 carriage of greater than 8 weeks had significantly higher persister levels than those with less  
69 than 8 weeks carriage, but did not address the underlying genetic mechanisms of persistence  
70 in this pathogen (5). To examine the genetic underpinnings of the persister phenotype in a  
71 clinical scenario, both a large, aligned patient cohort that places the bacteria under similar  
72 environmental stresses as well as isolate sampling at a resolution that captures the emergence  
73 and longevity of the phenotype are needed.

74

75 *P. aeruginosa* is the most frequent cause of chronic airway infections in patients with CF (7,  
76 8). Mutations in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene often  
77 result in inefficient mucociliary clearance of bacteria from the airways, creating opportunities  
78 for bacterial colonization (9, 10). Upon entering the host, environmental *P. aeruginosa* adapts  
79 to the CF lung environment, ultimately establishing an incurable airway infection (11, 12).  
80 Despite intensive antibiotic treatment from the first discovery of the bacterium in the lung,  
81 resistance emergence in the first years of infection is surprisingly low (13, 14). In the absence  
82 of clinically defined antibiotic resistance, survival of the bacteria is likely supported by small  
83 fractions of antibiotic tolerant subpopulations, among which persister cells may be clinically  
84 important (6, 15). While persister cells are stochastic phenotypic variants in any bacterial  
85 population, genetic changes in bacterial populations can result in a high persister (Hip) state,  
86 producing increased numbers of antibiotic tolerant cells following exposure to antibiotics (16).  
87 A previous investigation of the persister phenotype in young CF patients showed an increase  
88 in persister phenotype in early/late infection isolate pairs from 14 patients. In this study, 35  
89 longitudinal *P. aeruginosa* isolates taken from one child over a 96-month period showed

90 increased levels of persister cells over time as well as an accumulation of 68 mutations between  
91 the first and last isolate (6). However, the mutations in the single patient resembled those  
92 known to accumulate in other CF patients over infection rather than any mutations previously  
93 associated with the Hip phenotype in other, non-CF studies.

94


95 To acquire a high-resolution pan-cohort perspective of persister emergence, genetic  
96 mechanism, and impact in long-term infections, we have screened for Hip variants over a 10-  
97 year period in 40 young CF patients via 467 longitudinal isolates of *P. aeruginosa* collected  
98 from initial colonization onward. This unique isolate collection allows us to determine Hip  
99 prevalence and dynamics during each colonizing strain's transition from environmental isolate  
100 to persistent pathogen. We describe relationships between the Hip phenotype and the age of  
101 the isolate, as well as other adaptive traits in longitudinal infections. We show that the Hip  
102 phenotype is an independent and widespread trait. We further search for genetic changes  
103 associated with the Hip phenotype in independent clonal lineages within distinct patients,  
104 which may suggest adaptive routes to producing this phenotype. Finally, we show that the Hip  
105 phenotype generally accumulates over time in patients via several archetypal patterns, appears  
106 to contribute to long-term persistence of lineages, and increases the fitness of colonizing  
107 populations of *P. aeruginosa* in antibiotic-treated CF patient lungs.

108

## 109 **Results**

### 110 *The isolate collection*

111 We examined a collection of 467 *P. aeruginosa* airway isolates obtained from 40 young CF  
112 patients over a 10 year period while they were treated at the Copenhagen CF Centre (17). These  
113 patients represent a cohort aligned at the early infection stage and undergoing similar treatment  
114 regimens per CF Centre guidelines, with repeated culture of *P. aeruginosa* from their monthly  
115 sputum sampling within a time frame of 2-10 years (patient inclusion was on a rolling basis  
116 over the study period in order to capture all early colonization cases). Early isolates therefore  
117 represent bacteria that have not been exposed to substantial antibiotic treatment before the  
118 study start excepting rare cases of strain transmission from another patient. The bacterial CF  
119 isolates have been grouped into 53 genetically distinct clone types based on their genome  
120 sequences (17), and while the majority of patients retained a monoclonal infection during the  
121 entire course of infection, half (n=21, 53 %) were infected at least transiently with another  
122 clone type. To effectively account for these multi-clonal infections, clinical isolates are  
123 described by their patient-specific lineage, which is defined by the clone type and the patient



**Figure 1. High-throughput screening approach for isolates with a high persister (Hip) phenotype.** A large collection of *P. aeruginosa* clinical isolates were grown to stationary phase in quadruplicate wells for 2 experiments (4 technical replicates for each of 2 biological replicates - BRs). Each isolate was treated with 100 µg/ml of ciprofloxacin for 24 hours, while growth was assessed by plating on LB agar. Following antibiotic treatment, isolates were plated via serial dilution and survival was assessed. Each isolate was given a persister score based on consistent replicate survival following treatment. Isolates for which 2-4 replicates survived for each BR were given a score of 2-4, respectively, and were considered high persisters (Hip). Isolates with a respective score of 0-1 were considered low persisters (Lop).

124 of origin (75 lineages in total). Throughout this paper, we will also refer to ‘Time since first  
125 detection’ for each isolate, which represents the length of time between first detection of *P.*  
126 *aeruginosa* and subsequent isolations of the same clone type.

127

#### 128 *Identification of Hip by high-throughput screening*


129 We screened the collection of *P. aeruginosa* isolates for the propensity to survive in the  
130 presence of high concentrations of ciprofloxacin. This antibiotic was chosen because it is  
131 commonly used to treat early *P. aeruginosa* infections in CF patients, and it is bactericidal  
132 toward stationary phase *P. aeruginosa* (18, 19). Briefly, *P. aeruginosa* subcultures in micro-  
133 titer plates were grown for 48 hours until they reached stationary phase, after which they were  
134 challenged with a high concentration of ciprofloxacin (100 µg/ml) for 24 hours before survival  
135 was assessed (Fig. 1). Each isolate was screened eight times (technical quadruplicates  
136 performed in duplicate biological experiments) and scored based on the capacity to re-grow  
137 after treatment. An isolate was given a score of 0 if it failed to re-grow in any replicate, a score  
138 of 1 if it grew once in both biological duplicates, a score of 2 if it grew in half of the technical  
139 replicates on each plate, a score of 3 if it grew in at least three replicates on each plate, and a  
140 score of 4 if it grew in all replicates. We then defined high persister (Hip) isolates as those  
141 scoring between 2-4, and low persister (Lop) isolates as those scoring between 0-1. Using this  
142 scoring system, 120 isolates (25.7%) of the collection were defined as Hip (Fig. 2A). Isolates  
143 with a score of four made up the largest Hip group, while most of the Lop isolates were scored  
144 as zero, failing to grow in any replicate. To validate our high-throughput screening approach,  
145 we randomly selected 14 Hip isolates and compared their persister phenotypes with 8 Lop

146 control isolates as well as the laboratory strain  
 147 PAO1 using standard persister assays. The Hip  
 148 variants on average produced 1000-fold-more  
 149 CFU persisters than the Lop isolates (Fig. 2B).

150


151 As we were using ciprofloxacin as the antibiotic  
 152 to identify persister isolates, and most of our  
 153 patients have been treated with ciprofloxacin, we  
 154 also wanted to check if we were simply selecting  
 155 ciprofloxacin resistant isolates as opposed to  
 156 high persisters. Ciprofloxacin MICs were  
 157 determined using E-tests for isolates within the  
 158 collection. Most of the isolates were  
 159 characterized as susceptible based on the  
 160 European Committee on Antimicrobial  
 161 Susceptibility Testing (EUCAST) breakpoints  
 162 (Fig. 2C). Forty-eight susceptible isolates were  
 163 Hip variants while 92 Lop variants were resistant,  
 164 confirming that our screening approach  
 165 identified an antibiotic-resistant phenotype  
 166 separate from the persister phenotype.

167



**Figure 2. Persister screening results. (A)** Distribution of *P. aeruginosa* Hip (blue) and Lop (grey) isolates. **(B)** Validation assays were performed for Hip and Lop isolates. Colony forming units (CFU) per ml were determined following overnight treatment with 100  $\mu$ g/ml of ciprofloxacin. Each data point represents a single isolate and is the mean of 6 independent experiments. Bars represent SEM. **(C)** Hip and Lop persister isolates were classified as susceptible or resistant to ciprofloxacin according to EUCAST breakpoints ( $S \leq 0.5 \mu\text{g/ml}$ ,  $R > 0.5 \mu\text{g/ml}$ ) based on their MIC obtained via E-test.

168 *The persister phenotype is an independent trait*  
 169 As illustrated by the comparison between  
 170 persister class and response to ciprofloxacin, the  
 171 persister phenotype in our collection is not  
 172 arising in isolation from other adaptations.  
 173 Adaptations accumulate based on a  
 174 combinatorial response to both temporal and  
 175 fixed environmental factors unique to a given  
 176 patient. We and others have previously observed  
 177 that CF isolates adapt towards slow growth rates,  
 178 increased resistance to antibiotics, and  
 179 preference for a biofilm lifestyle (20–22), and it  
 180 has also been proposed that there is a specific  
 181 association between slowing growth rate and the  
 182 Hip phenotype (23). To probe interrelationships  
 183 with other phenotypes and compare the adaptive  
 184 background of the Hip and Lop variants, we used  
 185 a principle component analysis to evaluate the  
 186 distribution of Hip (blue diamonds) versus Lop  
 187 (gray circles) variants by multiple traits under  
 188 selection pressure in the CF lung. In Figure 3A,  
 189 we see that Hip variants group with isolates  
 190 exhibiting more adapted traits (increased  
 191 antibiotic MICs and slowing growth), but they  
 192 also appear across the full phenotypic space  
 193 alongside Lop isolates. The data ellipses  
 194 enclosing the approximated majority of each  
 195 population (68% of the population,  $t$   
 196 distribution) show that Hip variants (solid blue  
 197 ellipse) do not separate from Lop variants (solid  
 198 gray ellipse). We also specifically identify the  
 199 first Hip variant of each clone type infecting a  
 200 given patient (FirstHip – blue triangles, dashed  
 201 blue ellipse) in an attempt to assess these isolates’ other traits and state of adaptation at first



**Figure 3. High persisters in the multi-trait landscape.** Lop (gray circle), the earliest Hip (blue triangle) isolate(s) of each lineage, and the rest of the Hip isolates (blue diamond) were analyzed via principle component analysis with respect to their similarity with other infection-linked traits: growth rate (grLB), adhesion (adh), ciprofloxacin MIC (cip) and tobramycin MIC (tob)). FirstHip persisters were highlighted as Hip variants with mitigated effects of other accumulating mutations within the lineage to improve cross-lineage comparison. 438 isolates with complete trait sets are included. **(A)** Isolates exhibiting a persister phenotype do not consistently cluster with any other specific trait, while traits of Lop, FirstHip, and Hip isolates shift slightly from ‘naïve’ towards ‘adapted’ levels. We illustrate this using data ellipse enclosing samples approximately within the first standard deviation ( $t$  distribution, 68% of the set) for isolate sets characterized as Lop (gray ellipse), FirstHip (dashed blue ellipse), and Hip (blue ellipse). **(B)** Lop and Hip variants co-occur in time when evaluating five lengthy lineages characterized by periods of continuous Hip isolates (see Figure 3 – ‘Continuous’ set). First isolate (white fill), firstHip (triangle), and last isolate (dark fill) are demarcated by bolded marker outlines for all lineages.

202 appearance. These FirstHip isolates overlap substantially with both Lops and Hips, indicating  
203 much variation in the adaptive state. This variation could be due to likely lapses of time  
204 between the emergence of a Hip variant and its isolation as well as different adaptive  
205 trajectories with patients.

206

207 We theorized that patients with increased numbers of clustered Hip variants might be easier to  
208 interpret from a trait perspective, as Hip variants from the same patient-specific lineages would  
209 be both closely related genetically and exposed to the same treatment and immune-related  
210 selection pressures. We made a focused plot including only isolates associated with 5 patient-  
211 specific lineages that exhibit continuous periods of Hip variants (as classified in association  
212 with Figure 5) and have been sampled for more than 2 years since first detection (Figure 3B).  
213 Two of 5 lineages evolve in a directed fashion (dashed directional lines) based on time since  
214 detection (marker fill color). However, we do not see consistent separation of Hip variants from  
215 Lop variants within each lineage by other phenotypes, even within directionally evolving  
216 lineages – they co-occur across isolate age and direction of adaptation trajectory.

217

218 In a final related assessment, we contrast just the appearance of ciprofloxacin resistance versus  
219 Hip variant for all study lineages. In seven lineages (17%), a Hip isolate was detected before  
220 resistance emerged. In five lineages, resistance emerged in the absence of the Hip phenotype,  
221 and in an additional seven cases, a resistant isolate was detected before a Hip isolate. This  
222 supports that Hip and increased ciprofloxacin MIC are two independent phenotypes, which  
223 may be concurrently selected for over time as patients undergo treatment.


224

225 *Genetic evolution of the persister phenotype is not convergent across patient-specific lineages*

226 Sequencing and identification of genetic variations accumulating within each clone type have  
227 been previously performed for most of the isolates used in this study; furthermore, genes  
228 targeted in convergent evolution were identified by the significant enrichment of observed  
229 lineages with mutations in those genes compared to the number of lineages expected to have  
230 mutations in the same genes according to genetic drift (derived from a simulated evolution  
231 where lineages accumulate an equivalent number of mutations randomly for 1000 independent  
232 evolution simulations) (17). We split our dataset into Hip and Lop variants, and then performed  
233 this same observed versus expected lineage enrichment analysis for each population (see  
234 Materials and Methods for further details). A comparison of the significantly enriched mutated  
235 genes for Hip versus Lop populations allowed us to identify 9 candidate ‘hip’ genes targeted



236 in Hip variants when compared to Lop variants (Figure 4A). However, a maximum of 4 (15%)  
 237 Hip-containing lineages accumulated non-synonymous mutations in the same enriched ‘hip’  
 238 gene, which is a weak signal of convergent evolution within our 26 lineages with Hip variants  
 239 (hereafter referred to as Hip<sup>+</sup> lineages). Moreover, there was a surprising lack of the most  
 240 prominent ‘hip’ genes identified in *in vitro* studies and screens (Table S1).  
 241



**Figure 4. Survival following antibiotic treatment for clinical *P. aeruginosa* isolates harboring an intact and inducible copy of *rpoN*.** (A) Mutated genes enriched in Hip versus Lop isolates as assessed from a convergent evolution perspective accounting for lineage adaptation. Lineage enrichment ratio was calculated by dividing lineage-based gene mutation enrichment within Hip variants by that within Lop variants for each gene. Top Hip-linked genes were selected via the following criteria: greater than 2 lineages presenting mutations in that gene in the Hip population and a lineage enrichment ratio greater than 2. (B) Hip (blue bars) and Lop (grey bar) isolates with *rpoN* null mutations were grown for 24 hours then treated with ciprofloxacin (100  $\mu$ g/ml). Cultures were plated for survival following 24 hours of treatment, and colony forming units (CFU) were quantified. The expression of intact *rpoN* was induced via the addition of 0.02% arabinose (black bars). Data presented are the mean  $\pm$  SD,  $n \geq 3$ . Significance was determined using unpaired t-test (\*\*  $p < 0.01$ , \*  $p < 0.05$ , ns, not significant).

242 We experimentally evaluated the contribution of our top ‘hip’ gene, the sigma factor encoding  
 243 gene *rpoN* (Figure 4B). The gene encodes the RNA polymerase  $\sigma^{54}$  subunit, and is among the  
 244 general pathoadaptive genes previously identified in a subset of our clinical isolate collection  
 245 (17). The gene was mutated in 19 isolates from five distinct lineages, and of these 18 isolates  
 246 were also Hip (of 4 Hip<sup>+</sup> lineages). The specific *rpoN* mutant alleles in these isolates differ, but  
 247 in all cases premature translational stop codons were identified. To evaluate the impact of *rpoN*  
 248 mutations in affected mutants, two representative Hip variants (from the DK06-P4405 lineage)  
 249 with differing ciprofloxacin sensitivities (an MIC of .25 and 4, respectively) and the one Lop  
 250 isolate (from the DK32-P8203 lineage, MIC of 4) were selected for further characterization  
 251 (see Materials and Methods for more details). Each isolate was transformed with a plasmid  
 252 carrying an intact and inducible copy of *rpoN* (*prpoN*). When RpoN function was restored in  
 253 each of the Hip variants, the persister level was reduced for both (which also illustrated the

254 phenotype's independence from ciprofloxacin sensitivity level). In contrast, the Lop isolate did  
255 not show any changes in persister class when complemented with an intact copy of *rpoN*.  
256 Additionally, a PAO1 *rpoN* deletion mutant did not show a difference in persister phenotype  
257 when compared to wild-type PAO1 (data not shown). This highlights the complex role of  
258 lineage background on the persister phenotype – RpoN contributes to the persister phenotype,  
259 but lineage-based factors such as potentiating mutations and/or background genotypic features  
260 enable this contribution.

261


262 *Hip variants accumulate in patients over time (via diverse incidence patterns)*

263 The lack of strong genetic signatures differentiating Hip from Lop isolates motivated us to  
264 examine the temporal dynamics of high persister incidence. Our isolate collection is unique  
265 due to the coverage of each patient's initial colonization phase based on the patient's age and  
266 clinical history. In half of the patients, the earliest bacterial isolate is also the first-ever  
267 identified *P. aeruginosa* in the clinic. We can thus estimate the emergence of the Hip phenotype  
268 as *P. aeruginosa* adapts from a wild type-similar naïve state into an adapted persistent  
269 pathogen. Previous findings have indicated that the number of Hip variants from a lineage may  
270 increase over time as the bacteria adapt to the antibiotic pressure in the host, and that once a  
271 Hip isolate is observed, it is assumed to persist in the infecting population of the patient (5, 6).

272

273 To illustrate the range of persister dynamics we observe, we grouped each lineage by an array  
274 of descriptors. The lineage descriptors include Hip presence versus absence (Hip<sup>+</sup> vs Hip<sup>-</sup>),  
275 transience of the lineage (whether it appears for less than 2 years and less than half the length  
276 of a patient's infection and is afterwards replaced by another lineage), continuity of Hip  
277 variants (whether Hip variants are consistently present for at least 3 sampling dates in a row),  
278 dominance of Hip variants (whether Hip variants make up at least 2/3 of all collected isolates  
279 of a lineage), and whether a Hip variant initiates the lineage. Figure 5A shows the ordered  
280 distribution of the lineages in 10 different groups based on the distinct sets of descriptors,  
281 illustrating both the complexity of our samples and the diversity of lineage Hip dynamics. An  
282 assessment of the number of lineages in each group shows the following: 1) 29 of 75 lineages  
283 are Hip<sup>+</sup>, 2) 29 of 46 Hip<sup>-</sup> lineages are transient, while 3 Hip<sup>+</sup> lineages are transient, 3) 20 Hip<sup>+</sup>  
284 lineages have Hip variants appear in a seemingly stochastic manner after colonization is  
285 initiated versus 9 lineages that exhibit continuous periods and/or dominance of Hip variants,  
286 and 4) 10 lineages have initiating Hip<sup>+</sup> variants. Thus, the fraction (25.7%, Fig. 2) of total  
287 isolates with a Hip phenotype appears to be distributed over a subset of lineages (38.6%) in

288 both stable (continuous/dominant) and stochastic patterns of incidence, rather than present in  
 289 every evolving lineage.  
 290



**Figure 5. Persister incidence patterns from a lineage-based perspective.** (A) Lineages were classed according to several nested characteristics: transient versus non-transient lineages, Hip presence, continuous periods of isolated Hips, lineage-initiating Hips, and Hips dominating a lineage. Lineages representing each combination of traits are shown on the left while characteristic sets are identified and enumerated for the entire collection on the right. (B) Hip<sup>-</sup> (gray circles) and Hip<sup>+</sup> (blue diamonds) show the continuous patient count of Hip<sup>-</sup> patients versus Hip<sup>+</sup> patients for the prior year of colonization, while the accumulating count of Hip<sup>+</sup> patients from time 0 is shown by black diamonds. (C) Transient lineages (lineages of shorter than 2 years duration, less than 50% of total patient infection length, and which are followed by a new lineage's appearance) are significantly associated with the Hip<sup>-</sup> phenotype, while non-transient lineages are associated with the Hip<sup>+</sup> phenotype based on Pearson's chi-squared test. Transience-unclassifiable lineages of shorter than 2 years' duration at the end of a patient's collection period are shown for context.

291 Given the apparent variability in Hip presence and pattern, we examined the incidence of Hip  
 292 variants within patients over time as shown in Figure 5B. Here we plot the continuous counts


293 of patients exhibiting the Hip phenotype (Hip<sup>+</sup>) versus no Hip presence (Hip<sup>-</sup>) within the  
294 previous year (dashed lines), which illustrates both the number of patients assessed at a given  
295 colonization age and the increasing distribution of Hip<sup>+</sup> versus Hip<sup>-</sup> phenotypes over time. An  
296 inspection of the number of patients that have exhibited a Hip variant at least once by a certain  
297 age of colonization (solid line) shows that the number of patients with Hip variants increases  
298 over time, ultimately affecting 65% of the patients in our study cohort by 7 years of  
299 colonization. Next, we evaluated the relationship between lineage transience and Hip presence.  
300 In addition to our other classifications, we mark as ‘New’ the lineages present for less than 2  
301 years in a patient at the end of our study period since we cannot classify them as transient or  
302 non-transient without additional samples. Of the 56 remaining lineages, non-transient lineages  
303 are significantly associated with the Hip<sup>+</sup> lineage status, while transient lineages are  
304 significantly associated with the Hip<sup>-</sup> lineage status (Fig. 5C). Thus, a given patient often has  
305 multiple infecting lineages, but the Hip<sup>-</sup> lineages are much more likely to disappear over the  
306 course of infection. In summary, we find that despite variable incidence patterns, a clear  
307 majority of patients are infected by Hip<sup>+</sup> lineages, and these lineages have a significant  
308 persistence advantage in comparison to Hip<sup>-</sup> lineages over time, suggesting that the Hip  
309 phenotype carries a fitness increase in antibiotic-treated patients.

310

### 311 *Hip variants show increased fitness in patient-similar biofilms*

312 Having shown that Hip variants appear at high frequency in *P. aeruginosa* populations  
313 infecting CF airways, and that the Hip phenotype is dissociated from other adapting traits such  
314 as antibiotic resistance, we next asked if Hip variant bacteria are able to survive antibiotic  
315 treatment better than Lop bacteria with similar antibiotic susceptibilities and growth properties.  
316 This fitness test should be carried out in conditions as close as possible to those in which the  
317 Hip variants had been selected. We therefore simulated antibiotic treatment of CF patients in a  
318 recently developed biofilm Pharmacokinetic/Pharmacodynamic system, in which the bacteria  
319 are challenged with antibiotics in much the same way as in patients (24). We chose this model  
320 because *P. aeruginosa* often appears as biofilms in lungs of CF patients (25), because biofilms  
321 have been shown to harbor increased levels of persister cells (26), and because our model  
322 mimics the bacterial exposure to ciprofloxacin treatment as described for CF patients (24, 27).  
323 The isolates were chosen based on the criteria that 1) the respective times of isolation since  
324 their first detection in the CF lungs were similar, 2) they had similar MIC values for  
325 ciprofloxacin, 3) similar growth rates, and 4) belong to the same clone type. Two independent  
326 Hip/Lop pairs were differentially tagged with YFP (Hip) or CFP (Lop). All strains formed

327 biofilms with comparable biomasses in the flow-cell system (data not shown). Hip and Lop  
328 cells were then mixed 1:1 and allowed to form a biofilm. Representative images of one of the  
329 Hip/Lop biofilms are shown before (Fig. 6A) and after treatment with ciprofloxacin (Fig. 6B).  
330



**Figure 6. Fitness comparison of Lop and Hip isolates in biofilm conditions.** (A) Lop (tagged with CFP) and Hip (tagged with YFP) isolates were cocultured and allowed to form biofilms in a flow-cell model for 72 hours. Mixed biofilms were treated for 24 hours with ciprofloxacin (4  $\mu\text{g/ml}$ ). Propidium iodine (PI) was added to visualise dead cells (red). (B) Biomass was quantified for each population. Significant differences in biomass following treatment were determined using unpaired t-test (\*\*\*)  $p < 0.001$ .

331 The majority of Lop bacteria were located close to the glass substratum with the Hip population  
332 proliferating at the external surface of the biofilm, facing the liquid flow. The addition of  
333 ciprofloxacin preferentially killed the Lop population leaving the Hip population relatively  
334 unaffected by the antibiotic. COMSTAT analysis confirmed this changed population structure  
335 after ciprofloxacin addition (Fig. 5C). Similar results were observed for the second Hip/Lop  
336 pairing (data not shown). This documentation of a Hip associated fitness increase in an  
337 antibiotic containing environment is all the more striking, as it has been shown previously that  
338 ciprofloxacin treatment of flow-cell biofilms preferentially kills the surface sub-populations of  
339 micro-colonies (28) – yet the Hip cells on the colony surfaces survive much better than the  
340 internal Lop bacteria under treatment with ciprofloxacin.

341

## 342 Discussion

343 In this communication, we focus our interest on the clinical significance of the persister  
344 phenotype, mapping the prevalence of persisters in a large, aligned cohort of patients under  
345 intensive antibiotic treatment in a 10 year period (17, 27). Of over 450 *P. aeruginosa* isolates  
346 from the airways of 40 young CF patients (75 lineages in total), 26% of the isolates were scored  
347 as Hip using a high-throughput screening approach (Fig. 1). The validity of the screen was  
348 confirmed using traditional persister assays, suggesting our results provide a reliable reflection

349 of the frequency of occurrence of Hip isolates infecting CF patients (Fig. 2). We show that the  
350 isolates display different levels of persisters, in accordance with the variance of persister levels  
351 previously found between species and within strains (29–31). We have previously shown that  
352 most adaptive changes occur during the first few years of colonization, which matches our  
353 objective of searching for signs of increased fitness of Hip variants in patients treated  
354 continuously with antibiotics (20, 32). Here, we show that in a young CF patient cohort  
355 impacted by early longitudinal colonization by *P. aeruginosa* strains, Hip variants were  
356 sampled from 65% of the patients (N=26) during a 10-year observation window. Our analysis  
357 is a new and important comparative baseline for moving examinations of this clinically  
358 important phenotype towards effective surveillance, impact assessment, and eventual control  
359 in actual clinics.

360 In the early years of infection since the first detection of *P. aeruginosa*, the Hip phenotype  
361 appeared and disappeared over time (Fig. 5A). However, the number of patients that exhibit  
362 Hip variants increases over time (Fig. 5B), suggesting a selective advantage of this phenotype  
363 during the continuation of antibiotic therapy. In support of this suggestion, the majority of  
364 lineages that showed short-term colonization were made up of only Lop variants, which may  
365 in fact explain why they were unable to establish a persistent infection (Fig. 5C). These in-  
366 patient data support the hypothesis that the Hip phenotype may generally have increased fitness  
367 in the antibiotic containing lung environment. It is, however, very important to note that neither  
368 dominance nor continuous presence of Hip variants is observed very frequently (Fig. 5A). It is  
369 likely that fitness trade-offs and clonal interference impact on the fitness properties and the  
370 persistence level of Hip variants, including other fitness increasing mutations associated with  
371 antibiotics in the lung environment which accumulate in the bacterial populations (13).

372  
373 Multiple relationships between the Hip phenotype and other phenotypic traits such as growth  
374 rate and antibiotic resistance have been suggested in the literature. Reduced growth rates have  
375 been associated with high persister phenotypes in *E. coli* (23), but other studies point out that  
376 there is no correlation between the mean growth rates of isolates and the observed Hip  
377 phenotype (33–35). In our data, the ratio of Hip to Lop variants increases as traits adapt over  
378 time, and therefore the average growth rate of all Hip variants is reduced versus Lop variants;  
379 however, we did not observe a consistent link between reduced growth rates and Hip phenotype  
380 among the clinical isolates of *P. aeruginosa*. Drug-tolerant cells have also been proposed to  
381 facilitate evolution of true antibiotic resistance in *E. coli in vitro* (36). Intermittent antibiotic

382 exposure of a batch culture of *E. coli* first selected for mutant clones harboring tolerance  
383 mutations that increased the growth lag-time, during which tolerance to killing by ampicillin  
384 favored a subsequent selection for mutations that increased the MIC. Though *P. aeruginosa* in  
385 the CF lung is also exposed to fluctuating concentrations of antibiotic over the course of  
386 infection, we did not observe any obvious contingency in which Hip variants precede  
387 antibiotic-resistant mutants. In contrast, our data suggest that Hip variants and resistant clones  
388 evolve independently in patients under antibiotic selection pressure as has been suggested by  
389 comparative studies of lab strains (37) and that each of the phenotypes contributes to a  
390 continued population survival and hence establishment of long term infection. In general, the  
391 results described here suggest that Hip variants arise stochastically in infected patients treated  
392 with antibiotics, and that this phenotype is one of several making it possible for the bacteria to  
393 persist for extended periods of time. As Hip variants are present in the majority of our young  
394 patients by the end of the study period, the persister phenotype may be an early advantageous  
395 adaptation in contrast to the adaptive timeline of other traits (20); effective comparisons  
396 between stochastic persistence versus more fixed changes such as adjustment of growth rate or  
397 antibiotic susceptibility should be pursued in the future.

398 In many ways, persisters have been investigated in line with previous investigations of  
399 antibiotic resistance, which has resulted in a similar parallel search for genetic explanations for  
400 large variations in persister levels in bacterial populations treated with antibiotics. In a study  
401 of clinical isolates from patients with urinary tract infection, it was shown that a gain-of-  
402 function mutation in the HipA toxin is common among *E. coli* Hip isolates (38). The most  
403 common mutated genetic target we identified among *P. aeruginosa* Hip in the present study is  
404 the RpoN sigma factor. Restoring a functional RpoN in these clinical isolates caused a  
405 reduction in persisters, but knocking out this gene in a laboratory strain had no effect. This  
406 suggests that the RpoN mutation only confers a persister phenotype in the context of additional  
407 mutations acquired by the *P. aeruginosa* Hip strains, underscoring the complexity of evolution  
408 occurring in the lungs of CF patients. It is also noteworthy that we do not see the appearance  
409 of mutations, which previously have been associated with Hip phenotypes *in vitro*. As the  
410 RpoN mutation was present in only 15% of the Hip isolates in our large, longitudinal collection,  
411 we conclude that a Hip phenotype may derive from a diverse array of genetic changes, and it  
412 is likely that more than one mutation often determines the persister level in the respective  
413 bacterial populations. It is possible that this difference reflects the multiple and dynamic  
414 selection pressures *in vivo*, which challenge Hip variants in antibiotic-treated populations very

415 differently from what is the situation in steady state *in vitro* conditions with only one selective  
416 force. Since nearly all published data on persister cells derive from simple *in vitro* experiments,  
417 our results further suggest that extrapolations from such *in vitro* experiments to clinical  
418 scenarios of infection cannot be made.

419 Many studies have shown the increased survival of persister cells under antibiotic treatment  
420 (23, 39) and leveraged this in screens for genetic determinants of persistence, but few have  
421 evaluated the fitness of Hip versus Lop variants in direct competition experiments. Stepanyan  
422 et al. assessed the fitness tradeoffs of two Hip strains (one hospital strain and one *pilH* mutant)  
423 and two Lop *P. aeruginosa* strains via competitive growth of paired Hip-Lop strains at a 1:1  
424 ratio under periodic treatment with ofloxacin in Mueller-Hinton broth; Hip strains showed  
425 higher fitness under treatment conditions and the reverse in control conditions (29). In our  
426 study, we were able to test genotypically similar isolates from a similar point in infection with  
427 other selected traits in alignment in order to characterize the selective advantage of the Hip  
428 phenotype in a biofilm under clinically-replicating antibiotic exposure (24). In two independent  
429 Lop/Hip strain pairs assessed in a PD/PK flow chamber, we show that Hip cells survived  
430 ciprofloxacin treatment far better than Lop isolates, and we further find that this survival is  
431 potentially reliant on biofilm architecture. While homogeneous monoclonal *P. aeruginosa*  
432 biofilms treated with ciprofloxacin show preferential killing of bacteria in the top layers (13,  
433 28, 40), Lop bacteria are preferentially killed in the deeper layers of the biofilm. This further  
434 supports that the Hip phenotype is associated with increased survival in the presence of  
435 ciprofloxacin under conditions simulating the in-patient environment during antibiotic  
436 treatment. It is a striking observation that the *in vitro* biofilm fitness assessment shows efficient  
437 elimination of the Lop strain in presence of ciprofloxacin, whereas in the patient lungs Hip  
438 variants rarely outcompete Lop strains despite frequent treatments with ciprofloxacin as  
439 indicated by Fig. 3A. This suggests that in the patient, direct competition is limited by the many  
440 separate niches, the influence of the host and the large lung volume, which contrasts the flow-  
441 cell biofilm system. This slow competition explains how fully adapted strains like the DK01  
442 and DK02 strains simultaneously present in a single patient may require many months or even  
443 years to finally outcompete the other (32, 41, 42).

444

445 In summary, we have shown that Hip variants of *P. aeruginosa* emerge frequently in younger  
446 CF patients. The Hip phenotype provides a selective advantage for *P. aeruginosa* populations  
447 treated with antibiotics, which may explain how persisters contribute to the establishment of



448 chronic infection. Ultimately, our results provide the first window into the evolving landscape  
449 of persistence across a whole patient cohort; as pathogens increase their fitness in patients over  
450 time, they clearly deploy the high persister phenotype as an important component in their  
451 survival repertoire and can do so from the earliest stages of infection. It is still premature to  
452 conclude that the high-persister phenotype described here differs from what has been identified  
453 as Hip in *in vitro* experimental conditions, but we find a much broader bacterial repertoire for  
454 survival in antibiotic containing environments than so far anticipated. Hip variants do not seem  
455 to be mutated in genes previously found from *in vitro* experiments to associate with Hip or in  
456 any strongly conserved genetic route. *In vitro*, antibiotic-mediated selection is the only  
457 selective force, whereas, in the patients, antibiotics constitute one among several parallel  
458 selective forces. In support of this concept, Hip variants emerge alongside other beneficial trait  
459 adaptations but association with any single trait is inconsistent. We suggest that the difference  
460 in selection pressure when comparing *in vitro* and *in vivo* environmental conditions including  
461 antibiotics results in highly different evolutionary trajectories. With our investigation, we  
462 provide an important platform for broader clinically based studies and contribute important  
463 new context for monitoring and one day hopefully preventing the high persister phenotype in  
464 the clinic.

465

## 466 **Materials and Methods**

467 *Strain collection.* In total, we analyzed 467 *P. aeruginosa* airway isolates from young CF  
468 patients followed at the Copenhagen CF-clinic after excluding twenty-six isolates that had  
469 acquired mutS and/or mutL mutations resulting in hypermutability. Of these 467 isolates, 388  
470 isolates from 34 patients were described previously in Marvig et. al. (17) and the remaining  
471 isolates were taken from six previously undescribed patients. The isolates were collected and  
472 stored at the Department of Clinical Microbiology at Rigshospitalet, Copenhagen, Denmark,  
473 between 2002 and 2014. Of the patients included in this study, 35% were diagnosed as  
474 chronically infected with *P. aeruginosa* by the end of the study period. We defined chronicity  
475 based on the Copenhagen CF Centre definition, whereby either *P. aeruginosa* has been  
476 detected in six consecutive monthly sputum samples or fewer consecutive sputum samples  
477 combined with observation of two or more *P. aeruginosa*-specific precipitating antibodies (27,  
478 43). Intermittently colonized patients were defined as patients where at least one isolate of *P.*  
479 *aeruginosa* is detected, and normal levels of precipitating antibodies against *P. aeruginosa*  
480 were observed. Phenotyping data for 407 isolates of this strain collection (growth rate in LB,

481 adhesion in LB, and ciprofloxacin MIC) have been previously published (20). We include trait  
482 measurements for 60 additional isolates from the same phenotypic screens conducted as  
483 described below, and all available trait data is provided in Supplementary Dataset 1 along with  
484 the persister classification of each isolate and descriptive data such as patient origin, genotype,  
485 date of sampling, and SRA number.

486

487 *Minimum Inhibitory Concentrations (MIC)* for ciprofloxacin and tobramycin were determined  
488 using E-test methodology according to the manufacturer's recommendations (Liofilchem®,  
489 Italy).

490

491 *Generation times.* Bacteria were grown overnight in LB, diluted to an OD of 0.05 in 96 well  
492 plates containing 150  $\mu$ L of LB medium, and incubated for 20 hours at 37°C with constant  
493 shaking. OD 630 nm measurements were taken every 20 minutes using a microplate reader  
494 (Holm & Halby, Copenhagen, Denmark/Synergy H1). Generation times (Td) were determined  
495 on the best-fit line of a minimum of 3 points during exponential growth of the bacterial isolate.  
496 Growth rates ( $\text{hr}^{-1}$ ) were calculated using the formula  $\log(2)/\text{Td} \times 60$ .

497

498 *Adhesion.* Adhesion was measured via attachment assays in 96-well plates using NUNC peg  
499 lids and 96 well plates with 150 $\mu$ L Luria broth medium. OD<sub>600nm</sub> was measured after incubation  
500 for 20 hours at 37°C and subsequently, a “washing microtiter plate” with 180 $\mu$ L PBS was used  
501 to wash the peg lids and remove non-adhering cells. After transfer of the peg lids to a microtiter  
502 plate containing 160 $\mu$ L 0.01% crystal violet (CV), they were left to stain for 15 min. To remove  
503 unbound crystal violet, the lids were then washed again three times in three individual  
504 “washing microtiter plates” with 180 $\mu$ L PBS. Adhesion was measured by detaching adhering  
505 CV stained cells through washing the peg lids in a microtiter plate containing 180 $\mu$ L 99%  
506 ethanol. An ELISA reader was then used to measure the CV density at OD<sub>590nm</sub>. (Microtiter  
507 plates were bought at Fisher Scientific, NUNC Cat no. 167008, peg lids cat no. 445497).

508

509 *High-throughput screening for Hip mutants.* To determine the frequency at which *P.*  
510 *aeruginosa* Hip mutants emerge in CF patients, we screened 467 isolates for ciprofloxacin  
511 tolerance. Stock 96-well microtiter plates containing 4 technical replicates of each isolate  
512 stored in glycerol (25 % v/v) were prepared and stored at -80°C. Using a 96-well spot  
513 replicator, bacteria were transferred from the stock plates into sterile 96-well microtiter plates

514 containing 150  $\mu$ l of Lysogeny Broth (LB) media. Plates were incubated statically for 48 hours  
515 at 37°C until the bacteria reached the stationary phase of growth. To determine the initial  
516 viability of bacteria in each well, the replicator was used to spot bacteria onto LB agar plates.  
517 Subsequently, 100  $\mu$ g/ml of ciprofloxacin was added to each well and the microtiter plates were  
518 incubated statically for a further 20-24 hours at 37°C. Serial dilutions were performed in 96  
519 well microtiter plates containing 0.9 % NaCl using an automated fluid handling robot  
520 (Viaflo3844/ Integra Biosciences AG). Each dilution was spotted onto LB agar plates using  
521 the replicator and plates were incubated at 37°C for at least 24 hours. The growth of the bacteria  
522 was compared by counting colonies whenever possible and visually inspecting growth on the  
523 plates before and after antibiotic treatment. The experiment was performed in duplicate.

524

525 *Persister assay validation.* Persister assays were performed on 23 isolates to validate the high  
526 throughput persister screen. *P. aeruginosa* were inoculated in 3 ml of LB media in 10 ml tubes  
527 and incubated for 48 hours at 37°C with shaking at 250 rpm. Following incubation, each culture  
528 was serially diluted using sterile 0.9 % NaCl, plated onto LB agar and incubated at 37°C to  
529 determine the initial colony forming units (CFU). The remaining culture was treated with 100  
530  $\mu$ g/ml of ciprofloxacin and incubated for a further 24 hours at 37°C with shaking. Following  
531 treatment, cells were washed and diluted in sterile 0.9 % NaCl, spot plated onto LB agar and  
532 incubated for 24 hours at 37°C. Bacteria survival was measured by counting (CFU) per ml.

533

534 *Pharmacokinetic/Pharmacodynamic (PK/PD) flow chamber biofilm model.* For fitness  
535 experiments, we used a PK/PD biofilm model system combined with confocal laser-scanning  
536 microscopy. This system simulates the changing antibiotic concentrations in CF patients during  
537 intravenous dosing in addition to retaining a similar profile of antibiotic decay as the one taking  
538 place in CF patients (24). First, Hip and Lop isolates were differentially tagged with a yellow  
539 fluorescent protein (YFP) or cyan fluorescent protein (CFP) respectively (13). Flow chambers  
540 were inoculated with a 1:1 mixture of Hip and Lop bacteria (each isolate had an initial OD<sub>600</sub>  
541 of 0.5). Bacteria were incubated for one hour at 30 °C, then nutrient flow was applied to each  
542 chamber (40x diluted LB at a rate of 20 ml/h using a Watson Marlow 205S peristaltic pump).  
543 Biofilms were allowed to form for 72 hours, at which point flow was stopped and medium  
544 containing ciprofloxacin was added. Peak ciprofloxacin concentrations were calculated to be 4  
545 mg/L based on PK parameters generated from healthy patients and CF patients (44). The  
546 medium was pumped from the dilution flask through the antibiotic flask to the flow chambers  
547 at a constant rate calculated to mimic the elimination rate constant of the antibiotic for 24 hrs.

548 A confocal laser-scanning microscope (Zeiss LSM 510) equipped with an argon/krypton laser  
549 and detectors was used to monitor YFP (excitation 514 nm, emission 530 nm), CFP (excitation  
550 458 nm, emission 490 nm), and dead cells (propidium iodine, excitation 543 nm, emission 565  
551 nm). Multichannel simulated fluorescent projections (SFPs) and sections through the biofilms  
552 were generated using Imaris software (Bitplane AG, Switzerland). The images were later  
553 analyzed using COMSTAT (45). The PK/PD biofilm experiments were performed using two  
554 independent Hip/Lop isolate pairs. Pairs were taken from the same patient at a similar time  
555 since first detection and had similar growth rates and ciprofloxacin MICs (Table S2). The data  
556 presented are from 2 biological experiments with 4 independent images taken from each  
557 experiment.

558

559 *Lineage-based genetic analysis.* To generate a list of mutated genes associated with the Hip  
560 phenotype, we used previously generated whole-genome sequencing data and variant calling  
561 filtered to obtain nonsynonymous mutations that had accumulated within a lineage after the  
562 first isolate (17) to evaluate differential mutation patterns for Lop and Hip variants for 388  
563 sequenced isolates. To identify genes that were mutated more than would have been expected  
564 by drift/random mutation while accounting for lineage-based mutation accumulation over time,  
565 we adapted a statistical analysis of the relative mutation enrichment by lineage. After  
566 separating Lop and Hip variants, we compared the mutated-gene lineage enrichment ratios for  
567 each group - the number of lineages with observed mutation(s) in a given gene divided by the  
568 number of lineages expected to have mutations in that gene according to random mutation.  
569 This enrichment metric was obtained as follows for each group: we determined the observed  
570 number of lineages mutated (sum-obs) in each gene. Then we estimated the average number of  
571 lineages (avg-exp) that would have been mutated in each gene if mutations were spread out  
572 randomly over the PAO1 genome. Using a random-roulette algorithm, the number of genes  
573 that were observed to be mutated in a given lineage was spread out over the PAO1 genome for  
574 1000 iterations, providing a  $m_{\text{gene}}$  by  $n_{\text{iteration}}$  matrix of randomly mutated gene profiles for each  
575 lineage. For the same iteration  $n$  across all lineages, it was noted whether a given gene was  
576 mutated. This allowed us to determine an average number of lineages expected to be mutated  
577 over 1000 iterations. If a gene was hit by chance more than once in a single iteration, this would  
578 still only be denoted as one hit; this is in alignment with our observed mutation assessment,  
579 where multiple isolates could be hit in the same gene but we only noted whether or not the  
580 lineage was hit by unique mutations in the specific gene. After obtaining the relative

581 enrichment by lineage, a Poisson distribution was used to calculate the probability of the  
582 observed given random drift (expected). We also divided the lineage enrichment metric for  
583 genes mutated in Hip variants by that for Lop variants to obtain a lineage enrichment ratio to  
584 identify targeted genes particularly impactful in the evolution of the Hip population.

585

586 *Lineage set analysis and principle component analysis.* Analyses were conducted in RStudio  
587 v. 1.0.143 and R v. 3.4.0. Lineage set analysis was performed using UpSetR v. 1.3.3 in R.  
588 Principal component analysis was performed in R using ‘prcomp’ with centered and scaled  
589 phenotype data (Supplementary Dataset 1).

590

591 *Genetic manipulation.* We selected two Hip variants and one Lop variant with mutations in  
592 *rpoN* for validation of a *rpoN*-associated Hip phenotype. Each of the *rpoN* Hip variants had a  
593 single nucleotide deletion (T, nucleotide position 4993951 on PAO1 reference genome), which  
594 resulted in a frameshift following the 360<sup>th</sup> amino acid (aa) and premature truncation of the  
595 protein resulting in a 381aa polypeptide (RpoN has 497aa in the PAO1 reference genome). The  
596 Lop isolate had a single nucleotide polymorphism (C→T at nucleotide 112) producing a  
597 premature stop codon at the 38<sup>th</sup> aa. An *rpoN* expression plasmid was constructed for gene  
598 complementation experiments. The *rpoN* gene from PAO1 was PCR amplified using primers  
599 *rpoN* forward (5’ GAATTCATGAAACCATCGCTAGTCCTC 3’) and *rpoN* reverse  
600 (5’AAGCTTTCACACCAGTCGCTTGCGCTC 3’). The product was cloned into pHERD28T  
601 (46), using EcoRI and HindIII restriction sites and the resulting plasmid (*prpoN*) was used to  
602 transform *E. coli* OneShot Top10 (Thermo Fisher Scientific). The presence of the plasmid was  
603 verified by PCR and DNA sequencing then introduced into selected clinical *P. aeruginosa*  
604 isolates using biparental conjugation. Trimethoprim (TMP, 300 µg/ml) was used for selection  
605 and arabinose (0.02 %) was used for *rpoN* induction. Each isolate was transformed with empty  
606 pHERD28T and isolates with and without arabinose induction were included as controls.

607

## 608 **Acknowledgments**

609 This work was supported by Cystic Fibrosis Foundation Pilot and Feasibility Award to KL.  
610 HKJ was supported by The Novo Nordisk Foundation as a clinical research stipend  
611 (NNF12OC1015920), by Rigshospitalets Rammebevilling 2015-17 (R88-A3537), by  
612 Lundbeckfonden (R167-2013-15229), by Novo Nordisk Fonden (NNF15OC0017444) by  
613 RegionH Rammebevilling (R144-A5287) and by Independent Research Fund Denmark (DFF-

614 4183-00051). JAB was supported by postdoctoral fellowships from the Whitaker Foundation  
615 and the Cystic Fibrosis Foundation. We would like to thank Marilena Falcone, Ph.D. for  
616 supplying pHERD28T and technical guidance related to cloning.

617

### 618 **Conflict of interest**

619 The authors declare no competing financial interests.

620

### 621 **Author Contributions**

622 SM and KL designed the study. HKJ collected all the bacterial isolates. BM, DRC, and JH  
623 performed all experiments. JAB and LMS performed genetic and lineage analysis. All authors  
624 contributed to the writing of the manuscript. All authors approved the final version.

625

### 626 **Ethics approval**

627 The local ethics committee at the Capital Region of Denmark (Region Hovedstaden)  
628 approved the use of the stored *P. aeruginosa* isolates: registration number H-4-2015-FSP.

629

### 630 **Supplementary information**

631 **Table S1.** Comparison of persister genes identified in previous *P. aeruginosa* studies to  
632 mutated genes highlighted by our lineage analysis.

633 **Table S2.** Characteristics of Hip/Lop pairs analyzed using the PK/PD biofilm model.

634 **Supplementary Dataset 1.** Phenotypic dataset for all isolates.

635

636

### 637 **References**

- 638 1. Bigger JW (1944) Treatment of staphylococcal infections with penicillin. *Lancet* 2(6320):497–  
639 500.
- 640 2. Wood TK, Knabel SJ, Kwan BW (2013) Bacterial persister cell formation and dormancy. *Appl*  
641 *Environ Microbiol* 79(23):7116–7121.
- 642 3. Rowe SE, Conlon BP, Keren I, Lewis K (2016) Persisters: Methods for Isolation and  
643 Identifying Contributing Factors—A Review. *Methods Mol Biol* 1333:17–28.
- 644 4. Lewis K (2010) Persister Cells. *Annu Rev Microbiol*.  
645 doi:10.1146/annurev.micro.112408.134306.
- 646 5. LaFleur MD, Qi Q, Lewis K (2010) Patients with long-term oral carriage harbor high-persister  
647 mutants of *Candida albicans*. *Antimicrob Agents Chemother* 54(1):39–44.
- 648 6. Mulcahy LR, Burns JL, Lory S, Lewis K (2010) Emergence of *Pseudomonas aeruginosa*  
649 strains producing high levels of persister cells in patients with cystic fibrosis. *J Bacteriol*

- 650 192(23):6191–9.
- 651 7. Salsgiver EL, et al. (2016) Changing Epidemiology of the Respiratory Bacteriology of Patients  
652 With Cystic Fibrosis. *Chest* 149(2):390–400.
- 653 8. Parkins MD, Somayaji R, Waters VJ (2018) Epidemiology, Biology, and Impact of Clonal  
654 *Pseudomonas aeruginosa* Infections in Cystic Fibrosis. *Clin Microbiol Rev* 31(4):1–38.
- 655 9. Dalemans W, et al. (1991) Altered chloride ion channel kinetics associated with the  $\Delta F508$   
656 cystic fibrosis mutation. *Nature* 354(6354):526–528.
- 657 10. Boucher RC (2007) Cystic fibrosis: a disease of vulnerability to airway surface dehydration.  
658 *Trends Mol Med* 13(6):231–240.
- 659 11. Folkesson A, et al. (2012) Adaptation of *Pseudomonas aeruginosa* to the cystic fibrosis airway:  
660 an evolutionary perspective. *Nat Rev Microbiol* 10(12):841–51.
- 661 12. Gibson RL, Burns JL, Ramsey BW (2003) Pathophysiology and management of pulmonary  
662 infections in cystic fibrosis. *Am J Respir Crit Care Med* 168(8):918–951.
- 663 13. Frimodt-Møller J, et al. (2018) Mutations causing low level antibiotic resistance ensure  
664 bacterial survival in antibiotic-treated hosts. *Sci Rep* 8(1):1–13.
- 665 14. Burns JL, et al. (1999) Effect of Chronic Intermittent Administration of Inhaled Tobramycin  
666 on Respiratory Microbial Flora in Patients with Cystic Fibrosis. *J Infect Dis* 179(5):1190–  
667 1196.
- 668 15. van Gestel J, Vlamakis H, Kolter R (2015) Division of labor in biofilms: the ecology of cell  
669 differentiation. *Microbiol Spectr* 3(2):1–24.
- 670 16. Moyed HS, Bertrand KP (1983) *hipA*, a newly recognized gene of *Escherichia coli* K-12 that  
671 affects frequency of persistence after inhibition of murein synthesis. *J Bacteriol* 155(2):768–  
672 775.
- 673 17. Marvig RL, Sommer LM, Molin S, Johansen HK (2015) Convergent evolution and adaptation  
674 of *Pseudomonas aeruginosa* within patients with cystic fibrosis. *Nat Genet* 47(1):57–64.
- 675 18. Eliopoulos GM, Gardella A, Moellering RC (1984) In vitro activity of ciprofloxacin, a new  
676 carboxyquinoline antimicrobial agent. *Antimicrob Agents Chemother* 25(3):331–335.
- 677 19. Su HC, et al. (2010) The development of ciprofloxacin resistance in *Pseudomonas aeruginosa*  
678 involves multiple response stages and multiple proteins. *Antimicrob Agents Chemother*  
679 54(11):4626–4635.
- 680 20. Bartell JA, et al. (2019) Evolutionary highways to persistent bacterial infection. *Nat Commun*  
681 10(1):629.
- 682 21. La Rosa R, Johansen HK, Molin S (2018) Convergent Metabolic Specialization through  
683 Distinct Evolutionary Paths in *Pseudomonas aeruginosa*. *MBio* 9(2):e00269-18.
- 684 22. Winstanley C, O’Brien S, Brockhurst MA (2016) *Pseudomonas aeruginosa* Evolutionary  
685 Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. *Trends Microbiol*  
686 24(5):327–337.
- 687 23. Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S (2004) Bacterial persistence as a  
688 phenotypic switch. *Science* (80-) 305(5690):1622–1625.
- 689 24. Haagensen JAJ, Verotta D, Huang L, Spormann A, Yang K (2015) New in vitro model to

- 690 study the effect of human simulated antibiotic concentrations on bacterial biofilms. *Antimicrob*  
691 *Agents Chemother* 59(7):4074–4081.
- 692 25. Singh PK, et al. (2000) Quorum-sensing signals indicate that cystic fibrosis lungs are infected  
693 with bacterial biofilms. *Nature* 407:762–764.
- 694 26. Spoering AL, Lewis K (2001) Biofilms and planktonic cells of *Pseudomonas aeruginosa* have  
695 similar resistance to killing by antimicrobials. *J Bacteriol* 183(23):6746–6751.
- 696 27. Johansen HK, et al. (2004) Antibody response to *Pseudomonas aeruginosa* in cystic fibrosis  
697 patients: a marker of therapeutic success? - A 30-year cohort study of survival in Danish CF  
698 patients after onset of chronic *P. aeruginosa* lung infection. *Pediatr Pulmonol* 37(5):427–432.
- 699 28. Pamp SJ, Gjermansen M, Johansen HK, Tolker-Nielsen T (2008) Tolerance to the  
700 antimicrobial peptide colistin in *Pseudomonas aeruginosa* biofilms is linked to metabolically  
701 active cells, and depends on the *pmr* and *mexAB-oprM* genes. *Mol Microbiol* 68(1):223–240.
- 702 29. Stepanyan K, et al. (2015) Fitness trade-offs explain low levels of persister cells in the  
703 opportunistic pathogen *Pseudomonas aeruginosa*. *Mol Ecol* 24(7):1572–1583.
- 704 30. Stewart B, Rozen DE (2011) Genetic variation for antibiotic persistence in *Escherichia coli*.  
705 *Evolution (N Y)* 66(3):933–939.
- 706 31. Bink A, et al. (2011) Superoxide dismutases are involved in *Candida albicans* biofilm  
707 persistence against miconazole. *Antimicrob Agents Chemother* 55(9):4033–4037.
- 708 32. Yang L, et al. (2011) Evolutionary dynamics of bacteria in a human host environment. *Proc*  
709 *Natl Acad Sci* 108(18):7481–7486.
- 710 33. Fung DKC, Chan EWC, Chin ML, Chan RCY (2010) Delineation of a bacterial starvation  
711 stress response network which can mediate antibiotic tolerance development. *Antimicrob*  
712 *Agents Chemother* 54(3):1082–1093.
- 713 34. Keren I, Minami S, Rubin E, Lewis K (2011) Characterization and transcriptome analysis of  
714 *Mycobacterium tuberculosis* persisters. *MBio* 2(3):3–12.
- 715 35. Wakamoto Y, et al. (2013) Dynamic persistence of antibiotic-stressed *Mycobacteria*. *Science*  
716 (80-) 339(6115):91–95.
- 717 36. Levin-Reisman I, et al. (2017) Antibiotic tolerance facilitates the evolution of resistance.  
718 *Science (80- )* 355(6327):826–830.
- 719 37. Vogwill T, Comfort AC, Furió V, MacLean RC (2016) Persistence and resistance as  
720 complementary bacterial adaptations to antibiotics. *J Evol Biol* 29(6):1223–1233.
- 721 38. Schumacher MA, et al. (2015) HipBA-promoter structures reveal the basis of heritable  
722 multidrug tolerance. *Nature* 524(7563):59–66.
- 723 39. Spoering AL, Vulić M, Lewis K (2006) GlpD and PlsB participate in persister cell formation  
724 in *Escherichia coli*. *J Bacteriol* 188(14):5136–5144.
- 725 40. Haagensen JAJ, et al. (2007) Differentiation and distribution of colistin- and sodium dodecyl  
726 sulfate-tolerant cells in *Pseudomonas aeruginosa* biofilms. *J Bacteriol* 189(1):28–37.
- 727 41. Markussen T, et al. (2014) Environmental Heterogeneity Drives Within-Host Diversification  
728 and Evolution of *Pseudomonas aeruginosa*. 5(5):1–10.
- 729 42. Marvig RL, Johansen HK, Molin S, Jelsbak L (2013) Genome Analysis of a Transmissible



- 730 Lineage of *Pseudomonas aeruginosa* Reveals Pathoadaptive Mutations and Distinct  
731 Evolutionary Paths of Hypermutators. *PLoS Genet* 9(9):e1003741.
- 732 43. Høiby N, et al. (1977) *Pseudomonas aeruginosa* infection in cystic fibrosis - Diagnostic and  
733 prognostic significance of *Pseudomonas aeruginosa* precipitins determined by means of  
734 crossed immunoelectrophoresis. *Scand J Respir Dis* 58(2):65–79.
- 735 44. Touw DJ, Knox AJ, Smyth A (2007) Population pharmacokinetics of tobramycin administered  
736 thrice daily and once daily in children and adults with cystic fibrosis. *J Cyst Fibros* 6(5):327–  
737 333.
- 738 45. Heydorn A, et al. (2002) Statistical Analysis of *Pseudomonas aeruginosa* Biofilm  
739 Development: Impact of Mutations in Genes Involved in Twitching Motility, Cell-to-Cell  
740 Signaling, and Stationary-Phase Sigma Factor Expression. *Appl Environ Microbiol*  
741 68(4):2008–2017.
- 742 46. Qiu D, Damron FH, Mima T, Schweizer HP, Yu HD (2008) PBAD-based shuttle vectors for  
743 functional analysis of toxic and highly regulated genes in *Pseudomonas* and *Burkholderia* spp.  
744 and other bacteria. *Appl Environ Microbiol* 74(23):7422–6.
- 745