Multimorbidity and healthcare utilization: A register-based study in Denmark

Anne Frølich^{1&2*}, Nermin Ghith¹, Michaela Schiøtz¹, Ramune Jacobsen¹, Anders Stockmarr³

¹ The Research Unit for Chronic Conditions, Center for Clinical Research and Prevention,

Bispebjerg Frederiksberg Hospital, The Capital Region, Copenhagen, Denmark

² Institute for Public Health, University of Copenhagen

³ Department of Applied Mathematics and Computer Science, The Technical University of

Denmark, Copenhagen, Denmark

* Corresponding author

Email: Anne.Froelich@regionh.dk

Abstract

Background

People with multimorbidity have reduced functional capacity, lower quality of life, and higher mortality rates and use healthcare resources more intensively than healthy people or those with a single chronic condition. The aim of this study was to explore associations between multimorbidity and use of healthcare services and the impact of socioeconomic status on utilization of hospitalizations and bed days.

Methods

The study population included all individuals aged 16 years and older who lived in the Capital Region of Denmark on January 1st, 2012. Data on chronic conditions, use of healthcare services and demographics were obtained from Danish national administrative and health registries. Zero-inflated models were used to calculate anticipated annual use of hospitalizations and bed days.

Findings

The study population comprised 1,397,173 individuals; the prevalence of multimorbidity was 22%. Prevalence was inversely related to educational attainment. For people with multimorbidity, utilization of hospitalizations and bed days increased approximately linearly with the number of chronic conditions. However, a steep increase in utilization of bed days was observed between five and six or more chronic conditions. An educational gradient in hospitalization rates and use of bed days was observed regardless of the number of chronic conditions. Educational attainment was strongly associated with healthcare utilization.

Conclusion

40 Multimorbidity was associated with a significant increase in utilization of all healthcare services in
41 Denmark. In addition, a socioeconomic gradient was observed in utilization of hospitalizations and
42 bed days.

43

Introduction

Multimorbidity is often defined as the coexistence of two or more chronic conditions in the same person [1, 2]. People with multimorbidity have decreased functional competence [3], lower quality of life [4], higher mortality rates [5] and use healthcare resources more intensively than healthy people or those with just one chronic condition [6]. Most diseases and consequences of poor health are unequally distributed among socioeconomic population groups, and socioeconomic differences are obvious in the prevalence and consequences of multimorbidity [7, 8]. The prevalence of multimorbidity is increasing internationally. The overall prevalence of multimorbidity in Ontario, Canada increased in nearly all age groups, reflecting a 40% total increase between 2003 and 2009 from 17.4% to 24.3% [9]. A study from the Netherlands reported an increase in multimorbidity prevalence from 12.7% to 16.2% between 2004 and 2011 [10]. An American study showed that the prevalence of multimorbidity rose between 2000 and 2010 from 22% to 30%, a trend that was most pronounced among people younger than age 65 [11]. Expected continuing increases in the prevalence of multimorbidity are recognized as a major public health and healthcare challenge for modern societies [9].

To understand the healthcare challenges associated with multimorbidity, the impact of multimorbidity on healthcare utilization must be carefully assessed [12, 13]. However, detailed knowledge about how multimorbidity affects healthcare utilization is incomplete. A systematic review identified 35 studies investigating relationships between multimorbidity and healthcare utilization, healthcare costs, or both. All included studies showed a positive correlation between multimorbidity and at least one aspect of utilization (physician visits and hospital care) and costs (medications, out-of-pocket spending, and total healthcare costs) for

elderly populations [14]. The included studies were from the United States (23), Europe (5), Canada (4), Asia (2), and Australia (1). Many of these study populations were large enough to enable sophisticated statistical analyses, and four were large cross-sectional studies that included 1.13 million to 1.65 million people [15-17]. Regrettably, these large studies had varying inclusion criteria, and none reported sociodemographic data that are necessary for exploring possible associations between multimorbidity, healthcare utilization, and sociodemographic status. Synthesis of the studies was not possible due to ambiguous definitions and measurements of multimorbidity, as well as a multitude of healthcare utilization outcomes.

After publication of this systematic review, several large-scale studies [18-22] exploring the relationship between healthcare utilization and the number of chronic conditions demonstrated that additional factors affecting utilization include age [18, 19], gender [18, 19], impaired activities of daily life [20], and socioeconomic status [19]. Furthermore, the impact of multimorbidity on healthcare utilization has been shown to differ across individual factors, disease combinations, healthcare systems, and regions [22, 23].

The Danish healthcare system is based on universal coverage and principles of free and equal access to healthcare for all citizens. Information on the impact of multimorbidity on healthcare utilization in the Danish healthcare system is sparse. To the best of our knowledge, only a single study to date has demonstrated that hospitalizations, use of bed days and general practitioner (GP) visits, were significantly higher for patients with multimorbidity, compared with those who had no chronic conditions [24].

The aim of this study was to explore associations between multimorbidity and healthcare utilization and the impact of socioeconomic status on utilization of hospitalizations and bed days. The structure and content of Danish healthcare registers provide a unique

opportunity to quantify how these variables interact [25, 26], which we explored in a large-scale, cross-sectional, regional, register-based population study.

Methods

Study population and data sources

The study population included all individuals aged 16 years and older who lived in the Capital Region of Denmark on January 1st, 2012. The included 1,397,173 individuals represented approximately one-third of the entire Danish adult population. Data on chronic conditions, use of healthcare services, and demographics, including gender, age, and educational attainment, were obtained from Danish national administrative and health registries: the Danish National Patient Registry [27], the Danish National Prescription Registry [28], the Danish National Health Service Registry [29], and The National Diabetes Registry [30]. All data obtained from registries were merged at the level of individuals using their unique social security numbers. However, national registries do not provide data on conditions diagnosed in the primary care sector. In addition to data on diagnoses of chronic conditions available from the secondary healthcare sector, diagnostic algorithms developed by the Research Center for Prevention and Health at Glostrup University Hospital were used to identify primary sector diagnoses of 16 chronic conditions of interest for the entire population (Table 1) [6]. Details about the diagnostic algorithms are provided elsewhere [7]. Multimorbidity was defined as two or more chronic conditions occurring simultaneously in the same person.

114 **Table 1.** Chronic conditions included in definition of multimorbidity (N = 16)

- 115 Allergies
- 116 Hypertension
- 117 High cholesterol
- 118 Diabetes (type 1 and type 2)
- 119 Heart disease
- 120 Stroke
- 121 Back pain
- 122 Joint disease
- 123 Osteoarthritis
- 124 Osteoporosis
- 125 Chronic obstructive pulmonary disease
- 126 Cancer
- 127 Dementia
- 128 Anxiety
- 129 Long-term use of antidepressants
- 130 Schizophrenia

131

132 Because data on direct costs could not be linked to the study data, we identified
 133 utilization of hospitalizations and bed days, which are among the most expensive services in
 134 healthcare, as proxies for direct costs. Socioeconomic status was defined as highest
 135 educational achievement and grouped into four categories according to the length of
 136 education: none (primary school) (≤ 10 years), short (11-14 years), medium (14-17 years),
 137 and long (≥ 17 years) [31]. Finally, we also recorded healthcare utilization for emergency

department visits, outpatient visits, GP visits, out-of-hour GP visits, yearly control visits in general practice, and specialist visits during 2012.

Statistical analysis

Descriptive statistics were calculated for gender, age, and educational attainment by the number of chronic conditions. Means for each type of healthcare utilization were calculated. Logistic regression was used to calculate odds ratios (ORs) for healthcare utilization for individuals with multimorbidity versus individuals with none and with one chronic condition. ORs were calculated in both a raw form and adjusted for gender, age, and educational attainment.

As cost data was not linked to the present dataset we chose to investigate the effect of socioeconomic status based on the proxies for general healthcare utilization: hospitalizations and bed days. Hospitalizations and bed days were adjusted for emergency visits, out-patient visits GP visits, out-of-hours GP visits, yearly controls in general practice, private specialist visits, number of conditions, age, gender, cohabitation status, education attainment, and employment status.

This was accomplished by applying zero-inflated models to both proxies. The models were used to calculate anticipated annual use of hospitalizations and bed days within educational attainment groups, separately for each number of chronic conditions.

The decision to use zero-inflated models was prompted by the fact that many individuals do not use bed days or hospitalization [32]. To counter extreme values for some covariates, squared effects of all numerical covariates were included as explanatory variables. For the partial model (1) below, a squared effect of GP visits was not included due to convergence problems. For the partial models of the form (2) below, quadrupled effects of ambulatory visits and out-of-hours GP visits were included. The zero-inflated model was applied as follows.

First, the probability p of having at least one bed day/one hospitalization was modeled with a logistic regression model

$$\text{logit}(p_i) = \sum_{j=1}^{26} \alpha_j x_{ji}, i = 1, \dots, n, \quad (1)$$

where p_i is the probability of one or more hospitalizations or bed days for an individual i and x_j is the j th covariate derived from the specified explanatory variables. Subsequently, numbers of bed days and hospitalizations y were similarly modeled with a log-linear model,

$$\log(y_i) = \sum_{j=1}^{29} \beta_j x_{ji} + \varepsilon_i, i = 1, \dots, n_1, \quad (2)$$

where $n_1 \leq n$ is the number of individuals with at least one bed day or hospitalization. In the models (2), two observations of bed days and three observations of hospitalizations were omitted as outliers for the analysis of bed days and hospitalizations, respectively.

For scenarios in which a specific number of conditions and a specific level of education were assumed, the linear predictors $\hat{\eta} = \sum \hat{\alpha}_j x_j$ and $\hat{\lambda} = \sum \hat{\beta}_j x_j$ were calculated, with all other covariates kept at the empirical mean for the group with the specified number of conditions and level of education. Estimated responses were found as $\hat{y} = \text{logit}^{-1}(\hat{\eta}) \exp(\hat{\lambda} + \hat{\sigma}^2/2)$, where σ^2 is the variance of $\hat{\lambda}$, thus combining models (1) and (2). All analyses were carried out using R software version 3.5.0 (34).

Ethics approval:

Approval to conduct the study was obtained from the Danish Data Protection Agency. No informed consent was required.

Results

184 Among 1,397,173 individuals included in the study population, approximately half
185 (720,885; 52%) were women, the majority (927,568; 66%) had none or short education, and
186 the prevalence of multimorbidity was 22% (301,757). Table 2 shows the distribution of
187 gender, age, and educational attainment by the number of chronic conditions. The prevalence
188 of both one and two or more chronic conditions was significantly higher for women than for
189 men. Overall, multimorbidity was highest among individuals 65-84 years old, followed by
190 those who were 45-64 years old. Educational attainment was inversely related to
191 multimorbidity.

192

193

Table 2. Distributions of the study population (N = 1,397,173) by gender, age, and educational attainment and number of chronic conditions, N (%)

		Number of chronic conditions		
		0	1	≥ 2
		N = 809,920	N = 285,496	N = 301,757
Gender				
	Male	418,105 (62)	127,591 (19)	130,592 (19)
	Female	391,815 (54)	157,905 (22)	171,165 (24)
Age in years				
	16-24	170,486 (86)	25,768 (13)	1,770 (1)
	25-44	385,540 (77)	89,216 (18)	23,557 (5)
	45-64	205,947 (48)	111,147 (26)	110,949 (26)
	65-84	45,529 (19)	53,334 (22)	140,001 (59)
	>84	2,418 (7)	6,031 (18)	25,480 (75)
Education				
	None (≤ 10 years)	185,347 (53)	69,779 (20)	93,702 (27)
	Short (11-14 years)	325,241 (56)	123,439 (22)	130,420 (23)
	Medium (15-16 years)	132,531 (59)	49,576 (22)	41,536 (19)
	Long (≥ 17 years)	107,578 (65)	34,035 (21)	22,174 (14)
	Missing	59,233 (72)	8,667 (11)	13,925 (17)

Mean rates of healthcare utilization among people with multimorbidity were much higher than among people with no chronic conditions, by a factor of 1.73 to 9.67, depending on the type of utilization (Table 3). When comparing people with multimorbidity and those with one chronic condition, rates of healthcare utilization were 1.44 to 4.00 times higher for

those with multimorbidity. In both comparisons, the largest between-group difference was for yearly control visits in general practice. Unadjusted and adjusted ORs for all types of healthcare utilization were significantly higher ($p < 0.001$) among people with multimorbidity, compared with people with zero or one chronic condition. Unadjusted ORs for healthcare utilization for people with multimorbidity were 1.65 to 11.76 times higher than for people with no chronic conditions and 1.31 to 3.15 times higher than for people with one chronic condition (Table 3). Adjusted ORs for healthcare utilization for people with multimorbidity were 1.86 to 6.70 times higher than for people with no chronic conditions and 1.44 to 2.94 times higher than for people with one chronic condition (Table 3).

Table 3. Mean and odds ratios (ORs) for healthcare utilization among individuals with 0, 1, and ≥ 2 chronic conditions

	Number of chronic conditions						
	0	1	≥ 2	≥ 2 vs. 0		≥ 2 vs. 1	
	Mean			Unadjusted OR	Adjusted OR*	Unadjusted OR	Adjusted OR*
Hospitalizations	0.12	0.22	0.51	3.84 (3.80-3.88)	2.78 (2.74-2.83)	2.26 (2.23-2.29)	1.77 (1.75-1.80)
Bed days	0.31	0.82	2.65	3.84 (3.80-3.88)	2.78 (2.74-2.85)	2.26 (2.23-2.29)	1.77 (1.75-1.80)
Emergency visits	0.15	0.18	0.26	1.65 (1.63-1.66)	1.86 (1.83-1.88)	1.39 (1.37-1.41)	1.49 (1.47-1.52)
Outpatient visits	0.91	1.88	3.86	4.83 (4.79-4.87)	3.79 (3.76-3.84)	2.25 (2.23-2.29)	2.03 (2.01-2.05)
General practice visits	4.34	7.39	12.94	8.37 (8.31-8.44)	6.70 (6.64-6.76)	3.15 (3.13-3.18)	2.94 (2.91-2.27)
Out-of-hours general practice visits	0.28	0.38	0.56	1.66 (1.64-1.67)	2.23 (2.19-2.26)	1.31 (1.29-1.32)	1.59 (1.57-1.61)
Yearly control visits in general practice	0.03	0.12	0.29	11.76 (11.58-11.96)	5.66 (5.56-5.77)	2.86 (2.81-2.89)	1.96 (1.93-1.98)
Specialist visits	0.69	1.39	2.09	3.66 (3.63-3.69)	2.36 (2.34-2.39)	1.77 (1.75-1.79)	1.44 (1.43-1.45)

*Adjusted for gender, age, and educational attainment

Among people with multimorbidity, utilization of hospitalizations increased approximately linearly with the number of chronic conditions. In Fig. 1A, hospitalization rates are indicated with a black line; the reference regression line in red has a slope equal to the mean number of hospitalizations across the number of chronic conditions. The similarity between observed rates and the regression line indicates that each chronic condition corresponded to an average of approximately 0.24 hospitalizations per year. A similar pattern was observed for bed days (Fig. 1B). For people with five or fewer chronic conditions, the utilization of bed days was approximately proportional to the number of conditions. However, between five and six or more chronic conditions, a steep increase in utilization of bed days was observed. Among individuals with six or more chronic conditions, utilization was higher than the mean number of bed days per condition multiplied by the number of conditions.

Fig 1.

1A. Relationship between numbers of chronic conditions and hospitalizations.

The black line indicates the number of hospitalizations by the number of chronic conditions. The red line indicates the mean number of hospitalizations multiplied by the number of chronic conditions.

1B. Relationship between numbers of chronic conditions and bed days.

The black line indicates the number of bed days by number of chronic conditions. The red line represents the mean number of bed days multiplied by the numbers of chronic conditions.

Fig 2 and Fig 3 depict utilization of hospitalization and bed days stratified by educational attainment. An educational gradient in hospitalization rates was observed across

one to six or more chronic conditions (Fig 2). Hospitalizations were more frequent in individuals with shorter education, compared with those with longer education. For bed day utilization (Fig 3), individuals with no education exhibited the highest estimated utilization rates, regardless of the number of chronic conditions.

Fig 2. Number of chronic conditions associated with the modeled rate of hospitalizations by educational attainment levels

Black line, no education; red line, short education; green line, medium education; blue line, long education.

Fig 3. Number of chronic conditions associated with the modeled rate of bed days by educational attainment

Black line, no education; red line, short education; green line, medium education; blue line, long education.

DISCUSSION

This study is the first large-scale, register-based study investigating associations between multimorbidity and utilization of healthcare services in the secondary and primary sectors in Denmark. In this study we were able to obtain a medical diagnosis for all diseased adults in the Capital Region aged 16 and above using algorithmic diagnoses for 16 conditions. This contrasts with studies using register-based diagnoses that only included patients who had had a hospital admission or were affiliated with an outpatient clinic. Although the 16 selected conditions do not represent the full spectrum of chronic disease, they include highly prevalent chronic conditions. A

similar study from Scotland [33] included 40 conditions and revealed an age-stratified pattern of prevalence of chronic conditions nearly identical to that of the data reported here [7, 33]. This suggests that the 16 included diagnoses in this study encompass all conditions and combinations of conditions that are predominant at the population level.

The prevalence of multimorbidity in our study was 22% [7]. Females had a 10% overrepresentation among individuals with multimorbidity (Table 2). However, in this study the overrepresentation of women among multimorbid individuals cannot be explained by the longevity of women alone: The frequency of multimorbidity within each one-year age group was consistently higher for women than men from 68 years to 100 years of age, the difference increasing with age. This finding is consistent with previous results across healthcare systems and geography [34-36]. Eighty percent of the study population and 45% of individuals with multimorbidity were younger than age 65 (Table 2). The study of multimorbidity is often confined to adults aged 65 and up [19, 37-39]. However, younger individuals with multimorbidity who survive will age into this group and can be expected to have higher lifelong healthcare utilization. In fact, we observed that, compared to age-similar individuals without chronic conditions, younger multimorbid individuals had a relatively higher rate of bed day utilization than did older individuals with multimorbidity. This finding highlights the importance of understanding chronic illness and healthcare utilization of younger multimorbid persons.

Utilization vs. number of conditions

The consistency of increases in healthcare utilization for individuals with multimorbidity, compared to those without chronic conditions, is remarkable (Table 3). The use of all services was higher to a statistically significant degree. Excluding hospitalizations and bed days, the increase in utilization rates for individuals with multimorbidity ranged from 73% (emergency department visits) to 324% (outpatient visits), except for yearly control visits in general practice, which were

867% higher. Disregarding the latter due to very low utilization rates, the average increase in healthcare utilization was 180% for individuals with multimorbidity, compared to those with no chronic conditions. The corresponding average increase in utilization rates for multimorbid individuals, compared with those with one chronic disease, was 65%. Unadjusted ORs for healthcare utilization for individuals with multimorbidity, compared with those with no chronic conditions, were all statistically significant and ranged from 1.65 (95% confidence interval [CI], 1.63 – 1.66) for emergency department visits and 11.76 (95% CI, 11.58 – 11.96) for yearly control visits. When adjusted for gender, age, and educational attainment, utilization differences were less pronounced, ranging from 1.86 (95% CI, 1.83 – 1.88) for emergency department visits to 6.70 (95% CI, 6.64 – 6.76) for GP visits. A similar but less pronounced pattern was seen when comparing multimorbid individuals to those with one chronic disease (Table 3). These results demonstrate that the impact of multimorbidity on healthcare utilization applies to a range of services and varies relatively little. Healthcare utilization rates were 2-4 times higher than those for people without chronic conditions. The direct relationship between the number of chronic conditions and healthcare utilization is well-documented in the literature [14, 23, 24, 40, 41], but, to the best of our knowledge, the impact of multimorbidity on a broad spectrum of healthcare utilization has not been documented previously.

The approximately linear relationship between frequency of hospitalizations and the number of chronic conditions depicted in Fig. 1A also applies to the relationship between bed days and the number of chronic conditions in Fig. 1B for five or fewer conditions. When the number of conditions increases to six or more, the utilization of bed days increases by a factor greater than the impact of a single additional condition. The overall utilization pattern is that each condition corresponds to 0.24 hospitalizations and 0.92 bed days, while the length of each hospitalization is longer for individuals with six or more conditions. The slope of the curve for six or more conditions

increases twofold to 1.84 (Fig 1B). The fact that the regression coefficient no longer explains the frequency of bed days per chronic condition leads us to define individuals with six or more conditions as high utilizers; they make up 0.87% of the population but account for 7.29% of bed days. An earlier Danish study found that 5% of the population with chronic conditions accounted for 45% of healthcare expenses [42]. This finding is in line with a large US study that showed that 5% of high utilizers accounted for up to 47% of healthcare costs [43]. A recent German study reported two subgroups of high utilizers: the oldest patients who suffered from severe multimorbidity and younger elderly patients with psychiatric or psychosomatic conditions [40]. Future research is required to examine characteristics and utilization of the high utilizers identified in this study.

Fig 1A shows a decline in utilization between nine and ten or more conditions. A possible explanation is that individuals with a very high disease burden have higher mortality, while those who survive have lower healthcare utilization than expected. Only 43 individuals had ten or more diagnosed conditions, and small sample size effects may also contribute to this finding.

Socioeconomic status

We found that the prevalence of multimorbidity decreased with increasing educational attainment, revealing a pronounced and statistically significant inverse socioeconomic gradient. This is consistent with previous findings [7].

To study the impact of multimorbidity on healthcare costs, we adjusted data for proxy costs for different educational attainment groups with varying profiles in terms of age, gender, other healthcare utilization and level of multimorbidity. When adjusted for these effects, as described in the methods statistical section, a clear inverse social gradient in hospitalization utilization appeared (Fig. 2). Adjusted hospitalization utilization decreased as the level of educational attainment increased, generally irrespective of the number of chronic conditions. However, adjusted bed-day

utilization revealed a slightly different pattern. An educational gradient no longer appeared; curves for short, medium, and high educational attainment tended to overlap. However, the group with no educational attainment stood out by virtue of higher healthcare utilization than the other. One important feature is that the relative difference between utilization rates for individuals with no education and those in other educational groups appeared close to constant across the number of chronic conditions.

In general, the number of hospitalizations decreased with increasing level of educational attainment. For multimorbid individuals, the length of each hospitalization was longer for individuals without any education than for those with at least some education. The reason for this finding is unknown. It may be the case that chronic disease tends to be more severe among individuals without education and that longer hospitalizations are caused by non-chronic conditions not included in our study. In addition, one could posit that different spectra of chronic conditions occur for persons with and without education, but investigation of this supposition will require further research. Other factors affecting the higher healthcare utilization rate observed in people with lower education attainment include lower health literacy levels linked to lower education attainment [44]. Disease burden has been shown to be associated with lower education levels in type 2 diabetes, and this may apply to other chronic conditions [45]. Furthermore, people with none or low educational attainment often has weak social networks and proper discharge to home might demand coordinated preparation that rely, in part, on support from individual's own social support structure [46]. However, we currently lack a well-founded explanation for differences in the two adjusted proxies for general utilization of healthcare services.

Strengths and limitations

The major strength of our study is that it is a large-scale register-based study, including comprehensive information about chronic conditions, healthcare utilization, and educational

375 attainment of the complete population of the Capital Region of Denmark aged 16 years and above.
 376 Generally, data from the Danish national registers provide complete information about healthcare
 377 system contacts, are of high quality and reliability, and are used extensively in research [25, 47]. As
 378 a general population-based study, our findings reflect the actual situation in a real-world setting.
 379 Based on register data, our study was free of recall bias and there was no loss of follow up.

380 Several limitations of our study deserve consideration. The necessary use of diagnostic
 381 algorithms to identify patients with chronic conditions in the primary healthcare sector is an
 382 approximation of actual diagnoses. Although the diagnostic algorithms have been shown to be
 383 highly accurate [6], they are not clinically determined by physicians. The study was based on cross-
 384 sectional data collected during a single year, and the number of patients with specific chronic
 385 conditions may be underestimated [48, 49]. In addition, we were not able to include healthcare
 386 services provided in the municipalities. Scarce register data exist for municipality services; existing
 387 data are not well defined and thus not useful for research purposes. Inclusion of the data on
 388 utilization of community healthcare services would likely have helped to generate a more complete
 389 picture of healthcare utilization related to multimorbidity. Finally, 6% of our population did not
 390 have information on educational attainment. This information appeared to be missing at random,
 391 except for individuals aged 91 years and above; Danish administrative registers only contain
 392 information on education for individuals born since 1921 [31]. However, this group contained few
 393 individuals, and we estimated the effect of these missing data to be minimal and possible changes in
 394 parameter estimates therefore very small.

395 Comparison to other studies on multimorbidity in populations should be performed with
 396 care. Varying definitions of multimorbidity (i.e., two or more chronic conditions), included
 397 conditions, and data collection methods render comparisons difficult. However, these challenges
 398 may be overcome for large studies [33].

399

400

Conclusion

401

Multimorbidity is associated with a significant increase in utilization of all healthcare

402

services in Denmark. A socioeconomic gradient was observed in utilization of hospitalizations, and

403

socioeconomic effects in utilization of bed days. A steep increase in the utilization of bed days in

404

patients with six or more chronic conditions suggests a subpopulation of high utilizers that should

405

be explored in further studies.

406

407 **Acknowledgements**

408 We thank Jennifer Green for skillful editing.

409

410 **References**

- 411 1. Valderas JM, Starfield B, Sibbald B, Salisbury C, Roland M. Defining comorbidity:
412 implications for understanding health and health services. *Ann Fam Med*. 2009;7(4):357-63. doi:
413 10.1370/afm.983.
- 414 2. Mercer S, Furler J, Moffat K, Fischbacher-Smith D, Sanci L. Multimorbidity: technical
415 series on safer primary care. Geneva: World Health Organization; 2016.
- 416 3. Bayliss EA, Bayliss MS, Ware JE, Steiner JF. Predicting declines in physical function in
417 persons with multiple chronic medical conditions: what we can learn from the medical problem list.
418 *Health Qual Life Outcomes*. 2004;2(1):47. doi: 10.1186/1477-7525-2-47.
- 419 4. Williams JS, Egede LE. The association between multimorbidity and quality of life, health
420 status and functional disability. *Am J Med Sci*. 2016;352(1):45-52. doi:
421 10.1016/j.amjms.2016.03.004.
- 422 5. Emerging Risk Factors Collaboration. Association of cardiometabolic multimorbidity with
423 mortality. *JAMA*. 2015;314(1):52-60. doi: 10.1001/jama.2015.7008.
- 424 6. Robinson KM, Lau CJ, Jeppesen M, Vind AB, Glümer C. Kroniske sygdomme-hvordan
425 opgøres kroniske sygdomme?: Glostrup: Forskningscenter for Forebyggelse og Sundhed [Research
426 Center for Prevention and Health]; 2011.
- 427 7. Schiøtz ML, Stockmarr A, Høst D, Glümer C, Frølich A. Social disparities in the prevalence
428 of multimorbidity: a register-based population study. *BMC Public Health*. 2017;17(1):422. doi:
429 10.1186/s12889-017-4314-8.

- 430 8. Diderichsen F, Evans T, Whitehead M. The social basis of disparities in health. In: Evans T,
431 Whitehead M, Diderichsen F, Bhulya A, Wirth M, editors. Challenging inequities in health: from
432 ethics to action. New York: Oxford University Press; 2001. p. 12-23.
- 433 9. Pefoyo AJ, Bronskill SE, Gruneir A, Calzavara A, Thavorn K, Petrosyan Y, et al. The
434 increasing burden and complexity of multimorbidity. BMC Public Health. 2015;15:415. doi:
435 10.1186/s12889-015-1733-2.
- 436 10. van Oostrom SH, Gijzen R, Stirbu I, Korevaar JC, Schellevis FG, Picavet HSJ, et al. Time
437 trends in prevalence of chronic diseases and multimorbidity not only due to aging: data from
438 general practices and health surveys. PLoS One. 2016;11(8):e0160264. doi:
439 10.1371/journal.pone.0160264.
- 440 11. Freid VM, Bernstein AB, Bush MA. Multiple chronic conditions among adults aged 45 and
441 over: trends over the past 10 years. NCHS Data Brief. 2012;(100):1-8.
- 442 12. König H-H, Leicht H, Bickel H, Fuchs A, Gensichen J, Maier W, et al. Effects of multiple
443 chronic conditions on health care costs: an analysis based on an advanced tree-based regression
444 model. BMC Health Serv Res. 2013;13:219. doi: 10.1186/1472-6963-13-219.
- 445 13. Skinner HG, Coffey R, Jones J, Heslin KC, Moy E. The effects of multiple chronic
446 conditions on hospitalization costs and utilization for ambulatory care sensitive conditions in the
447 United States: a nationally representative cross-sectional study. BMC Health Serv Res. 2016;16:77.
448 doi: 10.1186/s12913-016-1304-y.
- 449 14. Lehnert T, Heider D, Leicht H, Heinrich S, Corrieri S, Lupp M, et al. Review: health care
450 utilization and costs of elderly persons with multiple chronic conditions. Med Care Res Rev.
451 2011;68(4):387-420. doi: 10.1177/1077558711399580.

- 452 15. Schneider KM, O'Donnell BE, Dean D. Prevalence of multiple chronic conditions in the
453 United States' Medicare population. *Health Qual Life Outcomes*. 2009;7:82. doi: 10.1186/1477-
454 7525-7-82. PubMed PMID: PMC2748070.
- 455 16. Wolff JL, Starfield B, Anderson G. Prevalence, expenditures, and complications of multiple
456 chronic conditions in the elderly. *Arch Intern Med*. 2002;162(20):2269-76. doi:
457 10.1001/archinte.162.20.2269.
- 458 17. Yu W, Ravelo A, Wagner TH, Barnett PG. The relationships among age, chronic conditions,
459 and healthcare costs. *Am J Manag Care*. 2004;10(12):909-16.
- 460 18. Glynn LG, Valderas JM, Healy P, Burke E, Newell J, Gillespie P, et al. The prevalence of
461 multimorbidity in primary care and its effect on health care utilization and cost. *Fam Pract*.
462 2011;28(5):516-23. doi: 10.1093/fampra/cmr013.
- 463 19. Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization and costs
464 in an elderly community-dwelling population: a claims data based observational study. *BMC Health*
465 *Serv Res*. 2015;15:23. doi: 10.1186/s12913-015-0698-2.
- 466 20. van den Bussche H, Schön G, Kolonko T, Hansen H, Wegscheider K, Glaeske G, et al.
467 Patterns of ambulatory medical care utilization in elderly patients with special reference to chronic
468 diseases and multimorbidity; results from a claims data based observational study in Germany.
469 *BMC Geriatr*. 2011;11:101. doi: 10.1186/1471-2318-11-54.
- 470 21. van Oostrom SH, Picavet HSJ, de Bruin SR, Stirbu I, Korevaar JC, Schellevis FG, et al.
471 Multimorbidity of chronic diseases and health care utilization in general practice. *BMC Fam Pract*.
472 2014;15:61. doi: 10.1186/1471-2296-15-61. PubMed PMID: PMC4021063.
- 473 22. Wang HHX, Wang JJ, Lawson KD, Wong SYS, Wong MCS, Li FJ, et al. Relationships of
474 multimorbidity and income with hospital admissions in 3 health care systems. *Ann Fam Med*.
475 2015;13(2):164-7. doi: 10.1370/afm.1757. PubMed PMID: PMC4369606.

23. McPhail SM. Multimorbidity in chronic disease: impact on health care resources and costs. Risk Manage Healthc Policy. 2016;9:143-56. doi: 10.2147/RMHP.S97248. PubMed PMID: PMC4939994.
24. Palladino R, Tayu Lee J, Ashworth M, Triassi M, Millett C. Associations between multimorbidity, healthcare utilisation and health status: evidence from 16 European countries. Age Ageing. 2016;45(3):431-5. doi: 10.1093/ageing/afw044.
25. Erlangsen A, Fedyszyn I. Danish nationwide registers for public health and health-related research. Scan J Public Health. 2015;43(4):333-9. doi: 10.1177/1403494815575193. PubMed PMID: 25759376.
26. Thygesen LC, Ersbøll AK. Danish population-based registers for public health and health-related welfare research: introduction to the supplement. Scan J Public Health. 2011;39(7 Suppl):8-10. doi: 10.1177/1403494811409654.
27. Lynge E, Sandegaard JL, Rebolj M. The Danish national patient register. Scan J Public Health. 2011;39(7 Suppl):30-3. doi: 10.1177/1403494811401482. PubMed PMID: 21775347.
28. Wallach Kildemoes H, Toft Sørensen H, Hallas J. The Danish national prescription registry. Scan J Public Health. 2011;39(7 Suppl):38-41. doi: 10.1177/1403494810394717.
29. Olivarius NF, Hollnagel H, Krasnik A, Pedersen PA, Thorsen H. The Danish national health service register. a tool for primary health care research. Dan Med Bull. 1997;44(4):449-53. PubMed PMID: 9377908.
30. Carstensen B, Kristensen JK, Marcussen MM, Borch-Johnsen K. The national diabetes register. Scan J Public Health. 2011;39(7 Suppl):58-61. doi: 10.1177/1403494811404278. PubMed PMID: 21775353.
31. Jensen VM, Rasmussen AW. Danish education registers. Scan J Public Health. 2011;39(7 Suppl):91-4. doi: 10.1177/1403494810394715. PubMed PMID: 21775362.

- 500 32. Lachenbruch PA. Analysis of data with excess zeros. *Stat Methods Med Res.*
501 2002;11(4):297-302. doi: 10.1191/0962280202sm289ra. PubMed PMID: 12197297.
- 502 33. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of
503 multimorbidity and implications for health care, research, and medical education: a cross-sectional
504 study. *Lancet.* 2012;380(9836):37-43. doi: 10.1016/S0140-6736(12)60240-2.
- 505 34. Abad-Díez JM, Calderón-Larrañaga A, Poncel-Falcó A, Poblador-Plou B, Calderón-Meza
506 JM, Sicras-Mainar A, et al. Age and gender differences in the prevalence and patterns of
507 multimorbidity in the older population. *BMC Geriatr.* 2014;14:75. doi: 10.1186/1471-2318-14-75.
- 508 35. Rzewuska M, de Azevedo-Marques JM, Coxon D, Zanetti ML, Zanetti ACG, Franco LJ, et
509 al. Epidemiology of multimorbidity within the Brazilian adult general population: evidence from
510 the 2013 national health survey (PNS 2013). *PLoS One.* 2017;12(2):e0171813. doi:
511 10.1371/journal.pone.0171813.
- 512 36. Alimohammadian M, Majidi A, Yaseri M, Ahmadi B, Islami F, Derakhshan M, et al.
513 Multimorbidity as an important issue among women: results of a gender difference investigation in
514 a large population-based cross-sectional study in West Asia. *BMJ Open.* 2017;7(5). doi:
515 10.1136/bmjopen-2016-013548.
- 516 37. Caughey GE, Ramsay EN, Vitry AI, Gilbert AL, Luszcz MA, Ryan P, et al. Comorbid
517 chronic diseases, discordant impact on mortality in older people: a 14-year longitudinal population
518 study. *J Epidemiol Community Health.* 2010;64(12):1036-42. doi: 10.1136/jech.2009.088260.
- 519 38. Jindai K, Nielson CM, Vorderstrasse BA, Quiñones AR. Multimorbidity and functional
520 limitations among adults 65 or older, NHANES 2005–2012. *Prev Chronic Dis.* 2016;13:E151. doi:
521 10.5888/pcd13.160174. PubMed PMID: PMC5094859.

- 522 39. Hopman P, Heins MJ, Rijken M, Schellevis FG. Health care utilization of patients with
523 multiple chronic diseases in the Netherlands: differences and underlying factors. *Eur J Intern Med.*
524 2015;26(3):190-6. doi: 10.1016/j.ejim.2015.02.006.
- 525 40. van den Bussche H, Kaduszkiewicz H, Schäfer I, Koller D, Hansen H, Scherer M, et al.
526 Overutilization of ambulatory medical care in the elderly German population?: an empirical study
527 based on national insurance claims data and a review of foreign studies. *BMC Health Serv Res.*
528 2016;16:129. doi: 10.1186/s12913-016-1357-y. PubMed PMID: PMC4831189.
- 529 41. Westert GP, Satariano WA, Schellevis FG, Van Den Bos GAM. Patterns of comorbidity and
530 the use of health services in the Dutch population. *Eur J Public Health.* 2001;11(4):365-72. doi:
531 10.1093/eurpub/11.4.365.
- 532 42. Lau CJ, Lykke M, Andreasen AH, Bekker-Jeppesen M, Buhelt LP, Robinson KM, et al.
533 Sundhedsprofil 2013-kronisk sygdom. Glostrup: Forskningscenter for Forebyggelse og Sundhed,
534 Region Hovedstaden; 2015.
- 535 43. Zulman DM, Pal Chee C, Wagner TH, Yoon J, Cohen DM, Holmes TH, et al.
536 Multimorbidity and healthcare utilisation among high-cost patients in the US Veterans Affairs
537 health care system. *BMJ Open.* 2015;5(4). doi: 10.1136/bmjopen-2015-007771.
- 538 44. van der Heide I, Wang J, Droomers M, Spreeuwenberg P, Rademakers J, Uiters E. The
539 relationship between health, education, and health literacy: results from the Dutch adult literacy and
540 life skills survey. *J Health Commun.* 2013;18(Suppl 1):172-84. doi:
541 10.1080/10810730.2013.825668.
- 542 45. Agardh EE, Sidorchuk A, Hallqvist J, Ljung R, Peterson S, Moradi T, et al. Burden of type
543 2 diabetes attributed to lower educational levels in Sweden. *Popul Health Metr.* 2011;9(1):60. doi:
544 10.1186/1478-7954-9-60.

- 545 46. Aartsen M, Veenstra M, Hansen T. Social pathways to health: On the mediating role of the
546 social network in the relation between socio-economic position and health. *SSM - population*
547 *health*. 2017;3:419-26. doi: 10.1016/j.ssmph.2017.05.006. PubMed PMID: 29349235.
- 548 47. Thygesen LC, Daasnes C, Thaulow I, Brønnum-Hansen H. Introduction to Danish
549 (nationwide) registers on health and social issues: structure, access, legislation, and archiving. *Scan*
550 *J Public Health*. 2011;39(7 Suppl):12-6. doi: 10.1177/1403494811399956.
- 551 48. van Oostrom SH, Picavet HSJ, van Gelder BM, Lemmens LC, Hoeymans N, van Dijk CE,
552 et al. Multimorbidity and comorbidity in the Dutch population: data from general practices. *BMC*
553 *Public Health*. 2012;12:715. doi: 10.1186/1471-2458-12-715.
- 554 49. Schram MT, Frijters D, van de Lisdonk EH, Ploemacher J, de Craen AJM, de Waal MWM,
555 et al. Setting and registry characteristics affect the prevalence and nature of multimorbidity in the
556 elderly. *J Clin Epidemiol*. 2008;61(11):1104-12. doi: 10.1016/j.jclinepi.2007.11.021.

557

558

559

Fig1

Fig2

Fig3