Preliminary evidence that maternal immune activation specifically increases diagonal domain volume in the rat brain during early postnatal development

Tobias C. Wood^{1§}, Michelle E. Edye^{2§}, Michael K. Harte², Joanna C. Neill²,
Eric P. Prinssen³ and Anthony C. Vernon^{*4, 5}

¹King's College London, Institute of Psychiatry, Psychology and Neuroscience,
Department of Neuroimaging, Centre for Neuroimaging Sciences, De
Crespigny Park, London, SE5 8AF, United Kingdom

²Division of Pharmacy and Optometry, School of Health Sciences, Faculty of
Medicine, Biology and Health, University of Manchester, Manchester, M13
9PT, UK

³Roche Innovation Centre Basel, Grenzacherstrasse 124, Basel, CH 4070,
Switzerland

⁴King's College London, Institute of Psychiatry, Psychology and Neuroscience,
Department of Basic and Clinical Neuroscience, Maurice Wohl Clinical
Neuroscience Institute, 5 Cutcombe Road, London, SE5 9RT, United Kingdom

⁵King's College London, MRC Centre for Neurodevelopmental Disorders, New
Hunt's House, Guy's Hospital Campus, London SE1 1UL, United Kingdom.

[§]These authors contributed equally to this publication

***Corresponding author:** Anthony C. Vernon (anthony.vernon@kcl.ac.uk);
King's College London, Institute of Psychiatry, Psychology and Neuroscience,
Department of Basic and Clinical Neuroscience, Maurice Wohl Clinical
Neuroscience Institute, 5 Cutcombe Road, London, SE5 9RT, United Kingdom

1 **Running title:** Diagonal domain volume increases following gestational poly

2 (I:C) exposure

3

4 **Keywords:** Maternal immune activation; poly (I:C); magnetic resonance

5 imaging; diagonal domain; rat.

6

7 **Word count**

8 Abstract: 247

9 Document: 3,692

10 Figures: 2

11 Tables: 1

12

1 **Abstract**

2 Maternal immune activation (MIA) is consistently associated with elevated risk
3 for multiple psychiatric disorders in the affected offspring. Related to this, an
4 important goal of our work is to explore the impact of MIA effects across the
5 lifespan. In this context, we recently reported the effects of poly (I:C)-induced
6 MIA at gestational day (GD)15, immediately prior to birth, at GD21 and again
7 at post-natal day (PD)21, providing a systematic assessment of plasma IL-6,
8 body temperature and weight alterations in pregnant rats following poly (I:C)
9 exposure and preliminary evidence for gross morphological changes and
10 microglial neuropathology in both male and female offspring at GD21 and
11 PD21. Here, we sought to complement and extend these data by
12 characterising in more detail the meso-scale impact of gestational poly (I:C)
13 exposure at GD15 on the neuroanatomy of the juvenile (PD21) rat brain using
14 high-resolution, *ex vivo* anatomical magnetic resonance imaging (MRI) in
15 combination with atlas-based segmentation. Our preliminary data suggest
16 subtle neuroanatomical effects of gestational poly (I:C) exposure (n=10)
17 relative to saline controls (n=10) at this time-point. Specifically, we report here
18 preliminary evidence for a significant increase in the relative volume of the
19 diagonal domain in poly (I:C) offspring ($p<0.01$; $q<0.1$), particularly in female
20 offspring. This occurred in the absence of any microstructural alterations as
21 detectable using diffusion tensor imaging (DTI). Longitudinal *in vivo* studies,
22 informed by the effect sizes from this dataset are now required to establish
23 both the functional relevance and cellular mechanisms of the apparent DD
24 volume increase.

1. Introduction

Epidemiological studies report associations between exposure to maternal immune activation (MIA) and increased risk for psychiatric illnesses, including schizophrenia (SZ) and autism spectrum disorder (ASD) in the affected offspring [1, 2]. Based on these findings, reverse-translational animal models of MIA were developed, particularly (but not exclusively) using the viral mimetic polyriboinosinic-polyribocytidylic acid (poly I:C) [3, 4]. Administration of poly (I:C) on specific gestational days (GD; range 9–17) induces a discrete inflammatory response in the dam and elicits behavioural and neurochemical abnormalities in the offspring that are of relevance for the aforementioned psychiatric illnesses, providing both causation for the epidemiological data and important mechanistic insights [3, 5-7]. On the other hand, differences with regard to the strain of rodent, dose and route of administration of poly (I:C) used to induce MIA are present in the literature [5, 8-11]. These methodological differences likely explain the variance in published data with regard to behavioural and *post-mortem* brain phenotypes in rodent MIA models [8, 9, 11, 12]. Reporting guidelines and painstaking methodological work to establish the sources of variation in rodent MIA models, such as the caging system used to house the animals, are critical steps forward to address this issue [8, 10]. There is also a clear unmet need for an early outcome measure with which MIA-exposed offspring that will develop a robust behavioural phenotype may be identified and further enhance reproducibility across laboratories [12].

1 To address this gap, small animal magnetic resonance imaging (MRI) is one
 2 experimental approach, which offers several advantages. First,
 3 neuroanatomical phenotypes defined using high resolution MRI in
 4 combination with advanced image processing techniques are quite robust [13,
 5 14]. Second, MRI provides whole brain coverage, eliminating the need for *a*
 6 *priori* hypotheses concerning implicated brain regions [15]. Third, MR imaging
 7 operates at mesoscopic resolution, which is technically translatable to human
 8 studies [16]. This is relevant in light of recent work examining the influence of
 9 maternal cytokine levels on brain structure and function in human offspring, in
 10 relation to the development of psychopathology [17, 18]. Fourth, non-invasive
 11 MR imaging in rodent models may be directly linked with behavioural assays
 12 and invasive *post-mortem* follow-up, to establish the neural correlates of
 13 behaviour and the underlying mechanism(s) [19]. The utility of this approach is
 14 exemplified by a number of landmark studies demonstrating that induction of
 15 MIA in rodents using a range of protocols is associated with consistent sex-
 16 specific deviations from the normative developmental trajectory of brain
 17 structure, function and neurochemistry [20-26]. As yet however, a
 18 reproducible non-invasive MRI biomarker measured in early life that is
 19 predictive of later dysfunction in MIA-exposed adult offspring has yet to
 20 emerge.

21

22 Related to this, an important goal of our work is to explore the developmental
 23 trajectory of MIA effects across the lifespan [27]. We have recently begun this
 24 process by exploring the effects of MIA prior to birth, at gestational day

(GD)21 and just prior to weaning, at post-natal day PD21 [27]. This initial study (see also commentary by Roderick and Kentner, 2019 [28]) provided a robust and systematic assessment of plasma IL-6, body temperature and weight alterations in pregnant rats following poly (I:C) exposure and preliminary evidence for gross morphological changes (e.g. brain to body weight ratio, brain weight) and neuropathology (Iba1+ microglia number and morphology) in male and female offspring at GD21 and PD21. The relevance of these changes is currently being confirmed by behavioural assessment [27]. Emerging evidence however, shows behavioural changes of relevance to psychiatric disorders in GD15 poly (I:C) exposed offspring, including an increase in ultra-sonic vocalisation in early life (PD6) and a deficit in sustained attention in adulthood (PD125) [29]. Here, we sought to complement and extend these initial results by characterising the impact of MIA on brain volume and microstructure using high-resolution, *ex vivo* anatomical magnetic resonance imaging (MRI) specifically at PD21. The rationale for this approach is two-fold. First it will provide a detailed, brain-wide assessment of early neuroanatomical differences between control and MIA-exposed offspring, using the protocol reported in [27]. Second, it will provide preliminary data (i.e. effect sizes) to inform the design of future longitudinal *in vivo* MRI studies, which are necessary to establish if early MRI-detectable brain changes have any functional relevance.

2. Methods

2.1 Animals

Animals used in the present study were taken from a large cohort of control and poly (I:C)-exposed Wistar rats generated at the University of Manchester. Full details of rat strain, sex and housing conditions are reported in detail elsewhere [27]. All procedures in this study were carried out in accordance with the UK Home Office, Animals (Scientific Procedures) Act 1986 and EU Directive 2010/63/EU. The University of Manchester Animal Welfare and Ethical Review Body (AWERB) approved all experimental protocols used in this study.

2.2 Maternal immune activation (MIA) and allocation of offspring for ex vivo MRI phenotyping

Induction of MIA was performed at the University of Manchester as described in detail elsewhere [27]. In brief, female Wistar rats were mated at 3 months of age and GD1 confirmed by the appearance of a vaginal plug. Several studies provide evidence that poly (I:C) treatment at GD15 in rats induces a maternal inflammatory response with development of relevant behavioural phenotype(s) in the affected offspring [24, 26, 30, 31]. Pregnant Wistar rats (293.0–428.7 g) therefore received a single intraperitoneal (i.p.) injection of either poly (I:C) (n=8; P9582, potassium salt, Sigma-Aldrich; Gillingham, Dorset, United Kingdom) at a dose of 10 mg/kg or 0.9% non-pyrogenic sterile saline as a control (n=8) on GD15. On PD1, pups were sex typed based on anogenital distance and then culled to litters of n=4 males and n=4 females. At PD21 all pups were sacrificed and n=1 male and n=1 female offspring

1 selected from 5 (out of the available 8) control and poly (I:C) litters, giving a
 2 total of n=10 (5 male, 5 female) offspring per experimental arm (control vs.
 3 poly (I:C)). In the absence of data concerning effects of MIA on brain volume
 4 in Wistar rats at PD21 and how this may vary by sex, a formal estimation of
 5 sample size by power calculation was not possible. Rather, this study was
 6 designed pragmatically to generate this data using the minimum number of
 7 animals, in line with UK and EU guidelines (see www.nc3rs.org.uk) and taking
 8 into account available MRI scanner time and financial resources. We therefore
 9 selected a group size of n=10, comprised of 5 males and 5 females, such that
 10 both experimental groups were matched for sex. This group size (n=10) is
 11 comparable with prior *in vivo* neuroimaging studies in rodent MIA models [20,
 12 24-26, 32]. As such, the results presented herein are preliminary given the
 13 limited sample size and should be interpreted accordingly (*see also*
 14 *discussion*).

15 16 2.3 Enzyme-linked immunosorbent assay

17
 18 To confirm successful MIA in the dams, IL-6 concentrations in maternal blood
 19 plasma (3 hours post-injection) were determined using a rat-specific ELISA
 20 DuoSet (R&D Systems, Abingdon, UK) as reported elsewhere [27]. In brief,
 21 absorbances were measured using a plate reader (MRX, Dynatech, UK) at
 22 room temperature and results were calculated from the standard curve using
 23 Prism software (v6.0, GraphPad, La Jolla, CA, USA). We present here only
 24 the IL-6 values (pg/ml) for the specific dams from which offspring were
 25 selected for inclusion in the MRI study.

26

2.4 Tissue preparation for ex vivo MRI

At PD21, offspring were culled by cardiac perfusion (0.9% saline followed by 4% paraformaldehyde) under terminal anaesthesia (sodium pentobarbital, 60 mg/kg i.p) and the brain tissue prepared for *ex vivo* MRI as described elsewhere [16]. In brief, fixed brain tissues were kept intact in the cranium and post-fixed for 24 hours in 4% PFA. Samples were then placed in 0.01M phosphate buffer containing 0.05% (w/v) sodium azide to allow tissue re-hydration prior to MR imaging. Samples were then shipped to KCL and stored at 4°C in this solution for 4 weeks prior to MR imaging.

2.5 MR image acquisition

A 7T horizontal small bore magnet (Agilent Technologies Inc. Santa Clara, USA) and a quadrature volume radiofrequency coil (39 mm internal diameter, Rapid Biomedical GmbH, GER) were used for all MRI acquisitions. Fixed brain samples were placed securely one at a time in a custom-made MR-compatible holder and immersed in proton-free susceptibility matching fluid (FluorinertTM FC-70; Sigma-Aldrich, UK). Samples were scanned in a random order, with the KCL operator (ACV) blinded to treatment group (CON or POL) by numerical coding of samples. Scanning was interspersed with phantoms to ensure consistent operation of the scanner. Two sets of MR images were acquired: a 3D Fast-Spin Echo (FSE) for structural analysis and a DTI protocol for microstructural analysis. The T2-weighted 3D FSE image had the following parameters: TE/TR=60/2000, echo train length = 8, matrix size =

1 192x128x192 and field of view (FOV) = 28.8x19.2x28.8 mm, yielding isotropic
 2 voxels of $150 \mu\text{m}^3$. Total scan-time was 1hr 44min. The DTI scans were
 3 acquired using a four shot Echo Planar Imaging (EPI) sequence with
 4 TE/TR=35/4000ms, matrix size = 128x96 and FOV = 25.6x19.2mm, with an
 5 in-plane resolution of $200 \mu\text{m}$ across 50, 0.5 mm-thick slices. A total of 30
 6 non-collinear diffusion directions were acquired, with four $b=0$ images with a
 7 target b value of 2000 s/mm. Total scan time was 1hr 22min. Reversed
 8 phase-encoded direction DTI images were additionally acquired to correct for
 9 eddy current distortions (8min). Total scan duration for the entire protocol was
 10 3hr 14min per brain.

11

12 *2.6 MR image processing*

13

14 After visual inspection of all MR images and elimination of those scans with
 15 artefacts, the final n values per group for statistical comparisons were:
 16 volume; CON, n=9 (4 male, 5 female) vs. POL, n=10 (5 male, 5 female); DTI:
 17 CON, n=10 (4 male, 4 female) vs. POL, n=10 (5 male, 5 female). The MR
 18 Images were converted to NIFTI format and processed using a combination of
 19 FSL [33], ANTs [34] and in-house C++ software utilizing the ITK library,
 20 available from <https://github.com/spinacist/QUIT>. The processing pipeline
 21 consisted of several steps, as described elsewhere [35]. The following
 22 operations were carried out in the native space of the acquired MR images.
 23 First, a Tukey filter was applied to the FSE images in k-space to remove high
 24 frequency noise and they were corrected for intensity inhomogeneity using the
 25 N4 algorithm [36]. Second, FSL *topup* and *eddy* were used to remove
 26 distortion and eddy current artefacts in the raw diffusion data [37, 38] using

1 acquired DTI data with a reversed phase-encode direction (*see section 2.5*).
2 The DTI parameter maps were then calculated using FSL *dtifit* and consisted
3 of Fractional Anisotropy (FA) and Mean Diffusivity (MD) [35]. Third, a template
4 image was constructed from the 3D FSE images of all subjects in the study
5 using the *antsMultivariateTemplateConstruction2.sh* script with cross-
6 correlation metric and SyN transform (n=19) [31]. This template was then
7 registered to an atlas image of the PD18 Wistar rat brain, again using a cross-
8 correlation metric and SyN transform [39]. Fourth, all subjects FSE images
9 were non-linearly registered to the study template using the
10 *antsRegistrationSyN.sh* script. Logarithmic Jacobian determinants were
11 calculated from the inverse warp fields in standard space to estimate apparent
12 volume change, and smoothed with a Gaussian filter at a full-width half-
13 maximum (FWHM) of 200 μm [40]. Fifth, each subject's DTI image was
14 registered to the same subject's FSE image using a SyN transform to account
15 for residual distortions and a Mutual Information metric to account for the
16 different contrast. This transform was then concatenated with those from the
17 FSE images to the templates to align them to the study template [35]. The DTI
18 images in the template space were also smoothed with a Gaussian filter with
19 full-width half-maximum of 200 μm . Sixth, a brain parenchyma mask was
20 created from the atlas labels by excluding Cerebrospinal Fluid (CSF) regions.
21 The inverse transforms from the atlas to the study template and from the
22 study template to each subject were applied to calculate the brain and atlas
23 based region of interest (ROI) volumes for each subject [35].

24

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

2.7 Statistical analysis

A subset of maternal plasma levels of IL-6, consisting of those values only from dams whose offspring were selected for MR imaging (n=5 CON dams vs. n=5 POL dams) were compared using unpaired t-test (2-tailed), with $\alpha=0.05$, using Prism software (v6.0; GraphPad Inc. La Jolla, CA, USA). For the MR image analysis, group-level differences in volume, FA (unit less) and MD ($\text{mm}^2 \text{s}^{-1}$) were assessed using atlas-based segmentation (ABS) [20], taking advantage of a publicly available high resolution MRI atlas of the Wistar rat brain at PD18 that is parcellated into 26 regions of interest (ROI) [39]. We analysed relative volumes since total brain volume accounts for the majority of inter-animal variation in the volume of individual brain structures [13, 41]. Volume data for each atlas ROI was therefore expressed as a percentage of total brain volume [20]. FA and MD values were analysed without this correction. After image registration we successfully extracted values for volume, FA and MD for 23/26 ROIs (missing ROIs: Pineal gland, pituitary gland, optic pathway). Total brain volume was calculated from the summation of each individual atlas ROI volumes [20]. Group-level differences between CON and POL-exposed animals in relative volume, FA and MD were assessed across these 23 ROI using a 2 x 2 ANOVA, with MIA (CON or POL) and sex (male or female) as between and within group factor respectively, using Prism software (v6.0; GraphPad Inc. La Jolla, CA, USA) with $\alpha=0.05$. The resulting *p*-values from the F-test contrasts (main effects of prenatal treatment, sex and prenatal treatment x sex interaction) were then corrected

for multiple comparisons (to account for Type I errors across the 23 ROI) using the False Discovery Rate (FDR) procedure, with the threshold set at 10% ($q < 0.1$). This threshold was chosen based on simulation studies showing that, at a 10% FDR, simulated volume losses/gains of 5–20% can be recovered with a true false positive rate of $< 0.2\%$ [42]. Therefore in addition to p -values, we report q -values, which are FDR-adjusted p -values. Effect sizes were calculated using either partial η^2 (ANOVA) or Cohen's d (t-test) as appropriate. Due to the small sample size, particularly for sex, we do not present any voxel-wise, whole brain analysis.

10

11 **3. Results**

12

13 *3.1 Maternal circulating IL-6 levels*

14

15 A statistically significant increase in the circulating levels of IL-6 in maternal plasma samples could be observed 3 hours post-injection in POL-injected dams as compared to controls ($t = 3.62$, $df = 14$; $p = 0.007$; Cohen's $d = 2.3$; **Figure 1**). There was however, a notable degree of variability in the plasma IL-6 levels within the POL-injected dams (range 480 to 2890 pg/ml).

20

21 *3.2 Atlas based segmentation analysis of volumetric and DTI datasets*

22

23 *3.2.1 Juvenile rat brain volume*

24

25 Whole brain volumes were significantly affected as a function of sex ($F(1,15) = 5.89$ $p = 0.03$). There were no significant effects of either MIA, or MIA x sex interaction (**Table 1**). Main effects of sex were also found for the relative volumes of 3/23 (13%) atlas ROIs, including the bed nucleus of the stria terminalis (BNST), the cerebellum and the ventricular system, although these

1 did not survive correction for multiple comparisons (**Table 1**). Nonetheless,
 2 these trend-level results are consistent with known sexually dimorphic regions
 3 in the rodent brain [43, 44]. A significant main effect of MIA was found for the
 4 relative volume of the diagonal domain (DD), with subtly increased volume in
 5 POL as compared to CON offspring (+4.4%; $p < 0.05$, $q < 0.1$; partial $\eta^2 = 0.44$;
 6 **Table 1**; **Figure 2**). The relative volumes of the olfactory structures and
 7 preoptic area were also affected by MIA exposure, albeit only at trend level
 8 ($p < 0.05$, $q > 0.1$; **Table 1**). Interactions between sex and MIA were found, again
 9 only at trend level, for the relative volumes of a further 3/23 (13%) of the atlas
 10 ROIs, including the DD, nucleus accumbens and anterior commissure
 11 ($p < 0.05$, $q > 0.1$; **Table 1**).

12 13 3.2.1 Juvenile rat brain microstructure (FA, MD)

14
15 There were no statistically significant effects of sex, MIA or MIA x sex
 16 interaction on either FA or MD, across any of the 23 ROIs in the Wistar PD18
 17 MRI atlas (*all $p > 0.05$; $q > 0.1$; data not shown*).

18 19 4. Discussion

20
21 The main findings of the current study are as follows. Exposure of pregnant
 22 Wistar rat dams to poly (I:C) (POL) at GD15 has no effect on the total brain
 23 volume (in mm^3) of juvenile offspring at PD21 relative to controls (CON).
 24 Irrespective of gestation treatment brain volume did vary as a function of sex,
 25 which is consistent with published data [43, 44]. When comparing the relative
 26 volumes of 23 brain regions between CON and POL offspring at PD21, we
 27 found only subtle neuroanatomical differences in the juvenile Wistar rat brain.

Specifically, we provide preliminary evidence for a significant increase in the relative volume of the diagonal domain (DD) in POL offspring with a robust effect size (partial $\eta^2 = 0.44$). This effect appeared to be driven primarily by female POL offspring, although this result did not survive FDR correction (MIA x sex interaction; $p < 0.05$; $q > 0.1$). Additional volumetric changes, albeit at trend-level only, were found for the relative volumes of the olfactory structures and pre-optic area. Finally, we found no significant differences, even at trend-level, for microstructural alterations as indexed by either FA or MD. Taken together, these data provide new information on the impact of MIA on the juvenile Wistar rat brain, complementing and extending our recently published work using this MIA protocol [27].

Before interpreting these findings, limitations of the current study should be noted. First, behavioural phenotyping of offspring generated using the MIA protocol used herein and in our prior study [27] is currently ongoing; as such the functional relevance of this MIA protocol still remains to be confirmed. Emerging evidence however, shows behavioural changes of relevance to psychiatric disorders in GD15 poly (I:C) exposed offspring, including an increase in ultra-sonic vocalisation in early life (PD6) and a deficit in sustained attention in adulthood (PD125) [29]. Nonetheless, we interpret our current results only within the narrow framework of further defining the impact of a maternal systemic poly (I:C) challenge of 10 mg/kg at GD15. Second, as already fully acknowledged (*see methods, section 2.2*) our sample size is small, precluding the full use of voxel-wise MR image analysis tools [20]. This

1 could account for the limited number of significant findings surviving multiple
2 comparisons correction, particularly with regard to MIA x sex interactions.
3 Equally, we cannot exclude the possibility that our negative results,
4 particularly where the DTI data are concerned, are not simply a reflection of
5 the small sample size. As already stated, the group sizes were chosen
6 pragmatically based on the absence of prior data regarding effect sizes,
7 ethical use of the minimum number of animals, access to the MRI scanner
8 and financial restraints. Our overall group size is also comparable to
9 previously published MRI studies in MIA models, but we acknowledge this is
10 likely underpowered to detect sex-specific effects [20, 24]. Taking these
11 factors into account we refer to all findings as preliminary, pending replication
12 in a larger cohort, which the data generated in the current study allows us to
13 accurately calculate. For example, using the effect size for the main effect of
14 MIA on relative DD volume (partial $\eta^2 = 0.44$), a power calculation (F-test,
15 repeated measures ANOVA, between factors) suggests that for a longitudinal
16 imaging study with 2 groups (CON and POL) and 3 time-points, at least n=16
17 (8 male, 8 female) offspring should be included in each group to achieve
18 $\alpha=0.05$ with 95% power (G*Power v3.1.9.2). Third, we chose to collect *ex vivo*
19 MR images, as opposed to *in vivo* MR images. The enhanced image quality
20 available with *ex vivo* data increases the statistical power to detect subtle
21 volume changes when performing cross-sectional comparisons of two groups
22 [13, 41]. Therefore, if the study objective is detection of a neuroanatomical
23 phenotype, without attention to its time course, then *ex vivo* imaging is
24 preferable, hence our decision to opt for this approach [13]. Balanced against

1 this argument is the fact that the sample preparation (perfusion and tissue
2 fixation) for *ex vivo* imaging may cause morphological disruption to the
3 tissues, which could affect interpretation of the data. This is particularly true
4 for the ventricular system, which may collapse post-perfusion, such that
5 group-level differences in ventricular volume *in vivo* are not preserved *ex vivo*
6 [45]. This is relevant as *in vivo* studies in MIA rat models do show differences
7 in ventricular volume [20, 24]. Total brain volume and that of most grey matter
8 structures also shrinks post-perfusion [16, 41, 46]. Prior work however,
9 including our own, suggests that major group-level differences in grey matter
10 volumes are preserved despite this shrinkage from *in vivo* to *ex vivo* and can
11 be confirmed *post-mortem* [16, 41]. Taken together, the choice of *ex vivo* MR
12 imaging is consistent with the aims of the current study, but the case for
13 longitudinal *in vivo* studies is also reinforced.

14

15 Accepting these limitations, a cautious interpretation of these preliminary data
16 suggests some interesting observations. We found no significant effects of
17 maternal POL exposure on the volume (or microstructure) of the rat
18 hippocampus, prefrontal cortex, corpus striatum or lateral ventricles at PD21.
19 This may suggest that MRI-detectable changes in these regions only emerge
20 with increasing post-natal age and are not present in the juvenile brain [20,
21 24]. Our preliminary findings highlighting the DD as a structure specifically
22 affected in the juvenile rat brain following MIA are potentially intriguing, based
23 on other lines of evidence. Efferent projections from the DD innervate several
24 brain regions, but the majority project to the hippocampus via the medial

1 septum, where they contribute to the modulation of hippocampal theta (θ)
 2 oscillations that are important for attention, spatial and working memory and
 3 sensory information processing [47-49]. It is noteworthy then that decreased θ
 4 rhythms are reported in the adult rat hippocampus following exposure to MIA
 5 at GD15, which are related to memory and sensory processing impairments in
 6 these animals [31, 50, 51]. Furthermore, increased number and activity of
 7 cholinergic neurons is reported in the basal forebrain (which includes the DD),
 8 in the offspring of poly (I:C) exposed dams in early post-natal life (PD1) [52].
 9 Similar findings of increased cholinergic neuron density and activity have also
 10 been reported in *post-mortem* human basal forebrain tissue from individuals
 11 aged <13 years old with a diagnosis of ASD [53]. Taken together, there is
 12 circumstantial evidence to suggest that our preliminary MRI finding of elevated
 13 DD volume could have functional and clinical relevance, which can be tested
 14 in future studies.

15

16 **5. Conclusion**

17

18 The findings of the current study provide new insights into the impact of
 19 gestational poly (I:C) exposure on the juvenile (PD21) Wistar rat brain,
 20 complementing and extending our work using this model [27]. The data, albeit
 21 preliminary due to the small sample size, suggests that the DD is the only
 22 structure significantly affected at this time-point, with increased volume in
 23 offspring exposed to MIA. Our data also provide publicly available data on
 24 effect sizes to aid the design of longitudinal *in vivo* MRI studies in this model.
 25 Such studies are now clearly required to confirm the functional relevance of

1 the observed DD volume changes with regard to adult behavioural dysfunction
2 and to establish the underlying cellular and molecular mechanisms.

3

4 **Author contribution statement:** ACV, JCN, and EPP conceived and
5 designed the study with input from MEE and MKH. All animal work, including
6 MIA, was carried out at the University of Manchester by MEE, supervised by
7 MKH, JCN and EPP. TCW optimised the MR pulse sequences used in this
8 study for high-resolution ex vivo MR image acquisition, which was carried out
9 at King's College London (KCL) by ACV. TCW and ACV carried out the
10 computational and statistical analyses, respectively, of the MR images and
11 atlas-based segmentation data at KCL. ACV drafted and revised the MS with
12 input and the approval of all authors. ACV, JCN and EPP provided financial
13 support for the study.

14

15 **Competing interests statement:** ACV acknowledges financial support for
16 this work from F. Hoffman La Roche Ltd and the Medical Research Council
17 (New Investigator Research Grant [NIRG], MR/N025377/1). The work [at
18 King's College, London] was also supported by the Medical Research Council
19 (MRC) Centre grant (MR/N026063/1). JCN acknowledges financial support for
20 this work at the University of Manchester from F. Hoffman La Roche Ltd, b-
21 neuro and the University of Manchester MRC Confidence in Concept Scheme.
22 The funders had no role in the decision to publish this work. MEE, TCW and
23 MKH declare no conflicts of interest. EPP is a full-time employee of F.
24 Hoffman La Roche Ltd.

1

2 **Data availability statement:** Data collected from this study including the ex
3 vivo MR images may be made freely available upon reasonable requests to
4 the corresponding author.

5

6 **Acknowledgements:** The authors would like to thank the British Heart
7 Foundation and Dr Po-Wah So (KCL) for supporting and managing the 7T
8 MRI scanner at the King's College Preclinical Imaging Unit (KCLPIU).

9

10

11

References

- [1] A.S. Brown, P.H. Patterson, Maternal infection and schizophrenia: implications for prevention, *Schizophr Bull*, 37 (2011) 284-290.
- [2] A.S. Brown, S. Vinogradov, W.S. Kremen, J.H. Poole, R.F. Deicken, J.D. Penner, I.W. McKeague, A. Kochetkova, D. Kern, C.A. Schaefer, Prenatal exposure to maternal infection and executive dysfunction in adult schizophrenia, *Am J Psychiatry*, 166 (2009) 683-690.
- [3] U. Meyer, Prenatal Poly(I:C) Exposure and Other Developmental Immune Activation Models in Rodent Systems, *Biol Psychiat*, 75 (2014) 307-315.
- [4] I. Knuesel, L. Chicha, M. Britschgi, S.A. Schobel, M. Bodmer, J.A. Hellings, S. Toovey, E.P. Prinssen, Maternal immune activation and abnormal brain development across CNS disorders, *Nat Rev Neurol*, 10 (2014) 643-660.
- [5] A.S. Brown, U. Meyer, Maternal Immune Activation and Neuropsychiatric Illness: A Translational Research Perspective, *Am J Psychiatry*, DOI 10.1176/appi.ajp.2018.17121311(2018) appi.ajp.201817121311.
- [6] S.D. Bilbo, C.L. Block, J.L. Bolton, R. Hanamsagar, P.K. Tran, Beyond infection - Maternal immune activation by environmental factors, microglial development, and relevance for autism spectrum disorders, *Exp Neurol*, 299 (2018) 241-251.
- [7] U. Meyer, J. Feldon, S.H. Fatemi, In-vivo rodent models for the experimental investigation of prenatal immune activation effects in neurodevelopmental brain disorders, *Neurosci Biobehav R*, 33 (2009) 1061-1079.
- [8] A.C. Kentner, S.D. Bilbo, A.S. Brown, E.Y. Hsiao, A.K. McAllister, U. Meyer, B.D. Pearce, M.V. Pletnikov, R.H. Yolken, M.D. Bauman, Maternal immune activation: reporting guidelines to improve the rigor, reproducibility, and transparency of the model, *Neuropsychopharmacology*, DOI 10.1038/s41386-018-0185-7(2018).
- [9] S. Smolders, T. Notter, S.M.T. Smolders, J.M. Rigo, B. Brone, Controversies and prospects about microglia in maternal immune activation models for neurodevelopmental disorders, *Brain Behav Immun*, 73 (2018) 51-65.
- [10] F.S. Mueller, M. Polesel, J. Richetto, U. Meyer, U. Weber-Stadlbauer, Mouse models of maternal immune activation: Mind your caging system!, *Brain Behav Immun*, 73 (2018) 643-660.
- [11] M. Careaga, S.L. Taylor, C. Chang, A. Chiang, K.M. Ku, R.F. Berman, J.A. Van de Water, M.D. Bauman, Variability in PolyIC induced immune response: Implications for preclinical maternal immune activation models, *J Neuroimmunol*, 323 (2018) 87-93.
- [12] M.L. Estes, A.K. McAllister, Maternal immune activation: Implications for neuropsychiatric disorders, *Science*, 353 (2016) 772-777.
- [13] J.P. Lerch, L. Gazdzinski, J. Germann, J.G. Sled, R.M. Henkelman, B.J. Nieman, Wanted dead or alive? The tradeoff between in-vivo versus ex-vivo MR brain imaging in the mouse, *Front Neuroinform*, 6 (2012) 6.
- [14] M.C. van Eede, J. Scholz, M.M. Chakravarty, R.M. Henkelman, J.P. Lerch, Mapping registration sensitivity in MR mouse brain images, *Neuroimage*, 82 (2013) 226-236.
- [15] C.J. Finlay, S. Duty, A.C. Vernon, Brain morphometry and the neurobiology of levodopa-induced dyskinesias: current knowledge and future potential for translational pre-clinical neuroimaging studies, *Front Neurol*, 5 (2014) 95.
- [16] A.C. Vernon, S. Natesan, M. Modo, S. Kapur, Effect of chronic antipsychotic treatment on brain structure: a serial magnetic resonance imaging study with ex vivo and postmortem confirmation, *Biol Psychiatry*, 69 (2011) 936-944.

- 1 [17] A.M. Graham, J.M. Rasmussen, M.D. Rudolph, C.M. Heim, J.H. Gilmore, M.
2 Styner, S.G. Potkin, S. Entringer, P.D. Wadhwa, D.A. Fair, C. Buss, Maternal
3 Systemic Interleukin-6 During Pregnancy Is Associated With Newborn Amygdala
4 Phenotypes and Subsequent Behavior at 2 Years of Age, *Biol Psychiatry*, 83
5 (2018) 109-119.
- 6 [18] M.D. Rudolph, A.M. Graham, E. Feczko, O. Miranda-Dominguez, J.M.
7 Rasmussen, R. Nardos, S. Entringer, P.D. Wadhwa, C. Buss, D.A. Fair, Maternal IL-
8 6 during pregnancy can be estimated from newborn brain connectivity and
9 predicts future working memory in offspring, *Nat Neurosci*, 21 (2018) 765-772.
- 10 [19] A.C. Vernon, W.R. Crum, J.P. Lerch, W. Chege, S. Natesan, M. Modo, J.D.
11 Cooper, S.C. Williams, S. Kapur, Reduced cortical volume and elevated astrocyte
12 density in rats chronically treated with antipsychotic drugs-linking magnetic
13 resonance imaging findings to cellular pathology, *Biol Psychiatry*, 75 (2014) 982-
14 990.
- 15 [20] W.R. Crum, S.J. Sawiak, W. Chege, J.D. Cooper, S.C.R. Williams, A.C. Vernon,
16 Evolution of structural abnormalities in the rat brain following in utero exposure
17 to maternal immune activation: A longitudinal in vivo MRI study, *Brain Behav*
18 *Immun*, 63 (2017) 50-59.
- 19 [21] E. Drazanova, J. Ruda-Kucerova, L. Kratka, K. Horska, R. Demlova, Z. Starcuk,
20 Jr., T. Kasparek, Poly(I:C) model of schizophrenia in rats induces sex-dependent
21 functional brain changes detected by MRI that are not reversed by aripiprazole
22 treatment, *Brain Res Bull*, 137 (2018) 146-155.
- 23 [22] R. Hadar, M.L. Soto-Montenegro, T. Gotz, F. Wieske, R. Sohr, M. Desco, C.
24 Hamani, I. Weiner, J. Pascau, C. Winter, Using a maternal immune stimulation
25 model of schizophrenia to study behavioral and neurobiological alterations over
26 the developmental course, *Schizophr Res*, 166 (2015) 238-247.
- 27 [23] N.V. Malkova, J.J. Gallagher, C.Z. Yu, R.E. Jacobs, P.H. Patterson, Manganese-
28 enhanced magnetic resonance imaging reveals increased DOI-induced brain
29 activity in a mouse model of schizophrenia, *Proc Natl Acad Sci U S A*, 111 (2014)
30 E2492-2500.
- 31 [24] Y. Piontkewitz, M. Arad, I. Weiner, Abnormal trajectories of
32 neurodevelopment and behavior following in utero insult in the rat, *Biol*
33 *Psychiatry*, 70 (2011) 842-851.
- 34 [25] J. Richetto, R. Chesters, A. Cattaneo, M.A. Labouesse, A.M.C. Gutierrez, T.C.
35 Wood, A. Luoni, U. Meyer, A. Vernon, M.A. Riva, Genome-Wide Transcriptional
36 Profiling and Structural Magnetic Resonance Imaging in the Maternal Immune
37 Activation Model of Neurodevelopmental Disorders, *Cereb Cortex*, 27 (2017)
38 3397-3413.
- 39 [26] A.C. Vernon, P.W. So, D.J. Lythgoe, W. Chege, J.D. Cooper, S.C. Williams, S.
40 Kapur, Longitudinal in vivo maturational changes of metabolites in the prefrontal
41 cortex of rats exposed to polyinosinic-polycytidylic acid in utero, *Eur*
42 *Neuropsychopharmacol*, 25 (2015) 2210-2220.
- 43 [27] K.N. Murray, M.E. Edye, M. Manca, A.C. Vernon, J.M. Oladipo, V. Fasolino, M.K.
44 Harte, V. Mason, B. Grayson, P.C. McHugh, I. Knuesel, E.P. Prinssen, R. Hager, J.C.
45 Neill, Evolution of a maternal immune activation (mIA) model in rats: early
46 developmental effects, *Brain Behav Immun*, DOI
47 10.1016/j.bbi.2018.09.005(2018).

- [28] R.C. Roderick, A.C. Kentner, Building a framework to optimize animal models of maternal immune activation: Like your ongoing home improvements, it's a work in progress, *Brain Behav Immun*, 75 (2019) 6-7.
- [29] H.G. Potter, A.T. Wang, S.T.M. Munni, H.M. Kowash, M.A. Burgess, T. Aarons, J.D. Glazier, J.C. Neill, R. Hager, Maternal immune activation affects offspring ultrasonic vocalisations and impairs cognition in rats, *J Psychopharmacol*, 32 (2018) A77.
- [30] D. Mattei, A. Djodari-Irani, R. Hadar, A. Pelz, L.F. de Cossio, T. Goetz, M. Matyash, H. Kettenmann, C. Winter, S.A. Wolf, Minocycline rescues decrease in neurogenesis, increase in microglia cytokines and deficits in sensorimotor gating in an animal model of schizophrenia, *Brain Behav Immun*, 38 (2014) 175-184.
- [31] A.R. Wolff, D.K. Bilkey, The maternal immune activation (MIA) model of schizophrenia produces pre-pulse inhibition (PPI) deficits in both juvenile and adult rats but these effects are not associated with maternal weight loss, *Behav Brain Res*, 213 (2010) 323-327.
- [32] Y. Piontkewitz, M. Arad, I. Weiner, Risperidone administered during asymptomatic period of adolescence prevents the emergence of brain structural pathology and behavioral abnormalities in an animal model of schizophrenia, *Schizophr Bull*, 37 (2011) 1257-1269.
- [33] M. Jenkinson, C.F. Beckmann, T.E. Behrens, M.W. Woolrich, S.M. Smith, *Fsl*, *Neuroimage*, 62 (2012) 782-790.
- [34] B.B. Avants, N.J. Tustison, G. Song, P.A. Cook, A. Klein, J.C. Gee, A reproducible evaluation of ANTs similarity metric performance in brain image registration, *Neuroimage*, 54 (2011) 2033-2044.
- [35] T.C. Wood, C. Simmons, S.A. Hurley, A.C. Vernon, J. Torres, F. Dell'Acqua, S.C. Williams, D. Cash, Whole-brain ex-vivo quantitative MRI of the cuprizone mouse model, *PeerJ*, 4 (2016) e2632.
- [36] N.J. Tustison, B.B. Avants, P.A. Cook, Y. Zheng, A. Egan, P.A. Yushkevich, J.C. Gee, N4ITK: improved N3 bias correction, *IEEE Trans Med Imaging*, 29 (2010) 1310-1320.
- [37] J.L. Andersson, S. Skare, J. Ashburner, How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging, *Neuroimage*, 20 (2003) 870-888.
- [38] J.L.R. Andersson, S.N. Sotiropoulos, An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging, *Neuroimage*, 125 (2016) 1063-1078.
- [39] E. Calabrese, A. Badea, C. Watson, G.A. Johnson, A quantitative magnetic resonance histology atlas of postnatal rat brain development with regional estimates of growth and variability, *Neuroimage*, 71 (2013) 196-206.
- [40] R.W. Cox, AFNI: software for analysis and visualization of functional magnetic resonance neuroimages, *Comput Biomed Res*, 29 (1996) 162-173.
- [41] D. Ma, H.E. Holmes, M.J. Cardoso, M. Modat, I.F. Harrison, N.M. Powell, J.M. O'Callaghan, O. Ismail, R.A. Johnson, M.J. O'Neill, E.C. Collins, M.F. Beg, K. Popuri, M.F. Lythgoe, S. Ourselin, Study the Longitudinal in vivo and Cross-Sectional ex vivo Brain Volume Difference for Disease Progression and Treatment Effect on Mouse Model of Tauopathy Using Automated MRI Structural Parcellation, *Front Neurosci-Switz*, 13 (2019).

- [42] C.R. Genovese, N.A. Lazar, T. Nichols, Thresholding of statistical maps in functional neuroimaging using the false discovery rate, *Neuroimage*, 15 (2002) 870-878.
- [43] L.R. Qiu, D.J. Fernandes, K.U. Szulc-Lerch, J. Dazai, B.J. Nieman, D.H. Turnbull, J.A. Foster, M.R. Palmert, J.P. Lerch, Mouse MRI shows brain areas relatively larger in males emerge before those larger in females, *Nature Communications*, 9 (2018).
- [44] C.E. Meyer, F. Kurth, S. Lepore, J.L. Gao, H. Johnsonbaugh, M.R. Oberoi, S.J. Sawiak, A. MacKenzie-Graham, In vivo magnetic resonance images reveal neuroanatomical sex differences through the application of voxel-based morphometry in C57BL/6 mice, *Neuroimage*, 163 (2017) 197-205.
- [45] J.Y. Zhang, Q. Peng, Q. Li, N. Jahanshad, Z.P. Hou, M.L. Jiang, N. Masuda, D.R. Langbehn, M.I. Miller, S. Mori, C.A. Ross, W.Z. Duan, Longitudinal characterization of brain atrophy of a Huntington's disease mouse model by automated morphological analyses of magnetic resonance images, *Neuroimage*, 49 (2010) 2340-2351.
- [46] H.E. Holmes, N.M. Powell, D. Ma, O. Ismail, I.F. Harrison, J.A. Wells, N. Colgan, J.M. O'Callaghan, R.A. Johnson, T.K. Murray, Z. Ahmed, M. Heggenes, A. Fisher, M.J. Cardoso, M. Modat, M.J. O'Neill, E.C. Collins, E.M. Fisher, S. Ourselin, M.F. Lythgoe, Comparison of In Vivo and Ex Vivo MRI for the Detection of Structural Abnormalities in a Mouse Model of Tauopathy, *Front Neuroinform*, 11 (2017) 20.
- [47] R.M. Yoder, K.C. Pang, Involvement of GABAergic and cholinergic medial septal neurons in hippocampal theta rhythm, *Hippocampus*, 15 (2005) 381-392.
- [48] E. Gould, N.J. Woolf, L.L. Butcher, Postnatal development of cholinergic neurons in the rat: I. Forebrain, *Brain Res Bull*, 27 (1991) 767-789.
- [49] C.Y. Huh, R. Goutagny, S. Williams, Glutamatergic neurons of the mouse medial septum and diagonal band of Broca synaptically drive hippocampal pyramidal cells: relevance for hippocampal theta rhythm, *J Neurosci*, 30 (2010) 15951-15961.
- [50] D.D. Dickerson, A.R. Wolff, D.K. Bilkey, Abnormal long-range neural synchrony in a maternal immune activation animal model of schizophrenia, *J Neurosci*, 30 (2010) 12424-12431.
- [51] G. Ducharme, G.C. Lowe, R. Goutagny, S. Williams, Early alterations in hippocampal circuitry and theta rhythm generation in a mouse model of prenatal infection: implications for schizophrenia, *PLoS One*, 7 (2012) e29754.
- [52] L. Pratt, L. Ni, N.M. Ponzio, G.M. Jonakait, Maternal inflammation promotes fetal microglial activation and increased cholinergic expression in the fetal basal forebrain: role of interleukin-6, *Pediatr Res*, 74 (2013) 393-401.
- [53] T.L. Kemper, M. Bauman, Neuropathology of infantile autism, *J Neuropathol Exp Neurol*, 57 (1998) 645-652.

1 Figures and table

2

3 **Figure 1.** Significant increases in circulating maternal IL-6 levels 3 hours post-
4 poly (I:C) (POL) injection (plasma samples from n=5 dams; 10 mg/kg i.p.
5 administered on GD15) as compared to saline-injected dams (CON; plasma
6 samples from n=5 dams; sterile saline, i.p.) determined using ELISA-based
7 assay. Data shown are IL-6 levels in pg/ml, * $p < 0.05$; 2-tailed student's t-test.

8

9

10 **Figure 2.** Prenatal exposure to POL at GD15 resulted in a significant main
11 effect of MIA ($p < 0.01$; $q < 0.1$; partial $\eta^2 = 0.44$) on the relative volume of the
12 diagonal domain (DD) with an increase in POL-exposed offspring compared to
13 controls at PD21. This increase is more apparent in the females POL offspring
14 as compared to the males. Data shown are relative volumes, expressed as
15 the percentage of total brain volume. ★ $p < 0.01$ main effect of MIA, corrected
16 for multiple comparisons at $q < 0.1$ using FDR procedure; Δ $p < 0.05$, MIA x sex
17 interaction at trend level significance ($p < 0.05$; $q > 0.1$). **A**, Anterior; **P**, posterior;
18 **L**, left, **R**, right.

Region of interest	2 x 2 ANOVA			Post-hoc test
	Sex (<i>p</i> , <i>q</i>)	MIA x Sex interaction (<i>p</i> , <i>q</i>)	MIA (<i>p</i> , <i>q</i>)	
Whole brain	<i>F</i> (1,15)=5.89; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=3.32; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.31; <i>p</i> >0.05 (<i>q</i> >0.1)	None
BNST	<i>F</i> (1,15)=4.97; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.16; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.01; <i>p</i> >0.05 (<i>q</i> >0.1)	None
Cerebellum	<i>F</i> (1,15)=4.40; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=1.25; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.44; <i>p</i> >0.05 (<i>q</i> >0.1)	None
Lateral ventricles	<i>F</i> (1,15)=4.34; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.09; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.01; <i>p</i> >0.05 (<i>q</i> >0.1)	None
Diagonal domain	<i>F</i> (1,15)=0.12; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=5.91; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=11.8; <i>p</i> <0.01 (<i>q</i> <0.1)	None
Olfactory structures	<i>F</i> (1,15)=0.13; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=1.07; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=6.11; <i>p</i> <0.05 (<i>q</i> >0.1)	None
Preoptic area	<i>F</i> (1,15)=0.02; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.03; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=4.77; <i>p</i> <0.05 (<i>q</i> >0.1)	None
Nucleus accumbens	<i>F</i> (1,15)=1.06; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=5.49; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=2.19; <i>p</i> >0.05 (<i>q</i> >0.1)	None
Anterior commissure	<i>F</i> (1,15)=0.41; <i>p</i> >0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=5.11; <i>p</i> <0.05 (<i>q</i> >0.1)	<i>F</i> (1,15)=0.10; <i>p</i> >0.05 (<i>q</i> >0.1)	None

Table 1. The results of a 2 x 2 ANOVA statistical analysis of whole and regional brain volumes at PD21 comparing CON and POL-exposed offspring. Only ROIs with *p*<0.05 (prior to multiple comparison correction) are shown. The *p*-values shown are derived from the F-tests of the main effects of either MIA or sex and MIA x sex interaction run across all 23 ROI. The *q*-values indicate statistical significance after FDR correction using the Benjamini-Hochberg procedure with a 10% threshold (*q*<0.1) to control for Type-I errors across the multiple ANOVA tests. Significant (*p*<0.05; *q*<0.01) F-tests are highlighted in red, whilst those in blue were significant at trend-level only (*p*<0.05; *q*<0.1).

1

2