The association between dietary habits and periodontal disease in young adult

women

Short title: Dietary habits and periodontal disease

Akemi Hosoda^{1,2}, Yuriko Komagamine^{1*}, Manabu Kanazawa¹, Yohei Hama¹,

Akiko Kojo², Shunsuke Minakuchi¹

¹Gerodontology and Oral Rehabilitation, Graduate School of Medical and Dental

Sciences, Tokyo Medical and Dental University, Tokyo, Japan

²Division of Medical Nutrition, Faculty of Healthcare, Tokyo Healthcare

University, Tokyo, Japan

*Corresponding author

Email: y.komagamine.gerd@tmd.ac.jp

Abstract

Background

Among middle-aged and elderly individuals, dietary habits have previously been reported to differ between patients with and without periodontal disease. However, in young adults, there are only a few reports that show a correlation between nutrient/food intake and periodontal disease. Moreover, no report has assessed the correlation between dietary habits measured by a self-administered diet history questionnaire (DHQ) and periodontal disease. Therefore, we assessed the correlation between dietary habits, determined using a DHQ, and periodontal disease in young adult women who are likely to develop a periodontal disease.

Methods

The participants were 120 healthy, non-smoking, female college students (mean age, 20.4 ± 1.1 years) from two universities who did not have any systemic disease. The participants were assessed for periodontal disease according to community periodontal index (CPI) and were divided into two groups. Subjects with a CPI code of 0, 1, or 2 were assigned to non-periodontal disease group (non-PD), and subjects with a CPI code of 3 or 4 were assigned to periodontal

disease group (PD). Dietary habits were assessed using a DHQ. In addition, physical status, level of difficulty in chewing food (dietary hardness), masticatory performance, and quality of life (QoL) were assessed.

Results

The PD group had a significantly lower nutrient intake of minerals, fat, water-soluble vitamins, and dietary fiber than the non-PD group. In terms of food groups, the PD group consumed significantly lesser amounts of green and yellow vegetables than the non-PD group. In addition, the PD group consumed significantly lesser amounts of hard foods than the non-PD group.

Conclusion

Young adult women with a periodontal disease had a significantly lower nutrient/food intake than young adult women without a periodontal disease.

1. Introduction

Periodontal disease is a chronic inflammatory disease that begins in the gingiva and spreads throughout the periodontal tissues, culminating to its destruction [1]. Periodontal disease occurs in young adulthood and progresses with age [2]. According to the results of the Survey of Dental Diseases 2016, the percentage of persons in Japan with periodontal pockets of ≥ 4 mm increased with age [3]. Moreover, since 17.6% individuals aged 15-24 years and 32.4% individuals aged 25-34 years have periodontal pockets of ≥ 4 mm, periodontal disease is now known to be common among the elderly and the young adults. Among middle-aged individuals, the prevalence of periodontal disease is higher in women than in men. The decrease in estrogen levels after menopause leads to a decrease in bone density, which causes the progression of alveolar bone resorption, a symptom of periodontal disease [4]. Therefore, with age, women are at a greater risk of developing periodontal disease than men [4].

It is important to prevent periodontal disease because it is known to affect systemic diseases such as cardiovascular disease [5], diabetes mellitus [6, 7], and metabolic syndrome [8] and premature/low-birth-weight delivery [9, 10]. Previous studies assessing the causes and etiology of periodontal disease

mainly focused on oral microbes and surrounding host environment [11, 12].

Previously reported environmental factors predisposing the host to periodontal disease include smoking habits, drinking habits [13], and dental hygiene [14].

In addition, the nutrients and food consumed have also been reported to be correlated with periodontal disease. Shariq Najeeb *et al.* suggested that decreased consumption of vitamins and minerals, particularly vitamin C, which has antioxidant activity, was highly associated with periodontal disease; however, vitamin C supplementation was not effective in treating periodontal disease [15]. Varun Kulkarni *et al.* showed a similar intake of particular nutrients in patients with and without periodontal disease. However, they pointed to a possible correlation between consumed nutrients and periodontal disease status [16]. In addition, Yoshihara *et al.* showed a negative correlation between the intake of green and yellow vegetables and periodontal disease prevalence. However, it has also been reported that a high intake of cereals, nuts, sugar, sweets, and candy is positively correlated with the presence of periodontal disease [17]. As described above, most of the previous studies on the effect of diet/nutrient targeted elderly patients, with very few studies targeting the young adults.

Our foods are divided into relatively hard foods and soft foods based on physical properties. The hardness of food is regarded as the difficulty involved in chewing, hereafter referred to as dietary hardness. Dietary hardness has been assessed as an estimate of masticatory muscle activity for the habitual diet [28]. A positive correlation between dietary hardness and chewing muscle activity has also been reported in a previous study [18].

Therefore, we aimed at assessing the correlation between dietary habits and periodontal disease in young adult women. In addition, we assessed the correlation between the intake of hard foods and periodontal disease.

2. Materials and Methods

2.1. Participants

One hundred and twenty-seven female college students from the Tokyo Medical and Dental University and the Tokyo Healthcare University were recruited for the study between February 2014 to March 2015.

The inclusion criteria were not having caries teeth and missing teeth excluding third molars. On the other hand, the exclusion criteria were in the following; a

smoker, taking gastrointestinal disease drugs, dry mouth, impaired oral movement and psychiatric illness.

After 6 smokers and 1 person taking gastrointestinal disease drugs were excluded, 120 subjects (mean age, 20.4 years) were finally enrolled in the study.

One dentist with ≥ 5 years of experience in clinical trials assessed the subjects for periodontal disease. Two other evaluators practiced the chewing gum and gummy jelly tests in advance so that their results matched. The Ethical Committee of the Tokyo Medical and Dental University (#1002) and Tokyo Healthcare University (#25-8) approved the study protocols. All participants provided written informed consent before enrolment in the study, and experiments were performed following the guidelines in the Helsinki Declaration on the use of human subjects for research.

2.2. Dietary assessment

Nutrient/food intake was measured using the self-administered diet history questionnaire (DHQ). The validity and reliability of the DHQ have previously been confirmed [19, 20]. The DHQ includes questions concerning the frequency of consumption, the semi-quantitative portions of 151 food and beverage items

consumed, general eating habits including skipping meals and eating irregularly, use of dietary supplements, and major cooking methods. An *ad hoc* of food composition in Japan was used to calculate the estimated dietary intake of energy, food group items, and nutrients [21]. The nutrient/food intake was converted to intake per 1,000 kcal of consumed energy. Since the total intake was expected to vary with body size, physical activity, and energy intake the nutrient/food intake was adjusted for total energy by using energy density (per 1,000 kcal) [22].

2.3. Estimation of dietary hardness

Dietary hardness was calculated using a previously described method [23]. Each food intake was multiplied by each masticatory muscle's activity for the habitual diet. Subsequently, dietary hardness was obtained by dividing the total sum of the above value-by the energy intake.

2.4. Periodontal disease variables

Each participant underwent assessments for periodontics according to the Community Periodontal Index (CPI) from code 0 to code 4 (code 0: health

periodontal conditions; code 1: gingival bleeding on probing; code 2: calculus and bleeding; code 3: periodontal pocket 4–5 mm; and code 4: periodontal pocket ≥ 6 mm) [24, 25]. Then, the participants were assigned to two groups, namely, non-periodontal disease (non-PD) group for CPI codes 0-2 and periodontal disease (PD) group for CPI codes 3-4.

2.5. Anthropometry

Age and self-reported body heights were obtained from the DHQ. The weight and percentage of body fat were measured using a body composition meter (Inner scan 50, TANITA corporation, Japan). Body mass index (BMI) was calculated as body weight (kilograms) divided by the square of body height (meters).

2.6. Measuring masticatory performance by using color-changeable

chewing gum and gummy jelly test

Mixing ability was measured using a color-changeable chewing gum (Masticatory Performance Evaluating Gum XYLITOL, Lotte Co., Ltd, Tokyo Japan). After mouth-rinsing with water for 15 seconds, the participants were then

allowed to chew the provided gum by free chewing at a rate of one stroke per second with an electric metronome. The participants chewed the gum for 60 strokes. Changes in color were assessed using a colorimeter (CR-13, Konica-Minolta sensing, Tokyo, Japan), and then the mean values of L^* , a^* , and b^* in the CIELAB color system were measured. Thereafter, the ΔE values were obtained using following the equation [26, 27].

$$\Delta E = \sqrt{(L^* - 72.3)^2 + (a^* + 14.9)^2 + (b^* - 33.0)^2}$$

Comminution test was measured using a test gummy jelly (UHA Mikakuto Co., Ltd, Japan) and a reported scoring method [28]. The participants chewed the test jelly on their preferred chewing side for a total of 30 strokes. Subsequently, the chewed test jellies were evaluated on a scale of one to ten by using a visual scoring method [29]. Higher scores represented better comminution.

2.7. Health-related QoL

The general health-related QoL of the participants was evaluated by using the Health-related quality of life scale, Short Form-8 (SF-8) [30, 31, 32].

2.8. Statistical analyses

Age, height, weight, BMI, body fat percentage, nutrient/food intake, dietary hardness, masticatory performance, and health-related QoL were compared between the two groups by using Student's *t*-test or Mann-Whitney U test. JMP Pro 11.0.0 (SAS) software was used for analyses. *P* < 0.05 was considered statistically significant. Except for physical status, dietary hardness, and QoL, all the other variables were log-transformed to achieve a normal distribution.

3. Results

3.1. Characteristics of participants

The mean age, body fat percentage, weight, height, and BMI of each group are shown in Table 1. The average age of the participants was 20.4 ± 1.1 years. Seventy-one subjects were assigned to the non-PD group, and 49 subjects were assigned to the PD group. There was no significant difference in age, height, weight, percentage body fat, and BMI between the two groups.

Table 1. The character of participants.

		Total		non-PD		PD		<i>p</i>
		Mean	SD	Mean	SD	Mean	SD	
Age ²⁾	yrs	20.4	1.1	20.4	1.1	20.3	1.1	0.10
Body height ²⁾	cm	158.9	5.8	158.4	5.3	159.0	6.3	0.91
Body weight ¹⁾	kg	51.2	6.9	51.2	7.4	51.1	6.2	0.89
BMI ²⁾	kg/m ²	20.3	2.4	20.4	2.6	20.2	2.0	0.78
Body fat ¹⁾	%	27.0	5.2	27.0	5.8	27.0	4.1	0.97

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

2) Mann-Whitney U test

3.2. Intake of nutrients

Table 2 shows the results of the intake of nutrients. A comparison of energy intake and protein, fat, and carbohydrate energy ratio did not show a significant difference between the two groups. The intake of minerals such as potassium, calcium, magnesium, and iron was significantly lower in the PD group. Moreover, the intake of fat-soluble vitamins, including vitamin A, beta-carotene equivalence, vitamin E, and vitamin K, was significantly lower in the PD group. The intake of water-soluble vitamins, including pantothenic acid (vitamin B5), vitamin B6, and folic acid (vitamin B9), were also significantly lower in the PD group.

202

Table 2. Nutrient intake in the PD and non-PD groups.

		Total		non-PD		PD		<i>p</i>
		Mean	SD	Mean	SD	Mean	SD	
Energy ¹⁾	kcal/day	1769	403	1806	431	1715	356	0.27
Protein ¹⁾	% energy	14.1	1.9	14.3	1.9	13.8	1.8	0.16
Fat ²⁾	% energy	30.6	5.3	31.1	5.1	29.8	5.6	0.30
Carbohydrate ¹⁾	% energy	53.1	5.8	52.6	5.5	54.0	6.3	0.21
Sodium ²⁾	mg/1000kcal	2198	582	2223	644	2161	484	0.86
Potassium ²⁾	mg/1000kcal	1142	248	1185	244	1080	242	0.04*
Calcium ¹⁾	mg/1000kcal	273	90	285	81	256	101	0.02*
Magnesium ¹⁾	mg/1000kcal	124	27	128	28	117	25	0.02*
Phosphorus ¹⁾	mg/1000kcal	527	93.0	539	91	509	94	0.06
Iron ¹⁾	mg/1000kcal	3.9	0.8	4.0	0.1	3.6	0.1	0.01*
Zinc ¹⁾	mg/1000kcal	4.1	0.5	4.2	0.5	4.0	0.5	0.11
Copper ²⁾	mg/1000kcal	0.57	0.10	0.59	0.11	0.55	0.09	0.09
Manganese ²⁾	mg/1000kcal	1.98	0.7	1.96	0.74	2.02	0.57	0.22
Vitamin A ¹⁾	µg/1000kcal	297	155	325	167	256	128	0.007**
Retinol ²⁾	µg/1000kcal	141	120	151	132	128	102	0.55
b-Carotene equivalents ¹⁾	µg/1000kcal	1841	1004	2068	1041	1515	855	0.001**
Vitamin D ¹⁾	µg/1000kcal	3.5	1.9	3.4	1.8	3.5	2.2	0.97
Vitamin E ¹⁾	mg/1000kcal	4.5	1.1	4.7	1.2	4.1	1.0	0.002**
Vitamin K ¹⁾	µg/1000kcal	147	74	162	83	125	53	0.004**
Thiamin ¹⁾	mg/1000kcal	0.46	0.10	0.47	0.10	0.44	0.12	0.13
Riboflavin ²⁾	mg/1000kcal	0.72	0.16	0.73	0.16	0.72	0.18	0.36
Niacin ¹⁾	mg/1000kcal	8.0	1.9	8.1	1.7	7.9	2.1	0.66
Vitamin B ₆ ¹⁾	mg/1000kcal	0.58	0.15	0.60	0.15	0.55	0.15	0.03*
Vitamin B ₁₂ ¹⁾	µg/1000kcal	3.0	1.2	3.0	1.2	3.0	1.3	0.75
Folic acid ¹⁾	µg/1000kcal	158	44	167	47	144	35	0.006**
Pantothenic acid ²⁾	mg/1000kcal	3.19	0.55	3.28	0.55	3.05	0.53	0.01*
Vitamin C ¹⁾	mg/1000kcal	50	22	52	22	48	22	0.25
Fatty acid ²⁾	g/1000kcal	29.63	5.40	30.16	5.29	28.85	5.52	0.39
Saturated ¹⁾	g/1000kcal	9.87	2.12	9.96	2.15	9.74	2.08	0.33
Monounsaturated ²⁾	g/1000kcal	12.02	2.63	12.21	2.52	11.73	2.79	0.38
Polyunsaturated ²⁾	g/1000kcal	7.45	1.60	7.67	1.55	7.12	1.63	0.12

n-3 Polyunsaturated ²⁾	g/1000kcal	1.28	0.36	1.30	0.35	1.25	0.38	0.50
n-6 Polyunsaturated ²⁾	g/1000kcal	6.27	1.31	6.48	1.26	5.95	1.31	0.06
Cholesterol ²⁾	mg/1000kcal	173	61	177	69	167	49	0.67
Total dietary fiber ¹⁾	g/1000kcal	6.7	1.8	7.1	1.7	6.1	1.8	< 0.001***
Soluble dietary fiber ¹⁾	g/1000kcal	1.7	0.5	1.8	0.4	1.6	0.6	0.003**
Insoluble dietary fiber ¹⁾	g/1000kcal	4.7	1.3	5.0	1.3	4.3	1.2	< 0.001***

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

2) Mann-Whitney U test

*P<0.05, **P<0.01, ***P<0.001

3.3. Intake of different food groups

The results of food intake per 1000 kcal of energy for each group are shown in Table 3. The intake of green and yellow vegetables and other vegetables was significantly lower in the PD group than in the non-PD group. Although there was no significant difference in the intake of seeds, nuts, and beans, their intake was lower in the PD group. In addition, the intake of sugar, animal fats, and beverages was higher in the PD group. When green and yellow vegetables and other vegetables were analyzed (Table 4), the intake of carrots (among green and yellow vegetables) and the intake of cabbage (among other vegetables) were significantly lower in the PD group than in the non-PD group. Furthermore, although there was no significant difference, the intake of broccoli, as a green

216 and yellow vegetable, and that of cucumber, eggplant, and lotus root, as a part

217 of the other vegetables, were lower in the PD group.

218

219

Table 3. Food intake per 1000 kcal in the PD and non-PD groups.

g/1,000kcal

	Total		non-PD		PD		<i>p</i>
	Mean	SD	Mean	SD	Mean	SD	
Cereals ²⁾	204.3	58.2	197.9	59.2	213.5	56.2	0.19
Nuts and seeds ²⁾	1.1	2.5	1.2	2.6	0.9	2.2	0.09
Potatoes ²⁾	13.9	8.4	14.8	8.4	12.7	8.4	0.06
Sugars ²⁾	6.2	3.0	6.1	2.6	6.3	3.6	0.86
Confectioneries ²⁾	51.4	25.1	51.3	25.3	51.6	25.1	0.89
Animal fat ²⁾	0.5	0.7	0.4	0.5	0.6	0.8	0.20
Vegetable fats and oil ²⁾	12.4	8.5	12.4	8.3	12.4	9.0	0.59
Pulses ¹⁾	38.2	41.1	42.9	48.4	31.5	26.2	0.17
Fruits ²⁾	45.6	38.1	46.7	41.6	43.9	32.6	0.93
Green and yellow vegetables ¹⁾	61.4	35.9	68.5	38.5	51.2	29.4	0.01*
Other vegetables ¹⁾	61.6	34.4	67.3	37.7	53.3	27.5	0.02*
Mushrooms ²⁾	7.3	7	7.3	6.5	7.4	8.7	0.21
Seaweeds ²⁾	7.4	7.3	7.6	6.7	7.0	8.2	0.57
Seasonings ²⁾	8.5	6.0	8.2	4.9	8.8	7.3	0.91
Alcoholic beverages ²⁾	25.3	50.8	25.9	52.7	25.6	48.5	0.58
Beverages ²⁾	383.7	241.4	357.5	244.0	421.7	234.8	0.09
Fish ¹⁾	28.9	15.2	28.4	15.3	29.5	15.2	0.84
Meat ¹⁾	43.3	18.9	44.4	17.1	41.7	21.3	0.17
Eggs ²⁾	19.3	13.5	20.2	15.1	18.1	10.9	0.76
Milk produces ²⁾	69.3	61.6	71.4	54.9	66.2	70.8	0.28

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

2) Mann-Whitney U test

*P<0.05

220

221

222

Table 4. Intake of green/yellow and other vegetables in the PD and non-PD groups.

g/1,000kcal

	Total		non-PD		PD		<i>p</i>
	Mean	SD	Mean	SD	Mean	SD	
Green and yellow vegetables							
Carrots ²⁾	8.1	5.9	9.1	5.9	6.7	5.8	0.02*
Pumpkins ²⁾	3.8	5.1	4.1	5.8	3.5	4.0	0.94
Tomatoes ¹⁾	13.0	12.5	14.5	13.6	10.9	10.3	0.13
Sweet peppers ²⁾	2.9	3.3	3.1	3.6	2.6	2.8	0.36
Broccoli ²⁾	6.6	9.3	7.7	10.2	5.1	7.8	0.06
Green leafy vegetables ²⁾	17.7	18.5	20.8	22.3	13.3	9.5	0.15
Other vegetables							
Cabbage ¹⁾	14.7	14.9	17.1	16.6	11.2	11.3	0.01*
Cucumbers ¹⁾	4.8	5.7	5.5	5.5	3.8	4.3	0.05
Lettuce ²⁾	6.3	5.8	7.3	6.5	5.0	4.2	0.13
Chinese cabbage ²⁾	4.4	7.4	4.4	7.5	4.4	7.2	0.73
Bean sprouts ²⁾	5.4	7.1	5.7	7.0	5.0	7.2	0.60
Radishes ²⁾	6.4	6.0	7.0	6.5	5.6	5.2	0.43
Onions ¹⁾	10.9	9.5	11.3	10.2	10.3	8.6	0.59
Cauliflower ²⁾	0.3	0.9	0.4	1.1	0.2	0.6	0.25
Eggplants ²⁾	3.4	5.4	3.6	4.0	3.1	7.1	0.06
Burdock ²⁾	1.7	2.3	2.0	2.8	1.2	1.1	0.12
Lotus root ²⁾	0.8	1.1	0.9	1.3	0.7	0.9	0.06

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

2) Mann-Whitney U test

*P<0.05

223

224 3.4. Dietary hardness

225 Masticatory muscle activity was analyzed as a measurement of dietary hardness

(Table 5). The results showed that the intake of hard foods was significantly lower in the PD group when compared with the non-PD.

Table 5. Hard food intake in the PD and non-PD groups.

m•V/1000kcal

	Total		non-PD		PD		<i>p</i>
	Mean	SD	Mean	SD	Mean	SD	
Intake of hard foods ¹⁾	204	30.4	208	31	197	30	0.046*

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

*P<0.05

3.5. Masticatory performance and health-related QoL

Masticatory performance and QoL are shown in Table 6. There was no significant difference in mixing ability, comminution ability, and QoL between the two groups.

Table 6. Masticatory performance and health related QoL.

	Total		non-PD		PD		<i>p</i>
	Mean	SD	Mean	SD	Mean	SD	
Masticatory Performance							
Color-changeable chewing gum ¹⁾	22.5	5.0	22.2	5.3	22.9	0.7	0.42
Gummy jelly test ²⁾	6.2	0.9	6.1	0.1	6.3	0.1	0.45
SF-8 score							
Physical component summary ²⁾	53.4	4.4	53.2	0.5	53.8	0.6	0.52
Mental component summary ²⁾	44.4	7.0	44.5	7.1	44.3	6.9	0.82

Total n=120; non-PD n=71; PD n=49

1) Student's t-test

2) Mann-Whitney U test

4. Discussion

In this study, the two groups showed significant differences in the intake of several nutrients.

In previous studies, subjects with periodontal disease significantly consumed less potassium and iron. In the present study, the intake of potassium and iron in the PD group was significantly lower than that in the non-PD group, which was consistent with that reported in previous studies. Magnesium is known to be involved in bone and tooth formation. It was reported that in 40-year-old individuals, serum magnesium concentration is associated with the

risk of periodontal disease [33]. The intake of magnesium was significantly lower in the PD group in the present study. We, therefore, think that there is a significant positive correlation between the presence of periodontal disease and low intake of magnesium in not only the elderly but also in young adults. Both animal and human studies have reported that a lack of calcium intake leads to alveolar bone resorption [34, 35] and a decrease in bone density [36], indicating that insufficient calcium intake might cause the progression of periodontal disease. In addition, it was shown that the risk of periodontal disease associated with calcium intake was observed in both men and women aged between 20 and 39 years (odds ratio; 1.99 and 1.84), only in men aged between 40 and 59 years, and in neither men nor women aged ≥ 60 years [37]. In the present study, the intake of calcium was significantly lower in the PD group. Therefore, we think that low calcium intake is more strongly correlated with the presence of periodontal disease in younger adults than in older adults.

In the present study, the intake of vitamin C was lower, although not significantly, in the PD group, while the intake of vitamins A, K, and E and beta-carotene were significantly lower. Among the elderly and smokers, the prevalence of periodontal disease was reported to be associated with vitamin C

265 intake. There was a weak but significant dose-response increase in the risk of
266 periodontal disease in subjects who had lower vitamin C intake [38]. Vitamin C is
267 known to suppress oxidative stress through its antioxidant activity [39]. In
268 addition, vitamins A and K [40] and vitamin E and beta-carotene [41] have
269 antioxidant activity. In previous studies, oxidative stress caused by reactive
270 oxygen species in the tissues surrounding the gingiva affects the onset of
271 periodontal disease [42]. Therefore, it was suggested that the antioxidant effect
272 of several vitamins contributed to the absence of periodontal disease in the
273 non-PD group. Low intake of folic acid is known to be related to increased
274 bleeding during probing [43, 44], which supports our findings that the intake of
275 folic acid was significantly lower in the PD group. In the present study, the intake
276 of pantothenic acid and vitamin B6, which are not known to have any correlation
277 with periodontal disease, was significantly lower in the PD group. Previous
278 reports have shown that decreased vitamin B complex intake reduced the
279 efficiency of access flap surgery for periodontal disease treatment [45].
280 However, the effect of each specific vitamin B in the complex on periodontal
281 disease treatment has not been shown yet. Based on our findings, it was
282 suggested that there is a positive correlation between the intake of pantothenic

acid and vitamin B6, among the vitamin B complex, and the absence of periodontal disease.

In a previous study, there was a significant positive correlation between the intake of whole grains and the absence of periodontal disease because whole grains contain dietary fiber [46]. Green and yellow vegetables and other vegetables like whole grains contain soluble and insoluble dietary fibers. Therefore, in this study, since the consumption of green/yellow vegetables and other vegetables was higher in the non-PD group than in the PD group, we think that dietary fibers play a protective role against the development of periodontal disease.

It was reported that the risk for periodontal disease among elderly persons with low intake of docosahexaenoic acid is approximately 1.5 times that among those with high intake [47]. However, in the present study, there was no significant difference in the intake of fatty acids between the two groups. N-3 polyunsaturated fatty acids such as docosahexaenoic acid and eicosapentaenoic acid exert anti-inflammatory effects by blocking intracellular signaling systems required for producing inflammatory cytokines such as IL-1, IL-6, and TNF- α [47]. Therefore, the immunological differences between younger

adults and the elderly might have caused the difference in the findings between the present study and previous studies.

This study showed that the PD group significantly consumed foods with lower dietary hardness when compared with the non-PD group. Yanagisawa, *et al.* classified foods into 10 ranks according to chewing muscle activity [18]. In the study, carrots and cabbage were ranked 8th out of 10 ($>1400 \mu V \cdot sec$) and carrots, cabbage, and lotus root were ranked 6th out of 10 ($>1000 \mu V \cdot sec$). We found that the non-PD group routinely consumed hard dietary foods when compared with the PD group. The quantity of saliva secreted has been reported to increase proportionally with an increase in chew number, chewing time, or chewing muscle activity [48]. Saliva washes away residual food particles and bacteria in the mouth. Saliva also contains lysozyme, amylase, peroxidase, and the proteins lactoferrin and IgA which possess antibacterial activity [49, 50]. Therefore, we think that eating foods with high dietary hardness (as seen in the non-PD group) protects against periodontal disease through oral washing and antibacterial activity resulting from increased saliva secretion.

In the present study, the participants in the PD group accounted for 40.8% of the total participants. The 2016 Survey of Dental Diseases reported that

women aged 20 to 24 with periodontal disease accounted for 16.7% of the total population [3]. Thus, the high prevalence of periodontal disease observed in our study population is likely because we did not comprehensively diagnose periodontal disease by using multiple diagnostic methods. Our diagnoses were based only on the CPI, which might have resulted in the observed high prevalence.

Presently, dietary or nutritional studies assessing how diet/nutrition can be helpful in the prevention or treatment of periodontal disease are lacking. Therefore, more studies are required to elucidate the connection between periodontal disease and diet/nutrition. In addition, longitudinal studies on dietary habits that effectively prevented or suppressed the progression of periodontal disease are warranted.

5. Conclusion

Under the limited conditions of the present study, the intake of minerals (potassium, calcium, magnesium, and iron), fat-soluble vitamins (vitamin A, beta-carotene, vitamin E, and vitamin K), water-soluble vitamins (pantothenic

acid, vitamin B6, and folic acid), and dietary fibers was significantly lower in young adult women with periodontal disease than in young adult women without periodontal disease.

Acknowledgments

The authors thank all the participants from the Tokyo Medical and Dental University and Tokyo Healthcare University.

References

1. Iwasaki M, Manz MC, Moynihan P, Yoshihara A, Muramatsu K, Watanabe R, et al. Relationship between saturated fatty acids and periodontal disease. J Dent Res. 2011; 90: 861-867.
2. Williams RC Jr, Mahan CJ. Periodontal disease and diabetes in young adults. J Am Med Assoc. 1960; 172: 776-778.

3. Ministry of Health, Labor and Welfare of Japan. Report on survey of dental diseases in Japan 2016, Table 21, p23. Available from: <https://www.mhlw.go.jp/toukei/list/dl/62-28-02.pdf> [accessed September 30, 2018]

4. Shapiro LF, Freeman K. The relationship between estrogen, estrogen receptors and periodontal disease in adult women. J Mich Dent Assoc. 2014; 96: 40-44.

5. Teeuw WJ, Slot DE, Susanto H, Gerdes VE, Abbas F, D'Aiuto F, et al. Treatment of periodontitis improves the atherosclerotic profile: a systematic review and meta-analysis. J Clin Periodontol. 2014; 41: 70-79.

6. Lalla E, Papapanou PN. Diabetes mellitus and periodontitis: a tale of two common interrelated diseases. Nat Rev Endocrinol. 2011; 7: 738-748.

7. Morita I, Inagaki K, Nakamura F, Noguchi T, Matsubara T, Yoshii S, et al. Relationship between periodontal status and levels of glycated hemoglobin. J

373 Dent Res. 2012; 91: 161-166.

374

375 8. Shimazaki Y, Saito T, Yonemoto K, Kiyohara Y, Iida M, Yamashita Y.

376 Relationship of metabolic syndrome to periodontal disease in Japanese women:

377 the Hisayama Study. J Dent Res. 2007; 86: 271-275.

378

379 9. Corbella S, Taschieri S, Francetti L, De Siena F, Del Frabro M.

380 Periodontal disease as a risk factor for adverse pregnancy outcomes: a

381 systematic review and meta-analysis of case-control studies. Odontology. 2012;

382 100: 232-240.

383

384 10. Chambrone L, Guglielmetti MR, Pannuti CM, Chambrone LA. Evidence

385 grade associating periodontitis to preterm birth and/or low birth weight: I. A

386 systematic review of prospective cohort studies. J Clin Periodontol. 2011; 38:

387 795-808.

388

389 11. Van Dyke TE, Dave S. Risk factors for periodontitis. J Int Acad

390 Periodontol 2005; 7: 3-7.

391

392 12. Cronin AJ, Claffey N, Stassen LF. Who is at risk? Periodontal disease
393 risk analysis made accessible for the general dental practitioner. Br Dent J.
394 2008; 205: 131-137.

395

396 13. Shimazaki Y, Saito T, Kiyohara Y, Kato I, Kubo M, Iida M, et al.
397 Relationship between drinking and periodontitis: the Hisayama Study. J
398 Periodontol. 2005; 76: 1534-1541.

399

400 14. Sakki TK, Knuuttila ML, Vimpri SS, Hartikainen MS. Association of
401 lifestyle with periodontal health. Community Dent Oral Epidemiol. 1995; 23:
402 155-158.

403

404 15. Najeeb S, Zafar MS, Khurshid Z, Zohaib S, Almas K. The role of
405 nutrition in periodontal health: an update. Nutrients. 2016; 8; pii: E530. Available
406 from <https://www.mdpi.com/2072-6643/8/9/530>

407

408 16. Kulkarni V, Bhatavadekar NB, Uttamani JR. The effect of nutrition on

periodontal disease: a systematic review. J Calif Dent Assoc. 2014; 42: 302-311.

17. Yoshihara A, Watanabe, Hanada N, Miyazaki H. A longitudinal study of the relationship between diet intake and dental caries and periodontal disease in elderly Japanese subjects. Gerodontology 2009; 26: 130-136.

18. Yanagisawa Y, Tamura A, Teramoto Y, Akasaka M. A classification of foods by the amount of masticatory action involved. Shoni Shikagaku Zasshi 1989; 27: 74-84.

19. Ainamo J, Barmes D, Beagrie G, Cutress T, Martin J, Sardo-Infirri J. Development of the World Health Organization (WHO) community periodontal index of treatment needs (CPITN). Int Dent J. 1982; 32: 281-291.

20. Cutress TW, Ainamo J, Sardo-Infirri J. The community periodontal index of treatment needs (CPITN) procedure for population groups and individuals. Int Dent J. 1987; 37: 222-233.

21. Hama Y, Kanazawa M, Minakuchi S, Uchida T, Sasaki Y. Properties of
a color-changeable chewing gum used to evaluate masticatory performance. J
Prosthodont Res. 2014; 58: 102-106.
22. Willett W and Stampfer MJ. Total energy intake: implications for
epidemiologic analy
23. Murakami K, Sasaki S, Takahashi Y, Uenishi K, Yamasaki M,
Hayabuchi H, et al. Hardness (difficulty of chewing) of the habitual diet in relation
to body mass index and waist circumference in free-living Japanese women
aged 18-22 y. Am J Clin Nutr. 2007; 86: 206-213.
24. Sasaki S, Yanagibori R, Amano K. Self-administered diet history
questionnaire developed for health education: a relative validation of the
test-version by comparison with 3-day diet record in women. J Epidemiol 1998;
8: 203-215.
25. Kobayashi S, Murakami K, Sasaki S, Okubo H, Hirota N, Notsu A, et al.

Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in Japanese adults. Public Health Nutr. 2011; 14: 1200-1211.

26. Ministry of Education, Culture, Sports, Science and Technology, Science and Technology Agency. Standard Table of Food Composition in Japan: Official Gazette Co-operation of Japan; 2015. Available from: http://www.mext.go.jp/en/policy/science_technology/policy/title01/detail01/1374030.htm

27. Tarkowska A, Katzer L, Ahlers MO. Assessment of masticatory performance by means of a color-changeable chewing gum. J Prosthodont Res. 2017; 61: 9-19.

28. Yasui S, Nokubi T, Yoshimuta Y, Nokubi F, Kusunoki C, Kida M, et al. Study on reliability for clinical application of a masticatory performance scoring method using test gummy jelly. J Masticat & Health Soc. 2012; 22: 11-17.

463

464 29. Nokubi T, Yoshimuta Y, Nokubi F, Yasui S, Kusunoki C, Ono T, et al.
465 Validity and reliability of a visual scoring method for masticatory ability using test
466 gummy jelly. Gerodontology. 2013; 30: 76-82.

467

468 30. Miyashita M, Yamaguchi A, Kayama M, Narita Y, Kawada N, Akiyama
469 M, et al. Validation of the Burden Index of Caregivers (BIC), a multi-dimensional
470 short care burden scale from Japan. Health Qual Life Outcomes. 2006; 4: 52.

471

472 31. Sugimoto M, Takegami M, Suzukamo Y, Fukuhara S, Kakehi Y.
473 Health-related quality of life in Japanese men with localized prostate cancer:
474 assessment with the SF-8. Int J Urol. 2008; 15: 524-528.

475

476 32. Fukuhara S, Suzukamo Y. Manual of the SF-8 Japanese Version.
477 Institute for Health Outcomes and Process Evaluation Research, Kyoto. 2004.

478

479 33. Meisel P, Schwahn C, Luedemann J, John U, Kroemer HK, Kocher T.
480 Magnesium deficiency is associated with periodontal disease. J Dent Res. 2005;

84: 937-941.

34. Aral K., Alkan BA., Saraymen R., Yay A., Şen A., Önder GÖ.
Therapeutic effects of systemic vitamin K2 and vitamin D3 on gingival
inflammation and alveolar bone in rats with experimentally induced periodontitis.
J Periodontol. 2015; 86: 666-673.

35. Gong A, Chen J, Wu J, Li J, Wang L, Goltzman D, et al.
1,25-Dihydroxyvitamin D deficiency accelerates alveolar bone loss independent
of aging and extracellular calcium and phosphorus. J Periodontol. 2018; 89:
983-994.

36. Song CH, Barrett-Connor E, Chung JH, Kim SH, Kim KS. Associations
of calcium and magnesium in serum and hair with bone mineral density in
premenopausal women. Biol Trace Elem Res. 2007; 118: 1-9.

37. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ.
Calcium and the risk for periodontal disease. J Periodontol. 2000; 71:

499 1057-1066.

500

501 38. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ.

502 Dietary vitamin C and the risk for periodontal disease. J Periodontol. 2000; 71:

503 1215-1223.

504

505 39. Washko P, Rotrosen D, Levine M. Ascorbic acid in human neutrophils. J

506 Biol Chem. 1993; 268: 15531-15535.

507

508 40. Khmelevskii IuV, Danilevskii NF, Borisenko AV, Poberezkina NV. Effect

509 of vitamins A, E, and K on the indices of the glutathione antiperoxide system in

510 gingival tissues in periodontitis. Vopr Pitan. 1985; 4: 54-56.

511

512 41. Moynihan PJ. The role of diet and nutrition in the etiology and

513 prevention of oral diseases. Bull World Health Organ. 2005; 83: 694–699.

514

515 42. Battino M, Bullon P, Wilson M, Newman H. Oxidative injury and

516 inflammatory periodontal diseases: the challenge of anti-oxidants to free radicals

and reactive oxygen species. Crit Rev Oral Biol Med. 1999;10: 458-476

43. Erdemir EO Bergstrom J. Effect of smoking on folic acid and vitamin B12 after nonsurgical periodontal intervention. J Clin Periodontol. 2007; 34: 1074-1081.

44. Esaki M, Morita M, Akhter R, Akino K, Honda O. Relationship between folic acid intake and gingival health in non-smoking adults in Japan. Oral Dis. 2010; 16: 96-101.

45. Neiva RF, Al-Shammari K, Nociti FH Jr., Soehren S, Wang HL. Effects of vitamin-B complex supplementation on periodontal wound healing. J. Periodontol. 2005; 76: 1084–1091.

46. Nielsen SJ, Trak-Fellermeier MA, Joshipura K, Dye BA. Dietary fiber intake is inversely associated with periodontal disease among US adults. J Nutr. 2016; 146: 2530-2536.

- 535 47. Iwasaki M, Yoshihara A, Moynihan P, Watanabe R, Taylor GW,
536 Miyazaki H. Longitudinal relationship between dietary ω -3 fatty acids and
537 periodontal disease. Nutrition 2010; 26: 1105-1109.
538
- 539 48. Yanagisawa Y, Wakabayashi T. The effect of difference in physical
540 property of cooked rice upon digestive speed of starch. J Masticat & Health Soc.
541 1991; 1: 45-52.
542
- 543 49. Tenovuo JO. Human saliva: Clinical chemistry and microbiology. CRC
544 press; 1998.
545
- 546 50. Edgar M, Dawes C, O'Mullane D. Saliva and oral health. 3th ed.
547 Stephen Hancocks Ltd; 2004.