1 **Chemical degumming increases larvae size and facilitates the**
2 **commercial production of Lumpfish (*Cyclopterus lumpus*) eggs**

3

4

5 Craig Pooley, Mia Berwick, & Carlos Garcia de Leaniz

6 Centre for Sustainable Aquatic Research

7 Swansea University, Swansea, SA2 8PP

8

9 **Abstract**

10 Many fishes produce adhesive eggs that confer protection from currents and
 11 predators in the wild, but that are more difficult to disinfect and aerate under
 12 aquaculture conditions. Removing egg adhesiveness ('degumming') has proved
 13 beneficial in the culture of many fish, and a recent gap analysis identified this as a
 14 potential way of increasing hatching success and minimize the risk of infectious
 15 diseases in the culture of lumpfish (*Cyclopterus lumpus*), a novel species to
 16 aquaculture. We tested the efficacy of the enzyme alcalase (0.02%, 0.2%, 2%) as a
 17 degumming agent for lumpfish eggs, and examined its effects on hatching success,
 18 survival, and larvae size under laboratory and commercial conditions. A five-minute
 19 exposure to 0.2% and 2% alcalase decreased chorion thickness by 14% and
 20 resulted in 61-75% degumming rates, without any negative effects on hatching rate,
 21 larval survival, or incidence of embryo malformations. Degummed eggs hatched
 22 earlier than controls and resulted in larger larvae, which may confer some benefits
 23 under aquaculture conditions. A cost-benefit analysis indicates that the benefits of
 24 egg degumming compensate the costs of chemical treatment under most conditions,
 25 and that the optimal alcalase concentration is around 0.2%. We therefore
 26 recommend egg degumming as a way of making the lumpfish industry more efficient
 27 and sustainable.

28

29 **Statement of relevance:** Improving the commercial production of lumpfish

30

31 **Keywords:** Adhesive eggs, Alcalase, Cleaner fish, Cost benefit, Egg degumming,
 32 Hatching

33

34 1. Introduction

35 Many fish species lay eggs that become adhesive in contact with the water, which is
 36 thought to be adaptive as it increases embryo survival (Rizzo et al., 2002). For
 37 example, some freshwater fish produce sticky eggs that cling to the substrate to
 38 avoid being washed away with the current, while other species may produce
 39 adhesive eggs to form discrete clumps that are easier to tend, or that stick to the
 40 vegetation to decrease predation risk or increase oxygen supply (Riehl &
 41 Patzner, 1998). However, the same adhesiveness that benefits eggs in the wild may
 42 hamper their viability under aquaculture conditions, where eggs are typically reared
 43 at very high densities and, if untreated, are prone to suffer from fungal infections and
 44 other diseases (Ringle et al, 1992).

45 The benefits of removing the adhesiveness of the eggs ('degumming') in fish
 46 farming have long been known (Rottman et al., 1991; Billard et al. 1995). These
 47 include improved aeration, reduced risk of fungal diseases, more effective
 48 disinfection, and less effort associated with egg count and husbandry (Ringle et al,
 49 1992). Egg degumming has proved beneficial for the culture of many freshwater
 50 fishes, including cyprinids (Linhart et al, 2003; El-Gamal & El-Greisy, 2008; Žibienė
 51 et al., 2017), tench (*Tinca tinca*, Linhart et al., 2000; Gela et al., 2003; Kujawa et al.,
 52 2010), walleye (*Sander vitreus*, Krise et al., 1986), pikeperch (*Sander lucioperca*,
 53 Demska-Zakes et al., 2005), barbels (Al Hazzaa & Hussein, 2003), sturgeons
 54 (Pšenička, 2016), and catfishes (Isaac & Fries, 1991; Ringle et al., 1992; Linhart et
 55 al., 2003; Asraf et al., 2013; Muchlisin et al., 2014; Kareem, et al., 2017). Although
 56 less common in marine aquaculture, egg degumming has also proved effective for
 57 anadromous and marine species (white sturgeon, *Acipenser transmontanus*; Kowtal
 58 et al, 1986; ballan wrasse, *Labrus bergylta*; Grant et al., 2016).

59 Naturally sticky fish eggs have been chemically treated with many different
60 degumming agents with various degrees of success, including urea (Kowtal et al.,
61 1986), milk (Billard et al., 1995; Linhart et al., 2000), pineapple juice (Thai & Ngo,
62 2004), clay (Siddique et al., 2016), talc (Linhart et al. 2002), tannic acid (Demska-
63 Zakes et al., 2005), sodium chloride (Kowtal et al., 1986) and sodium sulphite (Isaac
64 & Fries, 1991) among others. Glycoproteins and glycosaminoglycans present in the
65 egg chorion are the molecules responsible for making fish eggs sticky (Riehl &
66 Patzner, 1998), and for this reason enzymes that can break down peptide bonds,
67 such as papain (Isaac & Fries, 1991), trypsin (Linhart et al., 2002) and various endo-
68 proteases (Linhart et al., 2003; El-Gamal & El-Greisy, 2008), have proved
69 particularly effective as degumming agents. Among these, alcalase (a commercial
70 endo-protease with broad specificity obtained from the bacterium *Bacillus*
71 *licheniformis*) has been effective in both freshwater (Linhart et al., 2003) and marine
72 species (Grant et al., 2016). However, egg degumming also poses risks and could
73 easily damage the fragile fish embryo (Pšenička, 2016). Given that species differ
74 markedly in the structure and chemical composition of the egg chorion (Riehl &
75 Patzner, 1998), no single degumming method is likely to be suitable in all cases.

76 One novel species to Aquaculture that has adhesive eggs and whose
77 incubation in hatcheries could benefit from egg degumming is the lumpfish,
78 *Cyclopterus lumpus* (Powell et al., 2018a). Large scale commercial production of
79 lumpfish only started recently but has increased exponentially over the last few years
80 as the species is increasingly being used as cleaner fish to control parasitic sea lice
81 in salmon farming (Powell et al., 2018a). Different lumpfish populations have
82 different fecundities and size at maturation (Whittaker et al., 2018), but an analysis
83 identified egg degumming as a potential way of increasing hatching success and

84 minimizing the risk of infectious diseases during embryo development (Powell et al.,
85 2018a). Under natural conditions, lumpfish eggs become sticky following exposure to
86 divalent calcium and magnesium in seawater (Lönning et al., 1984). Following
87 fertilization, males create funnel-like depressions in the egg mass and puff water and
88 vigorously fan the eggs to facilitate gaseous exchange and the removal of
89 nitrogenous waste, such paternal care being maintained throughout the incubation
90 period (Powell et al., 2018b). In contrast, lumpfish eggs in hatcheries may be left to
91 incubate in large clumps or shaped into a thin monolayer and reared in upwelling
92 incubators (Powell et al., 2018a). Hatching success is very variable in culture and it
93 is not uncommon to lose entire egg batches due to poor oxygenation and
94 subsequent spread of bacterial and fungal diseases (Powell et al., 2018a). Since a
95 female can typically produce in excess of 100,000 eggs (Davenport, 1985; Powell et
96 al., 2018b), even a modest increase in hatching rate would yield a significant
97 improvement in the number of larvae produced.

98 As with ballan wrasse (Grant et al., 2016), a small-scale pilot trial showed
99 recently that alcalase could also be effective at degumming lumpfish eggs (Powell et
100 al., 2018a), but the side effects and most efficient concentrations were not
101 investigated. Thus, the aim of this study was to test the potential benefits of alcalase
102 treatment to remove the adhesive layer of the lumpfish eggs and examine any
103 potential impacts on embryo survival, timing of hatching, growth, and incidence of
104 malformations. To this end, two experiments were conducted, one under controlled
105 laboratory conditions, where accurate counts could be made of daily mortalities in
106 replicated mini-incubators, and one under commercially relevant conditions in order
107 to validate the results obtained in the laboratory.

108

109

110 **2. Materials and Methods**

111 *2.1. Source of eggs and artificial fertilization*

112 Wild lumpfish were obtained from commercial fishermen operating in Dorset (50° 44'
113 N; 2° 20' W) and Guernsey (49° 27' N; 2° 31' W) during 2016 and 2017 and were
114 transferred to a quarantine unit at the Centre for Sustainable Aquatic Research
115 (CSAR, Swansea University) where they were kept in 1.5 m³ circular tanks at 32 ppt
116 salinity, 13°C ± 0.3°C, and 8:16 L:D photoperiod. Broodstock were fed a mixture of
117 frozen blue mussels, squid, krill and whitebait.

118 Eggs were stripped from anaesthetized females (2-phenoxyethanol,
119 concentration 0.03 ml/litre) and milt was collected from euthanized males as
120 described in Powell et al. (2018a). Sperm motility was assessed by activation of a
121 small sample with seawater, followed by observation under the microscope at x40
122 magnification. Sperm was diluted with a milt extender (AquaBoost SpermCoat,
123 Cryogenetics AS, Norway) at a 1:1 ratio and maintained refrigerated at 4°C for up to
124 14 days. Artificial fertilization was carried out by adding 2-3 ml of extended milt to a
125 batch of eggs (~600 g), followed by gentle agitation for 1 min, activation with ozone-
126 treated seawater, further agitation for 2 min, and rinsing with clean seawater. Egg
127 hardening was completed after c. 30 min at 10°C.

128

129 *2.2. Laboratory (in vitro) degumming trial*

130 Three liquid alcalase (*Bacillus licheniformis*; 126741; VWR, UK) solutions were
131 prepared using 0.2 µm filtered seawater as a diluent to achieve concentrations of
132 0.02%, 0.2% and 2.0% along with a control treatment (0% alcalase). These were
133 maintained in the dark at 4°C for a maximum of 48h prior to use. Immediately

following sperm activation, and prior to egg hardening, 12 batches of fertilized eggs ($n = 104 \pm 10$ eggs) from a single female were exposed for 5 min to one of the four alcalase concentrations, gently inverting the vial to ensure uniform exposure for all eggs. Following alcalase exposure, eggs were allocated at random to 12 x 0.5 L upwelling vertical incubators constructed from 750 mL plastic funnels fitted with a 500 μm mesh “cradle” to hold the eggs. Flow was maintained at 1 Lmin⁻¹ (± 0.05 L), and water temperature was maintained at $8.0\text{ }^{\circ}\text{C} \pm 0.1\text{ }^{\circ}\text{C}$ during the 37-day duration of the trial. Eggs were disinfected with 30 ppm Pyceze (Novartis Pharmaceuticals UK Ltd) for 20 minutes every 5 days.

Eggs were inspected 48 h post-fertilisation (morula stage) to assess their viability and estimate fertilization success (Powell et al., 2018b). Degumming efficiency was assessed by counting the number of single eggs (i.e. those not in a clump, bound together by their adhesive chorion) and dividing by the total number of eggs in the sample. They were then photographed using a Leica DFC290 digital camera (Leica Microsystems Ltd, UK) mounted to a Nikon SMZ 800 Zoom Stereomicroscope at X20 magnification. Mean chorion thickness ($\pm 10\text{ }\mu\text{m}$) was obtained from photomicrographs of six eggs per replicate ($n=18$ per treatment) at 2-4 equidistant points of the egg, as described in Songe et al. (2016).

Hatched larvae were collected daily from individual incubators and preserved in 10% formalin (HT501128; Sigma-Aldrich, UK) to screen for embryo malformations. Hatching success was calculated by dividing the number of hatched larvae by the number of viable eggs (i.e. excluding non-viable eggs that failed to develop after 24 hrs). A sample of fixed larvae ($n=6$ per replicate, $n=18$ per treatment) were photographed and the standard length was calculated from the tip of the snout to the caudal peduncle (resolution $\pm 10\mu\text{m}$). The incidence of embryo deformities such as

spinal scoliosis (Gapasin et al., 1998; Merchie et al., 1997), and operculum and sucker deformities was recorded.

2.3. Degumming under commercial conditions

A 2% alcalase concentration was chosen for the large commercial trial, as this had shown promising results in a small-scale pilot study (Powell et al., 2018a). A solution was prepared by adding 10 ml of alcalase to a 20-litre beaker containing 490 ml of filtered sea water. Fertilised lumpfish eggs (590 ± 10 g) were added to the beaker and aerated. For the control treatment, the same amount of eggs was added to 500 ml of filtered sea water. After 5 minutes, eggs were rinsed with 1 litre of filtered sea water and transferred to 70 litre hopper incubators with a 20 L/min upwelling flow and a 500 μ m screen mesh to keep the eggs off the bottom. Salinity was maintained at 32 ppt and water temperature at 9.0 ± 2.0 °C. Eggs were treated with Pyceze every five days, as above. Approximately 30 eggs were sampled from each clump using a grid system to ascertain viability and degumming efficiency, as described above. During the peak of mass hatching, 100 larvae per batch were reared in triplicate in 30 litre tanks and monitored for eight weeks under the same commercial conditions (temperature 9.0 ± 2.0 °C) and fed enriched *Artemia* and dry diet. Mortalities were recorded daily for each replicate.

2.4. Statistical analysis

We used R v3.4.1 (R Core Team, 2017) for statistical analysis. Degumming efficiency, survival, and incidence of embryo malformations (all proportion data) were analysed via Generalized Linear Mixed-Effect Modelling (GLMM) using the R package “lme4” (Bates et al., 2015) and a binomial error distribution, with alcalase

concentration as a fixed factor and incubator ID as a random effect. Hatching success was modelled as a function of time and alcalase concentration as predictors, and incubator ID as a random effect, using the packages “*survival*” (Therneau, 2015) and “*FrailtyPack*” (Rondeau et al., 2012) to account for unmeasured heterogeneity. The effect of alcalase concentration on chorion thickness and larvae size at hatching was tested by Linear Mixed Modelling (LMM) using alcalase concentration as a fixed effect, and incubator identity as a random effect using the *lme4* and *lmerTest* (Kuznetsova et al., 2017) R packages. We compared models using untransformed and log10 transformed alcalase concentrations, and also treating alcalase concentration as a continuous or discrete predictor. Inspection of AIC values indicated that treating alcalase concentration as a factor resulted in the most plausible models.

2.5. Ethics Statement

This study adhered to the ARRIVE guidelines and was approved by Swansea University Animal Welfare Research Body (permit IP-1516-16)

3. Results

3.1. Efficacy of alcalase as a degumming agent

Alcalase was effective at separating (degumming) lumpfish eggs at concentrations of 0.2% and 2% for 5 minutes, but not at 0.02% (Figure 1, glmer: estimate = 1.97, SE = 0.226, Z value = 8.724, $P < 0.001$). Compared to controls, eggs exposed to alcalase concentrations of 0.2 and 2% displayed mean degumming rates of 61% (± 12) and 75% (± 21), respectively. The degumming effect of alcalase was confirmed by measurements of chorion thickness, which decreased significantly with increasing alcalase concentration (Figure 2; lmer; $F_{3,11.852} = 11.102$, $P < 0.001$). Thus, compared to controls (mean chorion thickness = $68.59 \pm 2.91 \mu\text{m}$), chorion thickness decreased by $8.77 \mu\text{m}$ (± 1.91) at 0.2% and by $10.24 \mu\text{m}$ (± 1.93) at 2% alcalase.

3.2 Effect of alcalase on hatching rate and timing of hatching

There was no significant difference in hatching success between controls and eggs exposed to alcalase, once tank effects were taken into account (Figure 3; frailty model treatment estimate = -0.084, SE = 0.056, Z = -1.48, $P = 0.138$; Exponential distribution

Loglik(model)= -2801.9, Loglik(intercept only)= -2806.4; $\chi^2 = 8.95$, df = 4.2, $P = 0.07$). However, alcalase treatment resulted in a shorter hatching period compared to controls ($F_{3,754} = 69.55$, $P < 0.001$), decreasing mean hatching time by -0.66 days (± 0.11) in the case of 0.02% alcalase ($t = -5.974$, $P < 0.001$) and by -1.59 days (± 0.11) in the case of 2% alcalase ($t = -14.097$, $P < 0.001$).

3.3 Effect of alcalase on larvae size and incidence of malformations

Embryo size increased with increasing alcalase concentration (Figure 4, lmer: estimate = 53.99, SE = 19.27, $t_{69.03} = 2.80$, $P = 0.006$). Compared to controls, eggs treated with 2% alcalase resulted in embryos that were $148.14 \pm 44.76 \mu\text{m}$ larger ($t_{68.98} = 3.31$, $P = 0.001$). No deformities were observed under the microscope in any of the larvae from any treatment.

3.4. Effects of alcalase under commercial conditions

Alcalase was also effective at degumming lumpfish eggs under large scale, commercial conditions (estimate = 0.531, SE = 0.230, Z value = 2.306, $P < 0.05$). Exposure of fertilised eggs to 2% alcalase for five minutes resulted in a mean degumming rate of 0.70 ± 0.06 , compared to 0.0 for controls. Alcalase treatment had no significant effect on larvae survival eight weeks post-hatch (estimate = -0.2285, SE = 0.1782, $t = -1.282$, $P = 0.2$), but eggs exposed to 2% alcalase were significantly larger (mean = $14.4 \text{ mm} \pm 0.14$) than controls (mean = $13.3 \text{ mm} \pm 0.15$; $F_{1,239} = 41.62$, $P < 0.001$), confirming the results of the *in vitro* laboratory trial.

3.5. Cost benefit analysis

During the commercial trial, we exposed fertilised lumpfish eggs at a ratio of one egg to one millilitre of alcalase solution. The alcalase solution was purchased at a cost of £46.45 for 500ml, but placing a large bulk order can probably obtain it c. 25% cheaper at £74.3 per Litre. Based on this bulk cost, and £10,000 fixed costs for the sourcing of wild broodstock, we estimated the benefit to cost ratio of incubating 1 million lumpfish eggs under different egg market prices (1-8 pence/egg), taking into account different hatching and degumming rates at different alcalase concentrations. The simulations (Figure 5) suggest that the optimal alcalase concentration is 0.2%

(over five minutes), and that benefits outweigh costs at 1.56 pence per egg which is the break even point.

4. Discussion

Without parental care in hatcheries, adhesive eggs tend to display reduced gaseous exchange rates, and this can make them more susceptible to opportunistic pathogens (Demska-Zakes et al., 2005). Our results show that alcalase can be effective at removing the adhesive layer of the lumpfish chorion, and that treated eggs result in larger larvae, without any evidence of adverse effects on larval survival or embryo malformations. These findings are similar to work carried out on ballan wrasse, where alcalase treatment achieved >69% degumming rate on a dosage-dependent fashion without any detrimental effects (Grant et al., 2016).

The advantages of egg degumming are various. Degummed eggs allow for more thorough disinfection, easier handling, more accurate egg counting and estimation of fecundity and survival rates, and potentially also more efficient use of space for egg incubation. This can in turn increase hatching success, reduce health risks, and allow for more efficient commercial production (Linhart et al., 2000; Linhart et al., 2003).

Our results indicate that although a five-minute exposure to 2% alcalase achieved the highest degumming rate, a concentration of 0.2% achieved similar results and can be more cost effective, based on a cost-benefit analysis. Little degumming, on the other hand, was observed at a concentration of 0.02% alcalase, although it is possible that increasing exposure time over 5 minutes may achieve better degumming rates, as shown for ballan wrasse (Grant et al., 2016). We exposed eggs to alcalase within ten minutes of artificial fertilisation, before the eggs

had hardened. Further research into the feasibility of degumming naturally spawned (hardened) eggs may be warranted, as naturally spawned clumps of eggs are often encountered in culture (Treasurer et al., 2018).

One unexpected consequence of alcalase treatment was a faster embryo development time and larger larvae. Although the reasons for this are unclear, it is possible that the thinner chorion of degummed eggs may have increased oxygen availability and allowed for more efficient use of energy. Oxygen availability is known to affect egg quality and embryo development in carp (Vodianitskiy et al., 2017), and in general increased oxygen results in larger embryos for many fish species (Braga Goncalves et al., 2015). Embryos developing within thick gelatinous egg masses often show retarded development in marine gastropods (Chaffee & Strathmann, 1984), and it is possible that the same may happen in lumpfish. Whatever the reasons, a significant size advantage was evident 8 weeks post-hatching, even though degummed eggs only hatched less than 2 days earlier than controls. The larger size of larvae originating from degummed eggs could be beneficial to industry, as larger larvae have a greater probability of surviving past the critical period during the transition from endogenous to exogenous feeding (Garrido et al., 2015).

Although a thick egg chorion can make it more difficult for water moulds, such as *Saprolegnia*, to penetrate the eggs (Songe et al., 2016), it can also favour settlement of water borne particulates that facilitate fungal and bacterial infection and can make eggs more vulnerable. Indeed, an adhesive chorion is thought to be particularly susceptible to viral, fungal and bacterial infection, and one of the benefits of achieving a thinner chorion through chemical degumming may be to reduce pathogen loads and facilitate egg prophylaxis under aquaculture conditions (Krise et

al., 1986). Egg degumming in lumpfish could also make it possible to use various types of upwelling incubators that maintain eggs in suspension, such as McDonald-type jars, pelagic egg jars, and Imhoff cones (i.e. Jensen et al., 2008). Their use could achieve a more uniform distribution of clean, oxygenated water with less chance of infection by water moulds. This is an area that warrants further attention because growth of opportunistic bacteria and water moulds on dead eggs can spread quickly and affect surrounding viable eggs (Davenport, 1983). Removing the adhesive layer can help to alleviate this problem as it would enable easier separation and removal of non-viable eggs. In this sense, several techniques are available for removing and counting eggs, including manual siphoning, use of flotation methods (i.e. Leitritz & Lewis, 1976) and use of automatic egg sorters, which could prove effective at separating large numbers of eggs if these are degummed and no longer form clumps. For example, using the method described by Coombs (1981), viable and non-viable lumpfish eggs could be separated using a column and a salinity gradient.

Conclusions

In summary, we describe (and validate under commercial conditions), a method for chemically removing the adhesive layer of lumpfish eggs immediately after artificial fertilization using the enzyme alcalase. A five minute exposure to 2% alcalase resulted in 70% degumming rate, 14% reduction in chorion thickness, and larger larvae without impacting on hatch rates or larvae survival up to 8 weeks after hatching. Cost-benefit analysis indicates that the costs of alcalase treatment are offset under most conditions and that a 0.2% concentration is the most cost-

effective. Egg degumming can facilitate the separation and removal of non-viable eggs, improves egg husbandry, and provides scope for using modified incubators that can reduce incubation space and decrease labour time compared to current methods.

Acknowledgments

This research was part-funded by Marine Harvest Scotland and the ERDF SMARTAQUA Operation. The funders played no role in the collection, analysis or interpretation of the data, the writing of the MS, or the decision to submit the article for publication. We are grateful to staff at CSAR for helping with the experimental set up and egg husbandry, and to Adam Powell and Maria Scolamacchia for setting an initial pilot trial.

References

- Al Hazzaa, R., & Hussein, A., 2003. Stickiness elimination of Himri Barbel (*Barbus lutes*, Heckel) eggs. Turkish Journal of Fisheries and Aquatic Sciences. 3(1), 47-50.
- Asraf, A., Muchlisin, Z.A., Siti-Azizah, M.N., 2013. Removal of eggs adhesiveness of African catfish (*Clarias gariepinus*) at different concentrations of urea solution. Aceh International Journal of Science and Technology, 2, 94-97.
- Billard, R., Cosson, J., Perchee, G., & Linhart, O., 1995. Biology of sperm and artificial reproduction in carp. Aquaculture, 129(1), 95-112.
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67, 1-48.
- Chaffee, C., Strathmann, R., 1984. Constraints on egg masses. I. Retarded development within thick egg masses. Journal of Experimental Marine Biology and Ecology, 84, 73 – 83. doi:10.1016/0022-0981(84)90231-4
- Coombs, S. H., 1981. A density-gradient column for determining the specific gravity of fish eggs, with particular reference to eggs of the mackerel *Scomber scombrus*. Marine Biology. 63(1), 101-106. doi:10.1007/bf00394667
- Davenport, J., 1983. Oxygen and the developing eggs and larvae of the lumpfish, *Cyclopterus lumpus*. Journal of the marine biological Association of the United Kingdom. 63(3), 633-640.
- Davenport, J., 1985. Synopsis of Biological Data on the Lumpsucker *Cyclopterus lumpus* (Linnaeus, 1758). FAO Fisheries Synopsis. 147, 1-41.
- Demska-Zakes, K., Zakes, Z., & Roszuk, J., 2005. The use of tannic acid to remove adhesiveness from pikeperch, *Sander lucioperca*, eggs. Aquaculture Research. 36(14), 1458-1464. doi:10.1111/j.1365-2109.2005.01370.x

366 El-Gamal, A.H.E., El-Greisy, Z.A., 2008. Effect of removal of egg adhesiveness on
367 hatchability and effect of different levels of salinity on survival and larval
368 development in common carp, *Cyprinus carpio*. Journal of Applied Sciences
369 Research. 4, 1935-1945.

370 Gapasin, R. S. J., Bombeo, R., Lavens, P., Sorgeloos, P., & Nelis, H., 1998.
371 Enrichment of live food with essential fatty acids and vitamin C: effects on
372 milkfish (*Chanos chanos*) larval performance. Aquaculture. 162(3-4), 269-
373 286.

374 Garrido, S., Ben-Hamadou, R., Santos, A., Ferreira, S., Teodósio, M., Cotano, U.,
375 Irigoien X., Peck, M.A., Saiz, E., Ré, P., 2015. Born small, die young:
376 Intrinsic, size-selective mortality in marine larval fish. Scientific Reports, 5,
377 17065.

378 Gela, D., Linhart, O., Flajshans, M., & Rodina, M., 2003. Egg incubation time and
379 hatching success in tench *Tinca tinca* (L.) related to the procedure of egg
380 stickiness elimination. Journal of Applied Ichthyology 19, 132-133.

381 Grant, B., Picchi, N., Davie, A., Leclercq, E., & Migaud, H., 2016. Removal of the
382 adhesive gum layer surrounding naturally fertilised ballan wrasse (*Labrus*
383 *bergylta*) eggs. Aquaculture. 456, 44-49.
384 doi:10.1016/j.aquaculture.2016.01.016

385 Isaac Jr, J., & Fries, L. T., 1991. Separation of channel catfish eggs in sodium sulfite
386 with and without papain. The Progressive Fish-Culturist. 53(3), 200-201.

387 Jensen, N.R., Ireland, S.C., Siple, J.T., Williams, S.R., Cain, K.D., 2008. Evaluation
388 of egg incubation methods and larval feeding regimes for North American
389 burbot. North American Journal of Aquaculture, 70, 162-170.

390 Kareem, O., Ajani, E., Akintunde, M., Olanrewaju, A., Oduntan, O., 2017. Effect of
391 different fertilization and egg de-adhesion methods on hatching and survival
392 of *Clarias gariepinus* (Burchell 1822) fry. Journal of FisheriesSciences.com,
393 11(1), 021-027.

394 Kowtal, G. V., Clark, W. H., & Cherr, G. N., 1986. Elimination of adhesiveness in
395 eggs from the white sturgeon, *Acipenser transmontanus*: chemical treatment
396 of fertilized eggs. Aquaculture. 55(2), 139-143.

397 Krise, W. F., Bulkowski-Cummings, L., Shellman, A.D., Kraus, K.A., & Gould, R.W.,
398 1986. Increased Walleye Egg Hatch and Larval Survival after Protease
399 Treatment of Eggs. The Progressive Fish-Culturist. 48(2), 95-100.
400 doi:10.1577/1548-8640(1986)48<95:IWEHAL>2.0.CO;2

401 Kujawa, R., Kucharczyk, D., & Mamcarz, A., 2009. The effect of tannin concentration
402 and egg unsticking time on the hatching success of tench *Tinca tinca* (L.)
403 larvae. Reviews in Fish Biology and Fisheries. 20(3), 339-343.
404 doi:10.1007/s11160-009-9136-z

405 Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B., 2017. lmerTest package:
406 tests in linear mixed effects models. Journal of Statistical Software. 82(13).

407 Leitritz, E., & Lewis, R. C., 1976. *Trout and Salmon Culture (Hatchery Methods)*.
408 Fish Bulletin. 1-198.

409 Linhart, O., Gela, D., Flajshans, M., Duda, P., Rodina, M., & Novak, V., 2000.
410 Alcalase enzyme treatment for elimination of egg stickiness in tench, *Tinca*
411 *tinca* L. Aquaculture. 191, 303-308.

412 Linhart, O., Rodina, M., Gela, D., Flajshans, M., & Kocour, M., 2003. Enzyme
413 treatment for elimination of egg stickiness in tench (*Tinca Tinca* L.),

European catfish (*Silurus Glanis* L.) and common carp (*Cyprinus Carpio* L.).
Fish Physiology and Biochemistry. 28, 507-508.

Linhart, O., Stech, L., Svarc, J., Rodina, M., Audebert, J. P., Grecu, J., & Billard, R.,
2002. The culture of the European catfish, *Silurus glanis*, in the Czech
Republic and in France. Aquatic Living Resources. 15, 139-144.

Lønning, S., Kjörsvik, E. and Davenport, J., 1984. The hardening process of the egg
chorion of the cod, *Gadhus morhua* L, and lumpsucker, *Cyclopterus lumpus*
L. Journal of Fish Biology. 24, 505–522.

Merchie, G., Lavens, P., & Sorgeloos, P., 1997. Optimization of dietary vitamin C in
fish and crustacean larvae: a review. Aquaculture. 155(1-4), 165-181.

Muchlisin, Z.A., Mastura, S., Asraf, A., Fadli, N., Hendri, A., Siti-Azizah, M.N., 2014.
A preliminary study to evaluate the effects of powder milk solution on the
eggs adhesiveness and fertilization rates of African catfish, *Clarias*
gariepinus. Aquaculture, Aquarium, Conservation & Legislation-International
Journal of the Bioflux Society (AACL Bioflux), 7(5), 412-418.

Powell, A., Keay, A.J., Pooley, C.L., Galloway, G., Scolamacchia, M., Frazao- Pires,
A., Scolding, J.W.S., Lloyd, R., Fletcher, D., Cockerill, D., 2015. Optimization
of captive breeding of the lumpfish (*Cyclopterus lumpus*) for sea lice control
in salmon farming. Proceedings of the Aquaculture Europe 2015, Rotterdam.
European Aquaculture Society. pp. 647.

Powell, A., Treasurer, J.W., Pooley, C.L., Keay, A.J., Lloyd, R., Imsland, A.K., Garcia
de Leaniz, C., 2018a. Use of lumpfish for sea-lice control in salmon farming:
challenges and opportunities. Reviews in Aquaculture, 10, 683–702.

437 Powell, A., Pooley, C., Scolamacchia, M., Garcia de Leaniz, C., 2018b. Review of
438 lumpfish biology. *In* Cleaner Fish Biology and Aquaculture Applications. ed.
439 J.W. Treasurer, pp. 98-121. 5M Publishing Ltd., Sheffield.

440 Pšenička, M., 2016. A novel method for rapid elimination of sturgeon egg stickiness
441 using sodium hypochlorite. *Aquaculture*. 453, 73-76.
442 doi:10.1016/j.aquaculture.2015.11.039

443 R Core Team, 2017. R: A language and environment for statistical computing. R
444 Foundation for Statistical Computing, Vienna, Austria. [http://www.R-](http://www.R-project.org/)
445 [project.org/](http://www.R-project.org/).

446 Riehl, R., & Patzner, R.A., 1998. Minireview: the modes of egg attachment in teleost
447 fishes. *Italian Journal of Zoology*. 65(S1), 415-420.

448 Ringle, J. P., Nickum, J. G., & Moore, A., 1992. Chemical separation of channel
449 catfish egg masses. *The Progressive Fish-Culturist*. 54(2), 73-80.

450 Rizzo, E., Sato, Y., Barreto, B.P., Godinho, H.P., 2002. Adhesiveness and surface
451 patterns of eggs in neotropical freshwater teleosts. *J. Fish Biol.* 61, 615–632.

452 Rondeau, V., Mazroui, Y., Mauguén, A., Diakite, A., Gonzalez, J.R., 2012.
453 FRAILTYPACK: An R package for General frailty models using a semi-
454 parametrical penalized likelihood estimation or a parametrical estimation,
455 2012. R package version 2.2-22.

456 Rottman, R. W., Shireman, J. V., & Chapman, F. A., 1991. Techniques for Taking
457 and Fertilizing the Spawn of Fish. Southern Regional Aquaculture Center,
458 SRAC Publication No. 426.

459 Siddique, M. A. M., Psenicka, M., Cosson, J., Dzyuba, B., Rodina, M., Golpour, A., &
460 Linhart, O., 2016. Egg stickiness in artificial reproduction of sturgeon: an
461 overview. *Reviews in Aquaculture*. 8(1), 18-29.

462 Songe, M. M., Willems, A., Sarowar, M. N., Rajan, K., Evensen, Ø., Drynan, K., ... &
463 Van West, P., 2016. A thicker chorion gives ova of Atlantic salmon (*Salmo*
464 *salar* L.) the upper hand against *Saprolegnia* infections. Journal of Fish
465 Diseases. 39(7), 879-888.

466 Thai, B. T., & Ngo, T. G., 2004. Use of pineapple juice for elimination of egg
467 stickiness of common carp (*Cyprinus carpio* L.). Asian Fisheries
468 Science. 17(1/2), 159-162.

469 Therneau, T. M., & Lumley, T., 2015. Package 'survival'. R Top Doc. 128.

470 Treasurer, J., Prickett, R., Zietz, M., Hempleman, C., Garcia de Leaniz, C., 2018.
471 Cleaner fish rearing and deployment in the UK., In Cleaner Fish Biology and
472 Aquaculture Applications. ed. J.W. Treasurer, pp. 376-391. 5M Publishing
473 Ltd., Sheffield.

474 Vodianitskiy, O. M., Potrokhov, O. S., & Zinkovskiy, O. G., 2017. Embryonic and
475 early postembryonic development of carp and activity of enzymes of the
476 energy and plastic metabolism under impact of water temperature
477 fluctuations. Hydrobiological Journal. 53(1), 76-86.

478 Whittaker, B.A., Consuegra, S., Garcia de Leaniz, C., 2018. Genetic and phenotypic
479 differentiation of lumpfish (*Cyclopterus lumpus*) across the North Atlantic:
480 implications for conservation and aquaculture. PeerJ 6, e5974.

481 Žibienė, G., Žibas, A., Švirinienė, L., 2017. The effects of tannic acid on the
482 effectiveness of egg fertilization and removing carp egg adhesiveness, In
483 Proceedings of the 8th International Scientific Conference Rural
484 Development 2017. ed. A. Raupelienė, pp. 857-861.

485
486

Figures

Figure 1. Effect of a 5-minute exposure to different alcalase concentrations on proportion (mean $\pm$ 95% CI) of degummed lumpfish eggs

Figure 2. Effect of alcalase treatment on thickness of the egg chorion. Boxplots show minimum, first quartile, median, third quartile, and maximum values. Extreme values are shown by closed circles.

Figure 3. Effect of alcalase treatment on hatching success and timing of hatching (cumulative proportion of hatched larvae).

Figure 4. Effect of alcalase treatment on size of lumpfish larvae (standard length). Shown are minimum, first quartile, median, third quartile, and maximum values. Extreme values are shown by closed circles.

Figure 5. Benefit-cost analysis for the production of 1 million lumpfish eggs using different alcalase concentrations and various egg sale prices (GBP pence/egg). Dotted line represents the estimated break-even point (1.56 pence/egg).

Cumulative % hatched

Concentration

— 0

····· 0.02

- - - 0.2

- - - 2

60

40

20

0

30

31

32

33

34

35

36

37

38

39

40

Days post fertilisation