

1 **Handling and Adequacy of Iodine at Household Level: Community Based Cross-**
2 **sectional Survey in Dega Damot District, West Gojjam Zone, Amhara Regional State,**
3 **Ethiopia**

4 **Aschalew Afework^{1*}, Wondemagegn Mulu², Almayehu Abate³, Abel Lule⁴**

5 **¹GAMBY Medical and Business College, Public Health Department, Bahir Dar,**
6 **Ethiopia,**

7 **E-mail: aschalew28@gmail.com**

8 ***Corresponding authors**

9 **²Bahir Dar University, College of Medicine and Health Sciences, Department of**
10 **Medical Laboratory Science**

11 **E-mail: wondem_32@yahoo.com**

12 **³Amhara Public Health Institute, Immunology department, Bahir Dar, Ethiopia,**

13 **E-mail: alexu2love@gmail.com**

14 **⁴GAMBY Medical and Business College, Public Health Department, Bahir Dar,**
15 **Ethiopia**

16 **E-mail: belolu.Tsi@gmail.com**

17

ABSTRACT

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

Introduction: Iodine is essential for regulation of physical growth and neural development. Although, fortification of iodine has been practiced decades before and iodized salt is available, handling and cultural food preparation may affect the content of iodine in the dishes. Moreover, Dega Damot is mountainous area that may lose its iodine via erosion. Therefore, this study aimed at determining the handling and adequacy of iodine in the salt in Dega Damot district, West Gojjam Zone, Ethiopia.

Method: Community based cross-sectional study was conducted from October 24 to November 15, 2017 on 802 households. Stratified multi-stage sampling was employed to select households. A total of 422 and 380 households from low land and high land, respectively took part in the study. For the interview either the husband or the spouse was selected randomly. Data on handling were collected by face-to-face interview using structured questionnaires. The concentration of iodine was determined using Rapid Test Kit. Descriptive statistics were used to describe relevant findings on the handling of iodized salt. Both bivariate and multivariate logistic regressions were analyzed to identify associated factors.

Result: From 802 samples tested, 37 (4.6%) had iodine greater than 15 parts per million. The majority (94.5%) of the respondents have been roasting their salt for 'Dikus' preparation whereas 91.1% of the households stored their salt in open containers. Salts with closed containers (AOR=3.22, CI=1.31-7.89), unroasted salts [AOR=5.23, CI=1.25- 22.11], good knowledge on handling [AOR=5.55, CI=1.64-18.77], salts from high land area [AOR=2.11, 95%CI=1.02-4.37], were significantly associated with adequacy of iodine

Conclusions: Adequate utilization of iodine was very low. Roasting of salt was common. These phenomena may be continued because 'Dikus' preparation is cultured in this population. We recommend the supplementation of packed iodized salt in the dishes

Key words: Handling, iodine content, Dega Damot, Dikus Preparation, roasted salt

43 INTRODUCTION

44 Iodine is needed for regulation of physical growth and neural development. Iodine deficiency is the major
45 cause of preventable mental retardation and its severity can range from mild mental blunting to cretinism [1].

46 Ethiopian public health institute 2016 survey reported that the national iodized salt utilization coverage was
47 89.2%. However, only about 26% of the surveyed households had salt that was adequately iodized (at ≥ 15
48 ppm). According to this report, the highest coverage of adequately iodized salt was in Tigray (55.2%) and
49 Somali (49.4%) regions, and the lowest in the regions of Gambela (9.5%), SNNPR (13.7%), and Amhara
50 (15%) [2]. The findings indicated that there is good national iodized salt coverage even if very low adequate
51 iodized salt utilization at household level. This might be due to improper handling of the iodized salt.
52 According to World Health Organization (WHO) and International Council for Control of Iodine Deficiency
53 Disorders (ICCIDD) standard, elimination of IDD will be possible if more than 90% of the households
54 consume adequately iodized salt [3].

55 Although Ethiopia, had set a national goal to eliminate IDD virtually by the year 2013 through universal salt
56 iodization (USI) and increase access to iodized salt among households up to 90% [4]. The 2016 Ethiopian
57 public health survey reported [3], only 26% adequate iodine coverage nationally. Moreover, there is great
58 difference between the adequacy of iodine consumption at house hold level in Ethiopia and other study
59 reports in different parts of the world. For example, 54.3% and 75.6% adequate iodine consumption at
60 house hold level reported in Pakistan [5] and Ghana, respectively [6].

61 Factors such as duration of storage, size of the crystals, impurities, and the ambient temperature of the
62 storage, humidity, cooking methods, time of adding and sunlight exposure influence the stability of iodized
63 salt at the household level include [7, 8]. But varieties of improper handling have been reported in different
64 studies. For example, a study done in Ghana, reported that 11.5% of households who stored salt in

65 containers without a lid [6]; and 3.5 % of the respondents in Gondar town wash the salt to remove its
66 impurities [9] and 64.9% of Assabi district add the salt during cooking [10].

67 Dega Damot district is one of the pocket areas of the country. Topographically, it consists of 35%
68 mountainous, 30% ups and downs, 20% valleys and 15% plains. This topographic characteristic might be
69 liable for erosion [11]. Thus, iodine may erode from the top area of the land. This characteristic of the soil
70 with inadequate iodine content salt consumption may worsen the iodine deficiency. Even if high utilization
71 of iodized salt in the country [2], improper handling might reduce the content of iodine in the dish and may
72 not be sufficient to fulfill the demand of the body.

73 In Dega Damot district, there is cultural food preparation called ‘Dikus’. Dikus is prepared primary as raw
74 material for the preparation of Wot (Ethiopian common cultural food). In this preparation, the iodized salt is
75 either heated in sunlight or roasted in dry oven. This might lead to evaporation of iodine from the salt and
76 might reduce the iodine content in the dish. Hence, the study determined the handling of salt and adequacy
77 of iodine at house level in Dega Damot district, Ethiopia.

78 **Materials and Methods**

79 **Study setting, design and period**

80 A community based cross-sectional study was conducted in Dega Damot District from October 24-
81 November 15, 2017. The District is located in West Gojjam Zone of Amhara National Regional State, 296
82 kilometers away from the capital city of Ethiopia. It is mountainous with high land and low land climate and
83 produce different crops. The most commonly used foods in the high land are potato, barely and peens while
84 lowland residents, commonly consumed maize, teff and barley foods. Erosion is very high in this area which
85 may leads to the loss of iodine. According to the information obtained from the zonal finance and economy
86 office based on the 2007 census, total population of the district is about 158, 667 with sex distribution of
87 78,115 males and 80,552 females.. It has 32 sub-districts and the average number of households in each sub-

88 district is estimated to be 830 with the average 6 members in each family [12]

89 **Sample size and Sampling procedure**

90 The sample size was determined using single population formula with the following assumptions: level of
91 confidence, 95%; margin of error, 5%; proportion of adequate iodine concentration from Lay Armachiho
92 District, 29.7% [13]. By taking design effect of 2.5 and 5% contingency for the non-response, the sample
93 size was 842. Stratified multi-stage sampling was used to select households from the source population. At
94 first stage, 6 sub-districts (3 from high land and 3 from low land) were selected by lottery method. At the
95 second stage, 6 villages (1 village from each district) were selected through lottery method. At the third
96 stage, cluster sampling was used to select households from each village. Hence, 421 households from the low
97 land and 421 from the high land were taken for this study. For the interview either the husband or the spouse
98 was selected randomly.

99 **Data Collection**

100 Demographic data and data for iodized salt handling were collected through face to face interview using
101 structured questionnaires. The questionnaire contains open and close ended questions and included a section
102 for observing the type of container used to store the salt and place of salt storage.

103 **Iodine determination**

104 A teaspoon of salts collected from each household. The salts were tested for the content of iodine using the
105 iodine rapid test kit (MBI Kits International). MBI KITS is improved iodized salt Field Test kit for salt
106 fortified with potassium iodide. The interpretation of the result was based on color comparison with its chart
107 (0, 1-15 and more than 15 parts per million, ppm). The cut-off proportion of 15 PPM and above was
108 considered as adequately iodized salt using the WHO reference indicators for monitoring of iodized salt in
109 the household level [14].

110 **Operational definitions**

111 **Adequate iodine in the salt:** If the iodine content of the tested salt was > 15 parts per million

112 **Low iodine content:** If the iodine content of the tested salt was \leq 15parts per million

113 **No iodine:** If the iodine content of the tested salt was 0 parts per million

114 **Good knowledge on handling:** If the respondent correctly answered greater than or equal to half of the
115 knowledge questions items

116 **Poor knowledge on handling:** If the respondent correctly answered less than half of the knowledge
117 questions item

118 **Data Analysis**

119 The data was analyzed using SPSS version 20. Descriptive statistics such as frequencies and mean were
120 used to describe the study participants in relation to relevant variables. The mean score of knowledge items
121 questions was calculated to measure the level of knowledge of the respondents. Both bivariate and
122 multivariate logistic regressions were used to identify associated factors. Odds ratio with 95% confidence
123 interval were used to identify the presence and strength of association. .P-value less than 0.05 were taken as
124 cut off point for statistical significance at the 95% confidence interval.

125 **Ethical Clearance**

126 Ethical clearance was approved from Amhara Public health institute, and permission letter was taken from
127 the district health office. After the purpose of the study was explained, verbal consent obtained from each
128 respondent and participation was based on their willingness. Participants were assured that their names or

129 personal identifiers are not included in the written questionnaire to ensure confidentiality. The findings were
130 reported to the concerned bodies.

131 **Results**

132 **Socio-demographic characteristics**

133 A total of 802 households were took part in the study. Of them, 422 (52.6%) were from the low land. The
134 mean ages of the respondents were 40-year-old (28-52). The average family size was six. Four hundred
135 eighty two (60.1%) of the respondents were females and five hundred thirty four (66.2%) were unable to read
136 and write. Details are depicted in table 1.

137 **Table1. Demographic Characteristics of the respondents in Dega Damot District, October- 2017**

	Categories	Frequency	Percent
Residence	High land	422	52.6
	Low land	380	47.4
Sex	Male	320	39.9
	Female	482	60.1
Age	18-27	92	11.5
	28-37	236	29.4
	38-47	268	33.4
	48-57	107	13.3
	>57	99	12.4
Religion	Orthodox	795	99.13
	Protestant	7	0.87
	Unable to read and write	534	66.6

Level of education	Literate	150	18.7
	1-8th	66	8.2
	9-12th	52	6.5
Family size	2-5	348	43.4
	6-9	412	51.4
	> 10	42	5.2
	Total		802

138

139 **Handling of Iodized Salt**

140 In this study, almost all (99.9) of the households had non-packed iodized salt. Of them, 85.8% were dried salts.
 141 Concerning frequency of buying, 534 (66.6%) of the households bought once a month and 96.5 percent of
 142 them buy the powdered one. The majority (91.1%) of the households stored their salt in an open container.
 143 Seven hundred twelve (88.8%) of them add the salt in the end stage of cooking. Most (94.5%) of the
 144 respondents had been roasting the salt while they prepare ‘Dikus’ (Table 2)

145 **Table2. Handling and utilization of iodized salt in Dega Damot district**

Characteristics	Categories	Frequency	Percentage
Type of salt	Packed	1	0.12
	Not packed	801	99.88
Condition of the salt during survey	Dry	688	85.8
	Moisture	114	14.2
Reason for not using packed iodized salt	Accessibility	104	12.98
	Cost	24	3.00
	Have no information	673	84.02
Amount of salt buy at	Less than 1 kg	76	9.5

a time	1-2 kg	539	67.2
	More than 3-5 kg	146	18.2
	More than 5kg	41	5.1
Frequency of buying	Less than once a month	79	9.8
	Once a month	534	66.6
	Once in 2-3 months	146	18.2
	Greater than 3 months	43	5.4
Type of salt bought	Powdered salt	774	96.5
	Both Crystalline salt and powdered	28	3.5
Storage material	open container	731	91.1
	closed container	71	8.9
Time to add salt	Middle	69	8.6
while cooking food	Late/end	712	88.8
	after cooking	21	2.6
Roasting on the oven	Yes	758	94.5
	No	44	5.5
Washing if impurities	Yes	36	4.5
	No	766	95.5
Salt to livestock	Same salt is given	759	94.6
	Buys different salt	43	5.4
Source of information	Radio	39	4.1
	Health Extension Workers	278	29.3
	Neighbors	631	66.6

146

Level of iodine concentration

147 From the total of 802 salts analyzed, 37 (4.6%) had adequate iodine content (greater than 15ppm) and 654
148 (81.6 %) had undetectable iodine content (Table 3)

149 **Table3. Level of iodine concentration at household in Dega Damot District , West Gojjam, Ethiopia,**

150

October 2017

151

Variable	Frequency	Percent
Iodine content of the salt		
0 parts per million	654	81.6
1-15 parts per million	111	13.8
>15 parts per million	37	4.6

152

153

Multivariate analysis on the level of iodine

154 On multivariate analysis, salts in closed container [AOR=3.22; 95%CI=1.31, 7.89] have 3 times more
155 likely to have concentration greater than 15ppm. Unroasted salts [AOR=5.23; 95%CI=1.25, 22.11] were
156 positively associated with iodine concentration greater than 15ppm. Having good knowledge on handling
157 [AOR=5.55; 95%CI=1.64, 18.77] significantly associated with iodine concentration greater than 15ppm.
158 Salts collected from the high land area [AOR=2.11; 95%CI=1.02, 4.37] were 2 times more likely to have
159 concentration greater than 15ppm. However, moisture content of the salt, educational status of the
160 respondents, income, family size and frequency of buying were not associated with concentration of
161 iodine salt in the salt (Table 4).

162

163

164

Table4. Multivariate analysis of factors that affect the iodine content of salt at Dega Damot

165

District, October 2017.

Variables	Iodine content		COR[95%CI]	AOR[95%CI]
	>15ppm	≤15ppm		
Storage Material				
Closed container	8	63	3.07 [1.35, 7.01]	3.22 [1.31, 7.89]
Open container	29	702	1.00	1.00
Roasted iodized salt				
No	35	588	5.27 [1.26, 22.12]	5.23 [1.25, 22.11]
Yes	2	177	1.00	1.00
Educational status of the respondents				
No educated individuals	28	506	0.71 [0.16, 3.06]	0.67 [0.15, 3.01]
Elementary education	7	208	1.17 [0.23, 5.78]	1.06 [0.20, 5.73]
Secondary educated individuals	2	51	1.00	1.00
Respondents Knowledge on Handling				
Poor	32	742	1.00	1.00
Good	5	23	5.04 [1.80, 14.12]	5.55 [1.64, 18.77]
Income				
Less than or equal to 11, 999	35	750	1.00	1.00
More than 12, 000	2	15	2.86 [0.63, 12.98]	1.69 [0.23, 12.57]
Salt collected site				
Low land area	13	409	1.00	1.00
High land area	24	356	2.12 [1.06, 4.23]	2.11 [1.02, 4.37]
Condition of the salt during survey				

Dry /normal	29	669	1.00	1.00
Moist salt	8	96	1.92[0.85, 4.33]	0.53 [0.23, 1.19]
Frequency of bought				
Less than once a month	4	75	1.00	1.00
Once a month	25	509	1.06 [0.37, 3.20]	1.16 [0.49, 3.59]
Once in 2-3 months	5	141	1.50 [0.39, 5.77]	2.06 [0.49, 8.56]
Greater than 3 months	3	40	0.71 [0.15, 3.33]	0.73 [0.14, 3.82]
Family size				
2-5	18	330	0.45 [0.06, 3.44]	0.43 [0.56, 3.34]
6-9	18	394	0.534 [0.07, 4.10]	0.52 [0.07, 3.99]
Greater than 9	1	41	1.00	1.00

166

167

DISCUSSION

168

169

170

171

172

173

174

175

176

177

178

A community based cross-sectional study attempted to assess handling and adequacy of iodine in the salt at the household level. Proper handling of the iodized salt at household level is a critical point to achieve the intended intervention of iodine fortification. In this study, only 4.6% of households had adequate iodine content in their salts. This finding is very far from WHO recommendation to eliminate iodine deficiency (3) and also with the Ethiopian national plan to eliminate IDD virtually by the year 2013 through adequate iodized salt utilization (4). Moreover, the present finding is significantly lower than 2016 Ethiopia Public health survey report [2]. This is an alarm to work at the community level in order to achieve the intended WHO goals .

Comparing with others, our study finding is by far lower than in Belgaum district, Indian [15], Pakistan [5] and Ghana [6] on which 33.3%, 54.3% and 44.8% of the household had adequate iodized salt, respectively. Moreover, the proportion of adequate iodine consumption in this population was lower than studies done in

179 Gondar [9] and Labo Assabi district [10]. The reason might be due to poor handling of the salt in the study
180 area or cultural food preparation led to high loss of iodine from the salt.

181 In the present study, 94.5% of respondents reported roasting the salt while preparing Dikus (the raw material
182 for the preparation of ‘Wot’ (common cultural food in Ethiopia)). This cultural food preparation led to the
183 common phenomena of roasting of salt in this population. Roasting is not common in other studies even if
184 exposure to sunlight is common practice [10, 17, 18]. This might be one of the reasons for the loss of iodine
185 from the salt. Moreover, almost all (99.9%) of the households used non-packed iodized salt. The result is
186 lower compared to studies done in Wolaita Sodo town [17] and Dera District, Ethiopia [18], the packed
187 iodized salt utilization was 11.1% and 12.49%, respectively. High proportion of roasting of salts and lower
188 utilization of packed iodized salt needs attention in this population.

189 In this study, the roasted salts were 5.2 times less likely to have adequate iodine than the counter parts. Even
190 if it is not identical to that of roasting, exposure to sunlight in the other similar studies were also associated
191 with inadequacy of iodine content. For example, not exposing salt to sunlight was positively associated with
192 utilization of adequately iodized salt in LaboAssabi District [11]. In a similar previous study at Gondar,
193 Ethiopia, those who did not expose salt to sunlight were 7.26 times more adequate iodized salt than those
194 who exposed salt to sunlight [10]. Moreover, exposure of salt to sunlight were about 4 times more likely to
195 have adequately iodized salt as compared to their counter parts in a similar study done in Wolaita Sodo
196 town[16]

197 In this study, respondents who had good knowledge on handling salts were 5.6 times more likely to have
198 iodine concentration greater than 15ppm. Comparatively, respondents who had good knowledge on handling
199 salts were 1.94 times more likely to have adequate iodine than those who do have poor knowledge in a study
200 conducted at Gondar town, Ethiopia [10]. Likewise, knowledge of the respondents was associated with

201 adequacy of iodine content in the salt in Dera District, Ethiopia [17]. However, knowledge about iodized salt
202 has no significant association with iodine content in LaboAssebi District, Ethiopia [11].

203 **Conclusion and recommendation**

204 The proportion of households with adequate iodine salt is very low. Roasting of the salt is a common practice
205 in this population. These phenomena may be continued in the future because ‘Dikus’ preparation is cultured
206 in this population. We recommend the utilization of packed iodized salt in the dish to attain the deficit due to
207 roasting of salts during Dikus preparation

208 **ACKNOWLEDGMENT**

209 We are grateful for GAMBY Medical and Business College in providing conducive environment for the
210 dissemination of the finding of this research. We would like to extend our appreciation to Amhara National
211 Regional State Health Bureau, Research and Technology Transfer Core Process for its ethical approval. Our
212 acknowledgment also goes to Dega Damot health office for its letter of permission. We acknowledge
213 UNICEF, Bahir Dar branch for materials supporting for the chemical analysis. We also thank the
214 respondents. Finally our deepest appreciation goes to the data collector: Tamirat Afework and Nibret Tilahun

215 **References**

- 216 1. National Guideline for Control and Prevention of Micronutrient Deficiencies in Ethiopia: Federal
217 Ministry of Health June 2004
- 218 2. Andinet Hailu. Ethiopian National Micronutrient Survey Report: September 2016. Ethiopian Public
219 Health Institute, Ministry of Health. 2016
- 220 3. UNICEF, ICCID, PAMM, and WHO, *Monitoring Universal Salt Iodization Program*,
221 PAMM/ICCIDD, Ottawa, Canada, 1995.

- 222 4. FMOH, *National Guideline for Control and Prevention of Micronutrient Deficiencies*, Edited by F.
223 H. Department, Federal Ministry of Health, Addis Ababa, Ethiopia, 2010.
- 224 5. Gul Nawaz Khan, Imtiaz Hussain, etal. A Study on the Household Use of Iodised Salt in Sindh and
225 Punjab Provinces, Pakistan: Implications for Policy Makers. *Journal of Pharmacy and Nutrition*
226 *Sciences*, 2012, 2, 148-154
- 227 6. Christiana Buxton, Benjamin Baguune. Knowledge and practices of people in Bia District, Ghana,
228 with regard to iodine deficiency disorders and intake of iodized salt. *Archives of Public Health*
229 2012, **70**:5
- 230 7. The entire salt sample lost iodine over the 12 month sampling period. High humidity leads to loss
231 of iodine in the salt (chemical engineering in canda)
- 232 8. Ritu Rana, Rita Singh Raghuvanshi. *Journal of food science technology*. 2013, 50(6) 1212 -1216
- 233 9. Hailay Gebrearegawi Gebremariam, Melkie Edris Yesuf, and Digsu Negese Koye. Availability of
234 Adequately Iodized Salt at Household Level and Associated Factors in Gondar Town, Northwest
235 Ethiopia. Volume 2013 (2013 SRN Public Health Volume 2013 (2013), Article ID 160582, 6 pages
236 <http://dx.doi.org/10.1155/2013/160582>
- 237 10. Meselech Regassa D, Tsedeke wolde H, and Befirdu mulatu J. utilization of adequately iodized salt
238 on prevention of iodine deficiency disorders at households level and associated factors in Labo
239 Assabi District, west Ethiopia
- 240 11. Dega Damot Woreda Agricultural Office, 2016
- 241 12. West Gojjam zone economic and development office
- 242 13. Molla Mesele, Getu Degu and Haimanot Gebrehiwot. Prevalence and associated factors of goiter
243 among rural children aged 6-12 years old in Northwest Ethiopia, cross -sectional study. *BMC*
244 *Public Health* 2014, 14:130. <http://www.biomedcentral.com/1471-2458/14/130>
- 245 14. UNICEF, ICCID, PAMM, and WHO, *Monitoring Universal Salt Iodization Program*,
246 PAMM/ICCIDD, Ottawa, Canada, 2004

- 247 15. Kamath, V. Bhat, R.S.P. Rao, D. Acharya, U. Kapil, M.S. Kotian, and D.S. Nayak. Prevalence of
248 goitre among school children in Belgaum District. Indian Journal of Pediatrics 2009; 76 (8) : 825-
249 828
- 250 16. Cherinet Abuye Yemane Berhane and Tessema.Ersumo. The Role of Changing Diet and Altitude on
251 Goitre Prevalence in Five Regional States in Ethiopia. East African Journal of Public Health. 2008;
252 5(3): 163-168
- 253 17. Wondimagegn Paulos Kumma, Yusuf Haji, Junayde Abdurahmen, and Yohannes Mehretie
254 Adinew. Factors Affecting the Presence of Adequately Iodized Salt at Home in Wolaita, Southern
255 Ethiopia: Community Based Study. Hindawi International Journal of Food Science. Volume 2018,
256 Article ID 4394908, 9 pages <https://doi.org/10.1155/2018/4394908>
- 257 18. Zelalem Alamrew Anteneh, Melese Engidayehu and Gedefaw Abeje. Iodine content of dietary salt
258 at household level and associated factors using Iodometric titration methods in Dera District,
259 Northwest Ethiopia. Anteneh et al. BMC Nutrition (2017) 3:83