Crab fisherman communities in north Brazil: a new high risk population for vampire bat rabies

Nailde de Paula Silva¹, Elane Araújo Andrade¹, Denis Cardoso², Ruth de Souza Guimarães¹, Mateus Borges Silva¹, Kelly Karoline Gomes Nascimento¹, Diego de Arruda Xavier¹, Isis Abel¹ *.

¹Laboratory of Epidemiology and Geoprocessing (EpiGeo), Veterinary Medicine Institute, Federal University of Pará (UFPA), Castanhal, Pará, Brazil.

² Veterinary Medicine Department, Federal University of Lavras (UFLA), Lavras, Minas Gerais, Brazil

*Corresponding author: isisabel@ufpa.br Phone: +55 (91) 3311-4719.

Abstract

An outbreak of human rabies transmitted by hematophagous bats occurred in 2018 in the state of Pará, Brazil, eastern Amazon, after 14 years with no record of the disease. It is necessary to understand the epidemiological characteristics of these attacks to protect the local population. This study aimed to characterize attacks of humans by vampire bats in the municipality of São João da Ponta, Pará state, Brazil, from 2013 to 2015. All individuals attacked by bats who sought medical care during the study period (n=5) were identified in the Notifiable Diseases Information System (SINAN) database and answered a questionnaire about the circumstances of the attack. Using snowball sampling, seed cases identified other individuals who were attacked in the same period but did not seek medical care (n=61), totalizing 66 people attacked in the same period. The interviewees

were male (92.4%), adults between 20 and 50 years old (69.6%) and had completed elementary education (86.3%). Most were rural residents (92.4%) and crab fishermen (79.3%). The interviewees (92.4%) identified the mangrove of the Mãe Grande de Curuçá extractive reserve as an area conducive to attacks by vampire bats, where groups of fishermen sometimes concentrate for days for crab fishing, often living in improvised dwellings without walls and covered by tarps or straw (88.8%). The wounds were single bites (71.2%) and were located on the lower limbs (93.9%). Overall, 42.4% of participants had been bitten more than four times throughout their life (range 1-23 attacks). Participants were unaware of the risk of contracting rabies by the bite (95.4%). Using São João da Ponta as a model, this study shows that bat attacks are an essentially occupational problem in the study region. Indeed, for each reported attack, there are 12.2 unreported cases. It is necessary to develop strategies to reach this population for prophylactic treatment.

Author Summary

Different from which occurs worldwide in relation to rabies transmission, in Amazon region, vampire bat is involved on direct transmission of rabies virus to humans when searching for bloodmeal. It is common in the state of Pará, Eastern Amazon, large areas inhabited near forests and mangroves. People living there use forest natural resources as a way of income and sustenance and these working conditions is what our study points out as an important factor for aggressions predisposition. Here this subject is shown as an occupational problem. This study also quantified for the first time underreported human's aggressions by bats in Amazon, using the snowball sampling, which valued the relationship between individuals to reach the target population. Based on these results,

rabies surveillance may direct actions for prevention and health education for these individuals, including changes in notifications forms and suggesting pre-exposure prophylaxis in vaccination calendar of the Brazilian Ministry of Health for these individuals exposed to the rabies virus.

Introduction

In general, the causal factors of rabies are manifold. The epidemiological cycle of the viral agent is complex and involves several mammalian hosts. Many researchers consider contact with the wild reservoir, human population mobility, and the social and cultural characteristics of a population as the main determinants for onset of the disease. The high impact of rabies transmission by vampire bats in the Amazon region is unquestionable [1-3].

Climate, seasonality, proximity to the forest, and the presence of livestock and natural prey provide conditions conducive to the proliferation of vampire bats in the Amazon. However, this in itself does not determine the occurrence of bites and transmission of rabies in humans. It is the relationships between humans and the environment that place human populations at epidemiological risk, which is also associated with the lack of medical care, poverty and low level education [4,5].

This relationship between humans and the environment is reflected in the use of the mangroves' natural resources by the families in that region. The municipality of São João da Ponta, Pará state (PA), is located in an Extractive Reserve (São João da Ponta RESEX) in the Eastern Amazon and is surrounded by other conservation units. The main economic activity of the municipality is swamp ghost crab (*Ucides cordatus*) fishing [6]. Informal conversations with the population indicated that many residents in the municipality had been attacked by bats but did not seek medical care.

According to the guidelines of the Health Surveillance Department of the Ministry of Health (SVS/MS), bat bites are subject to mandatory notification throughout Brazil. Every case of a bite or attack involving bats must be notified to health authorities using the Notifiable Diseases Information System (SINAN) and completing the human anti-rabies treatment notification form [7]. This form collects identification data of the

notifying unit and agent, patient identification, characteristics of the injury, characteristics of the animal responsible for the attack (in cases involving dogs or cats), place of residence and treatment regimen. This form must be completed at each visit and sent to the appropriate department for processing. Although this form has a lot of information to aid in rabies surveillance, it does not provide a record of the circumstances of the attack, which is a key for the surveillance of cases in which the attacking animal is a vampire bat.

To help in the recognition of areas where populations are susceptible to vampire bat attacks, and assuming that the population of the municipality of São João da Ponta is representative of the crab fishermen riverside communities in Amazon RESEX area study attacks by vampire bats in mangrove areas of the Eastern Amazon, Pará, Brazil.

Methods

Ethical considerations

This study was conducted with the authorization of the National Research Ethics Commission (Portuguese acronym: CONEP) of the National Health Council (Portuguese acronym: CNS; registration number CAAE: 49593315.1.0000.0018), the Brazilian Institute of the Environment and the Chico Mendes Institute for Biodiversity Conservation (Portuguese acronym: ICMBio; registration number 50344-1). All study participants provided written informed consent.

Description of the study area

The Salgado micro-region in Pará comprises 11 municipalities that are home to five Marine Extractive Reserves (RESEX): São João da Ponta RESEX, where the

municipality under study is located, Maracanã RESEX, Mocapajuba RESEX in São Caetano de Odivelas, Mestre Lucindo RESEX in Marapanim, and Mãe Grande de Curuçá RESEX. The latter covers practically the entire territory of the municipality of Curuçá-PA. The climate is hot and humid, with a rainy season (from January to June) and a dry season (from July to December) [8][9]. The vegetation is composed of moderately preserved mangrove forest with species such as *Rhizophora mangle*, *Avicennia germinans* and *Laguncularia racemosa* [10](Figure 1).

Data collection and analysis

Information of individuals attacked by bats in São João da Ponta municipality between 2013 and 2015 were from Notifiable Diseases Information System (SINAN). This period was selected to prevent the interviewee from having difficulty remembering the episode and answering the questions.

These individuals were visited in their homes, and during these visits, the study objectives were explained, and the informed consent form was signed. Subsequently, a semi-structured questionnaire was used to collect data on the circumstances of the attack, such as location and time of the attack, type of attack, time elapsed between attack and anti-rabies treatment, frequency of attack, professional occupation of the attacked individuals and structure of the household or shelter where the attack occurred.

A non-probabilistic snowball sampling technique was used to obtain information on individuals who were attacked by bats during the same period but did not seek medical care. The seeds were the cases reported in SINAN during the study period [11]. Individuals reached by this method were also interviewed and all locations indicated by the interviewees as the area of the attack were georeferenced.

The descriptive statistical analysis of the data was performed using SPSS v. 20. A geographic database was created with the coordinates of the locations of the attacks. Using the cartographic databases of the Brazilian Institute of Geography and Statistics (IBGE), the spatial distribution of cases was analyzed in ArcGIS™ 10.1. The relationships between the individuals reached by snowball sampling was visualized using R.

Results

Between 2013 and 2015, five residents of the São João da Ponta municipality sought anti-rabies treatment after being attacked by vampire bats and therefore were registered in SINAN. These individuals identified another 141 individuals who were attacked by bats in the same period. Of those, 61 were interviewed and the others 80 individuals were disregarded in the analysis because they no longer lived in the community or were bitten for more than three years or refused to answer the questions and we could not confirm the aggression. Therefore, for each individual who sought care, there were at list 12.2 cases that were not reported. The communication network between these individuals is illustrated in Figure 2. We can observe that there are individuals who have more indications. They are individuals better known in the community. In contrast, there are few individuals who know more individuals who have been attacked by bats. To these, besides being more connected with the others in the network, they have high "PageRank" (centrality measures that reflects how much is indicated by the others of the network and those that indicate also have greater popularity). So, we can admit that people have neighbors that have more indications than them, corroborated by the article of Feld (1991) [12]. One of these more connected individuals was a Community Health Agent, a professional that visit households daily as a municipality action of preventive medicine.

Among the victims, 10 (15.1%) reported that the attacks occurred in their homes or in areas near the São João da Ponta Extractive Reserve, whereas the majority (84.9%) were attacked in the mangrove areas of the Mãe Grande de Curuçá Extractive Reserve in Curuçá municipality, a neighboring city (Figure 3).

Nineteen sites were identified as locations of attacks: 14 in the Curuçá RESEX, four in the São João da Ponta RESEX and one in the Marapanim RESEX. The Cuimiri and Pacamorema beaches which are both located in Curuçá, accounted for the highest proportions of bites, 22.7% and 10.6% respectively. Figure 4 shows the route traveled by the crab fishermen between the locations of the attack and their city of origin, where they would seek anti-rabies treatment.

The attacked individuals were mostly male (92.4%) and adult (69.6%) and had less than 4 years formal school education (54.4%). Most lived in the rural area of the city (92.4%) and were crab fishermen (79.3%) (Table 1).

Table 1- Characteristics of individuals bitten by bats in the São João da Ponta municipality, Pará, Brazil.

VARIABLES	RESPONSE IN DESCENDING ORDER OF FREQUENCY (%)		
	1 ^a	2 ^a	3 ^a
CHARACTERISTICS OF GRAZING			
Ocupation	Crabfishing (79.3)	Student (12.1)	Farmer (6.1)
Sexo	Male (92.4)	Female (7.6)	-
Age	20 to 50 (69.9)	10 to 20 (28.8)	less than 10 (1.5)
CHARACTERISTICS OF THE PLACE OF AGGRESSION			
Type of home	Tent or straw taps (88.8)	Other (without wall) (10.5)	Masonry (with wall and ceiling) (7.5)
Are there animals at the scene of the aggression?	Wild (75.8)	Domestic (48.5)	-

How many people were at the scene of the aggression?	More than five (63.3)	Two to five (24.1)	One (4.5)
Was there any light source in place?	Yes (87.9)	No (12.1)	-
There is fresh water on site?	Yes (59.1)	No (40.9)	-
Where were you bitten??	Beach (56.1)	Mangrove (30.3)	“Tiso” (3.0)
In what circumstances have you been bitten?	Fishing (83.1)	At leisure (6.2)	Others (6.2)
At what time of year?	Seca (54.5)	Chuvosa (45.5)	-
FEATURES OF THE INJURY			
How many times have you been bitten in the last 3 years?	One (33.3)	More than five (25.8)	Two (21.2)
When was the last aggression?	1 to 3 anos (37.9)	2 to 6 months (34.8)	7 months to 1 any (16.7)
What was the amount of bites in the last attack?	More than tree (45.5)	One (33.3)	Two (21.2)
Type of Injury	Single (71.2)	Multiple (28.8)	-
Anatomical location of the wound	Lower member (80.8)	Head (12.8)	Upper limb (5.13)
Did you look for the health unit?	No (86.4)	Yes (13.6)	-
Do you believe that the bat bite causes any disease?	Yes (80.3)	No (19.7)	-
Do you know what rabies is?	No (94.5)	yes (5.5)	-

191

192 The interviewees reported that during crab fishing, they gather for days in
193 makeshift dwellings without walls and covered by tarp or straw (88.8%) (Figure 4). In
194 most cases, these shacks were set up in drier areas within the mangrove (30.3%) or on
195 beaches (56.1%), less than five kilometers from the forest (73.3%) and freshwater bodies
196 (45.4%). In addition, many fishermen reported that bats sheltered in a tree typical of the
197 mangrove area, commonly known as black mangrove (*Avicennia germinans*) (80.3%).
198 Most of the interviewees reported the presence of domestic animals like dogs (56.5%),
199 domestic birds (23.9%) and bovine (10.9%) where attacks occurred. Wild animals were

also observed (75.8%), with raccoon (*Procyon cancrivorus*) (26.1%), scarlet ibis (*Eudocimus ruber*), great egret (*Ardea alba*) (23.5%) and monkey (19.1%) being the most commonly reported species. Some interviewees had found lesions suggestive of bat attacks on dogs (42.9%) and domestic birds (10.6%).

On the nights that the attacks occurred, most of the individuals (63.6%) were in a group with more than five fishermen in the same shack (ranging from 2 to 12 people), and 33.3% reported that others were also attacked the same night. The shelters typically had a light source (87.9%) such as a lantern, a bonfire or an oil lamp that was usually extinguished by strong beach winds throughout the night.

When questioned about the frequency of attacks, most reported having been attacked more than four times. The number of attacks by bats during the life ranged from 1 to 18. In most cases (34.8%), the most recent episodes had occurred between 2 and 6 months prior to the interview, more frequently in the drier season (54.4%). In general, the injuries were single bites (71.2%) and were located on the lower limbs (80.8%). Among those who sought care (i.e., the seeds, 13.6%), all completed the prophylactic regimen with 5 doses of vaccine and serum.

Questions addressing respondents' perceptions regarding the risks associated with bat bites and their consequences revealed that most individuals did not use any type of personal protection equipment against bat attacks (63.6%). Those who claimed to use some type of protection used mosquito netting, "sapato de mangua" (resistant fabric wrapped around their feet), bonfires, or fishing nets around the shack as a physical barrier. Most (80.3%) were unaware of the risk of rabies transmission through a bat bite or even unaware of rabies (94.5%). Of those attacked who did not seek medical care, 66.7% reported "not caring about what happened", 13.6% did not know how to respond, and 7.6% mentioned the distance from the healthcare unit.

225

226 **Discussion**

227 This is the first study to quantify the underreporting of bat attacks in humans in
228 the Amazon using the snowball sampling method, which capitalized on relationships
229 between individuals to reach the target population. Network analysis was useful in
230 identifying the people most commonly mentioned by the community in relation to bat
231 attacks, and it can be applied to identify key individuals who can be the focus of more
232 intense health surveillance activities. People identified by the network can be great allies
233 in implementing interventions to change habits and attitudes that may hinder prevention.
234 Indeed, it can be useful to optimize Communities Health Agents job, when bitten people
235 active search is necessary.

236 Human behavior, in particular the lack of knowledge in this study population
237 regarding the predictable consequences of vampire bat bites hinders the use of SINAN
238 quantitative reports as a reliable data source to establish public health policies for rabies
239 transmission in these areas. This is because the number of people attacked by bats in this
240 region is much higher than that recorded in the system. Moreover, the reports do not
241 include the locations of attacks or the bat species involved. Although this characterization
242 was carried in municipality of Pará state, São João da Ponta, we believe that it is reflective
243 of the situation in other municipalities of the Salgado microrregion, which have similar
244 geographic and cultural characteristics.

245 Strategies for rabies prevention in vulnerable populations in Amazon regions have
246 been previously reviewed in the literature [1,3,4]. In fact, it has already been proposed
247 that the SINAN form be changed to include fields related to dog and cat behavior to
248 support animal rabies surveillance and prophylactic measures [13]. Although these
249 changes have not yet been incorporated, this study demonstrates the need to adapt the

form to include bat attacks, given that animal attacks relevant to rabies transmission in the Salgado microregion and the whole Amazon typically involve different species than in other regions [14]. The fact that these people are generally not bitten in their homes but, rather, in other circumstances demonstrates that the inclusion in the SINAN form of a field to report the location of attacks is necessary to better prevent vampire bat bites.

This study also shows for the first time that living conditions in the mangrove are important factors in bat attacks in the Amazon. Respondents reported that in these areas, attacks by bats have been occurring for a long time and have never resulted in death. In fact, there are no reports of human deaths from the rabies virus in the study region, and there were no reported cases of neurological syndromes in livestock between 2004 and 2013 [23]. However, there are no studies on the circulation of rabies virus in this micro-region and the possibility of inexistent or inadequate communication between healthcare and agricultural services cannot be ruled out [16]. Therefore, based on the evidence of human attacks reported here and the underreporting of bites, the study region cannot be considered as an area of controlled rabies transmission. When in doubt about whether the virus is circulating, it would be prudent to define a policy of immunization and health education for this population, given the history of rabies in Pará and the prevalent risk factors for rabies [24].

Travassos da Rosa et al. (2006) stated that the proximity of animal husbandry to the dwellings was a predisposing factor for the attacks on humans that occurred in 2004/2005 in Pará state. However, these characteristics were not observed in the present study. For the population, RESEX's are associated to the improvement of the quality of life of people that use forest resource for subsistence, being prohibited the creation of livestock in these areas, because it contradicts its rules of sustainable use [22]. These units were created for the protection of mangrove ecosystems, which constitute an area under

strong anthropic pressure, with increased exploitation of their natural resources [21]. Informal reports reveal that attacks on humans increased when bovine were removed from such areas. It is also possible that hunting activities by the native population are promoting a reduction in the supply of natural prey for bats, such as the crab-eating raccoon (*Procyon carnivores*) and nonhuman primates. These animals are targeted and slaughtered by fishermen because they feed on the crabs caught in their traps, thereby reducing their daily catch. Another relevant factor is the fact that the vegetation of the area is predominantly composed of natural fields and mangroves, where little food is available for the vampire bat [20].

The age group most affected in this study differs from that reported in previous studies. Whereas most of the victims in this study were male adults, Travassos da Rosa et al. (2006) reported children as the main victims of the rabies virus transmitted by vampire bats in the 2004-2005 period. In contrast to our study, the victims in the Travassos da Rosa study were bitten in their own residences, which commonly house families with individuals of various age groups. In this context, bats seem to prefer children. In the present study, the targets of attacks were clusters of adults who gathered for work, often associated with alcohol consumption. Individuals become passive and docile victims after falling asleep following intense work activity during the day. This is similar to the behavior of herd animals, which are favored targets of vampire bats, as they seek abundant prey and tend to remain in the same place for prolonged periods to feed [15,16].

This study also demonstrates for the first time the occupational nature of the bat attack problem in the region. The vulnerability of crab fishermen is closely tied to their work activity, as they are bitten primarily while working in the mangroves. Usually these individuals dwell in shacks (huts) built on upland soil inside the mangrove or on the

beaches. These dwellings are completely open, with just a tarp or straw roof, and are used for sleeping during the work period. Even students who were bitten reported that they were accompanying their parents to help with crab fishing during the school vacation.

Residents of the Deolândia, Guarajuba and Porto Grande communities accounted for most respondents. These communities are the main centers for swamp ghost crab fishing in the region, and ghost crabs are the main natural resource exploited as a source of income by these communities. Therefore, contact between humans and bats is inevitable, as crabs are the only source of income for many of these fishermen. However, the lack of individual protection equipment certainly contributes to the increased number of bat attacks in this population [2,4].

Conclusions

Using a single city as a model for investigating bat attacks in the Amazon, this study showed that, for each person who seeks anti-rabies treatment after being attacked by bats in the region, there are 12.2 people who do not seek care and can be easily reached by surveillance agencies through snowball sampling. We also found that the population most at risk is that of crab fishermen, suggesting that bat attacks are an occupational hazard. These results can help health surveillance agencies to establish measures for the prevention of human rabies in these individuals. They can also help minimize costs and increase the efficiency of public health measures, thus avoiding the emergence of rabies in the Salgado micro-region of Pará state.

Acknowledgments

We thank PAPQ/PROPESP-UFPA for financial support; the Secretaria Municipal Saúde de São João da Ponta for sharing the database; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the research grants and the researchers JG Barreto and CCG Moraes for their valuable contributions.

References

1. Kessels JA, Recuenco S, Navarro-Vela AM, Deray R, Vigilato M, Ertl H, et al. Pré-exposure rabies prophylaxis: a systematic review. Bulletin of the World Health Organization 2017; 95 (3): 210-219c.
2. Mendes WDS, Silva AAM, Neiva RF, Costa NM, Assis MS, Vidigal PMO, et al. An outbreak of bat-transmitted human rabies in a village in the Brazilian Amazon. Rev Saude Publica 2009; 43: 1075–1077. doi:10.1590/S0034-89102009005000073
3. Stoner-Duncan B, Streicker DG, Tedeschi CM Vampire Bats and Rabies: Toward an Ecological Solution to a Public Health Problem. PLoS Negl Trop Dis 2014; 8: 8–12. doi:10.1371/journal.pntd.0002867
4. Berger F, Desplanches N, Baillargeaux S, Joubert M, Miller M, Ribadeau-Dumas F, et al. Rabies Risk: Difficulties Encountered during Management of Grouped Cases of Bat Bites in 2 Isolated Villages in French Guiana. PLoS Negl Trop Dis 2013; 7(6): e2258. doi:10.1371/journal.pntd.0002258
5. Romero-Sandoval N, Escobar N, Utzet M, Feijoo-Cid M, Martin M. Sylvatic rabies and the perception of vampire bat activity in communities in the Ecuadorian Amazon. Cad Saude Publica 2014; 30: 669–674. doi:10.1590/0102-311X00070413

- 347 6. Glaser M, Diele K. Asymmetric outcomes: Assessing central aspects of the
348 biological, economic and social sustainability of a mangrove crab fishery, *Ucides*
349 *cordatus* (Ocypodidae), in North Brazil. *Ecol Econ* 2004; 49: 361–373.
350 doi:10.1016/j.ecolecon.2004.01.017
- 351 7. Brasil. Ministério da Saúde (2014). Portaria nº 1271 de 12 de dezembro de 2014.
352 Lista Nacional de Doenças de Notificação Compulsória. Brasília. Distrito Federal, Brasil.
- 353 8. INMET. Instituto Nacional de Meteorologia. Disponível em: www.inmet.gov.br.
354 Acesso em: 23/01/2018.
- 355 9. Instituto Brasileiro de Geografia e Estatística. Censo 2010. Disponível
356 em: www.ibge.gov.br. Acesso em: 03/02/2018.
- 357 10. Carneiro DB, Leitão CM, Sá LB, Menezes, MP. Plantas Nativas Úteis na Vila dos
358 Pescadores da Reserva Extrativista. *Acta Bot* 2010; 24: 1027–1033. doi:10.1590/S0102-
359 33062010000400017
- 360 11. Vinuto J. A amostragem em bola de neve na pesquisa qualitativa: um debate em
361 aberto. *Temáticas* 2014; 22 (44): 203-220.
- 362 12. Feld, SL. Why Your Friends Have More Friends Than You Do. *The American Journal*
363 *of Sociology* 1991; 96 (6): 1464-1477. doi: 10.1086/229693
- 364 13. Frias DFR, Carvalho AAB, Nunes JOR. Proposta de nova metodologia de apoio para
365 indicação racional de profilaxia antirrábica humana pós-exposição. *Arq. Cienc. Saúde*
366 *UNIPAR, Umuarama* 2016; v. 20, n. 1: 9-18. doi: 10.25110/arqsaude.v20i1.2016.4955
- 367 14. De Paula NS, Saraiva EA, Araújo IM, Nascimento KKG, Xavier DA, Santos KS,
368 et al. Characterization of rabies post-exposure prophylaxis in a region of the eastern

- 369 Amazon, state of Pará, Brazil, between 2000 and 2014. Zoonoses Public Health 2018;
370 65(4),395-403. doi:10.1111/zph.12444
- 371 15. Ito F, Bernard E, Torres RA What is for dinner? First report of human blood in
372 the diet of the hairy-legged vampire bat *Diphylla ecaudata*. Acta Chiropt 2016; 18: 509–
373 515. doi:10.3161/15081109ACC2016.18.2.017
- 374 16. Johnson N, Aréchiga-Ceballos N, Aguilar-Setien A Vampire bat rabies: Ecology,
375 epidemiology and control. Viruses 2014; 6: 1911–1928. doi:10.3390/v6051911
- 376 17. Taddei VA (1991). Distribuição do morcego vampiro, *Desmodus rotundus*
377 (*Chiroptera*, *Phyllostomidae*) no estado de São Paulo e a raiva dos animais domésticos.
378 Impresso Espec CATI, 107 páginas.
- 379 18. Mialhe P. Preferential prey selection by *Desmodus rotundus* (E. Geoffroy, 1810,
380 Chiroptera, Phyllostomidae) feeding on domestic herbivores in the municipality of São
381 Pedro - SP. Brazilian J Biol 2014; 74: 579–584. doi:10.1590/bjb.2014.0086
- 382 19. Magalhães HF de, Costa Neto EM, Schiavetti A. Saberes pesqueiros relacionados
383 à coleta de siris e caranguejos (Decapoda: Brachyura) no município de Conde, Estado da
384 Bahia. Biota Neotrop 2011; 11: 45–54. doi:10.1590/S1676-06032011000200005
- 385 20. Giarrizzo T and Krumme U. Temporal patterns in the occurrence of selected
386 tropical fishes in mangrove creeks: implications for the fisheries management in north
387 Brazil. Braz Arch Biol Technol 2009; 52 (3): 679-688
- 388 21. Ulrich Saint-Paul HS (2010). Mangrove Dynamics and Management in North
389 Brazil. Media SS& B, editor.
- 390 22. Brasil. Lei no 9.985, de 18 de julho de 2000. Regulamenta o art. 225, § 1o, incisos
391 I, II, III e VII da Constituição Federal, institui o Sistema Nacional de Unidades de

Conservação da Natureza e dá outras providências. Presidência da República. Casa Civil.

Subchefia para Assuntos Jurídicos. Diário Oficial [da] União, Brasília, DF, 19 jul. 2000.

23. Monteiro, FD. Raiva em animais de produção no estado do Pará, Brasil: um estudo descritivo, 2004-2013 (2014). Dissertação (Mestrado). Programa de pós-graduação em Saúde Animal da Amazônia. Universidade Federal do Pará, Castanhal-PA.

24. Travassos da Rosa ES, Kotait I; Barbosa TFS; Carrier ML, Brandão PE, Pinheiro AS, et al. Bat-transmitted human rabies outbreaks, Brazilian Amazon. Emergent Infect Dis 2006; 12: 1197–1202.

Figures Captions

Fig 1. Study area. Distribution of extractive reserves in the Salgado microregion, Pará, Brazil. The map was created specifically for this manuscript and was generated by ArcGis 10.1 (ESRI) based in Brazilian Institute of Geography and Statistics Database (<https://downloads.ibge.gov.br/>)

Fig 2. Network of relationships between the residents of the São João da Ponta municipality (Pará, Brazil) who were attacked by bats between 2013 and 2015. The nodes

represent the individuals who were attacked. The diameter of the nodes is proportional to the number of times the person was mentioned.

Fig 3. Spatial distribution of locations where residents of the Curuçá municipality were attacked by bats. The lines indicate the route traveled by the crab fishermen from their municipality of origin to the place where they camped for crab fishing. The map was created specifically for this manuscript and was generated by ArcGis 10.1 (ESRI) based in Brazilian Institute of Geography and Statistic Database (<https://downloads.ibge.gov.br/>)

Fig 4. Housing conditions of the individuals who were attacked. A) shack without walls and with a straw roof; B) Shack without walls and with a tarp roof; C and D) Siriubeira.

Figure

Figure

A

bioRxiv preprint doi: <https://doi.org/10.1101/590083>; this version posted March 26, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

B

Figure

Figure