Induction of p73, Δ 133p53, Δ 160p53, pAKT leads to neuroprotection via DNA repair by 5-LOX inhibition

Shashank Shekhar and Sharmistha Dey

Department of Biophysics All India Institute of Medical Sciences, New Delhi, India

Running title: p53 isoform and pAKT in neuroprotection

To whom correspondence should be addressed:

*Dr. Sharmistha Dey

Department of Biophysics

All India Institute of Medical Sciences

Ansari Nagar,

New Delhi - 110029, INDIA,

Tel:+91 - 11 - 2654 6435,

Fax:+91 - 11 - 2658 8663,

E-mail: sdey@aiims.edu

Induction of p73, Δ 133p53, Δ 160p53, pAKT leads to neuroprotection via DNA repair by 5-LOX inhibition

Shashank Shekhar and Sharmistha Dey

Department of Biophysics All India Institute of Medical Sciences, New Delhi, India

Abstract

Lipoxygenase-5 (5-LOX), protein is involved in the pathologic phenotype of AD which includes amyloid-plaque and tau hyperphosphorylation. This study aims to identify the mechanistic role in neuroprotection by peptide YWCS, the 5-LOX inhibitor in neurotoxic SH-SY5Y cell line developed by the treatment of $A\beta_{25-35}$. The cells were treated with $A\beta_{25-35}$ and with different doses of YWCS. The effect on cell survival pathways were determined by western blot using polyclonal anti body of p53, anti-Akt and anti-phosphorylated-Akt. Immunoprecipitation and mass spectroscopic studies were done to identify the altered proteins. Over expression of phosphorylated-Akt and 3 bands of p53 isoforms were observed which correspond to p73, Δ 133p53 and Δ 160p53 in the cells treated only with 80 μ M of YWCS compare to untreated cells. However, no alteration of total p53 and Akt were observed. The results exposed the novel mechanistic pathway of neuroprotection by 5-LOX inhibition, which is likely to be mediated by DNA DSB repair through p53 isoforms and PI3K/Akt pathway. Our finding has opened a new window in the therapeutic approach for the prevention of AD.

Key word: Alzheimer's disease, p73, Δ 133p53, Δ 160p53, DNA DSB repair

Introduction

Alzheimer's disease (AD) is the most common cause of dementia in the elderly. Due to increasing longevity and the lack of therapy, AD has become not only a major health problem but also imposes substantial social and economic burden worldwide[1]. The inflammatory process plays a key role in neurodegenerative disorder. The inflammatory molecule, 5-lipoxygenase (5-LOX), protein is involved in the pathologic phenotype of AD which includes A β amyloid deposition and tau hyperphosphorylation. Recent research from our lab has proposed the role of 5-LOX peptide inhibitor as neuroprotective molecule for AD[2], provides rescue to neuronal cells from amyloid induced proteotoxic stress/neurotoxicity. The 5-LOX peptide inhibitor reduced γ -secretase expression as well as tau hyperphosphorylation at threonine 181. The mechanistic detail of the 5-LOX inhibitor is still to be elaborated to understand how this inhibitor is providing rescue to neuronal cells from amyloid beta induced neurotoxicity. Some previous studies have shown that 5-LOX inhibitor accelerates the phosphorylation of Akt and thereby provides rescue to cells[3]. It is also reported that 5-LOX regulate p53 activity[4] and p53 isoform Δ 113p53/ Δ 133p53 promotes DNA double-strand break repair to protect cell from death and senescence in response to DNA damage[5]. In a recent finding it has been reported that the p53 isoform; p73 plays an important role in the DNA repair pathway in coordination with another isoform of low molecular weight; Δ 133p53[6]. The Δ 160p53 is a recently identified isoform of p53 and have role in senescence and DNA repair but the detailed function of it is still illusive[7]. The neuroprotective role for p53 was reported in an in-vivo model of tau-mediated neurodegeneration relevant to Alzheimer's disease and related disorders. All these findings are illustrated the importance of Akt, p53, p73, Δ 133p53 and Δ 160p53 in cell survival pathway. Their role in AD pathogenesis is not yet studied. Here, for the first time we have identified the

involvement of p73, $\Delta 133p53$ and $\Delta 160p53$ in amyloid beta induced neurotoxicity and the novel pathway of neuroprotection by YWCS peptide inhibitor of 5-LOX via p73, $\Delta 133p53$ and $\Delta 160p53$.

Materials and Methods

Solid phase Peptide Synthesis and processing

The peptides were synthesized by solid phase peptide synthesizer PS3 (Protein technology, USA) using Fmoc and Wang resin chemistry. The purity of the peptide was verified by analytical RP-HPLC as described earlier[8]. Briefly, the solvent used for the synthesis was dimethylformamide (DMF) and 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) was used as an activator. Fmoc was deprotected by 20% piperidine and wang resin was cleaved by Trifluoroacetic acid (TFA). The peptides were precipitated from dry ether.

Preparation of aggregated A β peptide

The A β_{25-35} peptide was dissolved in nuclease free water and then incubated for 5 days at 37°C and constant shaking in incubator shaker. The aggregation of peptide was confirmed by Thioflavin T (ThT) assay and the scanning electron microscopy.

Cells and treatment

SH-SY5Y cells were obtained from NCCS, Pune, India and maintained in Ham's F12 Nutrient media (Gibco) supplemented with 10% (v/v) fetal bovine serum and 1% antibiotic-anti-mycotic solution (Gibco). The cells used in the experiment were of passage number 30-33. Before starting the experiments, cells were authenticated by STR profiling (DNA forensic laboratory

Private limited, India). The cells were maintained at 37°C and 5% CO₂ under humidified condition. Cells were grown as monolayer.

SH-SY5Y cells were plated at a density of 1×10^6 cells per T25 flask and kept in CO₂ incubator at 37°C overnight to adhere. Cells were differentiated by the treated of 20 µM retinoic acid in fresh complete media [9,10] for 7 days where media was replaced every 2nd day. On 7th day, cells were treated with different concentrations of YWCS (20, 40 and 80µM) and Aβ₂₅₋₃₅ (20 µM) peptide for 72 h simultaneously. Aβ₂₅₋₃₅ and YWCS peptide were synthesized by solid phase peptide synthesis as described previously [8].

Western blot:

Cells were harvested and lysate was prepared in RIPA buffer (10mM Tris- HCl pH-8.0, 140mM NaCl, 1mM EDTA pH 8.0, 0.1% sodium deoxycholate, 1% Triton X100, 0.1% SDS, 0.1mM ethylene glycol tetra-acetic acid pH-8.0, 1mM Protease inhibitor, 1mM phenylmethanesulfonyl fluoride). The expression of p53, Akt and p-Akt were determined by western blotting as described previously [11]. Briefly, 30µg of total protein was separated on 12% SDS gel and protein was transferred on PVDF membrane (md, India). Membrane was blocked in 5%NFM and then incubated with primary antibodies of following proteins: p53 (1:500, Santa Cruz), Akt p-Akt, (1:1000, abchem). Membrane were developed with chemiluminescent substrate west Pico (Thermo scientific).

Immunoprecipitation

After treatment with YWCS cells were lysed in RIPA buffer as described above. Total protein (100µg) was incubated overnight at 4°C with 1µg of anti-human p53 polyclonal antibody in a

reaction buffer of 0.1 % BSA in PBS. The protein agarose beads (30μL pre washed with 0.1 % BSA) were added in the reaction and incubated for 4 h at 4°C. After 4 h beads were washed 3 times with wash buffer (10mM Tris-HCl pH 7.5, 1mM EDTA, 1mM EGTA, 150mM NaCl, 0.2 mM sodium orthovanadate, 1 mM PMSF). Beads were eluted with 0.2 M glycine pH 2.6 in 1:1 ratio. After elution all the fractions were pulled and neutralized by adding Tris-HCl pH 8.0.

Mass Spectroscopy

Immunoprecipitated samples were separated on 10 % SDS-PAGE. IgG was used as negative control. Then bands were carefully cut and sent to Sandor Life Sciences, India for the mass spectroscopy.

Results

Preparation of aggregated Aβ peptide

Thioflavin-T is a dye specific for detection of fibrillation of proteins. It has an excitation wavelength at 440nm and the emission wavelength at 480nm. ThT assay had shown high intensity after 4 days for its aggregation status (Figure 1 A). The results were further confirmed by scanning electron microscopy for the aggregation of peptide after incubation of 4 days (Figure 1 B and C).

Western blot for Akt and p-Akt

The western blot for Akt and p-Akt were carried out to check the effect of YWCS peptide on their expression and phosphorylation status. Our results suggest that the inhibition of 5-LOX with YWCS peptide had no effect on the Akt expression, while increased the phosphorylated Akt (Figure 2). This indicated that, YWCS peptide induced the phosphorylation of Akt protein. Some

recent literature reported that the p-Akt has inhibitory effect on apoptosis. Our study first time provides an evidence for phosphorylation of Akt by inhibiting 5-LOX with peptide inhibitor and thereby prevents neurotoxicity.

Western blot for p53

The western blot for p53 was performed with poly clonal antibody against human p53 since it has several isoforms and play diverse role in cell survival regulation. To our surprise we observed that there was no change in the expression of p53 protein upon treatment with the YWCS but 3 other bands were observed in the cells while treated with 80μM of YWCS. These bands were not present in any other treatment groups (Figure 3).

Identification of proteins which were expressed by the treatment of YWCS

Immunoprecipitation was performed followed by mass spectroscopy. Immunoblot of pull down showed the enrichment of the target proteins (Figure 4A). After immunoprecipitation, pull down was separated on SDS-PAGE and IgG was used as control (Figure 4B). Bands were then sent for mass spectroscopy. The mass result confirmed our hypothesis that the 5-LOX inhibition by peptide induced the expression of p53 isoforms p73, Δ133p53 and Δ160p53 (Figure 5 A, B and C). Our study first time demonstrated the role of p73, Δ133p53 and Δ160p53 in neuroprotection. The expression of these isoforms was induced in the 80μM peptide treated cells. These isoforms of p53 activates DNA double strand break repair. This study showed 5-LOX inhibitor activated these isoforms and prevented neurotoxicity via DNA double strand break repair pathway. This finding opens a new window of mechanistic pathway for 5-LOX inhibition and neuroprotection under oxidative stress and DNA damage.

Discussion:

AD is age associated disease, progressive with extracellular amyloid-beta ($A\beta$) deposits intracellular aggregates of hyperphosphorylated tau and neurofibrillary tangles. Ageing process normally arises DNA damage and in AD excessive DNA damage occurs due the $A\beta$ -induced oxidative stress. In our previous work, we found that inhibition of 5-LOX prevented the $A\beta$ induced neurotoxic effect in SH-SY5Y cells and downregulated the expression of γ -secretase components [2]. 5-LOX is a direct p53 target gene in humans. The p53 protein involved in DNA repair, down regulated in AD[12]. In the mammalian cell, p53 was found to be modulate the DNA repair and stimulate both removal of damaged bases and nucleotide reinsertion [13]. To ensure the effect on p53 through 5-LOX inhibition under neurotoxic condition in cells, we performed western blot with treated SH-SY5Y cells by 3 different concentrations (20, 40 and 80 μ M) of YWCS using p53 polyclonal antibody against human p53. No alteration of p53 protein was observed, but 4 different bands appeared only at high concentration (80 μ M) of peptide. This distinction leads us to search the protein bands by immunoprecipitation and mass spectroscopy, which revealed three isoforms of p53 corresponds to p73, Δ 133p53 and Δ 160p53. While going through literature we came to know that these isoforms play vital and diverse role in cell survival regulation. The isoforms of p53 are found to be pro-survival factor for DNA damage stress and their expression prevents apoptosis and promotes DNA-DSB repair. Some more recent studies have proposed the role of p53 isoforms in the cell survival such as Δ 133p53/ Δ 113p53 by repairing DNA damage in cells during low level of oxidative stress or reactive oxygen species (ROS). However, the full length p53 inhibits DNA-DSB repair. The present work found high expression of p53 isoforms in the $A\beta$ induced neurotoxic SH-SY5Y

cells treated by only at higher concentration of peptide YWCS in compare to untreated cells and thereby prevented neurotoxicity. However, no alteration of p53 was observed in treated cells.

The higher expression level of $\Delta 160p53$ and p73 isoforms was observed in treated SH-SY5Y cells with peptide (80 μ M YWCS). The internal promoter originates the $\Delta 133p53$ mRNA codes both the isoforms, $\Delta 133p53$ and $\Delta 160 p53$. Though $\Delta 160p53$ is the conserved isoform, very little information is available about it so far. Another study reported both p73 and $\Delta 133p53$ synergistically promote the expression of DNA repairing genes RAD51, LIG4 and RAD52 by homologous recombination (HR), non-homologous end joining (NHEJ) and single-strand annealing (SSA) and thereby promote DNA DSB repairing, which supports our study.

Inhibition of 5-LOX by YWCS peptide inhibitor was also upregulated the phosphorylation of Akt in SH-SY5Y cells. There is evidence of neuroprotective effects by stimulating PI3K/Akt signaling plays a pivotal role in neuronal survival [14]. Previous studies suggested that PI3K/Akt signaling was downregulated in the AD brain [15] and activation of this pathway showed to prevent A β -induced neuronal neurotoxicity. Akt has well established role in cell cycle control and in DNA repair by check point activation in late G2 phase.

This study first time reported the rescue of A β induced neurotoxicity by the treatment of 5-LOX inhibitor by activating the expression of $\Delta 160p53$, p73 and $\Delta 133p53$ in SH-SY5Y cells. The results reveal that the mechanistic pathway for the prevention of neurotoxic effect by targeting the inhibition of 5-LOX may proceed by DNA DSB repair, by stimulating p53 isoforms and PI3K/Akt pathway. Thus repairing the DNA damage ameliorates the AD pathologies. The results exposed the novel mechanistic pathway of neuroprotection by 5-LOX inhibition mediated by

DNA DSB repair through p53 isoforms and PI3K/Akt signaling pathway. Our finding has opened a new window in the therapeutic approach for the prevention of AD.

Acknowledgments: Authors acknowledge intramural grant, Research Section, All India Institute of Medical Sciences (A-422) for providing funds for the consumable items and CSIR, Government of India for the fellowship of Shashank Shekhar.

Conflict of Interest: Authors show no conflict of interest.

Role of the authors: SS performed the entire experiment and study concept. SD was responsible for the study concept, design and wrote the paper.

References:

1. Tarawneh R, Holtzman DM (2012) The clinical problem of symptomatic Alzheimer disease and mild cognitive impairment. *Cold Spring Harb Perspect Med* **2**: a006148.
2. Shekhar S, Yadav SK, Rai N, Kumar R, Yadav Y, Tripathi M, Dey AB, Dey S (2018) 5-LOX in Alzheimer's Disease: Potential Serum Marker and In Vitro Evidences for Rescue of Neurotoxicity by Its Inhibitor YWCS. *Mol Neurobiol* **55**: 2754–2762.
3. Tu X-K, Zhang H-B, Shi S-S, Liang R-S, Wang C-H, Chen C-M, Yang W-Z (2016) 5-LOX Inhibitor Zileuton Reduces Inflammatory Reaction and Ischemic Brain Damage Through the Activation of PI3K/Akt Signaling Pathway. *Neurochem Res* **41**: 2779–2787.
4. Catalano A, Rodilossi S, Caprari P, Coppola V, Procopio A (2005) 5-Lipoxygenase regulates senescence-like growth arrest by promoting ROS-dependent p53 activation. *EMBO J* **24**: 170–179.

5. Gong L, Gong H, Pan X, Chang C, Ou Z, Ye S, Yin L, Yang L, Tao T, Zhang Z, et al. (2015) p53 isoform $\Delta 113p53/\Delta 133p53$ promotes DNA double-strand break repair to protect cell from death and senescence in response to DNA damage. *Cell Res* **25**: 351–369.
6. Gong H, Zhang Y, Jiang K, Ye S, Chen S, Zhang Q, Peng J, Chen J (2018) p73 coordinates with $\Delta 133p53$ to promote DNA double-strand break repair. *Cell Death Differ* **25**: 1063–1079.
7. Marcel V, Perrier S, Aoubala M, Ageorges S, Groves MJ, Diot A, Fernandes K, Tauro S, Bourdon J-C (2010) $\Delta 160p53$ is a novel N-terminal p53 isoform encoded by $\Delta 133p53$ transcript. *FEBS Lett* **584**: 4463–4468.
8. Singh AK, Singh R, Naz F, Chauhan SS, Dinda A, Shukla AA, Gill K, Kapoor V, Dey S (2012) Structure based design and synthesis of peptide inhibitor of human LOX-12: in vitro and in vivo analysis of a novel therapeutic agent for breast cancer. *PLoS ONE* **7**: e32521.
9. Pählman S, Ruusala AI, Abrahamsson L, Mattsson ME, Esscher T (1984) Retinoic acid-induced differentiation of cultured human neuroblastoma cells: a comparison with phorbol-ester-induced differentiation. *Cell Differ* **14**: 135–144.
10. König G, Masters CL, Beyreuther K (1990) Retinoic acid induced differentiated neuroblastoma cells show increased expression of the beta A4 amyloid gene of Alzheimer's disease and an altered splicing pattern. *FEBS Lett* **269**: 305–310.
11. Shekhar S, Yadav Y, Singh AP, Pradhan R, Desai GR, Dey AB, Dey S (2018) Neuroprotection by ethanolic extract of *Syzygium aromaticum* in Alzheimer's disease like pathology via maintaining oxidative balance through SIRT1 pathway. *Exp Gerontol* **110**: 277–283.

12. Lovell MA, Gabbita SP, Markesbery WR (1999) Increased DNA oxidation and decreased levels of repair products in Alzheimer's disease ventricular CSF. *J Neurochem* **72**: 771–776.
13. Boesch P, Weber-Lotfi F, Ibrahim N, Tarasenko V, Cosset A, Paulus F, Lightowlers RN, Dietrich A (2011) DNA repair in organelles: Pathways, organization, regulation, relevance in disease and aging. *Biochim Biophys Acta* **1813**: 186–200.
14. Liu Y, Yang X, Lei Q, Li Z, Hu J, Wen X, Wang H, Liu Z (2015) PEG-PEI/siROCK2 Protects Against A β 42-Induced Neurotoxicity in Primary Neuron Cells for Alzheimer Disease. *Cell Mol Neurobiol* **35**: 841–848.
15. Liu Y, Liu F, Grundke-Iqbal I, Iqbal K, Gong C-X (2011) Deficient brain insulin signalling pathway in Alzheimer's disease and diabetes. *J Pathol* **225**: 54–62.

Figure legends:

Figure 1. Aggregation of A β_{25-35} peptide. (A) There was drastic change in the florescence intensity after 4 days of aggregation. (B) Electron microscopic image of A β_{25-35} peptide prior to aggregation. (C) Electron microscopic image of A β_{25-35} peptide after aggregation.

Figure 2. Western blot of expression of Akt and the phosphorylation of Akt at ser473 after treatment by YWCS in SH-SY5Y cells. There was no change in the level of Akt but there was phosphorylation of Akt only in 80 μ M of YWCS treated cells.

Figure 3. Western blot for p53. Expression of p73 and low molecular weight isoforms of p53 i.e. Δ 133p53 and Δ 160p53 in 80 μ M of YWCS treated cells. Beta actin was used as the loading control.

Figure 4. Immunoprecipitation (IP) of p53. (A) immunoblot of IP samples. (B) SDS page of Immunoprecipitate. IgG was used as negative control.

Figure 5. Mass spectroscopy of bands form IP. (A) Identified as p73, (B) identified as Δ 133p53 and (C) identified as Δ 160p53 isoforms of p53.

A **B** **C**

A**IP p53** **B****IP p53****IgG control**

(A) p73

(B) Δ133p53

(C) Δ160p53

Mascot Search Results

User : Samovar
 Email : samovar@pandor.co.in
 Search title :
 MS data file : peaklist.xml
 Database : P53_Human_Human_20180602 (678 sequences; 280684 residues)
 Timestamp : 2 Jun 2018 at 06:44:42 GMT
 Top Score : 76 for [sp00209.1375_0004](#), Reckase: full-length protein p73; AltName: full-length transcription factor; AltName: full-length-related p73

Mascot Score Histogram

Protein score is $-10 \cdot \log(P)$, where P is the probability that the observed match is a random event.
 Protein scores greater than 41 are significant ($p < 0.05$).

Protein Summary Report

Format As : Protein Summary [Help](#)
 Significance threshold $p < 0.05$ Max. number of hits 20
 Preferred taxonomy All entries
[Re-Search All](#) [Search Unmatched](#)

6/2/2018

Protein Summary Report (.Jdata/20180602/P005774.dat)

Mascot Search Results

User : Samovar
 Email : samovar@pandor.co.in
 Search title :
 MS data file : peaklist.xml
 Database : P53_Human_Human_20180602 (678 sequences; 280684 residues)
 Timestamp : 2 Jun 2018 at 10:29:08 GMT
 Top Score : 98 for [NP_001119587.1](#), cellular tumor antigen p53 isoform α (Homo sapiens)

Mascot Score Histogram

Protein score is $-10 \cdot \log(P)$, where P is the probability that the observed match is a random event.
 Protein scores greater than 41 are significant ($p < 0.05$).

Protein Summary Report

Format As : Protein Summary [Help](#)
 Significance threshold $p < 0.05$ Max. number of hits 20
 Preferred taxonomy All entries
[Re-Search All](#) [Search Unmatched](#)

6/2/2018

Protein Summary Report (.Jdata/20180602/P005778.dat)

Mascot Search Results

User : Samovar
 Email : samovar@pandor.co.in
 Search title :
 MS data file : peaklist.xml
 Database : P53_Human_Human_20180602 (678 sequences; 280684 residues)
 Timestamp : 2 Jun 2018 at 11:39:46 GMT
 Top Score : 93 for [NP_001119587.1](#), cellular tumor antigen p53 isoform α (Homo sapiens)

Mascot Score Histogram

Protein score is $-10 \cdot \log(P)$, where P is the probability that the observed match is a random event.
 Protein scores greater than 41 are significant ($p < 0.05$).

Protein Summary Report

Format As : Protein Summary [Help](#)
 Significance threshold $p < 0.05$ Max. number of hits 20
 Preferred taxonomy All entries
[Re-Search All](#) [Search Unmatched](#)