Identification of biomarkers and pathways to identify novel therapeutic targets in Alzheimer's disease: Insights from a systems biomedicine perspective

Md. Rezanur Rahman^{1,†,*}, Tania Islam^{2,†}, Toyfiquz Zaman¹, Md. Shahjaman³, Md. Rezaul Karim¹, Fazlul Huq⁴, Julian M.W. Quinn⁵, R. M. Damian Holsinger^{4,6}, and Mohammad Ali Moni^{4,6,*}

¹ Department of Biochemistry and Biotechnology, School of Biomedical Science, Khwaja Yunus Ali University, Sirajgonj, Bangladesh; rezanur12@yahoo.com(M.R.R.).

² Department of Biotechnology and Genetic Engineering, Islamic University, Kushtia, Bangladesh; taniaislam1304@gamil.com(T.I.).

³ Department of Statistics, Begum Rokeya University, Rangpur, Bangladesh; shahjaman_brur@yahoo.com (M.S.).

⁵ Bone Biology Division, Garvan Institute of Medical Research, Darlinghurst, NSW, Australia; j.quinn@garvan.edu.au(J.W.W.Q.).

⁴ Discipline of Pathology, School of Medical Sciences, The University of Sydney, Sydney, NSW, Australia; mohammad.moni@sydney.edu.au (M.A.M.).

⁵ Laboratory of Molecular Neuroscience and Dementia, Brain and Mind Centre, The University of Sydney, Camperdown, NSW, Australia; damian.holsinger@sydney.edu.au (R.M.D.H.).

[†]These two authors have made an equal contribution and hold joint first authorship for this work.

***Correspondence:**

Md. Rezanur Rahman, M.Sc

Lecturer,

Department of Biochemistry and Biotechnology

Khwaja Yunus Ali University,

Sirajgonj 6751, Bangladesh,

Tel: 88075163662-3, Fax: 88075163867, E-mail: rezanur12@yahoo.com

and

Dr. Mohammad Ali Moni

Research Fellow and Conjoint Lecturer

Discipline of Biomedical Science, School of Medical Sciences, Faculty of Medicine and Health,

The University of Sydney, Sydney, New South Wales 2006, Australia.

Tel: 61(02)93519458, Fax: +61 2 9351 9520, E-mail: mohammad.moni@sydney.edu.au

Abstract

Alzheimer's disease (AD) is a progressive neurodegenerative disease characterized by the accumulation of amyloid plaques and neurofibrillary tangles in the brain. However, there are no peripheral biomarkers available that can detect AD onset. This study aimed to identify such AD biomarkers through an integrative analysis of blood cell gene expression data. We used two microarray datasets (GSE4226 and GSE4229) comparing blood cell transcriptomes of AD patients and matched controls to identify differentially expressed genes (DEGs). Geneset and protein overrepresentation analysis, protein-protein interaction (PPI), DEGs-Transcription Factor interactions, DEGs-MicroRNAs interactions, protein-drug interactions, and protein subcellular localizations analyses were performed on DEGs common to the datasets. We identified 25 common DEGs between the two datasets. Integration of genome scale transcriptome datasets with biomolecular networks revealed hub genes (TUBB, ATF3, NOL6, UQCRC1, SND1, CASP2, BTF3, INPP5K, VCAM1, and CSTF1), common transcription factors (FOXC1, ZNF3, GEMIN7, and SMG9) and miRNAs (mir-20a-5p, mir-93-5p, mir-16-5p, let-7b-5p, mir-708-5p, mir-24-3p, mir-26b-5p, mir-17-5p, mir-4270, and mir-4441). These included 8 previously identified AD markers (BTF3, VCAM, FOXC1, mir-26b-5p, mir-20a-5p, mir-93-5p, let-7b-5p, mir-24-3p) and 13 novel AD biomarkers. Evaluation of histone modification revealed that hub genes and transcription factors possess several histone modification sites associated with AD. Protein-drug interactions revealed 10 compounds that affect the identified AD candidate markers, including anti-neoplastic agents (Vinorelbine, Vincristine, Vinblastine, Etoposide, Etoposide B, CYT997, and ZEN-012), a dermatological (Podofilox) and an immunosuppressive agent (Colchicine). The subcellular localization of markers varied, including nuclear, plasma membrane and cytosolic proteins. This study identified blood-cell derived molecular biomarker signatures that might be useful as peripheral biomarkers in AD. This study also identified drug and epigenetic data

associated with these molecules that may be useful in designing therapeutic approaches to ameliorate AD.

Keywords: systems biology; protein subcellular localization; epigenetics; protein-drug interaction; protein-protein interaction.

1. Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disease that gives rise to dementia and severe impairment of cognitive function in affected people, and is notably associated with the formation of extracellular amyloid plaques and intracellular accumulation of neurofibrillary tangles in the brain. It is the most common form of dementia, affecting 5.5 million individuals in the US alone, (Association A, 2017). Cognitive impairment is the main assessment method for patients, [2]. Sporadic AD is the most common form of the disease and generally affects individuals over the age of 65. Early diagnosis is rare, occurring in only a small fraction (1-5%) of patients. Early diagnosis of AD may improve management strategies for patients. However, there are no effective treatments for the disease. Therefore, research directed towards identifying AD biomarkers is needed not just for a better understanding of the development of AD but also as an important step in discovering new treatment strategies [3].

For differential diagnosis and AD patient assessments, neuroimaging techniques and cerebrospinal fluid biomarkers are widely used in clinical practice [4–6]. These methods are expensive, show low sensitivity and specificity and in the case cerebrospinal fluid, highly invasive. Thus, a simple blood-based test of dementia patients would be a valuable clinical tool for AD [7]. However, considering the dearth of useful peripheral blood protein or antigen biomarkers, despite much effort [8–11], blood-cell based transcript biomarkers may be a valuable approach [12–16]. Indeed, epigenetic factors are known to affect the pathogenesis of AD and have been accepted as playing significant roles in the development and progression of neurodegenerative diseases. Such epigenetic profiling, including studies of DNA methylation and histone modifications can be used to reveal the regulation of gene expression in various diseases [17]. Various external factors such as lifestyle, age, environment and health status result in epigenetic and transcriptome changes that may relate to AD

development [17]. However, epigenetic studies have yet to identify key mechanisms in AD development [17].

Dysregulation of the miRNAs has also been implicated in AD [18] and, consequently, miRNAs are increasingly being examined as possible biomarkers in or mediating factors in AD [19–21]. Thus, miRNA and epigenetic approaches have led to the use of systems biology approaches to elucidate key mediating or marker biomolecules in different complex diseases such as AD [20–23]. Accordingly, we have employed such an approach here. To identify critical genes, we used two peripheral blood microarray gene expression datasets derived from studies of AD patients. Using DEGs identified by this approach we performed functional annotations to identify important Gene Ontology and pathways enriched in these DEGs. We integrated the DEGs with interaction networks using the following approaches:

- (i) a PPI network of the proteins encoded by DEGs;
- (ii) DEG-TFs and DEGs-miRNAs interactions;
- (iii) protein-drug interaction networks to screen potential drugs with the binding affinity to target proteins estimated by molecular docking simulations;
- (iv) prediction of subcellular localization of proteins encoded by the DEGs was performed with the intent to provide insights into function.

Thus, we have used a comprehensive systems biology pipeline to explore molecular biomarker signatures of both mRNAs and miRNAs present in peripheral blood transcriptomes of AD patients. Additionally, we have also identified potential hub proteins, transcription factors (TFs) and pathways in blood that may provide potential biomarkers for early diagnosis of AD. These discoveries provide novel insights into the mechanisms underlying AD (Figure 1).

Figure 1: A multi-stage analysis methodology employed in this study. Transcriptomic datasets of blood from AD and normal individuals were collected from the NCBI-GEO database. The datasets were analyzed using GEO2R to identify DEGs. Functional enrichment analyses of DEGs were performed to identify significantly enriched pathways, Gene Ontology terms, and disease overrepresentation terms. Protein-protein interaction networks were reconstructed around DEGs. TF-target gene interactions and miRNA-target gene interactions were downloaded from JASPAR and miRTarbase databases, respectively. The protein-drug interaction analysis using the DrugBank database revealed candidate small molecular drugs that may interact with identified proteins. Significant pathways and GO terms, candidate biomarkers at protein and TFs levels, drug target, candidate drugs and drug-targeting sites were also identified.

2. Materials and Methods

2.1 Identification of Differentially Expressed Genes from Microarray High-throughput Datasets of Blood of AD Patients

The microarray gene expression high-throughput datasets in AD was obtained from the NCBI-GEO database [24]. This data was obtained from two separate studies of peripheral blood mononuclear cells that were collected from normal elderly control and AD subjects and analysed using human MGC cDNA microarrays; the results were deposited in NCBI-GEO database with the accession numbers GSE4226 and GSE4229 [25]. GSE4226 contained samples from 28 individuals in total, including 7 each of male and female controls and male and female AD patients, with the analysis originally published by Maes et al. [14] [PMID: 19366883], who also used DNA and blood protein studies in the search for new biomarkers. GSE4229 studied 40 blood samples (11 from male controls, 12 from female controls, 10 from male AD patients and 7 from female AD patients). The datasets were originally analyzed by Maes and colleagues using the NCBI-GEO2R online tool to identify DEGs in the AD compared to age matched controls. In our analysis, the gene expression dataset was first normalized by log2 transformation. Then, we analyzed the datasets in NCBI's GEO2R tool using Limma in hypothesis testing and the Benjamini & Hochberg correction to control the false discovery rate. The p-value<0.01 was regarded as the cut-off criteria to screen for significant DEGs. Venn analysis was performed using the online tool—Jvenn [26] to identify common target DEGs from the two datasets.

2.2 Gene Ontology and Pathway Enrichment Analysis

Gene over-representation analyses were performed via the bioinformatics resource Enrichr [27] to identify molecular function, biological process, and pathway annotations of the identified DEGs. In these analyses, the Gene Ontology (GO) terminology and KEGG

pathway database were used as annotation sources. The protein class over-representation was performed using the PANTHER database [28]. A p-value<0.05 was considered significant for all enrichment analyses.

2.3 Protein-protein Interaction Analysis

We used the STRING protein interactome database [29] to construct PPI networks of the proteins encoded by the DEGs. We set a medium confidence score of 400 for the construction of PPI networks since the number of DEGs is low. Network visualization and topological analyses were performed using NetworkAnalyst (Xia et al., 2015).

2.4 DEG-TFs Interaction Analysis

To identify regulatory TFs that control the DEGs at a transcriptional level, TF-target gene interactions were obtained using the JASPAR database [31] and were based on topological parameters (i.e., degree and betweenness centrality) using NetworkAnalyst [30].

2.5 DEGs-miRNA Interaction Analysis

To identify regulatory miRNAs that influence DEGs at the post-transcriptional level, miRNAs-target gene interactions were obtained from TarBase [32] and miRTarBase [33] based on topological parameters (i.e., degree and betweenness centrality) and analyzed using NetworkAnalyst [30].

2.6 Identification of Histone Modification Sites

Based on the recent systematic review on neurodegenerative disease aimed to identify most consistent epigenetic marks reported in methylations genes and histone modifications associated with AD [34], we used the human histone modification database (HHMD) to identify histone modification sites of the hub genes and TFs [35].

2.7 Cross-Validation of Candidate Biomarker Biomolecules

The AlzGene database contains 618 published genetic association studies of AD [36] and was used to evaluate the interactions between AD-associated cellular alterations in blood and AD-associated alterations collected in AlzGene database. The miRNAs were also crosschecked with blood-based miRNA signatures.

2.8 Protein-drug Interactions Analysis

Protein-drug interaction was analyzed using the DrugBank database (Version 5.0) [37] via NetworkAnalyst to identify potential drugs that could be used in the treatment of AD. The protein-drug interaction analysis revealed that TUBB had interactions with several drugs. Then, in order perform molecular docking analysis between TUBB and drugs, three dimensional (3D) crystal structure of the target proteins TUBB (PDB ID:1TUB) was obtained from protein data bank (PDB) [38]. Molecular docking analyses were performed using the protein-small molecule docking server SwissDock [39]. The docking score and best-fit pose were selected for each ligand/drug.

2.9 Prediction of Protein Subcellular Localization

We used the WoLF PSORT software package [40] to predict the subcellular localizations of the proteins encoded by the DEGs. The WoLF PSORT software predicts subcellular localizations based on amino acid sequence information.

3. Results

3.1 *Transcriptomic Signatures of AD*

The microarray datasets obtained from blood tissue of AD patients were analyzed and 25 mutually expressed core DEGs were identified between the two datasets. These core DEGs were considered as transcriptomic signatures for AD (Figure 2A) and classified into groups according to their functions and activities as transporters (4%), membrane traffic proteins (4%), hydrolases (24%), oxidoreductases (4%), cell junction proteins (4%), enzyme modulators (8%), transcription factors (12%), nucleic acid binding proteins (12%), calcium-binding proteins (4%) and cytoskeletal proteins (4%) (Figure 2B). A gene-set enrichment analysis showed that the DEGs were enriched in different biological processes, molecular functions and cellular components as summarized in Table 1. The molecular pathway enrichment analysis revealed that pathways previously identified in leukocyte transendothelial migration, oxidative phosphorylation, Parkinson's disease, cell adhesion molecules, non-alcoholic fatty liver disease, Alzheimer's disease and Huntington's disease were altered (Table 2).

Figure 2: (A) Identification of DEGs in microarray datasets GSE4226 and GSE4229. (A) The mutually expressed core DEGs identified between two datasets. (B) The protein class over-representation coded by the DEGs identified using the PANTHER database.

Table 1: Gene set enrichment analysis of the differentially expressed genes identified from microarray data of blood of AD patients

Category	GO ID	Term	P-value	Gene
Biological Process	GO:1902043	positive regulation of extrinsic apoptotic signaling pathway via death domain receptors	0.0081	ATF3
	GO:0045869	negative regulation of single stranded viral RNA replication via double stranded DNA intermediate	0.0111	INPP5K
	GO:0006508	proteolysis	0.0004	CELA3A;CAPN6;UQCRC1;CASP2
	GO:0070973	protein localization to endoplasmic reticulum exit site	0.0087	BCAP29
	GO:0090315	negative regulation of protein targeting to membrane	0.0087	INPP5K
	GO:0046838	phosphorylated carbohydrate dephosphorylation	0.0099	INPP5K
	GO:0045719	negative regulation of glycogen biosynthetic process	0.0087	INPP5K
	GO:0035810	positive regulation of urine volume	0.0099	INPP5K
	GO:0046855	inositol phosphate dephosphorylation	0.0099	INPP5K
	GO:0009308	amine metabolic process	0.0087	VCAM1
Molecular Function	GO:0008131	primary amine oxidase activity	0.0087	VCAM1
	GO:0016681	oxidoreductase activity, acting on diphenols and related substances as donors, cytochrome as acceptor	0.0111	UQCRC1
	GO:0052866	phosphatidylinositol phosphate phosphatase activity	0.0087	INPP5K
	GO:0008121	ubiquinol-cytochrome-c reductase activity	0.0111	UQCRC1

Cellular Component	GO:0005000	vasopressin receptor activity	0.0087	INPP5K
	GO:0004439	phosphatidylinositol-4,5-bisphosphate 5-phosphatase activity	0.0111	INPP5K
	GO:0106019	phosphatidylinositol-4,5-bisphosphate phosphatase activity	0.0111	INPP5K
	GO:0046030	inositol trisphosphate phosphatase activity	0.0136	INPP5K
	GO:0042577	lipid phosphatase activity	0.0149	INPP5K
	GO:0016641	oxidoreductase activity, acting on the CH-NH2 group of donors, oxygen as acceptor	0.0198	VCAM1
	GO:0097504	Gemini of coiled bodies	0.0099	GEMIN7
	GO:0005750	mitochondrial respiratory chain complex III	0.0185	UQCRC1
	GO:0042599	lamellar body	0.0198	KLK5
	GO:0048471	perinuclear region of cytoplasm	0.0113	CAPN6;NOCT; INPP5K
	GO:0032040	small-subunit processome	0.0440	NOL6
	GO:0000794	condensed nuclear chromosome	0.0428	NOL6
	GO:0000793	condensed chromosome	0.0500	NOL6
	GO:0000228	nuclear chromosome	0.0642	NOL6
	GO:0032587	ruffle membrane	0.0653	INPP5K
	GO:0005876	spindle microtubule	0.0606	CAPN6

Table 2: The molecular pathway enrichment in Alzheimer's disease.

Category	Pathways	Adj. p-value	Genes
KEGG	Ribosome	0.021	RPS9;RPL12;RPS5
BioCarta	Alternative Complement Pathway	0.02	C3
	Classical Complement Pathway	0.02	C3
	Lectin Induced Complement Pathway	0.02	C3
WikiPathways	Cytoplasmic Ribosomal Proteins	0.005	RPS9;RPL12;RPS5

3.3. Proteomic Signatures in AD

To reveal central proteins, a protein-protein interaction of the DEGs was constructed (Figure 3). Topological analysis revealed 10 central hub proteins (TUBB, ATF3, NOL6, UQCRC1, SND1, CASP2, BTF3, INPP5K, VCAM1, and CSTF1) (Table 3).

Figure 3: Protein-protein interaction network of the differentially expressed genes (DEGs) in Alzheimer's disease. The nodes indicate the DEGs and the edges indicate the interactions between two genes.

Table 3: Summary of hub proteins in Alzheimer's disease.

Symbol	Description	Feature
TUBB	Tubulin Beta Class	GTP binding and structural molecule activity
ATF3	Activating Transcription Factor 3	DNA binding transcription factor activity and sequence-specific DNA binding
NOL6	Nucleolar Protein 6	related to this gene include RNA binding
UQCRC1	Ubiquinol-Cytochrome C Reductase Core Protein 1	ubiquitin protein ligase binding and ubiquinol-cytochrome-c reductase activity
SND1	Staphylococcal Nuclease And Tudor Domain Containing 1	nucleic acid binding and transcription coregulator activity
CASP2	Caspase 2	enzyme binding and cysteine-type endopeptidase activity
BTF3	Basic Transcription Factor 3	this protein forms a stable complex with RNA polymerase IIB and is required for transcriptional initiation
INPP5K	Inositol Polyphosphate-5-Phosphatase K	vasopressin receptor activity and lipid phosphatase activity
VCAM1	Vascular Cell Adhesion Molecule 1	integrin binding and primary amine oxidase activity
CSTF1	Cleavage Stimulation Factor Subunit 1	RNA binding

3.4 The Regulatory Signatures of AD

We studied DEG-TF interactions and DEGs-miRNA interactions (Figure 4 and Figure 5) and detected central regulatory biomolecules (TFs and miRNAs) using topological parameters. Five TFs (FOXC1, ZNF3, GEMIN7, SMG9, and BCAP29) and 10 miRNAs (mir-20a-5p, mir-93-5p, mir-16-5p, let-7b-5p, mir-708-5p, mir-24-3p, mir-26b-5p, mir-17-5p, mir-4270, and mir-4441) were detected from the DEG-TF (Figure 4) and DEG-miRNA

interaction (Figure 5) networks respectively (Table 4). These biomolecules regulate genes at transcriptional and post-transcriptional levels.

Figure 4: Construction of regulatory networks of DEG-TF interactions.

Figure 5: Construction of regulatory networks of DEG-miRNA interactions.

Table 4: Summary of regulatory biomolecules (TFs, miRNAs) in Alzheimer's disease

Symbol	Description	Feature
FOXC1	Forkhead Box C1	DNA binding transcription factor activity and transcription factor binding
ZNF3	Zinc Finger Protein 3	nucleic acid binding and identical protein binding
GEMIN7	Gem Nuclear Organelle Associated Protein 7	Transport of the SLBP independent Mature mRNA and RNA transport.
SMG9	SMG9, Nonsense Mediated MRNA Decay Factor	Viral mRNA Translation and Gene Expression.
BCAP29	B Cell Receptor Associated Protein 29	B Cell Receptor Signaling Pathway (sino) and AKT Signaling Pathway
miRNAs		
mir-20a-5p	MicroRNA 20	Parkinson's Disease Pathway and cancer
mir-93-5p	MicroRNA 93	miRNAs involved in DNA damage response
mir-16-5p	MicroRNA 16	sudden infant death syndrome susceptibility pathways and in cancer
let-7b-5p	MicroRNA let-7b	MicroRNAs in cancer and metastatic brain tumor
mir-708-5p	MicroRNA 708	respiratory electron transport, ATP synthesis by chemiosmotic coupling, and heat production by uncoupling proteins.
mir-24-3p	MicroRNA 24	Peptide hormone metabolism and hepatitis C and hepatocellular carcinoma
mir-26b-5p	MicroRNA 26	Parkinson's Disease Pathway and cancer
mir-17-5p	MicroRNA 17	Parkinson's Disease Pathway and cancer
mir-4270	MicroRNA 4270	affiliated with the miRNA class
mir-4441	MicroRNA 4441	affiliated with the miRNA class

3.5 Epigenomic Signatures of AD: Histone Modification Sites

To provide insights regarding the epigenetic regulation of genes of interest, we studied possible histone modifications patterns. We observed that eight hub genes and TFs possessed one or more histone modification sites (Table 5).

Table 5: Histone modification patterns (obtained from HHMD) of novel hub genes and TFs with respect to the already known histone modification sites in neurodegenerative diseases.

Official symbol of Biomolecules	RefSeq ID	Histone modification sites already known in neurodegenerative diseases				
		H3K27	H3K4	H3K9	H3K9/H4K20	H4R3
<i>Hub protein</i>						
TUBB	NM_178014	✓	✓	✓	✓	✓
ATF3	NM_001030287	✓	✓	✓	✓	✓
NOL6	NM_139235	✓	✓	✓	✓	✓
UQCRC1	NM_003365	✓	✓	✓	✓	✓
SND1	NM_014390	✓	✓	✓	✓	✓
CASP2	NM_032983	✓	✓	✓	✓	✓
BTF3	NM_001037637	✓	✓	✓	✓	✓
INPP5K	NM_016532	✓	✓	✓	✓	✓
VCAM1	NM_080682	✓	✓	✓	✓	✓
CSTF1	NM_001324	✓	✓	✓	✓	✓
<i>TFs</i>						
FOXC1	NM_001453	✓	✓	✓	✓	✓
ZNF3	NM_017715	✓	✓	✓	✓	✓
GEMIN7	NM_001007269	✓	✓	✓	✓	✓
BCAP29	NM_001008407	✓	✓	✓	✓	✓

3.6 Cross-validation of Putative Biomarkers

A cross-check of the biomarkers identified in this study with the AlzGene database revealed no overlapped of the 25 DEGs with the AD-related biomarkers in AlzGene database. The present study did, however, identify miR-26a-5p, a miRNA that has previously been identified in a study of 12 blood-cell based miRNAs signatures of AD.

3.7 Protein-drug Interactions

In the protein-drug interaction network, TUBB protein was associated with the drugs Vinorelbine, Vincristine, Vinblastine, Podofilox, Colchicine, Epothilone D, Epothilone B, Cyt997, Ca4p, and Zen-012 (Figure 6). These identified drugs were categorized into different classes based on therapeutic classes (Figure 7A) and further classified according to their progress in the therapeutic pipeline (Figure 7B). Employing statistical significance thresholds of the drug-protein interactions and the possible role of the targeted protein in AD pathogenesis, a set of protein-drug interactions was selected and a series of molecular docking simulations were performed to analyze the binding affinities of identified drugs with their target protein (Table 6). The resultant energetic state and docking scores indicated the presence of thermodynamically feasible confirmations for all these interactions.

Figure 6: Protein-drug interaction network. The interactions between drugs and hub nodes (TUBB) are represented. The area of the node represents the degree of interaction in the network.

Table 5: Protein-drug interactions and their binding affinity by molecular docking statistics.

Target protein	Drug	Description	Full Fitness (kcal/mol)	Estimated ΔG (kcal/mol)
TUBB	Vinorelbine	Approved/Antineoplastic Agents	-4874.85	-8.43
	Vincristine	Approved/Antineoplastic Agents	-4848.09	-8.39
	Vinblastine	Approved/Antineoplastic Agents	-4757.89	-8.41
	Podofilox	Approved/Dermatologicals	-4947.47	-7.90
	Colchicine	Approved/Immunosuppressive Agents	-4906.52	-7.13
	Epothilone D	Experimental/Antineoplastic Agents	-5007.01	-7.29
	Epothilone B	Experimental/Antineoplastic Agents	-4782.27	-8.32
	CYT997	Investigational/Antineoplastic Agents	-5048.60	-7.14
	CA4P	Investigational/No available	-4951.83	-8.19
	ZEN-012	Investigational/Antineoplastic Agents	-	-

Figure 7: (A) Distribution of drugs into therapeutic chemical classes. (B) Classification of identified drugs according to the therapeutic pipeline.

3.8 Protein Subcellular Localization

Employing the WoLF PSORT software package we predicted the subcellular localization of proteins encoded by the DEGs in AD. The proportions of identified proteins in different subcellular compartments are displayed in Figure 8. Notably, nodes TUBB, SND1, CASP2 showed cytoplasm localization; ATF3, BTF3 were present in the nucleus; NOL6,

UQCRC1, INPP5K were localized in mitochondria; VCAM1 was localized in the plasma membrane; and CSTF1 is found in both cytoplasm and nucleus.

Figure 8: The distribution and percentages of the DEGs is shown to be the protein subcellular localization.

4. Discussion

The diagnosis of AD is currently performed based on neuropsychological evaluation and neuroimaging, but robust and specific biomarkers for diagnosis and prognosis of AD is an unmet challenge [11]. In this study, using a systems biology approach, we comprehensively investigated gene expression patterns evident in peripheral blood cells of AD patients and identified robust candidate biomarkers that may serve as potential biomarkers of AD. Our results may therefore provide significant insight into the process leading to AD.

Microarray data are extensively used in biomedical research and have become a major resource for identifying biomarker candidates [41]. Microarray gene expression profiling is widely used to identify DEGs in various diseases (Gov et al., 2017; Guo et al., 2017; Islam et al., 2018; Rahman et al., 2018). Analyzing gene expression patterns in the blood of AD patients revealed significant alterations in the expression profiles of 25 genes in two transcriptomic datasets. The over-representation analysis revealed AD and neurodegeneration associated molecular pathways (Oxidative phosphorylation and AD pathway) [45].

Integrative analysis of protein-protein interaction network can identify proteins that are key players in the mechanisms underlying disease [46]. Our study of the PPI network based on DEGs thus identified a number of key hub proteins (Table 2) which may indicate the onset and progression of neurodegenerative diseases. These hub proteins are associated with a number of different types of biological and pathological processes. According to the GeneCards database (<https://www.genecards.org/>), diseases associated with TUBB include cortical dysplasia with other brain malformations. ATF3-related pathway has been implicated in brain vascular damage [47] as well as amyloid deposition in AD [48], consistent with our observations. NOL6 has been associated with ribosome biogenesis which

affects the behavior of many cells types but has not previously been linked to AD (GeneCards database). Similarly, pathways involving UQCRC1 have been implicated include cell metabolism (including respiratory electron transport), but involvement in AD has yet to be reported. The multifunctional protein SND1 is deregulated in various cancers [49] and dysregulation of CASP2, a cysteine-aspartic acid protease (caspase) family member may play a role in AD. BTF3 is a blood biomarker of AD [50]. INPP5K is associated with the disease muscular dystrophy, and intellectual disability, but no association with AD has been reported to date. Hub protein CSTF1 is involved in RNA poly-adenylation but its function is otherwise largely unknown. VCAM1 is a cell adhesion molecule important both in leukocyte and neuronal function and its expression is associated with AD dementia [51]. In summary, these hub genes have disease associations (seen in dbGaP and OMIM data) which implies that they are involved in a range of pathogenic influences, but nevertheless most are poorly understood.

We also identified a number of TFs and miRNAs with altered levels of expression in AD patient blood cells. Differential expression of these molecules may provide significant information on dysregulation of gene expression in AD. Our previous study using microarray data obtained from brain also revealed FOXC1 as a dysregulated TF in AD (Rahman et al., 2018). A number of ZNF TFs have been implicated in the pathogenesis of neuronal diseases [52], but the role of ZNF3 in neurological disorders or AD is unknown. GEMN6 has been reported to be associated with AD [53], but this is poorly understood SMG9 is involved in nonsense-mediated mRNA decay but is not reported to have an AD association [54]. BCAP29 has been suggested as diagnostic biomarkers in peripheral blood gene expression analysis [55].

miRNAs are single stranded non-coding RNAs that can silence gene activities by targeting their transcripts [19] [56]. Dysregulated miRNAs have been reported in the

development of AD [19]. Since miRNAs may serve as biomarkers for diagnosis and therapeutic target for breakthrough treatment strategies in the AD [57]; therefore, we identified the miRNAs as regulatory component of the DEGs. mir-20a-5p identified in our study is associated with aging [58] and has also been proposed as peripheral blood biomarker in AD [10]. The upregulation of mir-93-5p has been reported as biomarkers in the serum of AD patients [10]. The mir-16-5p and mir-708-5p have also been identified as differentially expressed in cerebrospinal fluid cells in AD [59,60]. The let-7b-5p has previously been down-regulated in AD as a member of 12 blood based miRNA biomarker signature [61]. The dysregulation of mir-24-3p was also reported in lung cancer [10]. Decreased expression of mir-24-3p was identified as candidate that can discriminate AD from control CSF [62]. Mir-26b-5p is also down-regulated in AD as a member of 12 blood based miRNA biomarker signature [50]. Elevated levels of miR-26-b may thus contribute to the AD neuronal pathology, and this may be reflected also in blood cell functions dependent upon mir-26b-5p [63]. Mir-17-5p was found to be deregulated in circulatory biofluids in lung cancer [10] and was dysregulated in the AD [64]. In contrast, a role for mir-4270 and mir-4441 in the AD has not been reported previously.

To study the epigenetic regulation of the hub genes, we investigated histone modification sites found in the coding genes of biomolecules implicated with neurodegenerative disease [65], identifying a range of sites, summarized in Table 4. While these are also poorly understood it raises the possibility that their alteration may be a major means of regulation for these genes that warrant further investigation, and that these genes are under tight control by a number of regulatory processes. Given the importance of these hub genes, and the possibility that they may exert sufficient influence in pathogenic processes in AD (and AD-affected blood cells) we studied the protein-drug interactions to identify drugs that may influence them. A total 10 drugs were identified from the interaction network.

A number of molecular-modelling based methods are used in pharmaceutical research to investigate complex biological systems; in particular, molecular docking methods broadly used in rational drug design to elucidate the ligand conformation within binding sites of the target protein. These molecular docking methods estimate free energy by evaluating critical phenomena involved in the intermolecular recognition process in ligand-receptor binding [66]. Consequently, in the present study, we evaluated the binding mode of the ligands/drugs with the target protein TUBB and energetically stable conformations obtained from existing databases. In this way we discovered associations between the identified drugs and our putative AD markers that indicate that they may influence important pathways influencing disease progression, however it should be noted that the consequences of biomarker blockade is unclear from our work and require further investigations. We would also emphasize that our biomarkers are blood cell based and as yet we do not know how these related to brain biology. Nevertheless, many of these drugs have been evaluated for effects on tissues as part of their clinical evaluation and it may be useful to evaluate what is known of their effects on neural cells.

5. Conclusion

In the present study, we have analyzed the blood-based transcriptomics profiles using integrative multi-omics analyses to reveal systems level biomarkers at the protein (hub proteins, TFs) and RNA level (mRNAs, miRNAs). A number of key hub genes were significantly enriched in pathways involved in leukocyte transendothelial migration, oxidative phosphorylation, Parkinson's disease, cell adhesion molecules, non-alcoholic fatty liver disease, AD, and Huntington's disease. We also identified significant hub proteins TFs and miRNAs that regulate the function of the DEGs in the AD. These biomolecules may be considered candidate systems biomarkers at the protein level and RNA levels. The hub genes possess several histone modification sites of interest. Thus, we have identified potential

peripheral biomarkers for AD which can be detected as transcripts in blood cells, and these warrant clinical investigations in AD patients to evaluate their utility. The nature of these biomarkers and the pathways they participate in may reveal new aspects of AD development and progression since these biomarkers are evident in cells outside the central nervous system.

References

- [1] Association A. 2017 Alzheimer's disease facts and figures. *Alzheimer's Dement* 2017;13:325–73. doi:10.1016/j.jalz.2017.02.001.
- [2] Dunckley T, Beach TG, Ramsey KE, Grover A, Mastroeni D, Walker DG, et al. Gene expression correlates of neurofibrillary tangles in Alzheimer's disease. *Neurobiol Aging* 2006;27:1359–71. doi:10.1016/j.neurobiolaging.2005.08.013.
- [3] Li X, Jiang W, Zhang Y, Meng F, Lian B, Chen X, et al. Identification of active transcription factor and miRNA regulatory pathways in Alzheimer's disease. *Bioinformatics* 2013;29:2596–602. doi:10.1093/bioinformatics/btt423.
- [4] McKhann G, Knopman DS, Chertkow H, Hyman B, Jack CR, Kawas C, et al. The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging- Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 2011;7:263–9. doi:10.1016/j.jalz.2011.03.005.The.
- [5] Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on. *Alzheimer's Dement* 2011;7:270–9. doi:10.1016/j.jalz.2011.03.008.The.
- [6] Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM, et al. Toward defining the preclinical stages of Alzheimer's disease: Recommendations from the National Institute on Aging- Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's Dement* 2011;7:280–92. doi:10.1016/j.jalz.2011.03.003.Toward.

- [7] Lista S, Faltraco F, Prvulovic D, Hampel H. Blood and plasma-based proteomic biomarker research in Alzheimer's disease. *Prog Neurobiol* 2013;101–102:1–17. doi:10.1016/j.pneurobio.2012.06.007.
- [8] Schneider P, Hampel H, Buerger K. Biological Marker Candidates of Alzheimer's Disease in Blood, Plasma, and Serum. *CNS Neurosci Ther* 2009;15:358–74. doi:10.1111/j.1755-5949.2009.00104.x.
- [9] Shi L, Baird AL, Westwood S, Hye A, Dobson R, Thambisetty M, et al. A Decade of Blood Biomarkers for Alzheimer's Disease Research: An Evolving Field, Improving Study Designs, and the Challenge of Replication. *J Alzheimer's Dis* 2018;62:1181–98. doi:10.3233/JAD-170531.
- [10] Kumar S, Reddy PH. Are circulating microRNAs peripheral biomarkers for Alzheimer's disease? *Biochim Biophys Acta - Mol Basis Dis* 2016;1862:1617–27. doi:10.1016/j.bbadis.2016.06.001.
- [11] Sharma N, Singh AN. Exploring biomarkers for Alzheimer's disease. *J Clin Diagnostic Res* 2016;10:KE01-KE06. doi:10.7860/JCDR/2016/18828.8166.
- [12] Fehlbaum-Beurdeley P, Jarrige-Le Prado AC, Pallares D, Carriere J, Guihal C, Soucaille C, et al. Toward an Alzheimer's disease diagnosis via high-resolution blood gene expression. *Alzheimers Dement* 2010;6:25–38. doi:10.1016/j.jalz.2009.07.001.
- [13] Rye PD, Booij BB, Grave G, Lindahl T, Kristiansen L, Andersen H-M, et al. A novel blood test for the early detection of Alzheimer's disease. *J Alzheimers Dis* 2011;23:121–9. doi:10.3233/JAD-2010-101521.
- [14] Maes OC, Schipper HM, Chertkow HM, Wang E. Methodology for discovery of Alzheimer's disease blood-based biomarkers. *J Gerontol A Biol Sci Med Sci* 2009;64:636–45. doi:10.1093/gerona/glp045.

- [15] Han G, Wang J, Zeng F, Feng X, Yu J, Cao H-Y, et al. Characteristic transformation of blood transcriptome in Alzheimer's disease. *J Alzheimers Dis* 2013;35:373–86. doi:10.3233/JAD-121963.
- [16] Bai Z, Stamova B, Xu H, Ander BP, Wang J, Jickling GC, et al. Distinctive RNA expression profiles in blood associated with Alzheimer disease after accounting for white matter hyperintensities. *Alzheimer Dis Assoc Disord* 2014;28:226–33. doi:10.1097/WAD.000000000000022.
- [17] Wen KX, Milic J, El-Khodori B, Dhana K, Nano J, Pulido T, et al. The role of DNA methylation and histone modifications in neurodegenerative diseases: A systematic review. *PLoS One* 2016;11:1–31. doi:10.1371/journal.pone.0167201.
- [18] Maciotta S, Meregalli M, Torrente Y. The involvement of microRNAs in neurodegenerative diseases. *Front Cell Neurosci* 2013;7:1–17. doi:10.3389/fncel.2013.00265.
- [19] Femminella GD, Ferrara N, Rengo G. The emerging role of microRNAs in Alzheimer's disease. *Front Physiol* 2015;6:1–5. doi:10.3389/fphys.2015.00040.
- [20] Rahman MR, Islam T, Turanli B, Zaman T, Faruquee HM, Rahman MM, et al. Network-based approach to identify molecular signatures and therapeutic agents in Alzheimer's disease. *Comput Biol Chem* 2019;78:431–9. doi:10.1016/j.compbiolchem.2018.12.011.
- [21] Rahman MR, Islam T, Shahjaman M, Quinn JMW, Holsinger RMD, Moni MA. Identification of common molecular biomarker signatures in blood and brain of Alzheimers disease. *bioRxiv* 2019:482828. doi:10.1101/482828.
- [22] Rollo JL, Banihashemi N, Vafaei F, Crawford JW, Kuncic Z, Holsinger RMD. Unraveling the mechanistic complexity of Alzheimer's disease through systems

- biology. *Alzheimers Dement* 2016;12:708–18. doi:10.1016/j.jalz.2015.10.010.
- [23] Rahman MR, Islam T, Gov E, Turanli B, Gulfidan G, Shahjaman M, et al. Identification of Prognostic Biomarker Signatures and Candidate Drugs in Colorectal Cancer: Insights from Systems Biology Analysis. *Medicina (Kaunas)* 2019;55. doi:10.3390/medicina55010020.
- [24] Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI GEO: Archive for functional genomics data sets - Update. *Nucleic Acids Res* 2013;41:991–5. doi:10.1093/nar/gks1193.
- [25] Maes OC, Xu S, Yu B, Chertkow HM, Wang E, Schipper HM. Transcriptional profiling of Alzheimer blood mononuclear cells by microarray. *Neurobiol Aging* 2007;28:1795–809. doi:10.1016/j.neurobiolaging.2006.08.004.
- [26] Bardou P, Mariette J, Escudie F, Djemiel C, Klopp C. jvenn: an interactive Venn diagram viewer. *BMC Bioinformatics* 2014;15:293. doi:10.1186/1471-2105-15-293.
- [27] Kuleshov M V., Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* 2016;44:W90–7. doi:10.1093/nar/gkw377.
- [28] Mi H, Muruganujan A, Thomas PD. PANTHER in 2013: Modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res* 2013;41:377–86. doi:10.1093/nar/gks1118.
- [29] Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The STRING database in 2017: Quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res* 2017;45:D362–8. doi:10.1093/nar/gkw937.
- [30] Xia J, Gill EE, Hancock REW. NetworkAnalyst for statistical, visual and network-

- based meta-analysis of gene expression data. *Nat Protoc* 2015;10:823–44. doi:10.1038/nprot.2015.052.
- [31] Khan A, Fornes O, Stigliani A, Gheorghe M, Castro-Mondragon JA, Van Der Lee R, et al. JASPAR 2018: Update of the open-access database of transcription factor binding profiles and its web framework. *Nucleic Acids Res* 2018;46:D260–6. doi:10.1093/nar/gkx1126.
- [32] Sethupathy P, Corda B, Hatzigeorgiou AG. TarBase²: A comprehensive database of experimentally supported animal microRNA targets TarBase²: A comprehensive database of experimentally supported animal microRNA targets. *Rna* 2006;12:192–7. doi:10.1261/rna.2239606.scale.
- [33] Hsu S Da, Lin FM, Wu WY, Liang C, Huang WC, Chan WL, et al. MiRTarBase: A database curates experimentally validated microRNA-target interactions. *Nucleic Acids Res* 2011;39:163–9. doi:10.1093/nar/gkq1107.
- [34] Wen K-X, Milic J, El-Khodori B, Dhana K, Nano J, Pulido T, et al. The Role of DNA Methylation and Histone Modifications in Neurodegenerative Diseases: A Systematic Review. *PLoS One* 2016;11:e0167201. doi:10.1371/journal.pone.0167201.
- [35] Zhang Y, Lv J, Liu H, Zhu J, Su J, Wu Q, et al. HHMD: The human histone modification database. *Nucleic Acids Res* 2009;38:149–54. doi:10.1093/nar/gkp968.
- [36] Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE. Systematic meta-analyses of Alzheimer disease genetic association studies: The AlzGene database. *Nat Genet* 2007;39:17–23. doi:10.1038/ng1934.
- [37] Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res* 2018;46:D1074–82. doi:10.1093/nar/gkx1037.

- [38] Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, et al. SWISS-MODEL: Modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res* 2014;42:252–8. doi:10.1093/nar/gku340.
- [39] Grosdidier A, Zoete V, Michielin O. SwissDock, a protein-small molecule docking web service based on EADock DSS. *Nucleic Acids Res* 2011;39:270–7. doi:10.1093/nar/gkr366.
- [40] Horton P, Park K-J, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, et al. WoLF PSORT: protein localization predictor. *Nucleic Acids Res* 2007;35:W585-7. doi:10.1093/nar/gkm259.
- [41] Stewart JP, Richman S, Maughan T, Lawler M, Dunne PD, Salto-Tellez M. Standardising RNA profiling based biomarker application in cancer-The need for robust control of technical variables. *Biochim Biophys Acta* 2017;1868:258–72. doi:10.1016/j.bbcan.2017.05.005.
- [42] Guo Y, Bao Y, Ma M, Yang W. Identification of key candidate genes and pathways in colorectal cancer by integrated bioinformatical analysis. *Int J Mol Sci* 2017;18:1–15. doi:10.3390/ijms18040722.
- [43] Islam T, Rahman MR, Gov E, Turanli B, Gulfidan G, Haque MA, et al. Drug Targeting and Biomarkers in Head and Neck Cancers□: OMICS 2018;22:422–36. doi:10.1089/omi.2018.0048.
- [44] Gov E, Kori M, Arga KY. Multiomics Analysis of Tumor Microenvironment Reveals Gata2 and miRNA-124-3p as Potential Novel Biomarkers in Ovarian Cancer. *Omi A J Integr Biol* 2017;21:omi.2017.0115. doi:10.1089/omi.2017.0115.
- [45] Li H, Hong G, Lin M, Shi Y, Wang L, Jiang F, et al. Identification of molecular alterations in leukocytes from gene expression profiles of peripheral whole blood of

- Alzheimer's disease. *Sci Rep* 2017;7:1–10. doi:10.1038/s41598-017-13700-w.
- [46] Sevimoglu T, Arga KY. The role of protein interaction networks in systems biomedicine. *Comput Struct Biotechnol J* 2014;11:22–7. doi:10.1016/j.csbj.2014.08.008.
- [47] Aung HH, Altman R, Nyunt T, Kim J, Nuthikattu S, Budamagunta M, et al. Lipotoxic brain microvascular injury is mediated by activating transcription factor 3-dependent inflammatory and oxidative stress pathways. *J Lipid Res* 2016;57:955–68. doi:10.1194/jlr.M061853.
- [48] Zhu M, Allard JS, Zhang Y, Perez E, Spangler EL, Becker KG, et al. Age-Related Brain Expression and Regulation of the Chemokine CCL4 / MIP-1 A in APP / PS1 Double-Transgenic Mice. *J Neuropathol Exp Neurol* 2014;73:362–74.
- [49] Jariwal N, Rajasekaran D, Srivastava J, Gredler R, Akiel MA, Robertson CL, et al. Role of the staphylococcal nuclease and tudor domain containing 1 in oncogenesis (Review). *Int J Oncol* 2015;46:465–73. doi:10.3892/ijo.2014.2766.
- [50] Lunnon K, Sattlecker M, Furney SJ, Coppola G, Simmons A, Proitsi P, et al. A blood gene expression marker of early Alzheimer's disease. *J Alzheimer's Dis* 2013;33:737–53. doi:10.3233/JAD-2012-121363.
- [51] Huang C-W, Tsai M-H, Chen N-C, Chen W-H, Lu Y-T, Lui C-C, et al. Clinical significance of circulating vascular cell adhesion molecule-1 to white matter disintegrity in Alzheimer's dementia. *Thromb Haemost* 2015;114:1230–40. doi:10.1160/TH14-11-0938.
- [52] Cassandri M, Smirnov A, Novelli F, Pitolli C, Agostini M, Malewicz M, et al. Zinc-finger proteins in health and disease. *Cell Death Discov* 2017;3:17071. doi:10.1038/cddiscovery.2017.71.

- [53] Wong J. Altered Expression of RNA Splicing Proteins in Alzheimer's Disease Patients: Evidence from Two Microarray Studies. *Dement Geriatr Cogn Dis Extra* 2013;3:74–85. doi:10.1159/000348406.
- [54] da Costa PJ, Menezes J, Romão L. The role of alternative splicing coupled to nonsense-mediated mRNA decay in human disease. *Int J Biochem Cell Biol* 2017;91:168–75. doi:10.1016/j.biocel.2017.07.013.
- [55] Hsu WC, Denq C, Chen SS. A diagnostic methodology for Alzheimer's disease. *J Clin Bioinforma* 2013;3:1–13. doi:10.1186/2043-9113-3-9.
- [56] Goodall EF, Heath PR, Bandmann O, Kirby J, Shaw PJ. Neuronal dark matter: the emerging role of microRNAs in neurodegeneration. *Front Cell Neurosci* 2013;7:1–16. doi:10.3389/fncel.2013.00178.
- [57] Shaik MM, Tamargo IA, Abubakar MB, Kamal MA, Greig NH, Gan SH. The role of microRNAs in Alzheimer's disease and their therapeutic potentials. *Genes (Basel)* 2018;9:1–36. doi:10.3390/genes9040174.
- [58] Hackl M, Brunner S, Fortschegger K, Schreiner C, Micutkova L, Mück C, et al. miR-17, miR-19b, miR-20a, and miR-106a are down-regulated in human aging. *Aging Cell* 2010;9:291–6. doi:10.1111/j.1474-9726.2010.00549.x.
- [59] Burgos K, Malenica I, Metpally R, Courtright A, Rakela B, Beach T, et al. Profiles of extracellular miRNA in cerebrospinal fluid and serum from patients with Alzheimer's and Parkinson's diseases correlate with disease status and features of pathology. *PLoS One* 2014;9:e94839. doi:10.1371/journal.pone.0094839.
- [60] McKeever PM, Schneider R, Taghdiri F, Weichert A, Multani N, Brown RA, et al. MicroRNA Expression Levels Are Altered in the Cerebrospinal Fluid of Patients with Young-Onset Alzheimer's Disease. *Mol Neurobiol* 2018;10–2. doi:10.1007/s12035-

018-1032-x.

- [61] Leidinger P, Backes C, Deutscher S, Schmitt K, Mueller SC, Frese K, et al. A blood based 12-miRNA signature of Alzheimer disease patients. *Genome Biol* 2013;14:R78. doi:10.1186/gb-2013-14-7-r78.
- [62] Lusardi TA, Phillips JI, Wiedrick JT, Harrington CA, Lind B, Lapidus JA, et al. MicroRNAs in Human Cerebrospinal Fluid as Biomarkers for Alzheimer's Disease. *J Alzheimers Dis* 2017;55:1223–33. doi:10.3233/JAD-160835.
- [63] Absalon S, Kochanek DM, Raghavan V, Krichevsky AM. MiR-26b, upregulated in Alzheimer's disease, activates cell cycle entry, tau-phosphorylation, and apoptosis in postmitotic neurons. *J Neurosci* 2013;33:14645–59. doi:10.1523/JNEUROSCI.1327-13.2013.
- [64] Hu Y-B, Li C-B, Song N, Zou Y, Chen S-D, Ren R-J, et al. Diagnostic Value of microRNA for Alzheimer's Disease: A Systematic Review and Meta-Analysis. *Front Aging Neurosci* 2016;8:1–9. doi:10.3389/fnagi.2016.00013.
- [65] Pattaroni C, Jacob C. Histone methylation in the nervous system: functions and dysfunctions. *Mol Neurobiol* 2013;47:740–56. doi:10.1007/s12035-012-8376-4.
- [66] Ferreira LG, Dos Santos RN, Oliva G, Andricopulo AD. Molecular docking and structure-based drug design strategies. *Molecules* 2015;20:13384–421. doi:10.3390/molecules200713384.

Acknowledgment

We are grateful to our colleagues for their comments and suggestion about the manuscript.

Conflict of interest

The authors declare no conflict of interest.