Within-finger maps of tactile and nociceptive input in the human parietal cortex

Flavia Mancini^{1,2*}, Martin I. Sereno³, Min-Ho Lee⁴, Giandomenico D Iannetti², Irene Tracey⁴

¹*Computational and Biological Learning, Department of Engineering, University of Cambridge, Cambridge, UK*

²*Department of Neuroscience, Physiology and Pharmacology, University College London, London, UK*

³*Department of Psychology, San Diego State University, San Diego, USA*

⁴*Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK*

**Corresponding author:*

Dr Flavia Mancini

Computational and Biological Learning, Department of Engineering, University of Cambridge
Trumpington Street, Cambridge, CB2 1PZ

flavia.mancini@eng.cam.ac.uk

24 **Abstract**

25

26 The spatial representation of nociceptive input in the human parietal cortex is not fully
 27 understood. For instance, it is known that the primary somatosensory cortex (S1) contains a
 28 representation of nociceptive-selective input to different fingers, but it is unclear whether S1
 29 subdivisions contain finer-grained, within-finger maps of nociceptive input. It is also unknown
 30 whether within-finger maps of somatosensory input exist in intraparietal regions. Therefore,
 31 we used high-field 7T functional MRI to reveal within-finger maps of nociceptive and tactile
 32 inputs in the human parietal cortex. Although we replicated the previous findings of between-
 33 finger maps of nociceptive input spanning S1 subdivisions, we found weak and inconsistent
 34 evidence for within-finger maps of nociceptive input in S1 subdivisions. In the same subjects,
 35 we found mirrored, within-finger maps of tactile input in areas 3a, 3b, and 1. Importantly, we
 36 discovered a within-finger map of nociceptive input, but not of tactile input, in the human
 37 intraparietal area 1 (hIP1). In conclusion, our study indicates that the spatial representation
 38 of nociceptive input in the parietal cortex partly differs from that of tactile input and reports
 39 the discovery of a within-finger map of nociceptive input in hIP1.

40

41

42 **New & Noteworthy**

43

44 We report the discovery of a fine-grained map of nociceptive input to finger segments in the
 45 human intraparietal area 1.

46

47

48 Introduction

49

50 Human parietal areas contain topographic representations of innocuous somatosensory
51 input (Serenio and Huang 2014), which support spatial processing of tactile input,
52 sensorimotor transformations and goal-directed behaviour. Although these regions represent
53 predominantly non-nociceptive somatosensory information (Frot et al. 2013), they are also
54 the target of spinothalamic projections conveying nociceptive information (Dum et al. 2009;
55 Gingold et al. 1991; Kenshalo et al. 2000; Kenshalo et al. 1988; Kenshalo and Isensee 1983;
56 Schmahmann and Pandya 1990). However, the precise spatial organization of nociceptive
57 responses in the parietal cortex is not clear.

58

59 In particular, each of the four cytoarchitectonic areas (3a, 3b, 1, and 2) within the primary
60 somatosensory cortex (S1) contains a complete representation of mechanical, non-
61 nociceptive input to the hand and fingers, both along the *medial-to-lateral axis* (e.g., *between*
62 *finger representation*) and along the *anterior-to-posterior axis* (*within finger representation*)
63 (Kaas et al. 1979; Merzenich et al. 1978). The within-finger tactile map in each area of S1
64 mirrors the representation in the adjacent area; the base-to-tip finger representation is
65 posterior-to-anterior in area 3b but anterior-to-posterior in the adjacent area 1 (Kaas et al.
66 1979; Merzenich et al. 1978; Sanchez-Panchuelo et al. 2012).

67

68 Previous studies have shown that the human S1 contains somatotopic maps of nociceptive
69 inputs along its medio-lateral axis, spanning different cytoarchitectonic subdivisions: not only
70 the broad somatotopies for hand, face, and foot territories (Andersson et al. 1997; Bingel et
71 al. 2004), but also the between-fingers maps of nociceptive input (Mancini et al. 2012). It is
72 unknown whether and how nociceptive input is mapped along the anterior-to-posterior axis
73 of S1. Spinothalamic projections also target areas in the intra-parietal sulcus (IPS) and
74 superior parietal lobule (SPL) (Schmahmann and Pandya 1990). It is unknown whether IPS
75 contains a representation of somatosensory input to different finger segments. Therefore, we
76 used high-field functional MRI (fMRI) to determine whether within-finger maps of nociceptive
77 input exist in the parietal cortex, if they exist, to evaluate whether these maps are co-aligned
78 to within-finger maps of tactile input.

79

80

81 Methods

82

83 Participants

84

85 Fourteen healthy and right-handed subjects (3 women) aged 21–32 years (mean $\pm$ SD:
86 24.64 ± 3.39) participated in the study, after having given written informed consent. All
87 experimental procedures were approved by the Health Research Authority (REC reference:
88 11/SC/0249; IRAS projectID: 80049). Twelve participants undertook a within-finger mapping
89 protocol. Additionally, two participants undertook a between-finger mapping protocol for
90 replication of previous results obtained at lower magnetic field strengths (Mancini et al. 2012;
91 Mancini et al. 2018) and for validating our scanning and analyses methods.

92

93 Experimental design

94

Within-finger mapping. Twelve participants attended two scanning sessions, each lasting 75 minutes. The sessions were identical except for the modality of the somatosensory stimulus: mechanical and non-nociceptive stimulation of A β fibers in one session, nociceptive-selective stimulation of A δ /C fibers in the other session, in counterbalanced order across participants (see Supplemental Data for a description of the stimuli used: https://github.com/flamancini/parietalmaps/blob/master/supplemental_data_JNP19.pdf). Participants laid inside the scanner bore with their right hand open and supine, kept firmly in place with the aid of foam pads and a cushion. The stimulation was a travelling wave of somatosensory stimuli sweeping through the proximal segments of digits 2, 3, 4 and 5, the middle segments of digits 2-5, and the distal segments of digits 2-5. The different digits were stimulated in random order and only the proximal-to-distal stimulation was given in sequential order. To correct for systematic regional variations in the shape of the hemodynamic response function, we interleaved progressions in two directions (proximal-to-distal; distal-to-proximal). The first progression was counterbalanced among participants. For both modalities, 12 cycles of 42.67s trains of stimuli (interstimulus interval, 2.13s) were administered in each of the three runs. Each run both started and ended with 20s of rest. However, painful sensations elicited by high-energy laser pulses can typically last for more than a second. One participant withdrew from the study before completing the full protocol and was not included in any analysis.

Between-finger mapping. To validate our scanning and analyses methods, we mapped the between-finger representation of tactile stimuli in one participant and that of nociceptive stimuli in another participant. All procedures were identical to the 'within-finger mapping' except for the spatial progression of the stimuli. The stimulation was a travelling wave sweeping through random locations in d2, d3, d4, and d5. Each digit was stimulated sequentially, whereas the different segments of each digit were stimulated randomly (but covering the same surface of the within-finger mapping protocol). Again, we interleaved progressions in two directions (d2-to-d5; d5-to-d2).

MRI data acquisition

MRI data were acquired on a Siemens Magnetom 7 Tesla MRI scanner with a 32-channel head coil. Blood oxygenation level dependent (BOLD) fMRI was acquired using multislice gradient echoplanar imaging, with axial slices centred on the typical location of hand maps in S1 (voxel size 1.7mm isotropic, 281 volumes/run, 30 axial slices, flip angle 65°, TR 2000ms, TE 26ms, GRAPPA acceleration factor PE 2, bandwidth 1370Hz/Px, echo spacing 0.81ms). The slice coverage of our echoplanar images was centred on the hand-knob in S1 and included most of the parietal lobe and premotor cortex, but excluded most of the cingulate cortex and insular cortex. Task-fMRI data were collected during 6 runs (3 runs per session; 1 session per modality). The first 4 measurements of each functional run were discarded from all subsequent analyses.

We also collected a structural MP2RAGE scan (voxel size 1mm isotropic, 192 slices per slab, flip angle 14°, TR 5000ms, TI 1900ms, TE 2.84ms, GRAPPA acceleration factor PE 3, bandwidth 240Hz/Px, echo spacing 7 ms) and a gradient field map (voxel size 2mm isotropic, 64 slices, flip angle 39°, TR 620ms, TE 14.08ms, TE 25.1ms, bandwidth 734Hz/Px). We used the FreeSurfer program 'recon-all' (<https://surfer.nmr.mgh.harvard.edu/>) to reconstruct the cortical surface from the T1

structural image, after correcting for image intensity non-uniformity using the AFNI program '3dUniformize' (<https://afni.nimh.nih.gov>) and non-brain removal with 3dSkullStrip.

FMRI analyses

We performed two parallel sets of analyses: Generalized Linear Model analyses (GLM) in order to study the magnitude of the cortical response; phase mapping analyses in order to characterise the spatial organisation of parietal responses.

GLM analyses. The purpose of these analyses was to determine the magnitude of brain responses to tactile stimuli and nociceptive stimuli in S1 and IPS, irrespective of the somatotopic organization of these responses. Data processing was carried out using standard GLM analyses in FSL (www.fmrib.ox.ac.uk/fsl; supplemental data https://github.com/flamancini/parietalmaps/blob/master/supplemental_data_JNP19.pdf). fMRI data were subjected to exclusion in cases of visible spin history motion artefact as a result of sharp motion during more than one scan sessions (1 mm of absolute mean displacement in fewer than five volumes), as in previous studies (Kolasinski et al. 2016a; Kolasinski et al. 2016b); this happened in two participants, which were then excluded from further analyses. Thus, nine participants were included in further analyses. The two participants who undertook the between-finger stimulation paradigm were not included in univariate analyses, because we did not collect data from both modalities in each of the two participants. Task-based statistical parametric maps were computed separately for the tactile condition and nociceptive condition versus resting baseline. At group level, we used FLAME1+2 to test: (1) whether activations in the tactile condition were greater than baseline; (2) whether activations in the nociceptive condition were greater than baseline; (3) whether activations in the tactile condition were different than in the nociceptive condition. Z statistic images were thresholded non-parametrically using clusters determined by $Z > 2.3$ and a corrected cluster significance threshold of $p = 0.05$.

Phase-encoded analyses. The purpose of these analyses was to determine whether any parietal region showed a preferential response to the spatial frequency of the periodic finger stimulation; voxels sensitive to a given spatial frequency of stimulation respond maximally when the stimulus passes through the preferred location and decay as the stimulus moves away (Chen et al. 2017). Surface-based analyses were performed on unsmoothed data as previously described (Mancini et al. 2018) (see Supplemental Data for a detailed description of the analyses https://github.com/flamancini/parietalmaps/blob/master/supplemental_data_JNP19.pdf).

Results

Magnitude of cortical responses

At the group level, both tactile and nociceptive stimuli to the fingers activated clusters in S1, superior parietal lobule, and anterior intra-parietal cortex in the hemisphere contralateral to hand stimulation (Fig. 1, Supplemental Table I and II https://github.com/flamancini/parietalmaps/blob/master/supplemental_data_JNP19.pdf). Tactile stimuli significantly activated a S1 region in the hemisphere ipsilateral to hand stimulation (Fig. 1A). No brain region was activated more by nociceptive stimuli than tactile

stimuli. In contrast, tactile stimuli elicited greater responses in S1 than nociceptive stimuli, both in terms of response amplitude and spatial extent (Fig 1C). This difference is unlikely to be mediated by trivial factors, such as differences of perceived intensity or saliency of the two stimulus modalities. First, although S1 responses were lower for nociceptive stimuli than tactile stimuli, the average perceived intensity of pinprick pain was higher than the average perceived intensity of touch ($t_{23}=4.06$, $p=0.0004$; mean 'touch' intensity rating: 4.18, SD across participants: 1.03; mean 'pinprick pain' intensity rating: 6.24, SD across participants: 1.48). Second, the energy of the laser stimuli was set as the highest the participants agreed to endure for 3 runs of ~8 minutes of stimulation. Third, the laser energy was adjusted before each run to ensure that pain intensity remained at the same level of the first run (mean rating variability across runs: 0.38).

Somatotopic organization of cortical responses

First, we validated our 7T mapping protocols by replicating our previous findings of contralateral, between-finger maps of tactile and nociceptive input in S1 (Fig.2). As we already described (Mancini et al. 2012; Mancini et al. 2018), the between-finger map of both tactile and nociceptive stimuli progressed along the lateral-to-medial axis from index finger to little finger. Then, we went on searching for within-finger maps in S1 and IPS. At the group level (Fig.3A-C), we found mirrored maps of tactile input to finger segments across S1 subdivisions, in the hemisphere contralateral to hand stimulation. For reference, the estimated anatomical boundaries of S1 subdivisions were estimated using an automated and probabilistic surface-based parcellation. The ROIs were statistically thresholded and displayed on the inflated cortical surface of an average brain. The tactile finger map progressed, along the anterior-to-posterior axis, from base to fingertip in area 3a. The phase of the map reversed approximately at the border between area 3a and 3b, at the fingertip. We did not observe consistent phase reversals between areas 3b and 1, but we found a significant finger segments map in area 1, that progressed from tip to base of the fingers along the anterior-to-posterior axis (Fig.3B-C).

Nociceptive stimuli elicited a periodic response in areas 3b and 1 (Fig.3D-E), in a more ventro-lateral region than the one activated by tactile stimuli. The responses to nociceptive stimuli appeared to follow a similar spatial gradient than those to tactile stimuli, at group level. However, these activity clusters barely survived FDR correction (Fig.4F) due to large intra-individual variability. Therefore, we cannot exclude that these within-finger activations in S1 might be spurious. Nociceptive stimuli, but not tactile stimuli, elicited periodic responses in a region in the contralateral human intraparietal area 1 (hIP1) (Choi et al. 2006; Scheperjans et al. 2008), as shown in the group average maps (Fig.4A-C). The centroid of this nociceptive map was located at Talairach coordinates -43, -52, 37 (MNI coordinates -43, -55, 39). In the majority of subjects (Fig.4D), we found significant periodic responses to nociceptive stimulation of finger segments.

Discussion

Our study indicates that the somatotopic representation of nociceptive input in the parietal cortex differs from that of tactile input. In particular, we did not find robust evidence for the existence of within-finger maps of nociceptive-selective stimuli across S1 cytoarchitectonic

subdivisions, but we were able to successfully detect within-finger maps of tactile input in Brodmann areas 3a, 3b, and 1, in the hemisphere contralateral to hand stimulation (Fig.3). However, we confirmed the existence of between-finger maps of nociceptive input in the medial-to-lateral axis of S1 (Fig.2), in line with previous reports (Mancini et al. 2012; Mancini et al. 2018). Although nociceptive stimuli did not elicit consistent BOLD responses in contralateral (left) S1, they consistently activated areas in a region in the left intra-parietal area 1 (hIP1), which did not display a significant periodic response to tactile stimulation (Fig.3-4).

Within-finger maps of tactile input in S1

Only a few fMRI studies have investigated the within-finger somatotopy of mechanical input in the human S1 (Blankenburg et al. 2003; Overduin and Servos 2004; Sanchez-Panchuelo et al. 2012; Schweisfurth et al. 2011). To date, only one of these studies (Sanchez-Panchuelo et al. 2012) has been able to reliably identify the border of map reversals in individual subjects. In monkeys, the locations of reversals of within-finger somatotopic maps identified from electrophysiology and anatomy have consistently been found to coincide with cytoarchitectonic borders (Kaas et al. 1979; Merzenich et al. 1978; Nelson et al. 1980).

The within-fingers somatotopy of tactile input reported here is largely consistent with that reported by previous work; we found that the most anterior reversal was around the fingertips, at the estimated border between areas 3a and 3b (Fig. 4B-C). Although we did not observe, at the group level, a map reversal between areas 3b and 1, we found that the spatial arrangement of the tactile within-finger map in area 1 followed the one expected on the basis of previous work (Sanchez-Panchuelo et al. 2012) (Fig. 4B-C). At the group level, we did not observe a within-finger map in area 2.

Nociceptive responses in S1

It is clear from animal and human studies that a subset of S1 neurons responds to nociceptive input, but their functional relevance and spatial organization remains controversial (Vierck et al. 2013). Studies in primates support a possible nociceptive role for areas 3a and 1 (Kenshalo et al. 2000; Tommerdahl et al. 1996; Whitsel et al. 2009). In humans, nociceptive responses originating in areas 1–2 have been suggested from MEG and subdural EEG data (Inui et al. 2003; Kanda et al. 2000; Ogino et al. 2005; Ploner et al. 1999), but the involvement of area 3b in nociception remains controversial (Baumgartner et al. 2011; Kanda et al. 2000; Valeriani et al. 2004). Intracerebral recordings in the human S1 have shown that responses specific to nociceptive heat in area 3b were sparser and much less frequent than responses to non-nociceptive input (Frot et al. 2013). Our study suggests that the density of nociceptive units might be sparse or heterogeneously distributed in S1 along the anterior-to-posterior axis, and may account for weak and inconsistent within-finger representations.

Within-finger maps of nociceptive input in hIP1

Our study reveals a novel within-finger map of nociceptive input in hIP1 (Choi et al. 2006). The map was significant at both group and individual-subject level in the majority of participants (Fig.4). This sub-region of hIP1, or any other intraparietal subregion, was not significantly activated by tactile stimuli (Fig.3A). To our knowledge, no previous study has demonstrated the existence of within-finger maps of somatosensory input in the intraparietal

cortex, possibly because the parietal somatotopy of the hand is both spatially and temporally variable across individuals (Chen et al. 2017).

Nociceptive stimuli have frequently been reported to activate intra-parietal posterior parietal regions (Duerden and Albanese 2011; Tracey and Mantyh 2007) and these activations are usually interpreted as reflecting spatial attention (Lobanov et al. 2013; Oshiro et al. 2009; Oshiro et al. 2007; Peyron et al. 1999). It is possible that we were able to detect a within-finger map in hIP1 only in response to nociceptive input, but not tactile input, because nociceptive stimuli are inherently more salient (being aversive) than non-painful tactile stimuli (Legrain et al. 2011). Indeed, previous work has shown that the anterior intraparietal cortex is sensitive to bottom-up attention driven by stimulus saliency (Geng and Mangun 2009). Future studies are required to confirm the functional significance of the within-finger map of nociceptive input in hIP1.

Acknowledgments

This project has received funding from the Wellcome Trust (Strategic Award [102645/Z/13/Z](#)). The Wellcome Centre for Integrative Neuroimaging is supported by core funding from the Wellcome Trust ([203139/Z/16/Z](#)).

Figure legends

Figure 1. Cortical responses to tactile (**A**) and nociceptive stimuli (**B**), as quantified using GLM analyses. Panel **C** shows regions more activated by tactile stimuli than nociceptive stimuli, whereas panel **D** shows the overlay of cortical activity elicited by tactile and nociceptive stimuli. The displayed Z statistics were thresholded using clusters determined by $Z > 2.3$ and a corrected cluster significance threshold of $p = 0.05$.

Figure 2. Replication of previous evidence of between-finger maps of tactile stimuli (subject #13) and nociceptive stimuli (subject #14), as revealed by phase-encoded mapping analyses. The F-statistics of the signal at different phases of periodic finger stimulation (a travelling wave sweeping through the different fingers, from the index finger to the little finger) are rendered on the T1 volume and on the inflated cortical surface.

Figure 3. Surface-based, group average of cortical responses elicited by periodic within-finger tactile (**A-C**) and nociceptive (**D-F**) stimulation. The color-coding scheme used is shown on the top of the figure: red = proximal finger segments, blue = medial finger segments, green = distal finger segments. Panels **A**, **B**, **D** and **E** display the uncorrected F-statistics, thresholded at $p < 0.01$. Panels **C** and **F** show the F-statistics with an FDR-correction for p-values at level 0.05. Tactile stimuli, but not nociceptive stimuli, exhibit consistent within-finger maps in S1 subdivisions. The F-statistics is rendered onto an inflated average cortical surface (fsaverage) and overlaid the anatomical boundaries between areas 3a, 3b, 1 and 2 (estimated using Freesurfer's probabilistic surface-based parcellation and thresholded).

Figure 4. A-C. Surface-based, group average of within-finger maps of nociceptive stimuli ($n=9$) in human intraparietal area 1 (hIP1). The color-coding scheme is as follows: red = proximal finger segments, blue = medial finger segments, green = distal finger segments. Panels **A-B** display the uncorrected F-statistics, thresholded at $p < 0.01$, whereas panel **C** shows the F-statistics with an FDR-correction for p-values at level 0.05. For reference, in panel **C** the anatomical boundaries of area hIP1 are displayed (as derived from the probabilistic Juelich Histological Atlas and thresholded at 25%). **D.** Single-subject, within-finger maps of nociceptive input (one map per subject; the two subjects that did not show any significant response in this region are not displayed). The surface-based mask used to display the single-subject data corresponds to the boundaries of the group-average, within-finger map of nociceptive input displayed in panel **B**.

References

- Andersson JL, Lilja A, Hartvig P, Langstrom B, Gordh T, Handwerker H, and Torebjork E.** Somatotopic organization along the central sulcus, for pain localization in humans, as revealed by positron emission tomography. *Exp Brain Res* 117: 192-199, 1997.
- Baumgartner U, Vogel H, Ohara S, Treede RD, and Lenz F.** Dipole source analyses of laser evoked potentials obtained from subdural grid recordings from primary somatic sensory cortex. *J Neurophysiol* 106: 722-730, 2011.
- Bingel U, Lorenz J, Glauche V, Knab R, Glascher J, Weiller C, and Buchel C.** Somatotopic organization of human somatosensory cortices for pain: a single trial fMRI study. *NeuroImage* 23: 224-232, 2004.
- Blankenburg F, Ruben J, Meyer R, Schwiemann J, and Villringer A.** Evidence for a rostral-to-caudal somatotopic organization in human primary somatosensory cortex with mirror-reversal in areas 3b and 1. *Cerebral Cortex* 13: 987-993, 2003.
- Chen CF, Kreutz-Delgado K, Sereno MI, and Huang RS.** Validation of periodic fMRI signals in response to wearable tactile stimulation. *Neuroimage* 150: 99-111, 2017.
- Choi HJ, Zilles K, Mohlberg H, Schleicher A, Fink GR, Armstrong E, and Amunts K.** Cytoarchitectonic identification and probabilistic mapping of two distinct areas within the anterior ventral bank of the human intraparietal sulcus. *J Comp Neurol* 495: 53-69, 2006.
- Duerden EG, and Albanese MC.** Localization of pain-related brain activation: A meta-analysis of neuroimaging data. *Human Brain Mapping* 2011.
- Dum RP, Levinthal DJ, and Strick PL.** The spinothalamic system targets motor and sensory areas in the cerebral cortex of monkeys. *Journal of Neuroscience* 29: 14223-14235, 2009.
- Frot M, Magnin M, Mauguiere F, and Garcia-Larrea L.** Cortical representation of pain in primary sensory-motor areas (S1/M1)--a study using intracortical recordings in humans. *Hum Brain Mapp* 34: 2655-2668, 2013.
- Geng JJ, and Mangun GR.** Anterior intraparietal sulcus is sensitive to bottom-up attention driven by stimulus salience. *J Cogn Neurosci* 21: 1584-1601, 2009.
- Gingold SI, Greenspan JD, and Apkarian AV.** Anatomic evidence of nociceptive inputs to primary somatosensory cortex: relationship between spinothalamic terminals and thalamocortical cells in squirrel monkeys. *J Comp Neurol* 308: 467-490, 1991.
- Inui K, Wang X, Qiu Y, Nguyen BT, Ojima S, Tamura Y, Nakata H, Wasaka T, Tran TD, and Kakigi R.** Pain processing within the primary somatosensory cortex in humans. *Eur J Neurosci* 18: 2859-2866, 2003.
- Kaas JH, Nelson RJ, Sur M, Lin CS, and Merzenich MM.** Multiple representations of the body within the primary somatosensory cortex of primates. *Science* 204: 521-523, 1979.
- Kanda M, Nagamine T, Ikeda A, Ohara S, Kunieda T, Fujiwara N, Yazawa S, Sawamoto N, Matsumoto R, Taki W, and Shibasaki H.** Primary somatosensory cortex is actively involved in pain processing in human. *Brain Res* 853: 282-289, 2000.
- Kenshalo DR, Iwata K, Sholas M, and Thomas DA.** Response properties and organization of nociceptive neurons in area 1 of monkey primary somatosensory cortex. *J Neurophysiol* 84: 719-729, 2000.
- Kenshalo DR, Jr., Chudler EH, Anton F, and Dubner R.** SI nociceptive neurons participate in the encoding process by which monkeys perceive the intensity of noxious thermal stimulation. *Brain Res* 454: 378-382, 1988.
- Kenshalo DR, Jr., and Isensee O.** Responses of primate SI cortical neurons to noxious stimuli. *Journal of Neurophysiology* 50: 1479-1496, 1983.

396 **Kolasinski J, Makin TR, Jbabdi S, Clare S, Stagg CJ, and Johansen-Berg H.**
397 Investigating the Stability of Fine-Grain Digit Somatotopy in Individual Human Participants. *J*
398 *Neurosci* 36: 1113-1127, 2016a.
399 **Kolasinski J, Makin TR, Logan JP, Jbabdi S, Clare S, Stagg CJ, and Johansen-Berg H.**
400 Perceptually relevant remapping of human somatotopy in 24 hours. *Elife* 5: 2016b.
401 **Legrain V, Iannetti GD, Plaghki L, and Mouraux A.** The pain matrix reloaded: a salience
402 detection system for the body. *Progress in Neurobiology* 93: 111-124, 2011.
403 **Lobanov OV, Quevedo AS, Hadsel MS, Kraft RA, and Coghill RC.** Frontoparietal
404 mechanisms supporting attention to location and intensity of painful stimuli. *Pain* 154: 1758-
405 1768, 2013.
406 **Mancini F, Haggard P, Iannetti GD, Longo MR, and Sereno MI.** Fine-grained nociceptive
407 maps in primary somatosensory cortex. *Journal of Neuroscience* 32: 17155-17162, 2012.
408 **Mancini F, Wang AP, Shira MM, Isherwood ZJ, McAuley JH, Iannetti G, Sereno MI,**
409 **Moseley L, and Rae CD.** Preserved cortical maps of the body in Complex Regional Pain
410 Syndrome. *bioRxiv* 2018.
411 **Merzenich MM, Kaas JH, Sur M, and Lin CS.** Double representation of the body surface
412 within cytoarchitectonic areas 3b and 1 in "SI" in the owl monkey (*Aotus trivirgatus*). *J Comp*
413 *Neurol* 181: 41-73, 1978.
414 **Nelson RJ, Sur M, Felleman DJ, and Kaas JH.** Representations of the body surface in
415 postcentral parietal cortex of *Macaca fascicularis*. *J Comp Neurol* 192: 611-643, 1980.
416 **Ogino Y, Nemoto H, and Goto F.** Somatotopy in human primary somatosensory cortex in
417 pain system. *Anesthesiology* 103: 821-827, 2005.
418 **Oshiro Y, Quevedo AS, McHaffie JG, Kraft RA, and Coghill RC.** Brain Mechanisms
419 Supporting Discrimination of Sensory Features of Pain: A New Model. *The Journal of*
420 *Neuroscience* 29: 14924-14931, 2009.
421 **Oshiro Y, Quevedo AS, McHaffie JG, Kraft RA, and Coghill RC.** Brain mechanisms
422 supporting spatial discrimination of pain. *The Journal of neuroscience : the official journal of*
423 *the Society for Neuroscience* 27: 3388-3394, 2007.
424 **Overduin SA, and Servos P.** Distributed digit somatotopy in primary somatosensory cortex.
425 *NeuroImage* 23: 462-472, 2004.
426 **Peyron R, Garcia-Larrea L, Gregoire MC, Costes N, Convers P, Lavenne F, Mauguiere**
427 **F, Michel D, and Laurent B.** Haemodynamic brain responses to acute pain in humans:
428 sensory and attentional networks. *Brain* 122 (Pt 9): 1765-1780, 1999.
429 **Ploner M, Schmitz F, Freund HJ, and Schnitzler A.** Parallel activation of primary and
430 secondary somatosensory cortices in human pain processing. *J Neurophysiol* 81: 3100-
431 3104, 1999.
432 **Sanchez-Panchuelo RM, Besle J, Beckett A, Bowtell R, Schluppeck D, and Francis S.**
433 Within-digit functional parcellation of Brodmann areas of the human primary somatosensory
434 cortex using functional magnetic resonance imaging at 7 tesla. *J Neurosci* 32: 15815-15822,
435 2012.
436 **Scheperjans F, Eickhoff SB, Homke L, Mohlberg H, Hermann K, Amunts K, and Zilles**
437 **K.** Probabilistic maps, morphometry, and variability of cytoarchitectonic areas in the human
438 superior parietal cortex. *Cereb Cortex* 18: 2141-2157, 2008.
439 **Schmahmann JD, and Pandya DN.** Anatomical investigation of projections from thalamus
440 to posterior parietal cortex in the rhesus monkey: a WGA-HRP and fluorescent tracer study.
441 *J Comp Neurol* 295: 299-326, 1990.

442 **Schweisfurth MA, Schweizer R, and Frahm J.** Functional MRI indicates consistent intra-
443 digit topographic maps in the little but not the index finger within the human primary
444 somatosensory cortex. *Neuroimage* 56: 2138-2143, 2011.

445 **Sereno MI, and Huang RS.** Multisensory maps in parietal cortex. *Curr Opin Neurobiol* 24:
446 39-46, 2014.

447 **Tommerdahl M, Delemos KA, Vierck CJ, Jr., Favorov OV, and Whitsel BL.** Anterior
448 parietal cortical response to tactile and skin-heating stimuli applied to the same skin site. *J*
449 *Neurophysiol* 75: 2662-2670, 1996.

450 **Tracey I, and Mantyh PW.** The cerebral signature for pain perception and its modulation.
451 *Neuron* 55: 377-391, 2007.

452 **Valeriani M, Barba C, Le Pera D, Restuccia D, Colicchio G, Tonali P, Gagliardo O, and**
453 **Treede RD.** Different neuronal contribution to N20 somatosensory evoked potential and to
454 CO2 laser evoked potentials: an intracerebral recording study. *Clin Neurophysiol* 115: 211-
455 216, 2004.

456 **Vierck CJ, Whitsel BL, Favorov OV, Brown AW, and Tommerdahl M.** Role of primary
457 somatosensory cortex in the coding of pain. *Pain* 154: 334-344, 2013.

458 **Whitsel BL, Favorov OV, Li Y, Quibrera M, and Tommerdahl M.** Area 3a neuron
459 response to skin nociceptor afferent drive. *Cereb Cortex* 19: 349-366, 2009.

460

Tactile stimuli

Nociceptive stimuli

A

D

B

E

C

F