

Raptor genomes reveal evolutionary signatures of predatory and nocturnal lifestyles

Yun Sung Cho^{1†}, JeHoon Jun^{1†}, Jung A Kim², Hak-Min Kim^{3,4}, Oksung Chung¹, Seung-Gu Kang⁵, Jin-Young Park⁵, Hwa-Jung Kim⁵, Sunghyun Kim⁶, Hee-Jong Kim⁷, Jin-ho Jang⁷, Ki-Jeong Na⁸, Jeongho Kim⁹, Seung Gu Park³, Hwang-Yeol Lee¹, Andrea Manica¹⁰, David P. Mindell¹¹, Jérôme Fuchs¹², Jeremy S. Edwards¹³, Jessica A. Weber¹⁴, Christopher C. Witt¹⁴, Joo-Hong Yeo², Soonok Kim^{2*} and Jong Bhak^{1,3,4*}

¹Clinomics Inc., Ulsan, Republic of Korea.

²Biological and Genetic Resources Assessment Division, National Institute of Biological Resources, Incheon, Republic of Korea.

³Korean Genomics Industrialization Center (KOGIC), Ulsan National Institute of Science and Technology (UNIST), Ulsan, Republic of Korea

⁴Department of Biomedical Engineering, School of Life Sciences, Ulsan National Institute of Science and Technology (UNIST), Republic of Korea.

⁵Animal Resources Division, National Institute of Biological Resources, Incheon, Republic of Korea.

⁶Strategic Planning Division, National Institute of Biological Resources, Incheon, Republic of Korea.

⁷Chungnam Wild Animal Rescue Center, Kongju National University, Yesan, Republic of Korea.

⁸College of veterinary medicine, Chungbuk National University, Cheongju, Republic of Korea.

⁹Medical care team, Cheongju Zoo, Cheongju, Republic of Korea.

¹⁰Department of Zoology, University of Cambridge, Cambridge, UK.

¹¹Museum of Vertebrate Zoology, University of California, Berkeley, California, USA.

¹²Institut Systématique Evolution Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Paris, France.

¹³Chemistry and Chemical Biology, UNM Comprehensive Cancer Center, University of New Mexico, Albuquerque, New Mexico, USA.

¹⁴Museum of Southwestern Biology and Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA.*

Correspondence: sokim90@korea.kr; jongbhak@genomics.org

† Yun Sung Cho and JeHoon Jun contributed equally to this work.

Email addresses:

YSC: joys0406@gmail.com
JeHoonJ: junjh0701@gmail.com
JAK: jakim21@korea.kr
HMK: howmany2@gmail.com
OC: okokookk219@gmail.com
SGK: pfalcon@hanmail.net
JYP: birdkorea@korea.kr
Hwa-JungK: hwajung@korea.kr
SunghyunK: birdksh@korea.kr
Hee-JongK: heejjong@hanmail.net
Jin-hoJ: maverick57@hanmail.net
KJN: sigol@cbnu.ac.kr
JK: in-africa@korea.kr
SGP: seung9park@gmail.com
HYL: hyeol911@gmail.com
AM: am315@cam.ac.uk
DPM: dpmindell@gmail.com
JF: jeromefuchs@gmail.com
JSE: jeremy.scott.edwards@gmail.com
JAW: weberj.unm@gmail.com
CCW: cwitt@unm.edu
JHY: y1208@korea.kr
SoonokK: sokim90@korea.kr
JB: jongbhak@gmail.com

1 **Abstract**

2 **Background:** Birds of prey (raptors) are dominant apex predators in terrestrial communities,
3 with hawks (Accipitriformes) and falcons (Falconiformes) hunting by day, and owls
4 (Strigiformes) hunting by night.

5 **Results:** Here, we report new genomes and transcriptomes for 20 species of birds, including 16
6 species of birds of prey, and high-quality reference genomes for the Eurasian eagle-owl (*Bubo*
7 *bubo*), oriental scops-owl (*Otus sunia*), eastern buzzard (*Buteo japonicus*), and common kestrel
8 (*Falco tinnunculus*). Our extensive genomic analysis and comparisons with non-raptor genomes
9 identified common molecular signatures that underpin anatomical structure and sensory, muscle,
10 circulatory, and respiratory systems related to a predatory lifestyle. Compared with diurnal birds,
11 owls exhibit striking adaptations to the nocturnal environment, including functional trade-offs in
12 the sensory systems (e.g., loss of color vision genes and selection for enhancement of nocturnal
13 vision and other sensory systems) that are probably convergent with other nocturnal avian orders.
14 Additionally, we found that a suite of genes associated with vision and circadian rhythm were
15 differentially expressed between nocturnal and diurnal raptors, indicating adaptive expression
16 change during the transition to nocturnality.

17 **Conclusions:** Overall, raptor genomes showed genomic signatures associated with the origin and
18 maintenance of several specialized physiological and morphological features essential to be apex
19 predators.

20 **Keywords:** Raptor, *De novo* assembly, Comparative genomics, Evolutionary adaptation,
21 Predatory lifestyle, Nocturnality

1 **Background**

2 Birds of prey, also known as raptors, are key apex predators in nearly every terrestrial biotic
3 community. Species in this guild comprise a non-monophyletic set of three orders within the
4 core landbirds clade, and recent large-scale phylogenomic studies have led to the suggestion that
5 the common ancestor of this clade may have been a predator [1]. There are three main orders of
6 birds of prey: Strigiformes (true and barn owls), Falconiformes (falcons and caracaras), and
7 Accipitriformes (eagles, buzzards, hawks, kites, and vultures). Species in each of these three
8 raptor clades are obligate predators with adaptations for hunting, killing, and/or eating meat [2,
9 3]. Additionally, the common ancestor of owls evolved nocturnality, and most extant owl species
10 are nocturnal, a habit they share with two other avian orders for which we have genome
11 sequences (Caprimulgiformes and Apterygiformes). These independent transitions in lifestyle
12 provide an opportunity to test for patterns of genome evolution that are linked with being
13 raptorial and nocturnal, respectively [3-5].

14 Genomes have been published for more than 50 avian species, including nine birds of
15 prey (peregrine and saker falcons, bald, white-tailed, and golden eagles, turkey vulture, barn owl,
16 northern spotted owl, and burrowing owl) [3, 6-9]. However, the barn owl, white-tailed eagle,
17 and turkey vulture genomes were assembled at low-quality [6], and a detailed comparative
18 evolutionary analysis was performed only for the falcons [3]. Here, we report new high-quality
19 whole genome reference sequences of four raptor species (Eurasian eagle-owl [*Bubo bubo*] and
20 oriental scops-owl [*Otus sunia*] in Strigiformes, eastern buzzard [*Buteo japonicus*] in
21 Accipitriformes, and common kestrel [*Falco tinnunculus*] in Falconiformes) with a set of raptor
22 whole-genome and transcriptome data, extending the genomic coverage of raptors (Fig. 1,
23 Additional file 1: Figure S1 and Tables S1-S3). Our investigation revealed numerous genomic

1 signatures of evolution that are shared among the three raptor orders or that appear to be
2 associated with nocturnal adaptations of owls.

3

4 **Results and discussion**

5 **Raptor genome sequencing and assembly**

6 We applied whole-genome shotgun sequencing and *de novo* assembly strategies [10-12] to build
7 reference genomes of the four raptor species (Eurasian eagle-owl, oriental scops-owl, eastern
8 buzzard, and common kestrel). The extracted DNA samples from wild individuals were
9 sequenced at high coverage (>185×) using various insert sizes of short-insert and long-mate pair
10 libraries (Additional file 1: Tables S4 and S5). The four raptor genomes showed relatively higher
11 levels of genomic diversity compared to the previously assembled genomes of eagles and falcons
12 (Additional file 1: Figures S2 and S3). By assembling in various conditions and evaluating
13 assembly quality, we obtained the four raptor reference genomes at a high-quality, resulting in
14 scaffold N50 sizes from 7.49 to 29.92 Mb (Additional file 1: Tables S6-S9). Protein-coding
15 genes (~16,000 to 18,000 genes) for these four species were predicted by combining *de novo* and
16 homologous gene prediction methods with whole blood transcriptome data (Additional file 1:
17 Table S10). Roughly 9.2% of the raptor genomes were predicted as transposable elements
18 (Additional file 1: Table S11), consistent with the composition of other avian genomes [6].
19 Additionally, we sequenced the whole genome and blood transcriptome from another twelve
20 raptors (five owls, six accipitrids, and a falconid) and four non-raptor birds (Additional file 1:
21 Tables S12-S14), most of which were sequenced for the first time.

22

1 **Evolutionary analysis of raptors compared to non-raptor birds**

2 To identify the genetic basis of predation and nocturnality in raptors, we performed in-depth
3 comparative evolutionary analyses for 25 birds of prey (including ten nocturnal owls and 15
4 diurnal raptors) and 23 non-raptor bird species (including nocturnal brown kiwi [13] and chuck-
5 will's-widow [6], and other avian representatives genome-assembled at a high-quality; Fig. 2,
6 Additional file 1: Figure S4 and Tables S1, S2, and S15). Birds have evolved to employ many
7 different strategies to obtain food, and raptors are specialized for hunting [2, 3, 7]. Several
8 molecular signatures were shared by the three raptor orders, and the ancestral branches of these
9 orders each showed an expansion of gene families associated with regulation of anatomical
10 structure size, embryonic appendage morphogenesis, regulation of responses to stimulus and
11 wounding, and learning or memory functions ($P < 0.05$, Fisher's exact test; Additional file 1:
12 Tables S16 and S17). When comparing gene family sizes between the extant species, immune
13 system associated gene families were expanded in the birds of prey (Additional file 1: Table
14 S18).

15 To further examine the shared evolutionary adaptations related to avian predatory
16 lifestyles, we identified selection signatures shared by the three orders of birds of prey at the
17 gene sequence level; which possibly reflects their shared requirement for highly-developed
18 sensory systems, efficient circulatory and respiratory systems, and exceptional flight capabilities
19 necessary to capture prey [2-5, 7, 8]. Based upon d_N/d_S ratio calculation [14, 15], only *RHCE* and
20 *CENPQ* genes were commonly found as positively selected genes (PSGs) in the three raptor
21 ancestral branches of the Strigiformes, Accipitriformes, and Falconiformes (Additional file 2:
22 Datasheets S1-S3); consistent with the results from mammals demonstrating that adaptive
23 molecular convergence linked to phenotypic convergence is rare [12, 16]. In addition, we

1 identified three genes as positively selected in the ancestral branches of two raptor orders
2 (*SFTPA1* in the Strigiformes and Falconiformes; *TFF2* and *PARL* in the Strigiformes and
3 Accipitriformes). A lung surfactant protein encoded by *SFTPA1* play an essential role in the
4 defense against respiratory pathogens and normal respiration [17]. *TFF2* gene encodes a protein
5 that mediate gastric wound repair and inhibit gastric acid secretion [18]. Finally, we found that
6 148 genes showed accelerated d_N/d_S in the raptor ancestral branches (Additional file 1: Table
7 S19). Of these, *SLC24A1*, *NDUFS3*, and *PPARA* encode proteins that play roles in visual
8 transduction cascade, mitochondrial membrane respiratory chain, and lipid metabolism,
9 respectively [17, 19, 20]. Out of 50 collected beak development associated genes, 17, 17, and 18
10 genes (34 to 36%) showed accelerated d_N/d_S in the ancestral branches of Strigiformes,
11 Accipitriformes, and Falconiformes, respectively (Additional file 1: Table S20), hinting at
12 adaptation for enhanced beaks for killing and flesh-tearing [2, 3]. Of these, four genes (*BMP10*,
13 *GDF9*, *NABI*, and *TRIP11*) showed common acceleration signatures in the three raptor orders.

14 It has been suggested that genes with elevated frequencies of Guanine-Cytosine at the
15 third codon position (GC3) are more adaptable to external stresses, through providing more
16 targets for *de novo* methylation that affect the variability of gene expression [23]. Therefore, we
17 analyzed the GC3 content in the three raptor orders, and we found that regulation of nervous
18 system development, central nervous system neuron differentiation, and locomotion associated
19 genes showed high GC3 bias (Fig. 2c, Additional file 1: Figure S5, Table S21 and Additional file
20 2: Datasheet S6). In the highly conserved genomic regions (HCRs) among species belonging to
21 the same order [12], 79 functional categories were commonly enriched in the three raptor orders
22 (Additional file 1: Tables S22-S31). Among these categories, eye, sensory organ, muscle organ,
23 epithelium, and limb development functions were commonly conserved in the three raptor orders,

1 but not in Passeriformes (a control avian order in this analysis), suggesting that those functions
2 are important in raptors for their predatory lifestyle.

3

4 **Evolutionary analysis of nocturnal birds compared to diurnal birds**

5 Since several avian clades have adapted to a nocturnal lifestyle independently, the comparative
6 method can be used to identify genes underlying convergent phenotypes that are associated with
7 nocturnal adaptation [5]. Three nocturnal bird groups (the ancestral branch of owls, chuck-will's-
8 widow, and brown kiwi) shared an expansion of gene families associated with synapse
9 organization, cellular response to stimulus, and bile secretion functions ($P < 0.05$; Additional file
10 1: Tables S32, S33). As expected, gene families associated with vision were commonly
11 contracted in the nocturnal birds (Additional file 1: Tables S34 and S35). Specifically, gene loss
12 of the violet/ultraviolet-sensitive opsin *SWS1* (*OPN1SW*) was found in all of the nocturnal bird
13 genomes, as previously reported [4, 22]. The nocturnal birds also showed common selection
14 signatures likely linked to their adaptation to a nocturnal environment. A total 14 PSGs were
15 shared among the three nocturnal groups, and 98 PSGs were shared by at least two nocturnal bird
16 groups (Additional file 2: Datasheets S1, S4 and S5). The shared PSGs were over-represented in
17 detection of mechanical stimulus involved in sensory perception of sound, wound healing, and
18 skin development functions (Additional file 1: Table S36), although the enrichment did not pass
19 the false discovery rate criterion. Interestingly, at least one of two wound healing associated
20 genes (*TFF2* and *COL3A1*) [23, 24] was found to be positively selected in the nocturnal birds.
21 Additionally, six genes (*RHO*, *BEST1*, *PDE6B*, *RPE65*, *OPN4-1*, and *RRH*) involved in light
22 detection, and *RDH8* that is involved in retinol (vitamin A₁) metabolism [17, 25], showed

1 accelerated d_N/d_S in the nocturnal birds (Additional file 1: Table S37). It is well-known that
2 rhodopsin encoded by *RHO* is a light-sensitive receptor and thus enables vision in low-light
3 conditions [26]. Notably, *RHO* also showed a high level of GC3 biases in the nocturnal birds
4 (Additional file 2: Datasheet S7). Furthermore, *RPE65* encodes a protein that is a component of
5 the vitamin A visual cycle of the retina, while *PDE6B* plays a key role in the phototransduction
6 cascade and mutations in this gene result in congenital stationary night blindness. In addition,
7 melanopsin encoded by *OPN4-1* is a photoreceptor required for regulation of circadian rhythm
8 [17, 25]. We also found that only *SLC51A* gene possesses specific amino acid sequences to the
9 nocturnal birds (Additional file 1: Figure S6). *SLC51A*, also known as *OST- α* , is essential for
10 intestinal bile acid transport [27], and it has been suggested that the bile acids affect the circadian
11 rhythms by regulating the expression level of circadian clock associated gene families [28, 29].
12 Interestingly, burrowing owl (*Athene cunicularia*), which is known as one of diurnal/crepuscular
13 owls, showed a different sequence alteration pattern from the other nocturnal or diurnal birds in
14 *SLC51A* locus.

15

16 **Sensory adaptations to nocturnal environment**

17 Modifications of the major sensory systems (not only vision, but also olfaction, hearing, and
18 circadian rhythm) are among the most common changes that occur when shifting from a diurnal
19 to a nocturnal lifestyle [5]. Analysis of the major sensory systems in the nocturnal bird genomes
20 revealed evidence of highly developed senses for adaptation to nocturnality. First, vision system
21 associated genes showed significantly accelerated d_N/d_S in the three nocturnal birds compared to
22 diurnal birds ($P < 0.05$, Mann-Whitney U test; Fig. 3). Owls and chuck-will's-widow

1 (Caprimulgiformes) had the highest acceleration in vision-related genes. The total number of
2 functional olfactory receptors (ORs) was not larger in the nocturnal birds than that in the diurnal
3 birds. However, the numbers of γ -clade ORs in the nocturnal birds and γ -c-clade ORs in the owls
4 were significantly larger than others (after excluding two outlier species showing extensive γ -c-
5 clade ORs expansion, chicken and zebra finch; $P < 0.05$, Mann-Whitney U test; Fig. 3 and
6 Additional file 1: Table S38). The diversity of ORs is thought to be related to a detection range
7 of odors [30], and we found that the diversity of α -clade ORs was significantly higher in the
8 nocturnal birds (Additional file 1: Table S39). Additionally, the diversity in the γ -c-clade ORs
9 was much higher in the owls and brown kiwi (Apterygiformes) compared to their sister groups
10 (downy woodpecker in Piciformes and common ostrich in Struthioniformes, respectively),
11 suggesting that increased olfactory abilities evolved repeatedly under nocturnal conditions [5,
12 13]. Hearing system associated genes showed a relatively high-level of d_N/d_S ratio in the owls
13 and brown kiwi; interestingly, two vocal learning species (budgerigar in Psittaciformes and
14 Anna's hummingbird in Apodiformes) had the first and third most accelerated d_N/d_S for hearing
15 associated genes, which may be linked with their highly developed cognitive abilities [31, 32].
16 Circadian rhythm associated genes showed the first and second largest acceleration in the owls
17 and brown kiwi, but the lowest in chuck-will's-widow, suggesting that these independent
18 instances of adaptation to nocturnality occurred by different mechanisms [5]. Additionally, we
19 found that 33 hearing system and 18 circadian rhythm associated genes showed accelerated d_N/d_S
20 in the three nocturnal bird groups (Additional file 1: Table S40). Considered together, these
21 results suggest that selection to augment nocturnal vision and other sensory systems predictably
22 compensates for loss of color vision, supporting a functional trade-off of sensory systems in
23 nocturnal birds [4, 5, 13].

1 Changes in gene expression are thought to underlie many of the phenotypic differences
2 between species [33]. Therefore, we carried out cross-species comparison of gene expression
3 among the blood transcriptomes from 13 raptors (five owls, four accipitrids, and four falconids)
4 and five non-raptor birds. We found that several vision-associated genes [17, 25] were
5 differentially expressed in the raptor orders ($P < 0.05$, moderated t-test; Additional file 1: Figures.
6 S7 and S8, and Additional file 2: Datasheets S8-11). For example, *PDCL* (lowly-expressed) and
7 *WFS1* (highly-expressed) genes were differentially expressed specific to the owls. Interestingly,
8 we could also find several circadian rhythm-related genes that were differentially expressed
9 between the nocturnal and diurnal raptors. Three circadian rhythm-associated genes (*ATF4*,
10 *PER3*, and *NR1P1*) were lowly expressed and two genes (*BTBD9* and *SETX*) were highly
11 expressed in the owls, whereas *ATF4* and *SIRT1* in the falconids and *NR1P1* in the accipitrids
12 were highly expressed. These results likely indicate that selectively driven expression switches
13 contributed to nocturnal adaptation of owls [33].

14

15 **Conclusions**

16 Our study provides whole genome assemblies of Eurasian eagle-owl, oriental scops-owl, eastern
17 buzzard, and common kestrel, as well as a suite of whole-genome resequencing and
18 transcriptome data from birds of prey. This is the first in-depth genomics study comparing the
19 three raptor orders, and we identified a number of shared molecular adaptations associated with a
20 predatory lifestyle. Furthermore, compared with diurnal birds, owls and other nocturnal birds
21 showed distinct genomic features, especially in sensory systems. While functional studies of
22 candidate genes will be needed to understand the molecular mechanisms of adaptation, these

- 1 results provide a genome-wide description and gene candidates of adaptations that have allowed
- 2 each of these three raptor groups to evolve into diverse, ecologically dominant apex predators.
- 3

1 **Methods**

2 **Sample and genome sequencing**

3 All blood samples used for genome and transcriptome sequencing were collected from
4 individuals being euthanized during wound treatment of rescued animals, except blood samples
5 of *A. flammeus*, *O. semitorques*, and *P. ptilorhynchus* that were obtained from the live
6 individuals during medical check-up at the wildlife rescue center. Muscle tissues samples
7 collected in 2017 were obtained from the fresh carcasses.

8 To build reference genome assemblies of the four raptor species (Eurasian eagle-owl,
9 oriental scops-owl, eastern buzzard, and common kestrel), we constructed eleven genomic
10 libraries with various insert sizes (Illumina short insert and long mate-pair libraries) for each
11 species, according to the manufacturer's protocol. The libraries were sequenced using Illumina
12 HiSeq platforms. The remaining twelve raptor and four non-raptor bird samples were re-
13 sequenced using Illumina HiSeq platforms with a short-insert libraries. Blood transcriptomes of
14 ten raptors and four non-raptor birds were sequenced using Illumina HiSeq platforms according
15 to the manufacturer's instructions.

16

17 **Genome assembly and annotation**

18 To assemble the raptor genomes, PCR duplicated, sequencing and junction adaptor contaminated,
19 and low quality (Q20) reads were filtered out. The short-insert and long-mate library reads were
20 trimmed into 90bp and 50bp, respectively to remove low-quality bases at the ends of the reads.
21 The quality-filtered reads were used to assemble the four raptor genomes using the
22 SOAPdenovo2 software [10]. We applied various *K*-mer values (33, 43, 53, and 63) to obtain
23 fragments with long contiguity. In this process, oriental scops-owl genome was assembled poorly

1 when using SOAPdenovo2, probably because of its high level of genomic heterozygosity.
2 Therefore, we also assembled the four raptor genomes using Platanus assembler, which is more
3 efficient for highly heterozygous genomes [11]. To reduce the number of gaps in the scaffolds,
4 we closed the gaps using the short-insert library reads in two iterations. To correct base-pair level
5 errors, we performed two iterations of aligning the short-insert library reads to the gap-closed
6 scaffolds using BWA-MEM [34] and calling variants using SAMtools [35]. In this process,
7 homozygous variants were assumed as erroneous sequences from the assembly process, and thus
8 substituted for the correction purpose.

9 To select final high-quality reference assemblies for the four raptors, we annotated all
10 assemblies and evaluated quality of each assembly. We first searched the genomes for tandem
11 repeats and transposable elements using Tandem Repeats Finder (version 4.07b) [36], Repbase
12 (version 19.03) [37], RepeatMasker (version 4.0.5) [38], RMBlast (version 2.2.28) [39], and
13 RepeatModeler (version 1.0.7) [40]. The protein-coding genes were predicted by combining *de*
14 *novo* and homology-based gene prediction methods with the blood transcriptome data for each
15 assembly. For the homology-based gene prediction, we searched for avian protein sequences
16 from the NCBI database using TblastN (version 2.2.26) [41] with an *E*-value cutoff of 1E-5. The
17 matched sequences were clustered using GenBlastA (version 1.0.4) [42] and filtered by coverage
18 and identity of >40% criterion. Gene models were predicted using Exonerate (version 2.2.0) [43].
19 For the *de novo* gene prediction, AUGUSTUS (version 3.0.3) [44] was used with the blood
20 transcriptome for each species. We filtered out possible pseudogenes having premature stop-
21 codons and single exon genes that were likely to be derived from retro-transposition. The
22 assembly and gene annotation qualities were assessed by aligning independently *de novo*
23 assembled transcripts using the Trinity software [45] and by searching for evolutionary

1 conserved orthologs using BUSCO software [46]. By considering the assembly statistics (e.g.,
2 N50 values and assembled sequence length) and the completeness of the genome assembly, final
3 high-quality reference assemblies for the four raptors were obtained. Genome, transcriptome, and
4 protein sequences for other comparison species were downloaded from the NCBI database.
5 Genes with possible premature stop-codons were excluded in the comparative analyses. The
6 northern-spotted owl's genome and protein sequences were acquired from the Zenodo linked in
7 the published paper [8].

8

9 **Comparative evolutionary analyses**

10 Orthologous gene families were constructed for avian genomes using the OrthoMCL 2.0.9
11 software [47]. To estimate divergence times of the 25 avian representatives, protein sequences of
12 the avian single-copy gene families were aligned using the MUSCLE program [48]. The poorly
13 aligned regions from the alignments were trimmed using the trimAl software [49]. The
14 divergence times were estimated using the MEGA7 program [50] with the phylogenetic tree
15 topology of published previous studies [1, 6] and the TimeTree database [51]. The date of the
16 node between chicken and rock dove was constrained to 98 million years ago (MYA), chicken
17 and brown kiwi was constrained to 111 MYA and common ostrich and brown kiwi was
18 constrained to 50-105 according to the divergence times from TimeTree. To estimate divergence
19 times among the birds of prey, the date of the node between downy woodpecker and Eurasian
20 eagle-owl constrained to 61-78 MYA and common kestrel and budgerigar was constrained to 60-
21 80 MYA according to the divergence times from the previous studies [1, 6] and TimeTree. A
22 gene family expansion and contraction analysis for the ancestral branches of the three bird of
23 prey orders was conducted using the CAFÉ program [52] with a $P < 0.05$ criterion. The

1 significantly different gene family sizes of the present species were identified by performing the
2 Mann-Whitney U test ($P < 0.05$).

3 To identify selection at the gene sequence level, two orthologous gene sets were
4 compiled, as previously reported [3]: the single-copy orthologs among avian species and
5 representative genes from multiple-copy orthologs. The representative genes from multiple-copy
6 orthologs were selected, if all species' protein sequences are reciprocally best matched to a
7 chicken protein sequence using BLASTp with an E -value cutoff of $1E-5$. PRANK [53] was used
8 to construct multiple sequence alignments among the orthologs. The CODEML program in
9 PAML 4.5 was used to estimate the d_N/d_S ratio (non-synonymous substitutions per non-
10 synonymous site to synonymous substitutions per synonymous site) [14]. The one-ratio model
11 was used to estimate the general selective pressure acting among comparison species. The two-
12 ratio model (model=2) was used to ensure that the d_N/d_S ratio is difference between foreground
13 species (raptors and nocturnal birds, respectively) and other species. Additionally, the d_N/d_S
14 ratios for each order-level branch of raptors and nocturnal birds were used to confirm if the
15 foreground d_N/d_S ratio is not biased to a specific raptor and nocturnal bird order. The branch-site
16 test was also conducted [15]. Statistical significance was assessed using likelihood ratio tests
17 with a conservative 10% false discovery rate criterion.

18 We identified target species-specific amino acid sequences. To filter out biases derived
19 from individual-specific variants, we used all of the raptor re-sequencing data by mapping to the
20 Eurasian eagle-owl genome for Strigiformes, the eastern buzzard genome for Accipitriformes,
21 and the common kestrel genome for Falconiformes. The mapping was conducted using BWA-
22 MEM, and consensus sequences were generated using SAMtools with the default options, except
23 the “-d 5” option. When we identified the specific amino acid sequences, protein sequences of

1 other birds from the NCBI database were also compared. We also checked multiple-sequence
2 alignments manually to remove artifacts. To identify genetic diversity based on heterozygous
3 SNV rates, variants were also called using Sentieon pipeline [54] with the default options, except
4 the "--algo Genotyper" option. The heterozygous SNV rates were calculated by dividing the total
5 number of heterozygous SNVs by the length of sufficiently mapped (>5 depth) genomic regions.

6 To identify HCRs in the three raptor orders and Passeriformes, we scanned genomic
7 regions that show significantly reduced genetic variation by comparing variations of each
8 window and whole genome as previously suggested [12]. In the case of Passeriformes, whole
9 genome data of four Passeriformes species (medium ground-finch, white-throated sparrow,
10 common canary, and collared flycatcher) were mapped to the zebra finch genome assembly, and
11 then variants were identified using the same methods used for the three raptor orders. Genetic
12 variation was estimated by calculating the number of different bases in the same order genomes
13 within each 100 Kb window. *P*-value was calculated by performing Fisher's exact test to test
14 whether the genetic variation of each window is significantly different from that of whole
15 genome. Only adjusted *P*-values (*q*-values) [55] of <0.0001 were considered significant. The
16 middle 10 Kb of each significantly different window were considered as HCRs.

17 For functional enrichment tests of candidate genes, GO annotations of chicken, zebra
18 finch, turkey, flycatcher, duck, anole lizard, and human genomes were downloaded from the
19 Ensembl database [56] and used to assign the avian protein-coding genes with GO categories. A
20 KEGG pathway was assigned using KAAS [57]. Functional information of candidate genes was
21 retrieved from the GO, KEGG, UniProt [58], and GeneCards [17] databases.

22

23 ***De novo* transcriptome assembly and differentially expressed genes**

1 The blood transcriptome data were assembled using Trinity software. Contaminated transcripts
2 were searched for bacteria and fungi sequence from the Ensembl database using BLASTN, and
3 filtered by identity of > 95% and *E*-value cutoff of 1E-6 criteria. Coding sequence (CDS) were
4 predicted using TransDecoder [45, 59]. To identify differentially expressed genes, RNA reads
5 were aligned to the reference genome (species whole genome assembled) or the assembled
6 transcripts (species without reference genome) using TopHat2 software [60]. The number of
7 reads that were mapped to orthologous genes were counted using HTSeq- 0.6.1 software [61]
8 and then converted into RPKM (Reads per kilobase per million mapped reads) value. The RPKM
9 values were normalized with the Trimmed Mean of M-values (TMM) [62] correction using the R
10 package edgeR [63]. The significance of differential expression was calculated by the moderated
11 t-test [64] (eBayes function) using the R package limma ($P < 0.05$) [65].

12

13 **Sensory system and beak development associated gene analysis**

14 To compare the olfactory sense across avian clades, we collected a total of 215 chicken olfactory
15 receptor (OR) gene sequences (functional only) from a previously published paper [66]. These
16 ORs were then searched against the 25 avian species genomes using TblastN with default
17 parameters. For OR candidates lacking start/stop codons, we searched 90bp upstream to find
18 start codons and 90bp downstream to find stop codons. After collecting sequences for each
19 species, the CD-HIT program [67] was used to remove redundant sequences with an identity cut-
20 off of 100%. A Pfam [68] search against sequences using hmmer-3.1 program [69] with an *E*-
21 value cutoff of 1.0 was used to identify sequences that contained 7tm_4 domain. To assign OR
22 clades and filter out non-OR genes, the multiple sequence alignments and phylogenetic analysis
23 were conducted with previously clade-assigned OR and non-OR genes of human, anole lizard,

1 and chicken [70] using ClustalW2 program [71]. The remaining OR candidates were classified
2 into three categories: 1) intact genes with normal start and stop codons and longer than 215
3 amino acid sequences, thus can code for seven transmembrane domains; 2) partial genes without
4 start and/or stop codons; and 3) pseudogenes with frameshift mutations and/or premature stop
5 codons. OR genes have evolved by multiple duplications and display a large number of
6 pseudogenes, which makes the assembly of OR regions challenging and complicates the
7 annotation process of OR genes [5, 13, 72, 73]. To overcome these issues, we also calculated the
8 diversity of OR genes by Shannon entropy [74] using BioEdit [75] as previously suggested [5,
9 13]. Amino acid positions with above 20% of gaps were excluded, and entropy was averaged
10 across all amino acid positions.

11 The vision system associated genes were retrieved from previous studies [5, 13]. Hearing
12 associated genes were retrieved from the AmiGO database [76] using GO categories related to
13 hearing [5]. Circadian rhythm related genes were retrieved from the AmiGO database using
14 “biorhythm/circadian” as search keywords. For the beak development analysis, beak
15 development associated genes were retrieved from the falcon genome study [3]. The protein
16 sequences with the same gene name were aligned using ClustalW2 and manually inspected one
17 by one for quality. A total of 50 beak development associated genes shared by at least two
18 Strigiformes, two Accipitriformes, and two Falconiformes, and 402 sensory system associated
19 genes (64 genes for vision, 219 genes for hearing, and 133 genes for circadian rhythm) shared by
20 the brown kiwi, chuck-will’s-widow, and at least two Strigiformes were included for selection
21 constraint (the d_N/d_S ratio) analyses.

22

1 **Additional files**

2 **Additional file 1: Figures S1-S8, Tables S1-S40,** and Supplementary Methods.

3 **Additional file 2: Datasheet S1.** PSGs in the ancestral branch of Strigiformes. **Datasheet S2.**

4 PSGs in the ancestral branch of Accipitriformes. **Datasheet S3.** PSGs in the ancestral branch of

5 Falconiformes. **Datasheet S4.** PSGs in chuck-will's-widow genome. **Datasheet S5.** PSGs in

6 brown kiwi genome. **Datasheet S6.** List of genes showing a high level of GC3 biases in the

7 raptors. **Datasheet S7.** List of genes showing a high level of GC3 biases in the nocturnal birds.

8 **Datasheet S8.** Differentially expressed genes in Strigiformes. **Datasheet S9.** Differentially

9 expressed genes in Accipitriformes. **Datasheet S10.** Differentially expressed genes in

10 Falconiformes. **Datasheet S11.** Differentially expressed genes in raptors

11

12 **Abbreviations**

13 PSG: Positively selected gene; GC3: Guanine-Cytosine at the third codon position; HCR: Highly

14 conserved genomic region; OR: Olfactory receptor

15

16 **Acknowledgements**

17 Korea Institute of Science and Technology Information (KISTI) provided us with Korea

18 Research Environment Open NETwork (KREONET), which is the Internet connection service

19 for efficient information and data transfer. We thank ARA Jo for bird illustrations.

20

21 **Funding**

22 This work was supported by the grants from the National Institute of Biological Resources

23 (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea

1 (NIBR201403101, NIBR201503101, NIBR201703102). This work was also supported by the
2 Genome Korea Project in Ulsan (800 genome sequencing) Research Fund (1.180017.01) of
3 Ulsan National Institute of Science & Technology (UNIST).

4

5 **Availability of data and materials**

6 The Eurasian eagle-owl, oriental scops-owl, eastern buzzard, and common kestrel genomes have
7 been deposited at DDBJ/EMBL/GenBank under the accession numbers PYWY00000000,
8 PYXB00000000, PYWZ00000000, and PYXA00000000, respectively. The versions described in
9 this paper are the first versions, PYWY01000000, PYXB01000000, PYWZ01000000, and
10 PYXA01000000. All the DNA and RNA sequencing data have been deposited into the NCBI
11 Sequence Read Archive under the accession number SRP131743.

12

13 **Authors' contributions**

14 The birds of prey genome project was initiated by the National Institute of Biological Resources,
15 Korea. SoonokK, JHY, and JB supervised and coordinated the project. SoonokK, JB, and YSC
16 conceived and designed the experiments. JAK, SGK, JYP, Hwa-JungK, SunghyunK, Hee-JongK,
17 Jin-hoJ, KJN, and JK provided samples, advice and associated information. YSC, JeHoonJ,
18 HMK, OC, SGP, HYL, and JF conducted the bioinformatics data processing and analyses. YSC,
19 JeHoonJ, SoonokK, and JB wrote and revised the manuscript. AM, DPM, JF, JSE, JAW, and
20 CCW reviewed and edited the manuscript.

21

22 **Competing interests**

1 YSC, JeHoonJ, OC, and HYL are employees, and JB is a chief executive officer of Clinomics
2 Inc. JB, YSC, and HMK have an equity interest in the company. All other coauthors have no
3 conflicts of interest to declare.

4

5 **Ethics approval and consent to participate**

6 Permissions for the endangered species of Korea or animals listed as natural monument were
7 obtained from the Ministry of Environment (MOE) or from the Cultural Heritage Administration
8 (CHA), respectively (see Additional file 1: Table S3 for detailed sampling and permission
9 information). No animals were killed or captured as a result of these studies.

10

11 **Consent for publication**

12 Not applicable.

13

1 **References**

- 2 1. Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, et al. Whole-genome analyses resolve
3 early branches in the tree of life of modern birds. *Science*. 2014;346:1320–1331.
- 4 2. Fowler DW, Freedman EA, Scannella JB. Predatory functional morphology in raptors:
5 interdigital variation in talon size is related to prey restraint and immobilisation technique.
6 *PLoS One*. 2009;4:e7999.
- 7 3. Zhan X, Pan S, Wang J, Dixon A, He J, Muller MG, et al. Peregrine and saker falcon genome
8 sequences provide insights into evolution of a predatory lifestyle. *Nat Genet*. 2013;45:563–
9 566.
- 10 4. Wu Y, Hadly EA, Teng W, Hao Y, Liang W, Liu Y, et al. Retinal transcriptome sequencing
11 sheds light on the adaptation to nocturnal and diurnal lifestyles in raptors. *Sci Rep*.
12 2016;6:33578.
- 13 5. Le Duc D, Schöneberg T. Adaptation to nocturnality - learning from avian genomes.
14 *Bioessays*. 2016;38:694–703.
- 15 6. Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, et al. Comparative genomics reveals insights
16 into avian genome evolution and adaptation. *Science*. 2014;346:1311–1320.
- 17 7. Van Den Bussche RA, Judkins ME, Montague MJ, Warren WC. A Resource of Genome-Wide
18 Single Nucleotide Polymorphisms (Snps) for the Conservation and Management of Golden
19 Eagles. *J Raptor Res*. 2017;51:368–377.

- 1 8. Hanna ZR, Henderson JB, Wall JD, Emerling CA, Fuchs J, Runckel C, et al. Northern Spotted
2 Owl (*Strix occidentalis caurina*) Genome: Divergence with the Barred Owl (*Strix varia*) and
3 Characterization of Light-Associated Genes. *Genome Biol Evol.* 2017;9:2522–2545.
- 4 9. Mueller JC, Kuhl H, Boerno S, Tella JL, Carrete M, Kempenaers B. Evolution of genomic
5 variation in the burrowing owl in response to recent colonization of urban areas. *Proc Biol*
6 *Sci.* 2018;285:20180206.
- 7 10. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved
8 memory-efficient short-read de novo assembler. *Gigascience.* 2012;1:18.
- 9 11. Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, et al. Efficient de novo
10 assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome*
11 *Res.* 2014;24:1384–1395.
- 12 12. Kim S, Cho YS, Kim HM, Chung O, Kim H, Jho S, et al. Comparison of carnivore,
13 omnivore, and herbivore mammalian genomes with a new leopard assembly. *Genome Biol.*
14 2016;17:211.
- 15 13. Le Duc D, Renaud G, Krishnan A, Almén MS, Huynen L, Prohaska SJ, et al. Kiwi genome
16 provides insights into evolution of a nocturnal lifestyle. *Genome Biol.* 2015;16:147.
- 17 14. Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol.*
18 2017;24:1586–1591.
- 19 15. Zhang J, Nielsen R, Yang Z. Evaluation of an improved branch-site likelihood method for
20 detecting positive selection at the molecular level. *Mol Biol Evol.* 2005;22:2472–2479.

- 1 16. Foote AD, Liu Y, Thomas GW, Vinař T, Alföldi J, Deng J, et al. Convergent evolution of the
2 genomes of marine mammals. *Nat Genet.* 2015;47:272–275.
- 3 17. Safran M, Dalah I, Alexander J, Rosen N, Iny Stein T, Shmoish M, et al. GeneCards Version
4 3: the human gene integrator. Database (Oxford). 2010;2010:baq020.
- 5 18. Engevik K, Aihara E, Matthis A, Montrose M. TFF2, CXCR4 and EGF-R mediated gastric
6 wound repair in vitro in gastric organoids. *FASEB J.* 2017;31 Suppl 1:1043–1048.
- 7 19. Saada A, Vogel RO, Hoefs SJ, van den Brand MA, Wessels HJ, Willems PH, et al.
8 Mutations in NDUFAF3 (C3ORF60), encoding an NDUFAF4 (C6ORF66)-interacting
9 complex I assembly protein, cause fatal neonatal mitochondrial disease. *Am J Hum Genet.*
10 2009;84:718–727.
- 11 20. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome
12 proliferator-activated receptor α mediates the adaptive response to fasting. *J Clin Invest.*
13 1999;103:1489–1498.
- 14 21. Tatarinova TV, Alexandrov NN, Bouck JB, Feldmann KA. GC3 biology in corn, rice,
15 sorghum and other grasses. *BMC Genomics.* 2010;11:308.
- 16 22. Borges R, Khan I, Johnson WE, Gilbert MT, Zhang G, Jarvis ED, et al. Gene loss, adaptive
17 evolution and the co-evolution of plumage coloration genes with opsins in birds. *BMC*
18 *Genomics.* 2015;16:751.

- 1 23. Yu G, Zhang Y, Xiang Y, Jiang P, Chen Z, Lee W, et al. Cell migration-promoting and
2 apoptosis-inhibiting activities of Bm-TFF2 require distinct structure basis. *Biochem Biophys*
3 *Res Commun.* 2010;400:724–728.
- 4 24. Crane NJ, Brown TS, Evans KN, Hawksworth JS, Hussey S, Tadaki DK, et al. Monitoring
5 the healing of combat wounds using Raman spectroscopic mapping. *Wound Repair Regen.*
6 2010;18:409–416.
- 7 25. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:
8 tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.* 2000;25:25–29.
- 9 26. Zhao H, Ru B, Teeling EC, Faulkes CG, Zhang S, Rossiter SJ. Rhodopsin molecular
10 evolution in mammals inhabiting low light environments. *PLoS One.* 2009;4:e8326.
- 11 27. Dawson PA, Hubbert M, Haywood J, Craddock AL, Zerangue N, Christian WV, et al. The
12 heteromeric organic solute transporter α - β , Ost α -Ost β , is an ileal basolateral bile acid
13 transporter. *J Biol Chem.* 2005;280:6960–6968.
- 14 28. Govindarajan K, MacSharry J, Casey PG, Shanahan F, Joyce SA, Gahan CG. Unconjugated
15 bile acids influence expression of circadian genes: a potential mechanism for microbe-host
16 crosstalk. *PloS One.* 2016;11:e0167319.
- 17 29. Zhang F, Duan Y, Xi L, Wei M, Shi A, Zhou Y, et al. The influences of cholecystectomy on
18 the circadian rhythms of bile acids as well as the enterohepatic transporters and enzymes
19 systems in mice. *Chronobiol Int.* 2018;35:673–690.

- 1 30. Hasin-Brumshtein Y, Lancet D, Olender T. Human olfaction: from genomic variation to
2 phenotypic diversity. *Trends Genet.* 2009;25:178–184.
- 3 31. Khan I, Yang Z, Maldonado E, Li C, Zhang G, Gilbert MT, et al. Olfactory Receptor
4 Subgenomes Linked with Broad Ecological Adaptations in Sauropsida. *Mol Biol Evol.*
5 2015;32:2832–2843.
- 6 32. Emery NJ. Cognitive ornithology: the evolution of avian intelligence. *Philos Trans R Soc*
7 *Lond B Biol Sci.* 2006;361:23–43.
- 8 33. Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, et al. The evolution
9 of gene expression levels in mammalian organs. *Nature.* 2011;478:343–348.
- 10 34. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
11 *ArXiv.* 2013;1303:3997.
- 12 35. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
13 Alignment/Map format and SAMtools. *Bioinformatics.* 2009;25:2078–2079.
- 14 36. Benson G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.*
15 1999;27:573–580.
- 16 37. Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. Repbase
17 Update, a database of eukaryotic repetitive elements. *Cytogenet Genome Res.* 2005;110:462-
18 467.
- 19 38. Bedell JA, Korf I, Gish W. MaskerAid: a performance enhancement to RepeatMasker.
20 *Bioinformatics.* 2000;16:1040–1041.

- 1 39. RMBlast. <http://www.repeatmasker.org/RMBlast.html>. Accessed 16 Aug 2016.
- 2 40. Abrusán G, Grundmann N, DeMester L, Makalowski W. TEclass--a tool for automated
3 classification of unknown eukaryotic transposable elements. *Bioinformatics*. 2009;25:1329–
4 1330.
- 5 41. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
6 architecture and applications. *BMC Bioinformatics*. 2009;10:421.
- 7 42. She R, Chu JS, Wang K, Pei J, Chen N. GenBlastA: enabling BLAST to identify
8 homologous gene sequences. *Genome Res*. 2009;19:143–149.
- 9 43. Slater GS, Birney E. Automated generation of heuristics for biological sequence comparison.
10 *BMC Bioinformatics*. 2005;6:31.
- 11 44. Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: ab initio
12 prediction of alternative transcripts. *Nucleic Acids Res*. 2006;34:W435–W439.
- 13 45. rabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
14 transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*.
15 2011;29:644–652.
- 16 46. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing
17 genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*.
18 2015;31:3210–3212.
- 19 47. Li L, Stoeckert CJ Jr, Roos DS. OrthoMCL: identification of ortholog groups for eukaryotic
20 genomes. *Genome Res*. 2003;13:2178–2189.

- 1 48. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
2 Nucleic Acids Res. 2004;32:1792–1797.
- 3 49. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment
4 trimming in large-scale phylogenetic analyses. *Bioinformatics*. 2009;25:1972–1973.
- 5 50. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis
6 Version 7.0 for Bigger Datasets. *Mol Biol Evol*. 2016;33:1870–1874.
- 7 51. Hedges SB, Dudley J, Kumar S. TimeTree: a public knowledge-base of divergence times
8 among organisms. *Bioinformatics*. 2006;22:2971–2972.
- 9 52. Han MV, Thomas GW, Lugo-Martinez J, Hahn MW. Estimating gene gain and loss rates in
10 the presence of error in genome assembly and annotation using CAFE 3. *Mol Biol Evol*.
11 2013;30:1987–1997.
- 12 53. Löytynoja A, Goldman N. An algorithm for progressive multiple alignment of sequences
13 with insertions. *Proc Natl Acad Sci USA*. 2005;102:10557–10562.
- 14 54. Weber JA, Aldana R, Gallagher BD, Edwards JS. Sentieon DNA pipeline for variant
15 detection—software-only solution, over 20× faster than GATK 3.3 with identical results. *PeerJ*
16 *PrePrints*. 2016;4:e1672v2.
- 17 55. Storey JD. A direct approach to false discovery rates. *J R Stat Soc Series B Stat Methodol*.
18 2002;64:479–498.
- 19 56. Aken BL, Achuthan P, Akanni W, Amode MR, Bernsdorff F, Bhai J, et al. Ensembl 2017.
20 *Nucleic Acids Res*. 2017;45:D635–D642.

- 1 57. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. KAAS: an automatic genome
2 annotation and pathway reconstruction server. *Nucleic Acids Res.* 2007;35:W182–W185.
- 3 58. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*
4 2017;45:D158–D169.
- 5 59. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo
6 transcript sequence reconstruction from RNA-seq using the Trinity platform for reference
7 generation and analysis. *Nat Protoc.* 2013;8:1494–1512.
- 8 60. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate
9 alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome*
10 *Biol.* 2013;14:R36.
- 11 61. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput
12 sequencing data. *Bioinformatics.* 2015;31:166–169.
- 13 62. Robinson MD, Oshlack A. A scaling normalization method for differential expression
14 analysis of RNA-seq data. *Genome Biol.* 2010;11:R25.
- 15 63. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
16 expression analysis of digital gene expression data. *Bioinformatics.* 2010;26:139–140.
- 17 64. McCarthy DJ, Smyth GK. Testing significance relative to a fold-change threshold is a
18 TREAT. *Bioinformatics.* 2009;25:765–771.

- 1 65. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential
2 expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.*
3 2015;43:e47.
- 4 66. Steiger SS, Kuryshev VY, Stensmyr MC, Kempnaers B, Mueller JC. A comparison of
5 reptilian and avian olfactory receptor gene repertoires: species-specific expansion of group
6 gamma genes in birds. *BMC Genomics.* 2009;10:446.
- 7 67. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
8 nucleotide sequences. *Bioinformatics.* 2006;22:1658–1659.
- 9 68. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, et al. The Pfam protein
10 families database: towards a more sustainable future. *Nucleic Acids Res.* 2016;44:D279–D285.
- 11 69. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology search:
12 HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res.* 2013;41:e121.
- 13 70. Niimura Y. On the origin and evolution of vertebrate olfactory receptor genes: comparative
14 genome analysis among 23 chordate species. *Genome Biol Evol.* 2009;1:34–44.
- 15 71. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, et al. Multiple sequence
16 alignment with the Clustal series of programs. *Nucleic Acids Res.* 2009;31:3497–3500.
- 17 72. Niimura Y, Nei M. Evolution of olfactory receptor genes in the human genome. *Proc Natl*
18 *Acad Sci USA.* 2003;100:12235–12240.
- 19 73. Newman T, Trask BJ. Complex evolution of 7E olfactory receptor genes in segmental
20 duplications. *Genome Res.* 2003;13:781–793.

- 1 74. Shannon CE. The mathematical theory of communication. Bell Syst Tech J. 1948;27:379–
2 423.
- 3 75. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program
4 for Windows 95/98/NT. Nucleic Acids Symp Ser.1999;41:95–98.
- 5 76. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S. AmiGO: online access to
6 ontology and annotation data. Bioinformatics. 2009;25:288–289.
- 7






Fig. 1 Phylogeny and genomic data of birds of prey. The phylogenetic tree topology was adapted from the Avian Phylogenomics Project [1] and TimeTree database. The estimated divergence time from present (million years ago; MYA) is given at the nodes. Dark red indicates species with higher-quality (scaffold N50 length > 1 Mb) genome assemblies, light red indicates species with lower-quality genome assemblies, black indicates species for which the whole genome was resequenced, and grey indicates non-raptor species high-quality genome assemblies. * denotes birds of prey sequenced from this study. The white-tailed eagle (denoted with **) was previously assembled at low-quality and also whole genome resequenced from this study.



1
2 **Fig. 2** Relationship of birds of prey to other avian species. **a** Venn diagrams of orthologous gene
3 clusters in the birds of prey. Orthologous gene clusters were constructed using 25 avian genomes.
4 Only raptor gene clusters are displayed. **b** Gene expansion or contraction in avian species. The
5 numbers near order and species names indicate the number of gene families that have expanded
6 (+) and contracted (-) in each branch and species. Species in red are birds of prey. **c** Heatmap of
7 enriched Gene Ontology (GO) categories for raptor common GC3 biased genes. Bird icons from
8 left to right indicate Strigiformes, Accipitriformes, Falconiformes, and non-raptor birds. Z-scores
9 for the average of normalized GC3 percentages are shown as a yellow-to-black color scale.



1
2 **Fig. 3** A functional trade-off of sensory systems in nocturnal birds. **a** The phylogeny of the α
3 and γ olfactory receptor (OR) genes identified in 25 avian genomes. Only intact OR genes were
4 used. **b** Selection constraints on sensory systems. Values for α , γ , and γ -c ORs are the diversity
5 of ORs in each clade. For avian orders including two or more genomes (Strigiformes,
6 Accipitriformes, Passeriformes, Falconiformes, and Pelecaniformes), the average diversity
7 values were used. The diversity of α ORs in Piciformes and γ -c ORs in Psittaciformes were not
8 calculated as the number of identified OR genes were smaller than two. Values for vision,
9 hearing, and circadian rhythm are d_N/d_S ratios of each set of sensory system associated genes. For
10 avian orders including two or more genomes, d_N/d_S ratios of the ancestral branches were used.
11 Three avian orders in red are nocturnal.

12