5-HT_{2B} shapes the macrophage transcriptome via AhR

5-HT_{2B} serotonin receptor agonist BW723C86 shapes the macrophage gene profile via AhR and impairs monocyte-to-osteoclast differentiation

Short title: 5-HT_{2B} shapes the macrophage transcriptome via AhR

Concha Nieto¹, Ignacio Rayo¹, Mateo de las Casas-Engel¹, Elena Izquierdo¹, Bárbara Alonso¹, Miguel A. Vega¹, and Ángel L. Corbí^{1*}

¹ Myeloid Cell Laboratory, Centro de Investigaciones Biológicas, CSIC, Madrid, Spain.

Corresponding author: Dr. Ángel L. Corbí, Centro de Investigaciones Biológicas, CSIC.

Ramiro de Maeztu, 9. Madrid 28040; Phone: 34-91-8373112; FAX: 34-91-5627518;

E-mail: acorbi@cib.csic.es

Text word count: 3931

Abstract word count: 249

Figures: 7

Reference count: 128

5-HT_{2B} shapes the macrophage transcriptome via AhR

KEY POINTS

- The serotonin receptor 5-HT_{2B} modifies the human macrophage transcriptome through activation of the Aryl Hydrocarbon Receptor.
- BW723C86, an agonist used for 5-HT_{2B} activation *in vivo*, exerts 5-HT_{2B}-independent effects and limits monocyte osteoclastogenic potential.

5-HT_{2B} shapes the macrophage transcriptome via AhR

ABSTRACT

Peripheral serotonin (5-HT) exacerbates or limits inflammatory pathologies through interaction with seven types of 5-HT receptors (5-HT₁₋₇). As central regulators of inflammation, macrophages are critical targets of 5-HT, which promotes their anti-inflammatory and pro-fibrotic polarization primarily via the 5-HT₇-Protein Kinase A (PKA) axis. However, anti-inflammatory human macrophages are also characterized by the expression of 5-HT_{2B}, an off-target of anesthetics, anti-parkinsonian drugs and Selective Serotonin Reuptake Inhibitors (SSRI) that contributes to 5-HT-mediated pathologies. Since 5-HT_{2B} prevents mononuclear phagocyte degeneration in amyotrophic lateral sclerosis and modulates motility of murine microglial processes, we sought to determine the functional and transcriptional consequences of 5-HT_{2B} activation in human macrophages. Ligation of 5-HT_{2B} by the 5-HT_{2B}-specific agonist BW723C86, which exhibits antidepressant- and anxiolytic-like effects in animal models, significantly modified the cytokine profile and the transcriptional signature in macrophages. Importantly, 5-HT_{2B} agonist-induced transcriptional changes were partly mediated through activation of the Aryl hydrocarbon Receptor (AhR), a ligand-dependent transcription factor that regulates immune responses and the biological responses to xenobiotics. Besides, BW723C86 triggered transcriptional effects that could not be abrogated by 5-HT_{2B} antagonists and impaired monocyte-to-osteoclast differentiation by affecting the expression of negative (*IRF8*) and positive (*PRDM1*) regulators of osteoclastogenesis. Therefore, our results demonstrate the existence of a functional 5-HT_{2B}-AhR axis in human macrophages and indicate that the commonly used 5-HT_{2B} agonist BW723C86 exhibits 5-HT_{2B}-independent effects. The 5-HT_{2B}-AhR link extends the range of signaling pathways initiated upon 5-HT receptor engagement and identifies a point of convergence for endogenous and exogenous agents with ability to modulate inflammatory responses.

5-HT_{2B} shapes the macrophage transcriptome via AhR

5-HT_{2B} shapes the macrophage transcriptome via AhR

INTRODUCTION

Peripheral serotonin (5-hydroxytryptamine, 5-HT) influences the physiopathology of numerous tissues¹⁻³ as it controls vascular, heart and gastrointestinal functions^{4,5}, promotes cell proliferation⁶⁻⁹, regulates wound healing, and influences immune and inflammatory responses¹⁰ by modulating T lymphocyte¹¹ and myeloid cell functions¹²⁻¹⁶. In inflammatory pathologies, 5-HT contributes to Pulmonary Arterial Hypertension (PAH)¹⁷, atopic dermatitis¹⁸, systemic sclerosis¹⁹, inflammatory gut disorders²⁰⁻²⁵, cancer angiogenesis²⁶ and neuroendocrine neoplasms proliferation²⁷, whereas it reduces pathologic scores in collagen-induced arthritis²⁸. The existence of seven types of 5-HT receptors (5-HT₁₋₇), with different distribution and signaling properties²⁹, underlies these tissue-specific actions of 5-HT.

Macrophages promote the initiation and the resolution of inflammatory processes, with both processes being essential for maintaining tissue homeostasis³⁰⁻³². In fact, modulation of macrophage polarization is an attractive therapeutic target for chronic inflammatory pathologies, where the balance between pro-inflammatory and resolving macrophages is altered^{33,34}. Not surprisingly, macrophage effector functions are directly modulated by 5-HT^{17,22,26,35,36}, and also by Selective Serotonin Reuptake Inhibitors (SSRI)^{37,38}. We have previously demonstrated that 5-HT₇ is the major mediator of the anti-inflammatory actions of 5-HT on human macrophages³⁹ and that both 5-HT_{2B} and 5-HT₇ contribute to the maintenance of the anti-inflammatory state of M-CSF-dependent human macrophages⁴⁰.

The 5-HT_{2B} serotonin receptor is expressed in the central nervous system, where it exerts anti-depressant or anxiolytic-like effects⁴¹, and in periphery, where it plays a major role in heart development⁴². Pharmacologic and genetic studies indicate that 5-HT_{2B} is required for 5-HT-mediated pathologies like PAH^{17,43}, dermal, lung and liver fibrosis^{19,44,45}, cardiac

5-HT_{2B} shapes the macrophage transcriptome via AhR

hypertrophy^{46,47}, and mediates the valvular heart disease (VHD) and fibrosis associated with carcinoid syndrome⁴⁸. Importantly, anti-migraine drugs like methysergide and ergotamine^{49,50}, general anesthetics⁵¹, fenfluramine and conventional SSRIs⁵²⁻⁵⁵, and even the dopamine agonists and anti-parkinsonian drugs pergolide and cabergoline^{50,56,57}, exert potent off-target effects on 5-HT_{2B}⁵⁰. In fact, 5-HT_{2B} appears to be required for the therapeutic actions of SSRI⁵⁴. Since some of these 5-HT_{2B}-targeting drugs induce VHD, their use is restricted, and novel potential drugs are commonly screened for 5-HT_{2B} agonist activity.

5-HT_{2B} engagement enhances proliferation of numerous cell types^{9,58-67} via Gαq and Src phosphorylation, and through production of growth factors like insulin⁶⁸, TGFβ1, CTGF, FGF2^{45,60} and TGFα⁶⁹. In macrophages, 5-HT_{2B} prevents mononuclear phagocyte degeneration in amyotrophic lateral sclerosis⁷⁰, and *Htr2b*^{-/-} microglia exhibits a mild inflammatory state⁷¹, which is in line with the ability of 5-HT_{2B} antagonist SB204741 to impair the acquisition of human macrophage polarization-specific genes⁴⁰ and TGFβ1 expression by mouse Kupffer cells⁴⁵. However, although 5-HT_{2B} activation modifies inflammatory cytokine production by human monocytes⁷², the functional and transcriptional consequences of 5-HT_{2B} ligation in human macrophage remains to be determined. Since anti-inflammatory M-CSF-dependent macrophages express 5-HT_{2B}, we have assessed the specific role of 5-HT_{2B} in macrophage polarization. The use of the 5-HT_{2B} agonist BW723C86, used to evaluate the role of 5-HT_{2B} in preclinical models of depression⁵⁴ and anxiety^{73,74}, allowed us to demonstrate that 5-HT_{2B} shapes the macrophage transcriptome via Aryl hydrocarbon Receptor (AhR) activation and impairs the osteoclastogenic potential of human macrophages.

5-HT_{2B} shapes the macrophage transcriptome via AhR

METHODS

Generation of human monocyte-derived macrophages and cell culture. Buffy coats from anonymous healthy blood donors were provided by Comunidad de Madrid blood Bank. Human peripheral blood mononuclear cells (PBMC) were isolated from buffy coats over a Lymphoprep (Nycomed Pharma) gradient, and monocytes were purified by magnetic cell sorting using CD14 microbeads (Miltenyi Biotech). Monocytes (>95% CD14⁺ cells) were cultured at 0.5×10^6 cells/ml for 7 days in RPMI supplemented with 10% fetal calf serum (FCS, Biowest) (completed medium) at 37°C in a humidified atmosphere with 5% CO₂, and containing M-CSF (10 ng/ml) (ImmunoTools GmbH) to generate M-MØ monocyte-derived macrophages. M-CSF was added every two days. Before treatments, M-MØ were maintained in serum-free medium (Macrophage-SFM, Gibco) for 48 hours. For macrophage activation, cells were exposed to 5-HT_{2B} agonists for 6 hours and then treated with *Escherichia coli* 055:B5 LPS (10 ng/ml) for 18 h. The 5-HT_{2B} agonists BW723C86 (α -methyl-5-(2-thienylmethoxy)-1*H*-indole-3-ethanamine)^{75,76} and α -Methylserotonin (AMS)⁷⁷, and the 5-HT_{2B} antagonist SB204741⁷⁸, were purchased from Sigma-Aldrich. The 5-HT_{2B} agonist 6-APB (6-(2-aminopropyl)benzofuran) was obtained from Cayman and used at the indicated concentrations. The AhR agonist FICZ (Enzo) and antagonist CH223191 (Calbiochem) were used at 250 nM and 3 μ M, respectively. When indicated, SB204741 was used 1 hour before treatment with agonists. Monocyte-derived osteoclasts were generated by culturing monocytes for 12 days on glass coverslips with M-CSF (25 ng/ml) and Receptor Activator for NF- κ B Ligand (RANKL; 30 ng/ml) addition every 72 h and BW723C86 treatment 6 h before cytokine addition. Osteoclast generation was verified directly by phase contrast microscopy or after staining for Tartrate-Resistant Acid Phosphatase (TRAP) (Leukocyte Acid Phosphatase kit, Sigma-Aldrich). Human TNF α and CCL2 in M-MØ culture supernatants were measured using commercially available ELISA (BD Biosciences).

5-HT_{2B} shapes the macrophage transcriptome via AhR

Transfections and Reporter gene assays. Dual luciferase Cignal® Xenobiotic Response Element (XRE) Reporter assay kit (Qiagen), where the XRE-Luc construct harbours tandem repeats of the AhR-binding element (XRE), was used for the reporter gene assays. HepG2 cells

(80.000 cells/well) or M-MØ (1x10⁶ cells/well) were transfected with 1 µg of the XRE reporter construct using either Superfect (Qiagen) or VIROMER RED (Lipocalix), respectively. For normalizing transfection efficiency, transfected DNA included a 40:1 mixture of the XRE-specific firefly luciferase construct and a construct expressing *Renilla* luciferase from a constitutive promoter. After transfection, cells were cultured overnight before exposure to BW723C86 (10 µM), FICZ (250 nM), CH223191 (3 µM) or DMSO for 24 h, lysed, and firefly and *Renilla* luciferase activities were determined using the Dual-Luciferase® Reporter Assay System (Promega).

Small Interfering Ribonucleic Acid (siRNA) Transfection. To knockdown AhR expression, M-MØ (1 × 10⁶ cells) were transfected with AhR-specific siRNA (siAhR) (50 nM) (# s1199, Thermo-Fisher Scientific), using HiPerFect (Qiagen) using siC (#4390843, Thermo-Fisher Scientific) as negative control siRNA. Cells were allowed to recover from transfection in complete medium (18–24 h) and then treated for 6h with BW723C86 (10 µM) before analysis.

Quantitative real-time RT-PCR (qRT-PCR). Total RNA was extracted using the total RNA and protein isolation kit (Macherey-Nagel). RNA samples were retrotranscribed (High-Capacity cDNA Reverse Transcription kit, AB), and triplicates of amplified cDNA were analyzed on a Light Cycler® 480. Gene-specific oligonucleotides (Supplementary Table I)

5-HT_{2B} shapes the macrophage transcriptome via AhR

were designed using the Universal ProbeLibrary software (Roche Diagnostics). Results were expressed relative to the expression level of the endogenous reference gene *TBP* and using the $\Delta\Delta CT$ (cycle threshold) method for quantitation.

Microarray analysis. Global gene expression analysis was performed on RNA from three independent samples of untreated (control), BW723C86-treated (10 μM), SB204741-treated (1 μM), or SB204741+BW723C86-treated M-MØ. RNA was isolated using the RNeasy Mini kit (Qiagen) and used as hybridization probe on Whole Human Genome Microarrays (Agilent Technologies, Palo Alto, CA). Only probes with signal values >60% quantile in at least one condition were considered for the differential gene expression (DGE) and statistical analysis. Statistical analysis for DGE was carried out using empirical Bayes moderated paired t-test implemented in the *limma* package (<http://www.bioconductor.org>), and p values were adjusted for multiple hypotheses testing using the Benjamini-Hochberg method to control the false discovery rate ⁷⁹, with all procedures coded in R (<http://www.r-project.org>). Microarray data were deposited in the Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo/>) under accession no. GSE121825. Unsupervised hierarchical clustering was done on the mean expression level of genes significantly regulated by BW723C86 (adj $p < 0.002$) with the ClustVis webtool (<https://biit.cs.ut.ee/clustvis/>) ⁸⁰. Differentially expressed genes were analysed for annotated gene sets enrichment using ENRICH (<http://amp.pharm.mssm.edu/Enrichr/>) ^{81,82}, and enrichment terms considered significant with a Benjamini-Hochberg-adjusted p value < 0.05 . For gene set enrichment analysis (GSEA) ⁸³, the gene sets available at the website, as well as the previously defined “Pro-inflammatory gene set” and “Anti-inflammatory gene set” ⁸⁴ (GSE68061) were used.

5-HT_{2B} shapes the macrophage transcriptome via AhR

Statistical analysis. For comparison of means, and unless otherwise indicated, statistical significance of the generated data was evaluated using the Student paired t-test. In all cases, $p < 0.05$ was considered as statistically significant.

5-HT_{2B} shapes the macrophage transcriptome via AhR

RESULTS

The BW723C86 agonist modulates the transcriptional signature of human macrophages partially via 5-HT_{2B}. Since human anti-inflammatory M-MØ express 5-HT₇ and 5-HT_{2B} serotonin receptors⁴⁰ and clinically relevant drugs (anesthetics, SSRIs, antimigraine drugs) act via 5-HT_{2B}^{49,51-54}, we sought to determine the consequences of 5-HT_{2B} engagement in human macrophages using BW723C86, an 5-HT_{2B} agonist with 10-fold and 100-fold selectivity over human 5-HT_{2C} and 5-HT_{2A}, respectively⁸⁵, amply used as *in vivo*^{54,73,74,86-91} and *ex vivo*^{28,66,70}, and with potential therapeutic effects^{73,74,91}. Transcriptional profiling of BW723C86-treated M-MØ (Figure 1A) revealed that the agonist increased the expression of 357 annotated genes and downregulated the expression of 398 genes (*adj p*<0.002) (Figure 1B, Supplementary Tables II and III). Analysis of independent M-MØ samples confirmed the microarray data, as BW723C86 increased *NCF4* and reduced *ACSL1*, *EGR1* and *DUSP6* expression (Figure 1C). Gene Set Enrichment Analysis (GSEA) revealed that BW723C86 significantly modifies the expression of “Gα_{12/13} signaling”⁹² and “Amyotrophic Lateral Sclerosis” gene sets, in agreement with the signaling capability and pathological significance of 5-HT_{2B}⁷⁰ (Figure 1D), and affects biological processes like “GO Heart Valve Development” and “GO Regulation of Myeloid Leukocyte Differentiation” (Supplementary Figure 1A), in line with the known involvement of 5-HT_{2B} in heart development^{42,93,94} and macrophage differentiation⁴⁰. Further, terms associated with bone metabolism were also enriched in BW723C86-treated M-MØ (Supplementary Figure 1A). Interestingly, and unlike the 5-HT-mediated activation of 5-HT₇³⁹, BW723C86 significantly augmented the expression of genes associated to both pro-inflammatory and anti-inflammatory macrophage polarization^{84,95} (Supplementary Figure 1B), suggesting that engagement of 5-HT_{2B} or 5-HT₇ on human macrophages leads to different transcriptional outcomes.

5-HT_{2B} shapes the macrophage transcriptome via AhR

To identify genes whose expression is regulated by BW723C86 in a 5-HT_{2B}-dependent manner, the transcriptional effects of the 5-HT_{2B} agonist were also determined in the presence of SB204741 (BW723C86 + SB204741) (Figure 1A, Supplementary Table IV), a selective 5-HT_{2B} antagonist with 100-fold selectivity over 5-HT_{2C} and 5-HT_{2A}⁸⁵ that is widely used *in vivo*^{9,61,63,90,96}. We reasoned that the genes significantly modulated by BW723C86, but not by BW723C86+SB204741, would represent *bona fide* 5-HT_{2B}-regulated genes. In the presence of the antagonist SB204741, BW723C86 significantly (*adj p*<0.002) increased the expression of 155 annotated genes and downregulated the expression of 70 genes (Figure 1B, Supplementary Table V). Comparison of the BW723C86-induced transcriptional changes in the absence or presence of the antagonist (Figure 1E, F) revealed that the antagonist SB204741 prevents the BW723C86-mediated upregulation of 217 genes, and the BW723C86-mediated downregulation of 334 genes (Figure 1E, Supplementary Table VI). Therefore, 5-HT_{2B} mediates the augmented expression of 60% of the genes upregulated by BW723C86, and the diminished expression of 84% of the genes downregulated by BW723C86 (Figure 1E, Supplementary Table VI). Further, these results point towards a 5-HT_{2B}-independent effect of BW723C86, as it modulates the expression of 207 genes (140 upregulated, 67 downregulated) even in the presence of the 5-HT_{2B} antagonist. As representative examples, the upregulation of *CYP1B1* and *AHRR* by BW723C86 was abrogated by SB204741, whereas SB204741 did not alter the modulation of *ENG* or *APOL6* expression by BW723C86 (Figure 1F). Altogether, these results demonstrate that BW723C86 modifies the macrophage transcriptome through 5-HT_{2B}-dependent (SB204741-sensitive) and 5-HT_{2B}-independent mechanisms.

BW723C86 modifies the LPS-induced cytokine and gene profile. As GSEA revealed that BW723C86 downregulates the “TNFα Signaling via NFκβ” and “Inflammatory Response” gene sets (Figure 1D), we next assessed the ability of BW723C86 to alter macrophage

5-HT_{2B} shapes the macrophage transcriptome via AhR

responses to an inflammatory stimulus like LPS (Figure 2A). BW723C86 did not modify basal cytokine production by M-MØ (Figure 2B and not shown), while it significantly reduced TNF α production, and increased CCL2 release, from LPS-treated M-MØ (Figure 2B). A similar trend was observed with α -Methyl Serotonin (AMS), a less selective 5-HT_{2B} agonist⁷⁷, although its effects did not reach statistical significance (data not shown). Importantly, the effect of BW723C86 on cytokine production by LPS-activated M-MØ was blunted by the 5-HT_{2B} antagonist SB204741 (Figure 2B), thus confirming that BW723C86 modifies the LPS-induced cytokine profile via 5-HT_{2B} engagement. In fact, BW723C86 significantly impaired the acquisition of the characteristic transcriptome of LPS-stimulated M-MØ (GSE99056)⁹⁷, an effect that was abolished by the 5-HT_{2B} antagonist SB204741 (Figure 2C). Thus, engagement of 5-HT_{2B} by BW723C86 conditions human macrophages for altered responses to LPS.

BW723C86 upregulates the expression of AhR target genes via 5-HT_{2B}. Unsupervised hierarchical clustering and ENRICH analysis (<http://amp.pharm.mssm.edu/Enrichr/>)^{81,82} on BW723C86-regulated genes (adj $p < 0.002$) revealed an enrichment of the “Aryl Hydrocarbon Receptor Pathway_Homo sapiens_WP2873” and “Aryl Hydrocarbon Receptor_Homo sapiens_WP2586” gene sets (Figure 3A), suggesting that BW723C86 might influence the AhR signaling pathway. Indeed, GSEA also revealed an enrichment of the “Metabolism of xenobiotics by Cytochrome P450” gene set (Figure 3B), and of the “Ahr_UP” gene set, which contains genes upregulated by the Aryl Hydrocarbon receptor (AhR)⁹⁸, the major regulator of the biological responses to xenobiotics⁹⁹ (Figure 3B). In fact, the genes modulated by BW723C86 in an 5-HT_{2B}-dependent manner included paradigmatic AhR target genes like *AHRR* and *CYP1B1*¹⁰⁰, as well as genes upregulated by the AhR agonists Benzopyrene or

5-HT_{2B} shapes the macrophage transcriptome via AhR

VAF-347^{98,101} and whose regulatory regions are bound by AhR (<http://iregulon.aertslab.org>) (Supplementary Table VI).

Based on these findings, we assessed the ability of BW723C86 to upregulate the expression of xenobiotic response genes, and found that exposure to BW723C86 significantly upregulates AhR target genes (*AHRR*, *CYP1B1*, *SERPINB2*, *TIPARP*, *EREG*)^{98,100} (Figure 3C). Their upregulation was detected as early as 2 hours after addition of BW723C86 (Figure 3D), observed at BW723C86 concentrations between 0.1 μ M and 10 μ M (Figure 3E) and confirmed on 5-HT_{2B}-expressing human hepatoma HepG2 cells (Figure 3F). In agreement with microarray data, upregulation of AhR-target genes by BW723C86 was reduced by the 5-HT_{2B} antagonist SB204741 (Figure 3G). Therefore, the BW723C86 agonist upregulates the expression of AhR target genes in human macrophages in a 5-HT_{2B}-dependent manner.

AhR mediates the BW723C86-induced upregulation of AhR target genes in human macrophages. To evaluate whether BW723C86 leads to AhR activation, its transcriptional effects were determined after pharmacological or siRNA-mediated inhibition of AhR. The AhR antagonist CH223191¹⁰² prevented the BW723C86-induced upregulation of both *CYP1B1* and *TIPARP*, whose expression was increased by the AhR agonist FICZ¹⁰³ (Figure 4A). Besides, siRNA-mediated knockdown of AhR (Figure 4B,C) completely prevented the BW723C86-induced *CYP1B1* and *TIPARP* upregulation (Figure 4D). In agreement with the role of AhR in the BW723C86-induced upregulation of AhR-target genes, reporter gene experiments revealed that BW723C86 specifically enhances the transcriptional activity of AhR in M-MØ (Figure 4E), an effect that was abrogated by the AhR antagonist CH223191 (Figure 4E). Similar results were also obtained in the 5-HT_{2B}-expressing HepG2 cell line, where the BW723C86-induced AhR transcriptional activity was prevented by the 5-HT_{2B} antagonist SB204741 (Figure 4F) or the AhR antagonist CH223191 (Figure 4G). Therefore,

5-HT_{2B} shapes the macrophage transcriptome via AhR

engagement of the 5-HT_{2B} serotonin receptor by BW723C86 in human macrophages enhances the transcriptional activity of AhR and results in upregulated expression of AhR-target genes.

5-HT and other 5-HT_{2B} ligands influence the expression of BW723C86-regulated genes. To further support the involvement of 5-HT_{2B} in the ability of BW723C86 to promote AhR activation, the influence of 5-HT and other 5-HT_{2B} agonists on the expression of AhR target genes was determined. Albeit to a lower extent than BW723C86, the potent 5-HT_{2B} agonist 6-APB (10nM-1μM) upregulated *CYP1B1* expression in M-MØ (Figure 5A). Similarly, the 5-HT_{2B} agonist alpha-Methyl-Serotonin (AMS) enhanced macrophage *CYP1B1* expression in a dose-dependent manner, and its effect was abolished by the SB204741 antagonist (Figure 5B). Therefore, 5-HT_{2B} agonists upregulate AhR target gene expression in a 5-HT_{2B}-dependent manner in human macrophages. Next, we tested whether 5-HT has an impact on the BW723C86-induced transcriptional effects. As expected³⁹, 5-HT treatment upregulated the expression of *PDE2A* and *TREMI* but did not modify the expression of *CYP1B1* (Figure 5C). However, the BW723C86-mediated upregulation of *CYP1B1* was prevented in the presence of 100 μM 5-HT that, by contrast, did not affect the ability of BW723C86 to downregulate *OCTAMP* expression (Figure 5C). Therefore, 5-HT impairs the ability of the 5-HT_{2B} agonist BW723C86 to activate the AhR transcriptional activity in human macrophages.

5-HT_{2B}-independent effects of BW723C86. The identification of 207 genes whose regulation by BW723C86 is not affected by the 5-HT_{2B} antagonist SB204741 (Figure 1E,F) suggested the existence of 5-HT_{2B}-independent effects of BW723C86. As an example, the BW723C86-mediated downregulation of *OCSTAMP* was not altered by 5-HT or the 5-HT_{2B} antagonist. Along the same line, while the BW723C86-mediated changes in *CYP1B1* and *CSF1* expression was inhibited by the AhR antagonist CH223191, AhR inhibition had no influence

5-HT_{2B} shapes the macrophage transcriptome via AhR

of the downregulation of *EGR1* and *OCSTAMP* by BW723C86 (Figure 6). Altogether, these results confirmed the existence of transcriptional changes induced by BW723C86 that are AhR-independent and not prevented by the 5-HT_{2B} antagonist SB204741 (HT_{2B}-independent). ***BW723C86 blocks osteoclast differentiation.*** The ability of BW723C86 to downregulate the expression of *CSF1* (which encodes M-CSF), *EGR1* (which regulates osteoclastogenesis)^{104,105} and *OCSTAMP* (which drives osteoclast precursor fusion)¹⁰⁶ (Figure 6), together with the under-expression of the “GO Regulation of Osteoclast Differentiation” gene set in the BW723C86-M-MØ transcriptome (Supplementary Figure 1A), led us to test whether BW723C86 negatively affects the monocyte-to-osteoclast differentiation. The continuous presence of the agonist during M-CSF+RANKL-induced osteoclast differentiation significantly impaired the expression of *OCSTAMP* and *DCSTAMP*, which mediate osteoclast cell-cell fusion^{107,108}, as well as the expression of hallmark osteoclast markers like *CTSK*, *CALCR*, *CA2* and *TNFRSF11A* (which encodes the RANKL receptor) (Figure 7A). Therefore, BW723C86 inhibits the expression of genes involved in osteoclast differentiation, cellular fusion and bone resorption, a previously unnoticed action of this 5-HT_{2B} agonist.

To get further insight into the inhibitory action of BW723C86 on osteoclast differentiation, we next assessed whether the agonist modulates the expression of genes coding for factors driving osteoclastogenesis. The presence of BW723C86 along osteoclastogenesis did not modify the expression of *TRAF6*, *MITF*, *FOS* and *BCL6* (Figure 7B). However, BW723C86 significantly augmented the expression of *IRF8*, whose protein product negatively regulates osteoclast differentiation¹⁰⁷, and simultaneously provoked a significant reduction in the expression of *PRDMI*, whose encoded protein (Blimp1) represses the expression of negative regulators of osteoclast differentiation^{107,109,110} (Figure 7B). In fact, exposure of M-MØ to BW723C86 for 6 hours sufficed to significantly downregulate the expression of *PRDMI*

5-HT_{2B} shapes the macrophage transcriptome via AhR

(Supplementary Table II). In consonance with these results, the continuous presence of BW723C86 greatly impaired the generation of “ruffled border”-rich multinucleated cells during the M-CSF+RANKL-driven osteoclast differentiation, as seen by light microscopy (Figure 7C) and after TRAP staining (Figure 7D). Therefore, BW723C86 has an inhibitory effect on the ability of human monocytes to differentiate into osteoclasts at the transcriptional and morphological level, an effect that correlates with its ability to down-regulate *PRDMI* expression and to enhance the expression of *IRF8*.

5-HT_{2B} shapes the macrophage transcriptome via AhR

DISCUSSION

The polarization-specific expression of 5-HT_{2B}, an unintended target of commonly used drugs, prompted us to dissect the transcriptional consequences of 5-HT_{2B} activation in human macrophages through the use of the 5-HT_{2B} agonist BW723C86, which exhibits good selectivity for 5-HT_{2B}⁸⁵, and the 5-HT_{2B} antagonist SB204741, which exerts selectivity over 5-HT_{2C} and 5-HT_{2A}⁸⁵. We now provide evidence that 5-HT_{2B} engagement shapes the macrophage transcriptome partly via activation of AhR⁹⁹ and that BW723C86 impairs the osteoclastogenic differentiation of monocytes at the morphological and transcriptional level, thus emphasizing the role of 5-HT_{2B} in myeloid cell differentiation. The 5-HT_{2B}-AhR axis extends the range of signaling pathways initiated upon 5-HT receptor engagement and constitutes a novel point of convergence between endogenous and exogenous agents with capacity to modulate inflammatory responses.

The link between 5-HT_{2B} and AhR activation is compatible with 5-HT_{2B} favouring an anti-inflammatory polarization because AhR enhances IL-10 expression and limits pro-inflammatory cytokine expression in macrophages¹¹¹. The transcriptional profile promoted by BW723C86 in human macrophages fits with the known signaling ability of 5-HT_{2B}¹¹², as it shows a significant enrichment of genes involved in G protein signaling (Figure 1D). Like 5-HT, BW723C86 has been shown to induce hepatocyte proliferation via PLC activation, EGFR phosphorylation and subsequent mTOR, PI3K, ERK and PLA2 activation¹¹³, and these signaling molecules have been recurrently linked to 5-HT_{2B} activation¹¹². By contrast, the unexpected link of 5-HT_{2B} to AhR activation constitutes an addition to the plethora of cell-specific intracellular signaling pathways activated by 5-HT_{2B}, and its significance is emphasized by the finding that the 5-HT_{2B}-AhR link is also functional in the 5-HT_{2B}-expressing HepG2 hepatoma cell line. As a whole, our results point to AhR and AhR-target

5-HT_{2B} shapes the macrophage transcriptome via AhR

genes (e.g., *CYP1B1*, *AHRR*, *TIPARP*, ...) as effectors of the 5-HT_{2B} receptor in human macrophages.

The transcriptional effects of BW723C86 on human macrophages differ considerably from those triggered by 5-HT, most of which can be mimicked or abrogated by 5-HT₇ agonists or antagonists, respectively³⁹. Thus, although 5-HT_{2B} and 5-HT₇ are preferentially expressed by anti-inflammatory human macrophages⁴⁰ and promote a pro-fibrotic phenotype^{19,39,47,114,115}, their individual engagement leads to different transcriptional outcomes, with AhR-activating ability being restricted to 5-HT_{2B}. The failure of 5-HT to activate AhR in human macrophages might rely on the fact that 5-HT can bind to both 5-HT_{2B} and 5-HT₇, and can therefore initiate intracellular signaling from both receptors. Ligation of 5-HT₇ by 5-HT triggers protein kinase A (PKA)-dependent signaling and transcription in macrophages³⁹ and other cell types¹¹⁶, and results in the expression of PKA-regulated genes like *PDE2A*³⁹. Since PKA activation (or increased cAMP levels) represses AhR-dependent gene expression^{117,118}, it is therefore possible that the binding of 5-HT to 5-HT₇, and the resulting PKA activation^{39,116}, might prevent the activation of AhR secondary to 5-HT binding to 5-HT_{2B}. In line with this hypothesis, it is worth noting that *PDE2A* gene expression in macrophages is greatly upregulated after 5-HT binding to 5-HT₇³⁹, and that PDE2A interaction with the immunophilin-like protein XAP2 (a member of the molecular complex that retains unliganded AhR in the cytoplasm) inhibits dioxin-induced AhR nuclear translocation and transcription in hepatocytes¹¹⁹.

A recent report has illustrated that, in intestinal epithelial cells, 5-HT activates AhR and the expression of AhR target genes in a manner that is independent on 5-HT receptors but mediated by the serotonin transporter (SERT)¹²⁰. The BW723C86-mediated activation of

5-HT_{2B} shapes the macrophage transcriptome via AhR

AhR in macrophages (here reported) is unrelated to the 5-HT/SERT/AhR link because: 1) it can be inhibited by an antagonist of 5-HT_{2B}; 2) M-MØ are devoid of SERT expression⁸⁴; and 3) 5-HT does not upregulate *CYP1B1* expression in human monocyte-derived macrophages. However, the existence of these two pathways (5-HT/SERT/AhR in intestinal epithelial cells and BW723C86/5-HT_{2B}/AhR in macrophages) fits with the AhR-activating ability of tryptophan catabolites¹²¹ and supports the importance of 5-HT and related molecules in modulating AhR activity and, hence, inflammatory responses. In this regard, and besides a receptor for environmental toxins⁹⁹, AhR mediates vascular system development, immune system polarization and resolution of inflammatory responses^{122,123}, processes where 5-HT_{2B} has been also implicated^{16,70,71,93}. Thus, since 5-HT_{2B} agonists (and specially BW723C86) have been widely used *in vivo* to support the participation of 5-HT_{2B} in several animal models of disease^{73,74,86,87,89-91,124-126}, our results suggests the possibility that AhR might have influenced some of these previous results and might have even contributed to the pathological effects of deregulated 5-HT_{2B} expression or function. As an example, since AhR mediates IL-10 production and promotes immunological tolerance after macrophage capture of apoptotic cells¹²⁷, AhR might contribute to the anti-inflammatory effect of 5-HT_{2B} in mouse microglia⁷¹ and human monocytes⁷². On a related note, the assessment of the extent of the transcriptional action of BW723C86 evidenced that the effects of the agonist on the expression of more than 200 genes (including some encoding regulators of osteoclastogenesis) are not prevented by the 5-HT_{2B} antagonist SB204741 (Figure 1). This finding indicates the existence of 5-HT_{2B}-independent effects of BW723C86 and, in line with the above comments, suggest that some of the actions previously assigned to BW723C86 *in vivo* and *in vitro* might be actually unrelated to (independent on) the ability of BW723C86 to interact with 5-HT_{2B}.

5-HT_{2B} shapes the macrophage transcriptome via AhR

ACKNOWLEDGEMENTS

This work was supported by grants from Ministerio de Economía y Competitividad (SAF2014-52423-R and SAF2017-83785-R) to MAV and ALC, Grant 201619.31 from Fundación La Marató/TV3 to ALC, and Red de Investigación en Enfermedades Reumáticas (RIER, RD16/0012/0007), and cofinanced by the European Regional Development Fund “A way to achieve Europe” (ERDF). MCE and IR were funded by FPI predoctoral fellowship (BES-2009-021465 and BES-2015-071337, respectively) from Ministerio de Economía e Innovación.

AUTHORSHIP CONTRIBUTIONS:

CN, IR, MCE, EI and BA performed research and analyzed data; CN, IR, MCE, MAV and ALC designed the research and analyzed data; ALC wrote the paper.

DISCLOSURE OF CONFLICTS OF INTEREST

The authors declare no competing financial interests.

5-HT_{2B} shapes the macrophage transcriptome via AhR

REFERENCES

1. Lesurtel M, Soll C, Graf R, Clavien PA. Role of serotonin in the hepato-gastrointestinal tract: an old molecule for new perspectives. *Cell Mol Life Sci*. 2008;65(6):940-952.
2. Amireault P, Sibon D, Cote F. Life without peripheral serotonin: insights from tryptophan hydroxylase 1 knockout mice reveal the existence of paracrine/autocrine serotonergic networks. *ACS Chem Neurosci*. 2013;4(1):64-71.
3. Yano JM, Yu K, Donaldson GP, et al. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*. 2015;161(2):264-276.
4. Kaumann AJ, Levy FO. 5-hydroxytryptamine receptors in the human cardiovascular system. *Pharmacol Ther*. 2006;111(3):674-706.
5. Spiller R. Serotonin and GI clinical disorders. *Neuropharmacology*. 2008;55(6):1072-1080.
6. Nemecek GM, Coughlin SR, Handley DA, Moskowitz MA. Stimulation of aortic smooth muscle cell mitogenesis by serotonin. *Proc Natl Acad Sci U S A*. 1986;83(3):674-678.
7. Lesurtel M, Graf R, Aleil B, et al. Platelet-derived serotonin mediates liver regeneration. *Science*. 2006;312(5770):104-107.
8. Pakala R, Willerson JT, Benedict CR. Mitogenic effect of serotonin on vascular endothelial cells. *Circulation*. 1994;90(4):1919-1926.
9. Liu S, Miao R, Zhai M, et al. Effects and related mechanisms of serotonin on malignant biological behavior of hepatocellular carcinoma via regulation of Yap. *Oncotarget*. 2017;8(29):47412-47424.
10. Ahern GP. 5-HT and the immune system. *Curr Opin Pharmacol*. 2011;11(1):29-33.
11. Leon-Ponte M, Ahern GP, O'Connell PJ. Serotonin provides an accessory signal to enhance T-cell activation by signaling through the 5-HT₇ receptor. *Blood*. 2007;109(8):3139-3146.
12. Arzt E, Costas M, Finkielman S, Nahmod VE. Serotonin inhibition of tumor necrosis factor- α synthesis by human monocytes. *Life Sci*. 1991;48(26):2557-2562.
13. Cloez-Tayarani I, Petit-Bertron AF, Venters HD, Cavaillon JM. Differential effect of serotonin on cytokine production in lipopolysaccharide-stimulated human peripheral blood mononuclear cells: involvement of 5-hydroxytryptamine_{2A} receptors. *Int Immunol*. 2003;15(2):233-240.
14. Durk T, Panther E, Muller T, et al. 5-Hydroxytryptamine modulates cytokine and chemokine production in LPS-primed human monocytes via stimulation of different 5-HTR subtypes. *Int Immunol*. 2005;17(5):599-606.
15. Menard G, Turmel V, Bissonnette EY. Serotonin modulates the cytokine network in the lung: involvement of prostaglandin E₂. *Clin Exp Immunol*. 2007;150(2):340-348.
16. de Las Casas-Engel M, Corbi AL. Serotonin modulation of macrophage polarization: inflammation and beyond. *Adv Exp Med Biol*. 2014;824:89-115.

5-HT_{2B} shapes the macrophage transcriptome via AhR

17. Launay JM, Herve P, Callebort J, et al. Serotonin 5-HT_{2B} receptors are required for bone-marrow contribution to pulmonary arterial hypertension. *Blood*. 2012;119(7):1772-1780.
18. Morita T, McClain SP, Batia LM, et al. HTR7 Mediates Serotonergic Acute and Chronic Itch. *Neuron*. 2015;87(1):124-138.
19. Dees C, Akhmetshina A, Zerr P, et al. Platelet-derived serotonin links vascular disease and tissue fibrosis. *J Exp Med*. 2011;208(5):961-972.
20. Kim JJ, Khan WI. 5-HT₇ receptor signaling: improved therapeutic strategy in gut disorders. *Front Behav Neurosci*. 2014;8:396.
21. Levin AD, van den Brink GR. Selective inhibition of mucosal serotonin as treatment for IBD? *Gut*. 2014;63(6):866-867.
22. Kim JJ, Bridle BW, Ghia JE, et al. Targeted inhibition of serotonin type 7 (5-HT₇) receptor function modulates immune responses and reduces the severity of intestinal inflammation. *J Immunol*. 2013;190(9):4795-4804.
23. Guseva D, Holst K, Kaune B, et al. Serotonin 5-HT₇ receptor is critically involved in acute and chronic inflammation of the gastrointestinal tract. *Inflamm Bowel Dis*. 2014;20(9):1516-1529.
24. Ghia JE, Li N, Wang H, et al. Serotonin has a key role in pathogenesis of experimental colitis. *Gastroenterology*. 2009;137(5):1649-1660.
25. Li N, Ghia JE, Wang H, et al. Serotonin activates dendritic cell function in the context of gut inflammation. *Am J Pathol*. 2011;178(2):662-671.
26. Nocito A, Dahm F, Jochum W, et al. Serotonin regulates macrophage-mediated angiogenesis in a mouse model of colon cancer allografts. *Cancer Res*. 2008;68(13):5152-5158.
27. Svejda B, Kidd M, Timberlake A, et al. Serotonin and the 5-HT₇ receptor: the link between hepatocytes, IGF-1 and small intestinal neuroendocrine tumors. *Cancer Sci*. 2013;104(7):844-855.
28. Chabbi-Achengli Y, Coman T, Collet C, et al. Serotonin Is Involved in Autoimmune Arthritis through Th17 Immunity and Bone Resorption. *Am J Pathol*. 2016;186(4):927-937.
29. Barnes NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology*. 1999;38(8):1083-1152.
30. Jantsch J, Binger KJ, Muller DN, Titze J. Macrophages in homeostatic immune function. *Front Physiol*. 2014;5:146.
31. Sica A, Mantovani A. Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest*. 2012;122(3):787-795.
32. Martinez FO, Sica A, Mantovani A, Locati M. Macrophage activation and polarization. *Front Biosci*. 2008;13:453-461.
33. Schultze JL. Reprogramming of macrophages--new opportunities for therapeutic targeting. *Curr Opin Pharmacol*. 2016;26:10-15.
34. Schultze JL, Freeman T, Hume DA, Latz E. A transcriptional perspective on human macrophage biology. *Semin Immunol*. 2015;27(1):44-50.
35. Tsuchida Y, Hatao F, Fujisawa M, et al. Neuronal stimulation with 5-hydroxytryptamine 4 receptor induces anti-inflammatory actions via alpha7nACh receptors on muscularis macrophages associated with postoperative ileus. *Gut*. 2011;60(5):638-647.
36. Maehara T, Matsumoto K, Horiguchi K, et al. Therapeutic action of 5-HT₃ receptor antagonists targeting peritoneal macrophages in post-operative ileus. *Br J Pharmacol*. 2015;172(4):1136-1147.

5-HT_{2B} shapes the macrophage transcriptome via AhR

37. Kalkman HO, Feuerbach D. Antidepressant therapies inhibit inflammation and microglial M1-polarization. *Pharmacol Ther.* 2016;163:82-93.
38. Nazimek K, Strobel S, Bryniarski P, Kozłowski M, Filipczak-Bryniarska I, Bryniarski K. The role of macrophages in anti-inflammatory activity of antidepressant drugs. *Immunobiology.* 2016.
39. Dominguez-Soto A, Usategui A, Casas-Engel ML, et al. Serotonin drives the acquisition of a profibrotic and anti-inflammatory gene profile through the 5-HT₇R-PKA signaling axis. *Sci Rep.* 2017;7(1):14761.
40. de las Casas-Engel M, Dominguez-Soto A, Sierra-Filardi E, et al. Serotonin skews human macrophage polarization through HTR_{2B} and HTR₇. *J Immunol.* 2013;190(5):2301-2310.
41. Zmudzka E, Salaciak K, Sapa J, Pytko K. Serotonin receptors in depression and anxiety: Insights from animal studies. *Life Sci.* 2018.
42. Nebigil CG, Choi DS, Dierich A, et al. Serotonin 2B receptor is required for heart development. *Proc Natl Acad Sci U S A.* 2000;97(17):9508-9513.
43. West JD, Carrier EJ, Bloodworth NC, et al. Serotonin 2B Receptor Antagonism Prevents Heritable Pulmonary Arterial Hypertension. *PLoS One.* 2016;11(2):e0148657.
44. Fabre A, Marchal-Somme J, Marchand-Adam S, et al. Modulation of bleomycin-induced lung fibrosis by serotonin receptor antagonists in mice. *Eur Respir J.* 2008;32(2):426-436.
45. Ebrahimkhani MR, Oakley F, Murphy LB, et al. Stimulating healthy tissue regeneration by targeting the 5-HT_{(2)B} receptor in chronic liver disease. *Nat Med.* 2011;17(12):1668-1673.
46. Jaffre F, Bonnin P, Callebort J, et al. Serotonin and angiotensin receptors in cardiac fibroblasts coregulate adrenergic-dependent cardiac hypertrophy. *Circ Res.* 2009;104(1):113-123.
47. Jaffre F, Callebort J, Sarre A, et al. Involvement of the serotonin 5-HT_{2B} receptor in cardiac hypertrophy linked to sympathetic stimulation: control of interleukin-6, interleukin-1 β , and tumor necrosis factor- α cytokine production by ventricular fibroblasts. *Circulation.* 2004;110(8):969-974.
48. Gustafsson BI, Hauso O, Drozdov I, Kidd M, Modlin IM. Carcinoid heart disease. *Int J Cardiol.* 2008;129(3):318-324.
49. Segelcke D, Messlinger K. Putative role of 5-HT_{2B} receptors in migraine pathophysiology. *Cephalalgia.* 2016.
50. Roth BL. Drugs and valvular heart disease. *N Engl J Med.* 2007;356(1):6-9.
51. Matsunaga F, Gao L, Huang XP, Saven JG, Roth BL, Liu R. Molecular interactions between general anesthetics and the 5HT_{2B} receptor. *J Biomol Struct Dyn.* 2015;33(1):211-218.
52. Zhang S, Li B, Lovatt D, et al. 5-HT_{2B} receptors are expressed on astrocytes from brain and in culture and are a chronic target for all five conventional 'serotonin-specific reuptake inhibitors'. *Neuron Glia Biol.* 2010;6(2):113-125.
53. Li B, Zhang S, Zhang H, et al. Fluoxetine-mediated 5-HT_{2B} receptor stimulation in astrocytes causes EGF receptor transactivation and ERK phosphorylation. *Psychopharmacology (Berl).* 2008;201(3):443-458.
54. Diaz SL, Doly S, Narboux-Neme N, et al. 5-HT_{(2)B} receptors are required for serotonin-selective antidepressant actions. *Mol Psychiatry.* 2012;17(2):154-163.
55. Setola V, Dukat M, Glennon RA, Roth BL. Molecular determinants for the interaction of the valvulopathic anorexigen norfenfluramine with the 5-HT_{2B} receptor. *Mol Pharmacol.* 2005;68(1):20-33.

5-HT_{2B} shapes the macrophage transcriptome via AhR

56. Zanettini R, Antonini A, Gatto G, Gentile R, Tesei S, Pezzoli G. Valvular heart disease and the use of dopamine agonists for Parkinson's disease. *N Engl J Med.* 2007;356(1):39-46.
57. Schade R, Andersohn F, Suissa S, Haverkamp W, Garbe E. Dopamine agonists and the risk of cardiac-valve regurgitation. *N Engl J Med.* 2007;356(1):29-38.
58. Georgia S, Bhushan A. Pregnancy hormones boost beta cells via serotonin. *Nat Med.* 2010;16(7):756-757.
59. Kim H, Toyofuku Y, Lynn FC, et al. Serotonin regulates pancreatic beta cell mass during pregnancy. *Nat Med.* 2010;16(7):804-808.
60. Svejda B, Kidd M, Giovinazzo F, et al. The 5-HT_{2B} receptor plays a key regulatory role in both neuroendocrine tumor cell proliferation and the modulation of the fibroblast component of the neoplastic microenvironment. *Cancer.* 2010;116(12):2902-2912.
61. Soll C, Jang JH, Riener MO, et al. Serotonin promotes tumor growth in human hepatocellular cancer. *Hepatology.* 2010;51(4):1244-1254.
62. Liang C, Chen W, Zhi X, et al. Serotonin promotes the proliferation of serum-deprived hepatocellular carcinoma cells via upregulation of FOXO3a. *Mol Cancer.* 2013;12:14.
63. Asada M, Ebihara S, Yamanda S, et al. Depletion of serotonin and selective inhibition of 2B receptor suppressed tumor angiogenesis by inhibiting endothelial nitric oxide synthase and extracellular signal-regulated kinase 1/2 phosphorylation. *Neoplasia.* 2009;11(4):408-417.
64. Tharayil VS, Wouters MM, Stanich JE, et al. Lack of serotonin 5-HT_{2B} receptor alters proliferation and network volume of interstitial cells of Cajal in vivo. *Neurogastroenterol Motil.* 2010;22(4):462-469, e109-410.
65. Wouters MM, Gibbons SJ, Roeder JL, et al. Exogenous serotonin regulates proliferation of interstitial cells of Cajal in mouse jejunum through 5-HT_{2B} receptors. *Gastroenterology.* 2007;133(3):897-906.
66. Collet C, Schiltz C, Geoffroy V, Maroteaux L, Launay JM, de Vernejoul MC. The serotonin 5-HT_{2B} receptor controls bone mass via osteoblast recruitment and proliferation. *Faseb J.* 2008;22(2):418-427.
67. Chabbi-Achengli Y, Launay JM, Maroteaux L, de Vernejoul MC, Collet C. Serotonin 2B receptor (5-HT_{2B} R) signals through prostacyclin and PPAR- α /delta in osteoblasts. *PLoS One.* 2013;8(9):e75783.
68. Bennet H, Mollet IG, Balhuizen A, et al. Serotonin (5-HT) receptor 2b activation augments glucose-stimulated insulin secretion in human and mouse islets of Langerhans. *Diabetologia.* 2016;59(4):744-754.
69. Naito K, Moteki H, Kimura M, Natsume H, Ogihara M. Serotonin 5-HT_{2B} Receptor-Stimulated DNA Synthesis and Proliferation Are Mediated by Autocrine Secretion of Transforming Growth Factor- α in Primary Cultures of Adult Rat Hepatocytes. *Biol Pharm Bull.* 2016;39(4):570-577.
70. El Oussini H, Bayer H, Scekcic-Zahirovic J, et al. Serotonin 2B receptor slows disease progression and prevents degeneration of spinal cord mononuclear phagocytes in amyotrophic lateral sclerosis. *Acta Neuropathol.* 2016;131(3):465-480.
71. Kolodziejczak M, Bechade C, Gervasi N, et al. Serotonin Modulates Developmental Microglia via 5-HT_{2B} Receptors: Potential Implication during Synaptic Refinement of Retinogeniculate Projections. *ACS Chem Neurosci.* 2015;6(7):1219-1230.

5-HT_{2B} shapes the macrophage transcriptome via AhR

72. Palmqvist N, Siller M, Klint C, Sjodin A. A human and animal model-based approach to investigating the anti-inflammatory profile and potential of the 5-HT_{2B} receptor antagonist AM1030. *J Inflamm (Lond)*. 2016;13:20.
73. Kennett GA, Bright F, Trail B, Baxter GS, Blackburn TP. Effects of the 5-HT_{2B} receptor agonist, BW 723C86, on three rat models of anxiety. *Br J Pharmacol*. 1996;117(7):1443-1448.
74. Kennett GA, Trail B, Bright F. Anxiolytic-like actions of BW 723C86 in the rat Vogel conflict test are 5-HT_{2B} receptor mediated. *Neuropharmacology*. 1998;37(12):1603-1610.
75. Kursar JD, Nelson DL, Wainscott DB, Cohen ML, Baez M. Molecular cloning, functional expression, and pharmacological characterization of a novel serotonin receptor (5-hydroxytryptamine_{2F}) from rat stomach fundus. *Mol Pharmacol*. 1992;42(4):549-557.
76. Baxter GS. Novel discriminatory ligands for 5-HT_{2B} receptors. *Behav Brain Res*. 1996;73(1-2):149-152.
77. Ismaiel AM, Titeler M, Miller KJ, Smith TS, Glennon RA. 5-HT₁ and 5-HT₂ binding profiles of the serotonergic agents alpha-methylserotonin and 2-methylserotonin. *J Med Chem*. 1990;33(2):755-758.
78. Forbes IT, Jones GE, Murphy OE, Holland V, Baxter GS. N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea: a novel, high-affinity 5-HT_{2B} receptor antagonist. *J Med Chem*. 1995;38(6):855-857.
79. Hochberg Y, Benjamini Y. More powerful procedures for multiple significance testing. *Stat Med*. 1990;9(7):811-818.
80. Metsalu T, Vilo J. ClustVis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Res*. 2015;43(W1):W566-570.
81. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res*. 2016;44(W1):W90-W97.
82. Chen EY, Tan CM, Kou Y, et al. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics*. 2013;14:128.
83. Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545-15550.
84. Gonzalez-Dominguez E, Dominguez-Soto A, Nieto C, et al. Atypical Activin A and IL-10 Production Impairs Human CD16⁺ Monocyte Differentiation into Anti-Inflammatory Macrophages. *J Immunol*. 2016;196(3):1327-1337.
85. Maroteaux L, Ayme-Dietrich E, Aubertin-Kirch G, et al. New therapeutic opportunities for 5-HT₂ receptor ligands. *Pharmacol Ther*. 2017;170:14-36.
86. Miwa H, Koseki J, Oshima T, et al. Impairment of gastric accommodation induced by water-avoidance stress is mediated by 5-HT_{2B} receptors. *Neurogastroenterol Motil*. 2016;28(5):765-778.
87. Belmer A, Quentin E, Diaz SL, et al. Positive regulation of raphe serotonin neurons by serotonin 2B receptors. *Neuropsychopharmacology*. 2018;43(7):1623-1632.
88. Doly S, Valjent E, Setola V, et al. Serotonin 5-HT_{2B} receptors are required for 3,4-methylenedioxymethamphetamine-induced hyperlocomotion and 5-HT release in vivo and in vitro. *J Neurosci*. 2008;28(11):2933-2940.
89. Martin RS, Martin GR. Investigations into migraine pathogenesis: time course for effects of m-CPP, BW723C86 or glyceryl trinitrate on appearance of Fos-like

5-HT_{2B} shapes the macrophage transcriptome via AhR

- immunoreactivity in rat trigeminal nucleus caudalis (TNC). *Cephalalgia*. 2001;21(1):46-52.
90. Walker EA, Brown EK, Jr., Sterious SN. In vivo Schild regression analyses using nonselective 5-HT_{2C} receptor antagonists in a rat operant behavioral assay. *Psychopharmacology (Berl)*. 2007;193(2):187-197.
91. Kennett GA, Ainsworth K, Trail B, Blackburn TP. BW 723C86, a 5-HT_{2B} receptor agonist, causes hyperphagia and reduced grooming in rats. *Neuropharmacology*. 1997;36(2):233-239.
92. Deraet M, Manivet P, Janoshazi A, et al. The natural mutation encoding a C terminus-truncated 5-hydroxytryptamine 2B receptor is a gain of proliferative functions. *Mol Pharmacol*. 2005;67(4):983-991.
93. Hutcheson JD, Setola V, Roth BL, Merryman WD. Serotonin receptors and heart valve disease--it was meant 2B. *Pharmacol Ther*. 2011;132(2):146-157.
94. Goldberg E, Grau JB, Fortier JH, Salvati E, Levy RJ, Ferrari G. Serotonin and catecholamines in the development and progression of heart valve diseases. *Cardiovasc Res*. 2017;113(8):849-857.
95. Sierra-Filardi E, Puig-Kroger A, Blanco FJ, et al. Activin A skews macrophage polarization by promoting a proinflammatory phenotype and inhibiting the acquisition of anti-inflammatory macrophage markers. *Blood*. 2011;117(19):5092-5101.
96. Janssen W, Schymura Y, Novoyatleva T, et al. 5-HT_{2B} receptor antagonists inhibit fibrosis and protect from RV heart failure. *Biomed Res Int*. 2015;2015:438403.
97. Riera-Borrull M, Cuevas VD, Alonso B, et al. Palmitate Conditions Macrophages for Enhanced Responses toward Inflammatory Stimuli via JNK Activation. *J Immunol*. 2017;199(11):3858-3869.
98. Sparfel L, Pinel-Marie ML, Boize M, et al. Transcriptional signature of human macrophages exposed to the environmental contaminant benzo(a)pyrene. *Toxicol Sci*. 2010;114(2):247-259.
99. Mulero-Navarro S, Fernandez-Salguero PM. New Trends in Aryl Hydrocarbon Receptor Biology. *Front Cell Dev Biol*. 2016;4:45.
100. Lo R, Matthews J. High-resolution genome-wide mapping of AHR and ARNT binding sites by ChIP-Seq. *Toxicol Sci*. 2012;130(2):349-361.
101. Lawrence BP, Denison MS, Novak H, et al. Activation of the aryl hydrocarbon receptor is essential for mediating the anti-inflammatory effects of a novel low-molecular-weight compound. *Blood*. 2008;112(4):1158-1165.
102. Zhao B, Degroot DE, Hayashi A, He G, Denison MS. CH223191 is a ligand-selective antagonist of the Ah (Dioxin) receptor. *Toxicol Sci*. 2010;117(2):393-403.
103. Rannug A, Rannug U, Rosenkranz HS, et al. Certain photooxidized derivatives of tryptophan bind with very high affinity to the Ah receptor and are likely to be endogenous signal substances. *J Biol Chem*. 1987;262(32):15422-15427.
104. Kukita T, Kukita A, Harada H, Iijima T. Regulation of osteoclastogenesis by antisense oligodeoxynucleotides specific to zinc finger nuclear transcription factors Egr-1 and WT1 in rat bone marrow culture system. *Endocrinology*. 1997;138(10):4384-4389.
105. Srivastava S, Weitzmann MN, Kimble RB, et al. Estrogen blocks M-CSF gene expression and osteoclast formation by regulating phosphorylation of Egr-1 and its interaction with Sp-1. *J Clin Invest*. 1998;102(10):1850-1859.
106. Yang M, Birnbaum MJ, MacKay CA, Mason-Savas A, Thompson B, Odgren PR. Osteoclast stimulatory transmembrane protein (OC-STAMP), a novel protein induced by RANKL that promotes osteoclast differentiation. *J Cell Physiol*. 2008;215(2):497-505.

5-HT_{2B} shapes the macrophage transcriptome via AhR

107. Park JH, Lee NK, Lee SY. Current Understanding of RANK Signaling in Osteoclast Differentiation and Maturation. *Mol Cells*. 2017;40(10):706-713.
108. Novack DV, Teitelbaum SL. The osteoclast: friend or foe? *Annu Rev Pathol*. 2008;3:457-484.
109. Miyauchi Y, Ninomiya K, Miyamoto H, et al. The Blimp1-Bcl6 axis is critical to regulate osteoclast differentiation and bone homeostasis. *J Exp Med*. 2010;207(4):751-762.
110. Nishikawa K, Nakashima T, Hayashi M, et al. Blimp1-mediated repression of negative regulators is required for osteoclast differentiation. *Proc Natl Acad Sci U S A*. 2010;107(7):3117-3122.
111. Kimura A, Naka T, Nakahama T, et al. Aryl hydrocarbon receptor in combination with Stat1 regulates LPS-induced inflammatory responses. *J Exp Med*. 2009;206(9):2027-2035.
112. Millan MJ, Marin P, Bockaert J, Mannoury la Cour C. Signaling at G-protein-coupled serotonin receptors: recent advances and future research directions. *Trends Pharmacol Sci*. 2008;29(9):454-464.
113. Nebigil CG, Etienne N, Schaerlinger B, Hickel P, Launay JM, Maroteaux L. Developmentally regulated serotonin 5-HT_{2B} receptors. *Int J Dev Neurosci*. 2001;19(4):365-372.
114. Nebigil CG, Jaffre F, Messaddeq N, et al. Overexpression of the serotonin 5-HT_{2B} receptor in heart leads to abnormal mitochondrial function and cardiac hypertrophy. *Circulation*. 2003;107(25):3223-3229.
115. Elangbam CS. Drug-induced valvulopathy: an update. *Toxicol Pathol*. 2010;38(6):837-848.
116. Raymond JR, Mukhin YV, Gelasco A, et al. Multiplicity of mechanisms of serotonin receptor signal transduction. *Pharmacol Ther*. 2001;92(2-3):179-212.
117. Oesch-Bartlomowicz B, Oesch F. Role of cAMP in mediating AHR signaling. *Biochem Pharmacol*. 2009;77(4):627-641.
118. Oesch-Bartlomowicz B, Huelster A, Wiss O, et al. Aryl hydrocarbon receptor activation by cAMP vs. dioxin: divergent signaling pathways. *Proc Natl Acad Sci U S A*. 2005;102(26):9218-9223.
119. de Oliveira SK, Hoffmeister M, Gambaryan S, Muller-Esterl W, Guimaraes JA, Smolenski AP. Phosphodiesterase 2A forms a complex with the co-chaperone XAP2 and regulates nuclear translocation of the aryl hydrocarbon receptor. *J Biol Chem*. 2007;282(18):13656-13663.
120. Manzella C, Singhal M, Alrefai WA, Saksena S, Dudeja PK, Gill RK. Serotonin is an endogenous regulator of intestinal CYP1A1 via AhR. *Sci Rep*. 2018;8(1):6103.
121. Mezrich JD, Fechner JH, Zhang X, Johnson BP, Burlingham WJ, Bradfield CA. An interaction between kynurenine and the aryl hydrocarbon receptor can generate regulatory T cells. *J Immunol*. 2010;185(6):3190-3198.
122. Veldhoen M, Hirota K, Westendorf AM, et al. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature*. 2008;453(7191):106-109.
123. Quintana FJ, Basso AS, Iglesias AH, et al. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. *Nature*. 2008;453(7191):65-71.
124. Gunther S, Maroteaux L, Schwarzwacher SW. Endogenous 5-HT_{2B} receptor activation regulates neonatal respiratory activity in vitro. *J Neurobiol*. 2006;66(9):949-961.
125. Diaz SL, Maroteaux L. Implication of 5-HT(2B) receptors in the serotonin syndrome. *Neuropharmacology*;61(3):495-502.

5-HT_{2B} shapes the macrophage transcriptome via AhR

126. Diaz SL, Maroteaux L. Implication of 5-HT(2B) receptors in the serotonin syndrome. *Neuropharmacology*. 2011;61(3):495-502.
127. Shinde R, Hezaveh K, Halaby MJ, et al. Apoptotic cell-induced AhR activity is required for immunological tolerance and suppression of systemic lupus erythematosus in mice and humans. *Nat Immunol*. 2018;19(6):571-582.
128. Cuevas VD, Anta L, Samaniego R, et al. MAFB Determines Human Macrophage Anti-Inflammatory Polarization: Relevance for the Pathogenic Mechanisms Operating in Multicentric Carpotarsal Osteolysis. *J Immunol*. 2017;198(5):2070-2081.

5-HT_{2B} shapes the macrophage transcriptome via AhR

FIGURE LEGENDS

Figure 1. The 5-HT_{2B} agonist BW723C86 modifies the gene signature of human M-MØ. A.

Experimental design of the gene profiling experiments. **B.** Number of annotated genes whose expression is significantly (*adj p*<0.002) upregulated (UP) or downregulated (DOWN) in M-MØ after exposure to BW723C86 (BW, 6h) in the absence (BW *vs* Control, CNT) or presence of SB204741 (SB) (BW+SB *vs* SB). **C.** Expression of the indicated genes in non-treated (-) or BW723C86-treated (6h) (+) M-MØ. Results are expressed as the mRNA level of each gene relative to the *TBP* mRNA level in the same sample. Mean and SEM of seven independent experiments is shown (*, *p* < 0.05; **, *p* < 0.01). **D.** GSEA analysis of the *t* statistic-ranked list of genes obtained from the BW723C86-treated M-MØ *versus* control M-MØ limma analysis, using the Molecular Signatures Database available at the GSEA webpage (<http://software.broadinstitute.org/gsea/index.jsp>)⁸³. Size and FDR *q-values* for the shown gene sets are shown at the top of the panel. **E.** Number of SB204741-sensitive and SB204741-insensitive annotated genes within the group of genes significantly (*adj p*<0.002) upregulated (BW723C86>Control, left circle) or downregulated (BW723C86<Control, right circle) by BW723C86. **F.** Volcano plot representation of the microarray data. Gene expression profiles of BW-treated M-MØ versus control M-MØ (left panel) and SB+BW-treated M-MØ versus SB-treated M-MØ (right panel) are plotted according to the log₂ fold change (X axis) and log₁₀ adjusted *p*-value (Y axis). In both plots, the shaded area includes the genes with *adj p*<0.05. Some of the differentially expressed genes in each case are indicated (Note the distinct relative position of *CYP1B1* and *AHRR* in both plots).

Figure 2. The 5-HT_{2B} agonist BW723C86 conditions human macrophages for altered LPS-

stimulated cytokine and gene expression. A. Experimental design to assess the effect of BW723C86 and SB204741 on LPS-stimulated cytokine production and gene expression in M-

5-HT_{2B} shapes the macrophage transcriptome via AhR

MØ. B. Production of CCL2 and TNF α from LPS-stimulated (18h) M-MØ that had been non-treated (-) or exposed (6 h) to BW723C86, SB204741 or both (+), as determined by ELISA. Mean and SEM of 10/12 independent experiments are shown (*, $p < 0.05$; ***, $p < 0.001$). **C.** Expression of the indicated genes in LPS-stimulated (4h) M-MØ that had been non-treated (-) or exposed (6 h) to BW723C86, SB204741 or both (+), as determined by qRT-PCR. Results are expressed as the mRNA level of each gene relative to the level of *TBP* mRNA in the same sample. Mean and SEM of six independent experiments is shown (*, $p < 0.05$; ***, $p < 0.001$).

Figure 3. BW723C86 upregulates the expression of AhR target genes in a 5-HT_{2B} - dependent manner. **A.** Non-supervised hierarchical clustering using the ClustVis webtool (<https://biit.cs.ut.ee/clustvis/>)⁸⁰ on the mean expression level of genes significantly (adj $p < 0.002$) regulated by BW723C86. For the first cluster, the pattern of expression of three representative genes and the ENRICH data (<http://amp.pharm.mssm.edu/Enrichr/>)^{81,82} on the significant enrichment of specific signaling pathway gene sets, are shown. **B.** Gene set enrichment analysis on the “t statistic-ranked” list of genes obtained from the BW723C86-treated M-MØ versus untreated-M-MØ limma analysis, using the indicated gene sets from the GSEA webpage (<http://software.broadinstitute.org/gsea/index.jsp>)⁸³ (upper panel) or retrieved from⁹⁸ (lower panel). **C.** Relative expression of the indicated genes in non-treated (-) and BW723C86-treated M-MØ (6h) (+) (n=7-11; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). **D.** Relative expression of *CYP1B1* and *TIPARP* in non-treated (-) and M-MØ treated with BW723C86 (+) for the indicated periods of time (n=3; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). **E.** Relative expression of *CYP1B1* in non-treated (0) or M-MØ treated for 6h with the indicated concentrations of BW723C86 (n=3; *, $p < 0.05$). **F.** Relative expression of *CYP1B1* in non-treated (-) or HepG2 hepatoma cells treated with BW723C86 (+) for the indicated

5-HT_{2B} shapes the macrophage transcriptome via AhR

periods of time (n=3 *, $p < 0.05$; **, $p < 0.01$). **G.** Relative expression of *CYP1B1*, *AHRR* and *TIPARP* in non-treated (-) or M-MØ treated for 6h with BW723C86, SB204741 or both (+) (n=5; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). **(D-G)** Results are shown as the expression of each gene after the different treatments and relative to its expression in control (untreated) samples. In all cases, mean and SEM are shown.

Figure 4. *CYP1B1* upregulation by BW723C86 is dependent on 5-HT_{2B} and requires AhR activation. **A.** Relative expression of *CYP1B1* and *TIPARP* in M-MØ either untreated (-) or treated for 6h with BW723C86 (BW) in the absence or presence (+) of the AhR antagonist CH223191. M-MØ were treated in parallel with the AhR agonist FICZ (6h) for control purposes. Results are shown as the expression of each gene after the different treatments and relative to its expression in control (untreated) samples. In all cases, mean and SEM are shown (n=4; *, $p < 0.05$; ***, $p < 0.0005$). **B-C.** *AHR* mRNA (**B**) and AhR protein level (**C**) in M-MØ transfected with control siRNA (siC) or an AhR-specific siRNA (siAhR). 18 hours after transfection, M-MØ were washed and lysed for qRT-PCR (**B**) or Western blot (**C**). Six independent experiments were performed for *AHR* mRNA determination by qRT-PCR (**B**) (n=6; ***, $p < 0.001$), and Western blots from two independent knockdown experiments are shown in (**C**). **D.** Relative expression of *CYP1B1* and *TIPARP* in M-MØ transfected with control siRNA (siC) or AhR-specific siRNA (siAhR) and either left untreated (-) or treated for 6h with BW723C86 (+) (n=6; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). In **A-D**, results are shown as the expression of each gene after the different treatments and relative to its expression in control (untreated) samples. In all cases, mean and SEM are shown. **E.** AhR-dependent transcriptional activity in XRE-Luc transfected M-MØ left untreated (-) or treated with BW723C86 (14h) in the absence or in the presence of the AhR antagonist CH223191, added 1 hour before agonist addition. Mean and SEM of the AhR-dependent luciferase

5-HT_{2B} shapes the macrophage transcriptome via AhR

activity of five independent experiments is shown (n=5; *, $p < 0.05$; **, $p < 0.01$). **F.** AhR-dependent transcriptional activity in XRE-Luc-transfected HepG2 cells left untreated (-) or treated with BW723C86 (16h) in the absence or in the presence of the indicated concentrations of the 5-HT_{2B} antagonist SB204741 (n=4; ***, $p < 0.001$). **G.** AhR-dependent transcriptional activity in XRE-Luc-transfected HepG2 cells left untreated (-) or treated with BW723C86 (16h) in the absence or presence of the AhR antagonist CH223191, which was added 1 hour before agonist addition. HepG2 cells were treated with FICZ for control purposes (n=4; ***, $p < 0.001$). In **F-G**, results are shown as the mean and SEM of the Relative AhR-dependent luciferase activity of each sample (referred to the luciferase activity measured in untreated samples).

Figure 5. AhR activation in human macrophages is also triggered by other 5-HT_{2B} ligands and is prevented by serotonin. **A.** Relative expression of *CYP1B1* in M-MØ treated for 6h with DMSO (vehicle, -), BW723C86 (+) or with increasing concentrations of APB (10 nM, 50 nM, 100 nM or 1 µM) (n=9). **B.** Relative expression of *CYP1B1* in non-treated (-) or M-MØ treated for 6h with BW723C86 (+) or with increasing concentrations of α-Methyl 5HT (AMS) (10 nM, 20 nM, 50 nM, 100µM or 500 µM) (left panel) (n=6), and either in the absence or presence of SB204741 using 100µM or 500 µM AMS (right panel) (n=4). **(A-B)** Results are shown as the expression of each gene after the different treatments and relative to its expression in control (untreated) samples, and mean and SEM are shown ($p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). **C.** Relative expression of the indicated genes in M-MØ either untreated (-) or treated for 6h with 10 µM BW723C86, 5-HT (10 µM or 100 µM) or both (n=4). Shown is the expression of each gene after the different treatments and relative to its expression in control (untreated) samples. Mean and SEM are shown ($p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$).

5-HT_{2B} shapes the macrophage transcriptome via AhR

Figure 6. BW723C86 also regulates the macrophage transcriptome in a 5-HT_{2B}-independent manner. Relative expression of the indicated genes in non-treated (-) or M-MØ treated for 6h with BW723C86 after a previous treatment (1h) with either SB204741 or CH223191 (n=5; ***, $p < 0.001$). Results are shown as the expression of each gene after the different treatments and relative to its expression in control (untreated) samples. In all cases, mean and SEM are shown.

Figure 7. BW723C86 prevents the generation of osteoclasts from human peripheral blood monocytes. A-B. Expression of the indicated genes in monocyte-derived osteoclasts generated after a 12 day-culture with M-CSF and RANKL in the absence (-) or presence of BW723C86 (+) (n=4; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$). Results are shown as the expression of each gene in BW723C86-treated cells and relative to its expression in control (untreated) samples. In all cases, mean and SEM are shown. **C-D.** Phase-contrast microscopy (**C**) and TRAP-staining (**D**) of untreated (CONTROL, left panels) or BW723C86-treated monocytes cultured for 12 days in the presence of M-CSF and RANKL for differentiation into osteoclasts (right panels). Two independent experiments are shown in panel C.

Supplementary Figure 1. A. Selected GSEA on the “t statistic-ranked” list of genes obtained from the BW723C86-treated M-MØ versus untreated-M-MØ limma analysis, using the C5 collection within the Molecular Signatures Database (MSigDB) gene sets (<http://software.broadinstitute.org/gsea/index.jsp>). **B.** GSEA on the “t statistic-ranked” list of genes obtained from the BW723C86-treated M-MØ versus untreated-M-MØ limma analysis, using the “Proinflammatory gene set” (left) and “Anti-inflammatory gene set” (right) previously defined^{84,95,128}.

5-HT_{2B} shapes the macrophage transcriptome via AhR

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7