Molecular changes in *Mesembryanthemum crystallinum* guard cells underlying the C₃ to CAM transition

Wenwen Kong^{1,2,3*}, Mi-Jeong Yoo^{2*}, Jerald D. Noble^{4,5}, Theresa M Kelley^{2,5}, Jing Li¹, Matias Kirst^{4,5}, Sarah M. Assmann⁶, Sixue Chen^{2,5,7}

¹ College of Life Sciences, Northeast Agricultural University, Harbin, China;

² Department of Biology, University of Florida (UF), Gainesville, FL, USA;

³ Guangdong Province Key Laboratory for Plant Epigenetics, College of Life Science and Oceanography, Shenzhen University, 518060 Shenzhen, China;

⁴ School of Forest Resources and Conservation, University of Florida, Gainesville, FL, USA;

⁵ Genetics Institute, University of Florida, Gainesville, FL, USA;

⁶ Department of Biology, Pennsylvania State University, University Park, Pennsylvania, USA;

⁷ Interdisciplinary Center for Biotechnology Research, UF, Gainesville, FL, USA

*Equal contribution

Short running title: **C₃ to CAM transition of ice plant guard cells**

Correspondence:

Professor Matias Kirst, e-mail: mkirst@ufl.edu

Professor Sarah M. Assmann, e-mail: sma3@psu.edu

Professor Sixue Chen (lead contact), e-mail: schen@ufl.edu

ABSTRACT

Crassulacean acid metabolism (CAM) is a specialized type of photosynthesis: stomata close during the day, enhancing water conservation, and open at night, allowing CO₂ uptake. *Mesembryanthemum crystallinum* (common ice plant) is a facultative CAM species that can shift from C₃ photosynthesis to CAM under salt or drought stresses. However, the molecular mechanisms underlying the stress induced transition from C₃ to CAM remain unknown. Here we determined the transition time from C₃ to CAM in *M. crystallinum* under salt stress. In parallel, single-cell-type transcriptomic profiling by 3'-mRNA sequencing was conducted in guard cells to determine the molecular changes in this key cell type during the transition. In total, 495 transcripts showed differential expression between control and salt-treated samples during the transition, including 285 known guard cell genes, seven CAM-related genes, 18 transcription factors, and 185 other genes previously not found to be expressed in guard cells. *PEPC1* and *PPCK1*, which encode key enzymes of CAM photosynthesis, were up-regulated in guard cells after seven days of salt treatment, indicating that guard cells themselves can transition from C₃ to CAM. This study provides important information towards introducing CAM stomatal behavior into C₃ crops to enhance water use efficiency.

Keywords: common ice plant, guard cell, water use efficiency, salt stress, C₃ to CAM transition

Summary statement

We determined the timing of salt induced transition of common ice plant from C₃ to CAM and identified transcriptomic changes during the transition. The data support the notion that guard cells themselves can transition from C₃ to CAM.

1. INTRODUCTION

Global climate change is causing an increase in frequency of extreme drought and heat events and reducing fresh water and arable land for agriculture (Kaushal *et al.* 2017; Ziska *et al.* 2016). The impacts of extreme weather and drought are exacerbated by the demands of a growing human population, predicted to reach nine billion by 2050 (Borland *et al.* 2014). Therefore, improving the water-use efficiency (WUE) of agricultural crops is crucial to sustain productivity under increasing abiotic stresses and to expand cultivation to marginal lands, thereby enhancing food security.

In crassulacean acid metabolism (CAM), a specialized type of photosynthesis, stomata close during the day and open at night. CO₂ is taken up at night and phosphoenolpyruvate (PEP) is converted to oxaloacetate by phosphoenolpyruvate carboxylase (PEPC). Oxaloacetate can be subsequently transformed into malate by malate dehydrogenase and transported into the vacuole. During the day, the organic acids are exported from the vacuoles and decarboxylated to produce PEP or pyruvate and release CO₂ for light-driven carboxylation via ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) in the Calvin cycle (Owen & Griffiths 2013). The PEP is recycled as the substrate to assimilate CO₂ during the night or used for synthesis of carbohydrates (Borland *et al.* 2014). By shifting atmospheric CO₂ uptake to nighttime, when evapotranspiration rates are drastically reduced compared to the day, CAM plants achieve 3- to 6-times higher WUE than C₄ and C₃ plants (Nobel 1996). Therefore, introducing CAM into C₃ crops could greatly improve WUE and drought tolerance of C₃ plants.

Mesembryanthemum crystallinum (common ice plant) is a facultative CAM plant – it can shift between C_3 and CAM. When *M. crystallinum* is grown under non-stress conditions, it can complete its life cycle solely with C_3 photosynthesis (Adams *et al.* 1998). However, under stress conditions, such as water-deficit, salinity or high light, *M. crystallinum* can perform all the physiological features of CAM (Winter & Holtum 2005; Winter & Ziegler 1992). The shift from C_3 to CAM in *M. crystallinum* may be mediated by a calcium-dependent signaling pathway. Pretreating leaves with a calcium chelator (ethyleneglycol-bis(aminoethyl ether)-N,N-tetraacetic acid) inhibits stress-induced transcription of *PEPC1*, NAD-glyceraldehyde-3-phosphate dehydrogenase gene 1 (*GapC1*) and cytosolic NAD-malate dehydrogenase gene 1 (*Mdh1*), which are all important for CAM (Taybi & Cushman 1999). In addition, increased activity of the antioxidative stress system (e.g., superoxide dismutase) caused by salinity and high irradiance (Hurst *et al.* 2004), or H_2O_2 applied to *M. crystallinum* roots (Surowka *et al.* 2016), can also induce the C_3 to CAM transition. The inducibility of CAM and the biochemistry of C_3 and CAM in the same cells (unlike C_4 photosynthesis) make *M. crystallinum* an excellent system to study the mechanisms underlying the C_3 to CAM transition.

Previous studies have revealed omics level changes in *M. crystallinum* in response to salt stress treatment. The first microarray study by Cushman *et al.* (2008) used five-week-old plants (as C_3), and plants plus 14 days of salt stress (as CAM). A total of 1457 genes showed more than two-fold changes in mRNA steady-state levels between control (C_3) and salt treatment (CAM). Many of the differentially regulated genes are involved in CAM-related C_4 acid carboxylation/decarboxylation, glycolysis/gluconeogenesis, starch metabolism, protein degradation, transcriptional activation, signaling, stress response, and transport (Cushman *et al.* 2008). Using

next-generation sequencing, Tsukagoshi and co-workers (2015) identified 53,516 cDNA contigs from *M. crystallinum* roots and provided a transcriptome database (Tsukagoshi *et al.* 2015). They found that ABA responsive genes, a sodium transporter (*HKT1*), and peroxidase genes exhibited opposite responses to 140 mM NaCl treatment in *Arabidopsis* (C3) and ice plant (CAM) (Tsukagoshi *et al.* 2015). Oh and co-workers (2015) constructed a reference transcriptome containing 37,341 transcripts from control and salt-treated epidermal bladder cells (EBCs) of *M. crystallinum*, and 7% of the transcripts related to ion transport and signaling were salt stress responsive. At the small RNA level, Chiang *et al.* (2016) used roots of 3-day-old *M. crystallinum* seedlings and found 135 conserved miRNAs belonging to 21 families. The expression of mcr-miR159b and 166b, predicted to target transcription factors such as MYB domain protein 33, homeobox-leucine zipper family protein (HD-ZIP), and TCP4, were induced by salt treatment and mcr-miR319 expression was repressed. The proteome, ionome, and metabolome of *M. crystallinum* epidermal bladder cells have also been investigated (Barkla & Estrella 2015; Barkla *et al.* 2016).

The above studies identified numerous candidate genes, miRNAs, proteins and metabolites, which may inform efforts to improve plant salt tolerance. However, they have focused on either the steady-state levels of the C₃ and/or CAM photosynthetic modes or the stress tolerance of *M. crystallinum*, not the transition from C₃ to CAM. Therefore, the molecular mechanisms underlying this stress-induced transition remain unknown. Here we determined the critical transition time from C₃ to CAM in *M. crystallinum* by measuring several key attributes, including titratable acidity, stomatal aperture, gas exchange, CAM-related enzyme activity, and CAM-related gene expression. Since reversed stomatal movement behavior is essential for CAM

development, we tested the hypothesis that guard cells themselves undergo transition from C3 to CAM using single-cell-type transcriptomics.

2. MATERIALS AND METHODS

2.1. Plant material and growth conditions

Mesembryanthemum crystallinum seeds were germinated on vermiculite moistened with 0.5× Hoagland’s solution (Hoagland & Arnon 1950). One-week old seedlings were transferred to 32 ounce containers. The plants were grown in a growth chamber under $200 \mu\text{mol m}^{-2} \text{s}^{-1}$ white light with a 12h (26°C) day /12h (18°C) night cycle, and watered daily with 0.5× Hoagland’s solution with micronutrients (50 ml/per plant). Four-week-old plants were treated by irrigating with 0.5× Hoagland’s solution containing 500 mM NaCl, in which the salt was provided as a source of stress to induce CAM (Cushman *et al.* 2008). Control plants were continuously watered with 0.5× Hoagland’s solution. The third pairs of leaves from the control and salt-treated plants were collected on day 5, day 6 and day 7 following the initiation of the salt treatment. Unless stated otherwise, three biological replicates from three different sets of plants were used.

2.2. Leaf nocturnal acidification

Leaf titratable acidity was measured as previously described (Cushman *et al.* 2008) with minor modifications. Leaves were harvested from control and salt-treated plants at the end of the night period (8 am). Fresh weight of the collected leaves was measured, followed by snap freezing in liquid N₂ and storage at −80°C until analysis. Frozen leaves were ground to a fine powder using a mortar and pestle, followed by further homogenization in 80% methanol at a ratio of 7.5 ml/gram of fresh weight at room temperature. The homogenate was allowed to warm to room temperature.

Aliquots of the methanol extracts were titrated against 5 mM NaOH to a neutral endpoint (pH = 7.0), using a pH meter. Leaf titratable acidities were expressed as $\mu\text{mol H}^+ \text{g}^{-1}$ fresh weight.

2.3. Net CO₂ Exchange

Gas exchange measurements were performed at day 5, day 6 and day 7 of salt treatment. Net CO₂ uptake was measured using a LI-6800 system (LI-COR Inc., Lincoln, NE, USA), and parameters were calculated using the manufacturer software. Conditions for measuring net CO₂ uptake, stomatal conductance, and transpiration rates were: photon flux density 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$, chamber temperature 18°C/night and 26°C/day, flow rate 500 $\mu\text{mol s}^{-1}$, relative humidity 50%, and 400 ppm CO₂ reference. The chamber covered approximately 7 cm² area of the ice plant leaf to measure net gas exchange. For each time point at least five plants were used. Rates of net CO₂ uptake, stomatal conductance, and transpiration were retrieved from the LI-6800 data.

2.4. Stomatal aperture assay

The epidermis of control and salt-treated plant leaves were removed using 10 cm wide tape at 4 am and 4 pm, during the CAM transition process. After peeling, minor contamination from adhering mesophyll cells was removed by scratching the epidermis using a scalpel blade, and clean peels were directly used for imaging stomatal apertures. For stomatal aperture measurement, 60 - 80 stomata were randomly selected, and the sample identity was blinded during the measurement. Stomatal length and width were measured as described (Savvides *et al.* 2012), and stomatal aperture was estimated by the length/width ratio.

2.5. CAM-related gene expression profiling via quantitative real-time PCR

To identify the C₃ to CAM transition time points, we assessed the expression of 28 CAM-related genes using quantitative real-time PCR (Table S1). Total RNA was isolated from leaf tissue using a CTAB method (Doyle & Doyle, 1987), and the quality and quantity were analyzed using a NanoDrop 1000 spectrophotometer (Thermo Fischer Scientific, San Jose, CA, USA). Five micrograms of total RNA were treated with RNase-free DNase I (New England Biolabs, Ipswich, MA, USA) to remove genomic DNA. cDNA was synthesized using a ProtoScript[®] First Strand cDNA Synthesis Kit (New England Biolabs, Ipswich, MA, USA) according to the manufacturer's manual. The cDNA was diluted to a final concentration of 50 ng/μl, and 5 μl was used in a 20 μl PCR reaction with 1 μl of each 10 μM primer and 10 μl of VeriQuest SyBr with Fluorescein kit (Affymetrix, Santa Clara, CA, USA). For each reaction, three technical replicates were included for each biological replicate. All amplifications were carried out on a CFX96 Touch[™] Real-Time PCR Detection System (Bio-Rad, Hercules, CA, USA). The product size was confirmed by 2% agarose gel electrophoresis and the specificity of the amplicon was confirmed by melting curve analysis. Data were analyzed and quantified with the Bio-Rad CFX Manager software. The transporter gene *PLASMA MEMBRANE INTRINSIC PROTEIN 1;2* (*PIP1;2*) was used as an internal PCR standard, as its expression is constant during the diurnal cycle in leaves from control and salt-treated *M. crystallinum* (Vera-Estrella *et al.* 2012). The relative expression levels were calculated by the $2^{-\Delta\Delta C_T}$ method (Livak & Schmittgen 2001).

2.6. PEPC enzyme activity

For PEPC enzyme activity assays, 50 mg frozen leaf tissue was ground in liquid nitrogen to a fine powder, and extracted with 0.5 ml enzyme extraction buffer (200 mM HEPES-NaOH (pH 7.0), 5 mM dithiothreitol (DTT), 10 mM MgCl₂ and 2% (w/v) PVP-40). The homogenized

mixture was then centrifuged at 14,000 rpm, 4°C for 20 minutes and the supernatant was retained. To measure the activity of PEPC, 50 µl supernatant, 150 µl extraction buffer and 2.8 ml assay solution (100 mM HEPES-NaOH (pH 8.0), 5 mM DTT, 5 mM MgCl₂, 10 mM NaHCO₃, 2.5 mM phosphoenolpyruvate (PEP), 0.2 mM NADH and 0.0034 units/µl malate dehydrogenase (MDH)) were used. Absorbance at 340 nm was recorded for at least 5 min. PEPC enzyme activity was calculated as described previously (Chu *et al.* 1990).

2.7. Guard cell enrichment and construction of RNA-Seq libraries

For guard cell enrichment, parallel samples from control and salt-treated plants were collected at 12 am and 12 pm from day 5 to day 7 using a tape-peel method (Lawrence *et al.* 2018). Briefly, the abaxial epidermis was directly peeled off using Scotch Transparent tape, and adherent mesophyll cells were removed from the epidermis with a scalpel. After washing in basic solution (0.55 M sorbitol, 0.5 mM CaCl₂, 0.5 mM MgCl₂, 0.5 mM ascorbic acid, 10 µM KH₂PO₄, 5 mM 4-morpholineethanesulfonic acid (MES), pH 5.5 adjusted with 1 M KOH), the tapes with adherent epidermis were incubated with a cell wall digesting enzyme solution (0.7% cellulase R-10 (Yakult Honsha Co., Ltd, Tokyo, Japan), 0.025% macerozyme R-10 (Yakult Honsha Co., Ltd, Tokyo, Japan), 0.1% (w/v) polyvinylpyrrolidone-40 (Calbiochem, Billerica, Massachusetts, USA), and 0.25% (w/v) bovine serum albumin (Research Products International Corp., Mt Prospect, Illinois, USA) in 55% basic solution on a reciprocal shaker for 30 min at room temperature. Digested peels were washed three times with basic solution and quickly blotted dry on a filter paper, then immediately frozen in liquid nitrogen and stored at -80°C until RNA extraction. For each time point, three independent biological replicates were collected using three different sets of 120 plants. Total RNA was isolated using a CTAB method (Doyle & Doyle,

1987). After eliminating any genomic DNA contamination with RNase-free DNase I (New England Biolabs, Ipswich, MA, USA), mRNA was purified using a Dynabeads[®] mRNA Purification Kit (Thermo Fischer Scientific, San Jose, CA, USA). RNA-seq libraries were constructed using the QuantSeq 3' mRNA-Seq Library Prep Kit (Lexogen, Vienna, Austria) according to the manufacturer's instructions. RNA sequencing was performed on an Illumina NextSeq 500 with 75 bp single end reads at the NextGen DNA Sequencing Core of the University of Florida, Gainesville, FL.

2.8. RNA-Seq data analysis

For gene expression quantification, a reference transcriptome was constructed using sequences from ice plant bladder cell (www.lsugenomics.org/data-from-our-group) (Oh *et al.* 2015) and root (dandelion.liveholonics.com/pothos/Mcr/data/reference/Mcr.transcript.fasta) (Tsukagoshi *et al.* 2015), sequences downloaded from NCBI, and Expressed Sequence Tags (ESTs) previously published (Cushman *et al.* 2008). Sequences from each reference were concatenated into a single fasta file to serve as a reference transcriptome. Because the *de novo* assemblies and transcripts were accessed from multiple sources, it was not known which transcripts were identical, or alternatively spliced isoforms, due to lack of a reference ice plant genome. To address the redundancy of the sequences in the reference and to collapse putative isoforms into a single transcript, Cap3 (Huang & Madan 1999) was used with default parameters (-a 20 -b 20 -c 12 -d 200 -e 30 -f 20 -g 6 -h 20 -i 40 -j 80 -k 1 -m 2 -n -5 -o 40 -p 90 -r 1 -s 900 -t 300 -u 3 -v 2 -w NA -x cap -y 100 -z 3). Assembled contigs representing putative collapsed transcript isoforms and singlet transcripts that were not homologous to other transcripts were kept as transcript references for alignments. These transcripts represent putative gene sequences. To

exclude multiple isoforms of the same gene, the single longest contig was used for each gene. A single transcript representing each gene was used because downstream analysis of differential transcript expression was conducted at the gene level rather than the isoform level, as 3' end sequencing or short read sequencing in general is not suitable for isoform level quantification. Low quality bases/reads were removed from the sequence data with Trimmomatic (Bolger *et al.* 2014) with parameters HEADCROP:0 LEADING:3 TRAILING:3 SLIDING-WINDOW:4:20 MINLEN:18. Reads were mapped to our reference transcriptome with RSEM (Li & Dewey 2011) version 1.2.31. Gene expression was measured as the number of reads that aligned to a given transcript (counts). Gene counts for each sample were consolidated into a matrix and imported into EdgeR (McCarthy *et al.* 2012) to conduct differential expression analysis. Differentially expressed (DE) transcripts were identified by comparing control versus salt treatment groups or day versus night groups from day 5, day 6, day 7, night 5, night 6 and night 7 sampling groups using a 2-fold change and an adjusted *p*-value threshold of 0.05. The RNA-Seq data have been deposited at the National Center for Biological Information (NCBI) Sequence Read Archive (SRA) under the accession number SRX3878746.

3. Results

3.1. Determination of the C₃ to CAM transition timing

3.1.1. Titratable acidity

Because of the nocturnal CO₂ assimilation in CAM plants, high levels of malate accumulate in vacuoles of mesophyll cells before dawn, which increases cellular acidity. Changes of leaf titratable acidity measured at the start (8 pm) and end (8 am) of the dark period were used as a measure of CAM induction in *M. crystallinum* from day 4 to day 14 after 500 mM salt treatment.

Salt-treated plants, measured from days 21 and 28, were used as positive controls because they use the CAM mode of photosynthesis only (Cushman *et al.* 2008; Vera-Estrella *et al.* 2012). A modest increase of overnight acidity was observed on day 6 in leaves of the salt-treated plants. The difference between treated and control plants became statistically significant on day 7, after which titratable acidity steadily increased (Figure 1A). It was noteworthy that there were no visible morphological differences between the control and salt-treated plants at day 7 (Figure 1B). Based on the nocturnal acid accumulation, we inferred that the C₃ to CAM transition occurred between days 5 to 7 of salt treatment; thus, we collected samples from day 5 (no transition), day 6 and day 7 plants for follow-up analyses.

3.1.2. Net CO₂ exchange and stomatal movement

At day 5, control and salt-treated plants showed no difference in gas exchange parameters. However, day 6 salt-treated plants showed increased net CO₂ uptake in the night and much lower net CO₂ uptake during the day, compared to control plants (Figure 2A). Salt-treated plants from day 7 showed decreased net CO₂ uptake during the day and positive net CO₂ uptake in the night, significantly different from control plants (Figure 2A).

To further confirm the C₃ to CAM transition time-points, stomatal aperture was measured in the control and salt-treated plants during the night (at 4 am) and the day (at 4 pm). These two time-points were chosen based on the result of net CO₂ exchange (Figure 2A). As shown in Figure 2B, the control plants showed C₃-type stomatal movement, i.e., they closed stomata during the night and opened stomata during the day. In contrast, the salt-treated plants showed an inversion in stomatal movement, starting on day 5. Stomatal aperture of the salt-treated plants increased

consistently during the nights of day 6 and day 7 (Figure 2B). The gas exchange and stomatal movement results also support that the C_3 to CAM transition of *M. crystallinum* occurs between day 5 to day 7 of the salt treatment.

3.1.3. CAM-related gene expression profiles

To further validate the C_3 to CAM transition, CAM-related genes were selected for analysis based on previous studies (Brilhaus *et al.* 2016; Cushman *et al.* 2008; Vera-Estrella *et al.* 2012). In total, 28 genes were selected, including PEP carboxylase (*PEPC*, the enzyme responsible for dark CO_2 fixation) (Osmond 1978), PEPC kinase (*PPCK1*) (Dittrich 1976), 4-alpha-glucanotransferase or disproportionating enzyme (*GTF1*, involved in starch degradation) (Cushman *et al.* 2008), circadian clock associated 1 (*CCA1*) (Abraham *et al.* 2016), phosphoglycerate kinase (*PGK3*) (Brilhaus *et al.* 2016; Cushman *et al.* 2008), and ABA insensitive 1 (*ABII*, involved in guard cell movement) (Abraham *et al.* 2016). From day 5 to day 7, control and salt-treated leaves were collected at 2 am and 4 am because two core CAM genes, *PEPC* and *PPCK1*, were reported to exhibit the highest expression levels at these time-points (Dodd *et al.* 2002). As shown in Figure 3A, *PEPC* exhibited a > 5-fold increase in transcript abundance at the two time points of day 6 in salt-treated plants compared to control plants and kept increasing up to > 200-fold at day 7. *PPCK1* showed a similar expression profile as *PEPC*. *ABII* and *GTF1* also showed higher transcript abundances at the two time points of day 6 in the salt-treated plants, and the levels kept increasing at day 7. *ABII* is an important ABA signaling component. The increased *ABII* transcript abundance may contribute to the development of CAM (Figure 2B). *CCA1*, a key regulator of circadian rhythm in plants, showed a significant increase at 4 am of day 7 in salt-treated samples relative to control plants. *PGK3* showed

significant decreases at the two time points of day 7 in the salt-treated samples. All six genes exhibited similar expression patterns at day 7 in the salt-treated plants as those reported in previous studies (Brilhaus *et al.* 2016; Cushman *et al.* 2008). Most of other 22 genes analyzed showed similar expression patterns as the above six genes at day 7 in the salt-treated plants (Table S2).

3.1.4. PEPC enzyme activity

PEPC activity in the dark is much higher in CAM plants than in C_3 plants (Chu *et al.* 1990). PEPC activity changes were measured in the samples collected at 2 am and 4 am of day 5 to day 7. As shown in Figure 3B, PEPC enzyme activity was significantly enhanced at the two day 7 time points in the salt-treated plants, as compared to the control. Taken together, the physiological data on titratable acidity, CO_2 exchange, PEPC activity and stomatal movement, as well as the transcription of key CAM-related genes, demonstrate that the transition from C_3 to CAM photosynthesis occurs between days 5 and 7.

3.2 Transcriptomics of *M. crystallinum* guard cells during the C_3 to CAM transition

To investigate the regulatory mechanism(s) underlying the transition to inverse stomatal opening in *M. crystallinum*, we profiled the single-cell type transcriptome of guard cells using RNA-Seq during days 5, 6 and 7 at 12 am (night) and 12 pm (day). A total of 197,790,866 raw reads were acquired. Removal of low quality reads with Trimmomatic (Bolger *et al.* 2014) resulted in 188,147,736 high quality reads. By using previously published microarray and transcriptome data (Cushman *et al.* 2008; Oh *et al.* 2015; Tsukagoshi *et al.* 2015), we created a reference transcriptome, to which our short reads were mapped. In total, 43,165 different transcripts

(including isoforms) were expressed in *M. crystallinum* guard cells (Counts Per Million (CPM) $\geq$ 10 in at least two biological replicates) (Table 1). Among these transcripts, 10,628 transcripts were not matched to the reference plant *Arabidopsis thaliana* transcripts (based on BLAST e-value ≤ 0.001 , similarity ≥ 70).

Significantly differentially expressed (DE) transcripts were defined as those with at least a 2-fold difference between control and salt-treated samples and an adjusted *p*-value < 0.05 . A total of 495 transcripts showed significant changes at one or more time points by comparing the salt-treated and control plants; 369 of these DE transcripts have homologs in *Arabidopsis*. Among the 369 DE transcripts, *PEPCI* was found to increase at 12 am (night) of day 7 in the salt-treated samples (Table 2). This increased expression of *PEPCI* in guard cells correlates with our real-time PCR result in leaves (Figure 3).

Gene Ontology (GO) enrichment analysis of the 369 DE transcripts was performed using R package “clusterProfiler” (DOI: 10.18129/B9.bioc.clusterProfiler). The DE transcripts were enriched for the category “response to stimulus”, especially “response to abiotic stimulus” and “response to stress”. In terms of molecular function, the encoded proteins were enriched in catalytic activity (Figure 4). It is interesting to note that there was little overlap between the DE transcripts in the day samples (Figure 5A) or night samples (Figure 5B) at days 5, 6 and 7, suggesting different changes took place in the course of the early stages of transition from C_3 to CAM. When the DE genes from day and night samples were compared, only about 10% of the DE genes were shared (Figure 5B). A small number of genes showed opposite change patterns under the day versus the night conditions (Table S3). For example, a papain-like cysteine

protease showed high expression in control samples collected at night in day 5. At days 6 and 7, its expression levels started to decrease in control night and significantly increase in the day of salt-treated samples (Table S3). How this cysteine protease play a role in the CAM transition is not known. Nevertheless, this result suggests that the GC transcriptome is diurnally regulated during development of CAM (as are stomatal apertures).

3.3 Abundance changes of previously identified guard cell transcripts during the C₃ to CAM transition

To identify the CAM-related DE transcripts in our study, we compared these 369 DE transcripts with Cushman et al. (2008)'s microarray data, which identified 56 CAM-related genes and non-CAM isogenes in *M. crystallinum* that displayed maximal inverse expression between CAM-performing and C₃ leaves. Seven out of the 56 genes were retrieved from our DE transcripts, including *PEPC1* (comp24500_c0_seq1), *PPCK1* (Contig5856), *BETA-AMYLASE 5* (Contig3509, *AMYB5*), *CARBONIC ANHYDRASE 2* (Contig20069, *CAH2*), *FRUCTOSE-BISPHOSPHATE ALDOLASE 2* (Contig12312, *FBA2*), *PHOSPHOGLYCERATE MUTASE* (Mcr017048.026, *PGM*) as well as another *PHOSPHOENOLPYRUVATE CARBOXYLASE* family protein gene (Contig20312) (Table 2). Among these seven genes, only *PPCK1* and *AMYB5* were not found in previous guard cell studies (Table 2). *PPCK1*, *PEPC1*, as well as another *PEPC* family gene, were up-regulated in our day 7 night samples, similar to the real-time PCR result on leaf tissue (Figure 3). The up-regulation of *PGM*, *CAH2*, *FBA2* and *AMYB5* in guard cells (Table 2) was also found in *M. crystallinum* leaves in a previous study (Cushman et al., 2008). However, in day 7 samples, *PGM* encoding a phosphoglycerate mutase up-regulated in leaves of the CAM plants (Cushman et al., 2008), was down-regulated in guard cells (Table 2).

Overall, this analysis corroborates that guard cells themselves are transitioning to CAM because they exhibit many of the transcriptomic changes expected of cells undergoing that process. Additionally, guard cells also have their own transcriptional program during the transition.

3.4 Transcription factor changes during the C₃ to CAM transition

It has been reported that in response to internal or external environment changes, transcription factors (TFs) exhibit more rapid expression changes than the bulk of the regulated genes (Jiao *et al.* 2003). Thus, the expression profiles of TF genes may in some way reflect the subsequent transcription activities regulated by them. In total, 18 TFs were identified among the 369 DE transcripts, and 14 of them were observed in other guard cell studies (Table 2). Based on the annotation from the *Arabidopsis* homologs, four TFs (Mcr002150.001, Mcr008625.003, Contig18172 and Contig9771) were identified in previous studies as responsive to salt or water deprivation stress in leaves or shoots (Table S4) (Ding *et al.* 2013; Seo & Park 2011; Yanhui *et al.* 2006). Since CAM mode in *M. crystallinum* is induced by abiotic stresses, such as salt and drought stress, these TFs may be general stress response genes and the others may have important roles in the C₃ to CAM transition.

To evaluate how gene expression changes during the transition process, we grouped genes with similar pattern of expression using k-means clustering. Eleven clusters were retrieved, of which cluster 5 contained the greatest number of transcripts (Figure 6). In cluster 5, several genes showed differential expression profiles in the day samples, but the majority of transcripts showed increases in expression during the day 5 night. Only one gene, *ARGININE/SERINE-RICH SPLICING FACTOR 35* (Mcr012474.005) showed decreased expression during the day 7 night.

In cluster 10, 20 of 24 transcripts have *Arabidopsis* homologs, and 17 of them were present in other guard cell studies (Figure 6). In this cluster, there are no differences in any samples during the day time, while all of them showed increases during the day 7 night (Figure 6). This cluster contains two key players in the CAM mode, *PEPC1* and *PPCK1*. It also includes three TFs: *AGAMOUS-LIKE 8* (AT5G60910), *HOMEODOMAIN 7* (AT2G46680) as well as *BEL1-LIKE* *HOMEODOMAIN 7* (AT2G16400) (Table 2).

Discussion

In this study, the C₃ to CAM transition was induced by 500 mM salt treatment, and the transition time points were supported by several physiological parameters and molecular marker expression profiles (Figures 1-3). Furthermore, the cell-type specific transcriptome of guard cells was characterized during the C₃ to CAM transition (Figures 4-6). Results are discussed here in light of the molecular mechanisms occurring in guard cells during the transition.

Although there have been many steady-state studies of C₃ and/or CAM (e.g., Cushman *et al.* 2008; Davies & Griffiths 2012; Tsukagoshi *et al.* 2015), determination of the critical transition time-point was not reported before. The results of this study suggest that the C₃ to CAM induction in *M. crystallinum* takes place within a short period of time (from day 5 to day 7), but that the full transition of photosynthesis to CAM is slow (another 2-4 weeks) (Figures 1, 2). Therefore, to determine the mechanisms underlying the C₃ to CAM transition, we cannot rely on the analysis of one-time point. For the transcriptomic analysis of guard cells, we targeted six-time points from day 5 to day 7 of salt treatment.

414 We generated a reference transcriptome based on previously sequenced ESTs (Cushman *et al.*
415 2008), assembled transcripts (Oh *et al.* 2015; Tsukagoshi *et al.* 2015), and individual genes from
416 *M. crystallinum* reported at NCBI. The genome size of *M. crystallinum* is 390 Mbp (Ha *et al.*
417 2014), with an estimate of 30,000 to 35,000 genes (De Rocher *et al.* 1990; Meyer *et al.* 1990).
418 We constructed more than 180 thousand contigs, and identified 40,757 different transcripts
419 (including isoforms) that were expressed in our samples. Among them, 10,628 unique transcripts
420 in guard cells (no isoforms) were found to have homologs in Arabidopsis. Based on the previous
421 studies of Arabidopsis guard cell transcriptomes (Bates *et al.* 2012; Bauer *et al.* 2013; Leonhardt
422 *et al.* 2004; Pandey *et al.* 2010; Wang *et al.* 2011) and proteome (Zhao *et al.* 2008), about 30%
423 of guard cell genes of *A. thaliana* had homologs expressed in *M. crystallinum* guard cells under
424 our salt stress condition. From the RNA-Seq data, we identified 495 DE transcripts that showed
425 significant changes at one or more time points of transition, of which 369 have homologs in
426 Arabidopsis. Among these 369 DE transcripts, there were 199 up-regulated and 178 down-
427 regulated transcripts in response to the salt treatment (Table 1). Both up- and down-regulated
428 transcripts were enriched in “response to stress” and “cellular carbohydrate metabolic process”,
429 suggesting a subset of genes involved in the two biological processes was employed in the
430 course of transition. Key CAM molecular marker genes such as *PPCK1*, *PEPC1* and *GTF1*, as
431 well as another *PEPC* family gene were up-regulated in the night 7 samples (Table 2, Table S4).
432 This new observation suggests that guard cells switch from C₃ to CAM; the timing of the
433 differential expression in guard cells is also consistent with our physiological determination of
434 the C₃ to CAM transition time points (Figures 1-3). Another four CAM-related genes were also
435 identified: *PGM*, *CAH2*, *FBA2* and *AMYB5*. Except for *PGM*, the other three genes showed
436 similar expression profiles to those reported in leaves of CAM *M. crystallinum* (Table 2)

(Cushman *et al.* 2008). *PGM*, which encodes a phosphoglycerate mutase was reported to be up-regulated in the plants performing CAM (Cushman *et al.* 2008), while in our study it was down-regulated in the day 7 samples. This may be due to difference in materials used since we targeted guard cells only, while Cushman and co-workers (2008) utilized the whole leaf in their study. Interestingly, *PGM* was shown to play an important role in guard cell functions. A null *pgm* mutant displayed defects in blue light-, abscisic acid-, and low CO₂-regulated stomatal movements in *Arabidopsis* (Zhao & Assmann 2011). Another difference between our study and that of Cushman *et al.* (2008) is the time of sample collection: our samples were harvested during the C₃ to CAM transition, while the previous study used leaves from plants performing complete CAM photosynthesis (Cushman *et al.* 2008). Therefore, this disparity indicates that the guard cells regulate transcription differently from those of leaves, either in response to salt stress or during transition and after transition to the CAM mode.

Notably, 18 TFs were identified among the 369 DE transcripts, and 14 of them were also detected in previous guard cell studies (Table 2). Among these 18 TFs, six of them were down-regulated, while 12 were up-regulated during the transition compared to the control guard cells from plants undergoing C₃ photosynthesis. Four TFs (Mcr002150.001, Mcr008625.003, Contig18172, and Contig9771) may be related to salt or water deprivation stress response. These TFs are key players in the regulatory networks underlying plant responses to abiotic stresses and development processes (Coelho *et al.* 2018; Gollack *et al.* 2014; Hoang *et al.* 2017). Among them, *Mcr010456.002* (*At2g42280.1*), which encodes a basic helix-loop-helix (bHLH) TF, was implicated in stomatal movement through activation of genes encoding inwardly rectifying K⁺ channels (Takahashi *et al.* 2013). In addition, from the *k*-means clustering result (Figure 4), 26

DE transcripts (including the two CAM genes, *PEPC1* and *PPCK1*, present in cluster 10) were up-regulated in samples collected during the night of day 7 (Table S4), suggesting these genes were induced in the initial CAM guard cells. The significantly changed genes (including TFs) that are co-regulated with known CAM genes (e.g., *PPCK1* and *PEPC1*) can be expected to play important roles in the C_3 to CAM transition process. In ice plants performing C_3 photosynthesis, light increases leaf conductance and also promotes stomatal opening in isolated epidermal peels, while in plants performing CAM, stomatal opening in epidermal peels becomes unresponsive to light (Figure S1). This result and the RNA-Seq data from the isolated stomatal guard cells corroborate previous studies in facultative CAM species (Lee & Assmann 1992; Tallman *et al.* 1997), and demonstrate the presence in the guard cells themselves of molecular switches for the CAM inverse stomatal behavior, separate from mesophyll cells.

In summary, we induced CAM transition from C_3 in *M. crystallinum* by 500 mM NaCl treatment and successfully determined the timing of the C_3 to CAM transition. Furthermore, we characterized the guard cell transcriptomic changes during the critical transition process. The presence and the diel changes of CAM marker genes in stomatal guard cells indicate the guard cells themselves can transit from C_3 to CAM. Many candidate genes (including TFs) were identified. Functional studies of these candidate genes in guard cells of either ice plant or the reference plant *Arabidopsis* are important future directions. In addition, these results indicate that efforts focused solely on engineering the mesophyll to introduce CAM into other species for improving WUE and stress tolerance may fail. Engineering both the mesophyll cells and guard cells is likely to be necessary.

483

484 ACKNOWLEDGMENTS

485 We thank Christopher Krieg, Emily Sessa, Tianyi Ma, and Zepeng Yin for their help in obtaining
486 data with the LI-6800, and Christopher Dervinis for his technical assistance in the RNA-Seq
487 experiments. We also thank Dr. John Cushman from University of Nevada for providing the
488 *Mesembryanthemum crystallinum* seeds and protocol for growing the plants. This work was
489 supported by the University of Florida internal funds, with additional partial support from NSF
490 Plant Genome Research Program Award #1444543, and NSF REU grant # 1560049 for the
491 support to an undergraduate student Theresa M Kelley for carrying out part of this research.

492

493 AUTHOR CONTRIBUTIONS

494 MK, SA and SC designed the research project. WK, MY and TK performed the experiments.
495 MY and JN analyzed the transcriptomics data. JL helped with data interpretation and manuscript
496 editing. MK and MY made the figures, and wrote the manuscript draft. All the authors read and
497 approved the manuscript. SC finalized the manuscript for submission.

498

499

500 REFERENCES

501 Abraham P.E., Yin H., Borland A.M., Weighill D., Lim S.D., De Paoli H.C., Engle N., Jones
502 P.C., Agh R., Weston D.J., Wullschleger S.D., Tschaplinski T., Jacobson D., Cushman
503 J.C., Hettich R.L., Tuskan G.A. & Yang X. (2016) Transcript, protein and metabolite
504 temporal dynamics in the CAM plant Agave. *Nature Plants* 2, 16178.

505 Adams P., Nelson D.E., Yamada S., Chmara W., Jensen R.G., Bohnert H.J. & Griffiths H. (1998)
506 Growth and development of *Mesembryanthemum crystallinum* (Aizoaceae). *New*
507 *Phytologist* 138, 171-190.

508 Barkla B.J. & Vera-Estrella R. (2015) Single cell-type comparative metabolomics of epidermal
509 bladder cells from the halophyte *Mesembryanthemum crystallinum*. *Frontiers in Plant*
510 *Science* 6.

511 Barkla B.J., Vera-Estrella R. & Raymond C. (2016) Single-cell-type quantitative proteomic and
512 ionomic analysis of epidermal bladder cells from the halophyte model plant
513 *Mesembryanthemum crystallinum* to identify salt-responsive proteins. *BMC Plant*
514 *Biology* 16, 110.

515 Bates G.W., Rosenthal D.M., Sun J.D., Chattopadhyay M., Peffer E., Yang J., Ort D.R. & Jones
516 A.M. (2012) A comparative study of the *Arabidopsis thaliana* guard-cell transcriptome
517 and its modulation by sucrose. *PLoS One* 7.

518 Bauer H., Ache P., Lautner S., Fromm J., Hartung W., Al-Rasheid K.A.S., Sonnewald S.,
519 Sonnewald U., Kneitz S., Lachmann N., Mendel R.R., Bittner F., Hetherington A.M. &
520 Hedrich R. (2013) The stomatal response to reduced relative humidity requires guard
521 cell-autonomous ABA synthesis. *Current Biology* 23, 53-57.

522 Bolger A.M., Lohse M. & Usadel B. (2014) Trimmomatic: a flexible trimmer for Illumina
523 sequence data. *Bioinformatics* 30, 2114-2120.

524 Borland A.M., Hartwell J., Weston D.J., Schlauch K.A., Tschaplinski T.J., Tuskan G.A., Yang
525 X.H. & Cushman J.C. (2014) Engineering crassulacean acid metabolism to improve
526 water-use efficiency. *Trends in Plant Science* 19, 327-338.

- 527 Brilhaus D., Brautigam A., Mettler-Altmann T., Winter K. & Weber A.P. (2016) Reversible
528 burst of transcriptional changes during induction of crassulacean acid metabolism in
529 *Talinum triangulare*. *Plant Physiology* 170, 102-122.
- 530 Chiang C.P., Yim W.C., Sun Y.H., Ohnishi M., Mimura T., Cushman J.C. & Yen H.E. (2016)
531 Identification of ice plant (*Mesembryanthemum crystallinum* L.) microRNAs using RNA-
532 Seq and their putative roles in high salinity responses in seedlings. *Frontiers in Plant*
533 *Science* 7, 1143.
- 534 Chu C., Dai Z., Ku M.S. & Edwards G.E. (1990) Induction of crassulacean acid metabolism in
535 the facultative halophyte *Mesembryanthemum crystallinum* by abscisic acid. *Plant*
536 *Physiology* 93, 1253-1260.
- 537 Coelho C.P., Huang P., Lee D.Y. & Brutnell T.P. (2018) Making roots, shoots, and seeds: IDD
538 gene family diversification in plants. *Trends in Plant Science* 23, 66-78.
- 539 Cushman J.C., Tillett R.L., Wood J.A., Branco J.M. & Schlauch K.A. (2008) Large-scale mRNA
540 expression profiling in the common ice plant, *Mesembryanthemum crystallinum*,
541 performing C₃ photosynthesis and Crassulacean acid metabolism (CAM). *Journal of*
542 *Experimental Botany* 59, 1875-1894.
- 543 Davies B.N. & Griffiths H. (2012) Competing carboxylases: circadian and metabolic regulation
544 of Rubisco in C₃ and CAM *Mesembryanthemum crystallinum* L. *Plant, Cell and*
545 *Environment* 35, 1211-1220.
- 546 De Rocher E.J., Harkins K.R., Galbraith D.W. & Bohnert H.J. (1990) Developmentally regulated
547 systemic endopolyploid in succulents with small genomes. *Science* 250, 99-101.

548 Ding Y., Liu N., Virlouvet L., Riethoven J.J., Fromm M. & Avramova Z. (2013) Four distinct
549 types of dehydration stress memory genes in *Arabidopsis thaliana*. *BMC Plant Biol* 13,
550 229.

551 Dittrich P. (1976) Nicotinamide adenine dinucleotide-specific "malic" enzyme in *Kalanchoe*
552 *daigremontiana* and other plants exhibiting crassulacean acid metabolism. *Plant*
553 *Physiology* 57, 310-314.

554 Dodd A.N., Borland A.M., Haslam R.P., Griffiths H. & Maxwell K. (2002) Crassulacean acid
555 metabolism: plastic, fantastic. *J Exp Bot* 53, 569-580.

556 Doyle J.J. & Doyle J.L. (1987) A rapid DNA isolation from small amount of fresh leaf tissue.
557 *Phytochemical Bulletin* 19, 11-15.

558 Golldack D., Li C., Mohan H. & Probst N. (2014) Tolerance to drought and salt stress in plants:
559 unraveling the signaling networks. *Frontiers in Plant Science* 5.

560 Ha J., Bernard W., Yim W.C., Albion R.L., Schlauch K.A., Yin H. & Cushman J.C. (2014) Draft
561 genome sequence of the common ice plant (*Mesembryanthemum crystallinum* L.) a
562 facultative crassulacean acid metabolism (CAM) and halophytic plant model. In: *Plant*
563 *and Animal Genome XXII Conference*, San Diego, CA.

564 Hoagland D. & Arnon D. (1950) The water culture method of growing plants without soil. In:
565 *California Agricultural Experiment Station Circular* pp. 1-32. College of Agriculture,
566 University of California, Berkeley, San Francisco, CA.

567 Hoang X.L.T., Nhi D.N.H., Thu N.B.A., Thao N.P. & Tran L.S.P. (2017) Transcription factors
568 and their roles in signal transduction in plants under abiotic stresses. *Current Genomics*
569 18, 483-497.

570 Huang X. & Madan A. (1999) CAP3: A DNA sequence assembly program. *Genome Research* 9,
571 868-877.

572 Hurst A.C., Grams T.E.E. & Ratajczak R. (2004) Effects of salinity, high irradiance, ozone, and
573 ethylene on mode of photosynthesis, oxidative stress and oxidative damage in the
574 C₃/CAM intermediate plant *Mesembryanthemum crystallinum* L. *Plant, Cell and*
575 *Environment* 27, 187-197.

576 Jiao Y., Yang H., Ma L., Sun N., Yu H., Liu T., Gao Y., Gu H., Chen Z., Wada M., Gerstein M.,
577 Zhao H., Qu L.J. & Deng X.W. (2003) A genome-wide analysis of blue-light regulation
578 of Arabidopsis transcription factor gene expression during seedling development. *Plant*
579 *Physiology* 133, 1480-1493.

580 Kaushal S.S., Gold A.J. & Mayer P.M. (2017) Land use, climate, and water resources - global
581 stages of interaction. *Water* 9, 1-10.

582 Lee D.M. & Assmann S.M. (1992) tomato responses to light in the facultative crassulacean acid
583 metabolism species, *Portulacaria afra*. *Physiologia Plantarum* 85, 35-42.

584 Leonhardt N., Kwak J.M., Robert N., Waner D., Leonhardt G. & Schroeder J.I. (2004)
585 Microarray expression analyses of Arabidopsis guard cells and isolation of a recessive
586 abscisic acid hypersensitive protein phosphatase 2C mutant. *Plant Cell* 16, 596-615.

587 Li B. & Dewey C.N. (2011) RSEM: accurate transcript quantification from RNA-Seq data with
588 or without a reference genome. *BMC Bioinformatics* 12, 323.

589 Livak K.J. & Schmittgen T.D. (2001) Analysis of relative gene expression data using real-time
590 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402-408.

591 McCarthy D.J., Chen Y. & Smyth G.K. (2012) Differential expression analysis of multifactor
592 RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res* 40, 4288-
593 4297.

594 Meyer G., Schmitt J.M. & Bohnert H.J. (1990) Direct screening of a small genome: estimation of
595 the magnitude of plant gene expression changes during adaptation to high salt. *Molecular*
596 *Genomics and Genetics* 224, 347-356.

597 Nobel P. (1996) High productivity of certain agronomic CAM Species. In: *Crassulacean Acid*
598 *Metabolism. Biochemistry, ecophysiology and evolution* (eds K. Winter & J.A.C. Smith),
599 pp. 255-265. Springer-Verlag, Berlin.

600 Oh D.H., Barkla B.J., Vera-Estrella R., Pantoja O., Lee S.Y., Bohnert H.J. & Dassanayake M.
601 (2015) Cell type-specific responses to salinity - the epidermal bladder cell transcriptome
602 of *Mesembryanthemum crystallinum*. *New Phytologist* 207, 627-644.

603 Osmond C.B. (1978) Crassulacean acid metabolism: a curiosity in context. *Annual Review of*
604 *Plant Physiology* 29, 379-414.

605 Owen N.A. & Griffiths H. (2013) A system dynamics model integrating physiology and
606 biochemical regulation predicts extent of crassulacean acid metabolism (CAM) phases.
607 *New Phytologist* 200, 1116-1131.

608 Pandey S., Wang R.S., Wilson L., Li S., Zhao Z.X., Gookin T.E., Assmann S.M. & Albert R.
609 (2010) Boolean modeling of transcriptome data reveals novel modes of heterotrimeric G-
610 protein action. *Molecular Systems Biology* 6.

611 Savvides A., Fanourakis D. & van Ieperen W. (2012) Coordination of hydraulic and stomatal
612 conductances across light qualities in cucumber leaves. *Journal of Experimental Botany*
613 63, 1135-1143.

614 Seo P.J. & Park C.M. (2011) Signaling linkage between environmental stress resistance and leaf
615 senescence in Arabidopsis. *Plant Signal Behav* 6, 1564-1566.

616 Surowka E., Dziurka M., Kocurek M., Goraj S., Rapacz M. & Miszański Z. (2016) Effects of
617 exogenously applied hydrogen peroxide on antioxidant and osmoprotectant profiles and
618 the C₃-CAM shift in the halophyte *Mesembryanthemum crystallinum* L. *Journal of Plant*
619 *Physiology* 200, 102-110.

620 Takahashi Y., Ebisu Y., Kinoshita T., Doi M., Okuma E., Murata Y. & Shimazaki K. (2013)
621 bHLH transcription factors that facilitate K⁺ uptake during stomatal opening are
622 repressed by abscisic acid through phosphorylation. *Science Signaling* 6, ra48.

623 Tallman G., Zhu J.X., Mawson B.T., Amodeo G., Nouhi Z., Levy K. & Zeiger E. (1997)
624 Induction of CAM in *Mesembryanthemum crystallinum* abolishes the stomatal response
625 to blue light and light-dependent zeaxanthin formation in guard cell chloroplasts. *Plant*
626 *and Cell Physiology* 38, 236-242.

627 Taybi T. & Cushman J.C. (1999) Signaling events leading to crassulacean acid metabolism
628 induction in the common ice plant. *Plant Physiology* 121, 545-556.

629 Tsukagoshi H., Suzuki T., Nishikawa K., Agarie S., Ishiguro S. & Higashiyama T. (2015) RNA-
630 seq analysis of the response of the halophyte, *Mesembryanthemum crystallinum* (ice plant)
631 to high salinity. *PLoS One* 10, e0118339.

632 Vera-Estrella R., Barkla B.J., Amezcua-Romero J.C. & Pantoja O. (2012) Day/night regulation
633 of aquaporins during the CAM cycle in *Mesembryanthemum crystallinum*. *Plant, Cell*
634 *and Environment* 35, 485-501.

635 Wang R.S., Pandey S., Li S., Gookin T.E., Zhao Z.X., Albert R. & Assmann S.M. (2011)
636 Common and unique elements of the ABA-regulated transcriptome of *Arabidopsis* guard
637 cells. *BMC Genomics* 12.

638 Winter K. & Holtum J.A. (2005) The effects of salinity, crassulacean acid metabolism and plant
639 age on the carbon isotope composition of *Mesembryanthemum crystallinum* L., a
640 halophytic C₃-CAM species. *Planta* 222, 201-209.

641 Winter K. & Ziegler H. (1992) Induction of crassulacean acid metabolism in
642 *Mesembryanthemum crystallinum* increases reproductive success under conditions of
643 drought and salinity stress. *Oecologia* 92, 475-479.

644 Yanhui C., Xiaoyuan Y., Kun H., Meihua L., Jigang L., Zhaofeng G., Zhiqiang L., Yunfei Z.,
645 Xiaoxiao W., Xiaoming Q., Yunping S., Li Z., Xiaohui D., Jingchu L., Xing-Wang D.,
646 Zhangliang C., Hongya G. & Li-Jia Q. (2006) The MYB transcription factor superfamily
647 of *Arabidopsis*: expression analysis and phylogenetic comparison with the rice MYB
648 family. *Plant Mol Biol* 60, 107-124.

649 Zhao Z.X., Assmann S.M. (2011) The glycolytic enzyme, phosphoglycerate mutase, has critical
650 roles in stomatal movement, vegetative growth, and pollen production in *Arabidopsis*
651 *thaliana*. *J Exp Bot* 62, 5179-5189.

652 Zhao Z.X., Zhang W., Stanley B.A. & Assmann S.M. (2008) Functional proteomics of
653 *Arabidopsis thaliana* guard cells uncovers new stomatal signaling pathways. *Plant Cell*
654 20, 3210-3226.

655 Ziska L., Crimmins A., Auclair A., DeGrasse S., Garofalo J.F., Khan A.S., Loladze I., Pérez de
656 León A.A., Showler A., Thurston J. & Walls I. (2016) Food safety, nutrition, and
657 distribution. In: *The Impacts of Climate Change on Human Health in the United States: A*

658 *Scientific Assessment* (eds A. Crimmins, J. Balbus, J.L. Gamble, C.B. Beard, J.E. Bell, D.
659 Dodgen, R.J. Eisen, N. Fann, M.D. Hawkins, S.C. Herring, L. Jantarasami, D.M. Mills, S.
660 Saha, M.C. Sarofim, J. Trtanj, & L. Ziska), pp. 189-216. U.S. Global Change Research
661 Program, Washington, DC.

662

663

664

665

Table 1 The number of differentially expressed transcripts in comparison of control and salt-treated samples

Contrast	# Contigs ^a	# transcripts differentially expressed between control and salt-treated samples (% ^b)	# transcripts down-regulated in salt (% ^b)	# transcripts upregulated in salt (% ^b)
Day 5 control vs. Day 5 salt	31,560	64 (0.20)	35 (0.11)	29 (0.09)
Day 6 control vs. Day 6 salt	26,954	65 (0.24)	37 (0.14)	28 (0.10)
Day 7 control vs. Day 7 salt	30,378	91 (0.30)	26 (0.09)	65 (0.21)
Night 5 control vs. Night 5 salt	34,367	235 (0.68)	75 (0.22)	160 (0.47)
Night 6 control vs. Night 6 salt	29,618	38 (0.13)	19 (0.06)	19 (0.06)
Night 7 control vs. Night 7 salt	28,462	81 (0.28)	37 (0.13)	44 (0.15)

^a shows the number of transcripts have at least one expected transcripts predicted from RSEM (RNA-Seq by Expectation-Maximization)

^b is calculated by dividing the number of transcripts differentially expressed by the number of transcripts^a

Table 2 Functional categorization of selected transcripts differentially expressed in guard cells in response to salt stress. Significant change is expressed as a Log2 value of fold change of salt/control samples. For the full list of transcripts, please refer to Table S4.

Transcript ID	Annotation	Log ₂ FC	Time point	Gene function ^a
Contig12312	Fructose-bisphosphate aldolase 2 (<i>FBA2</i>)	-5.23	Night 7	CAM
Mcr017048.026	Phosphoglycerate mutase, 2,3-bisphosphoglycerate-independent (<i>PGM</i>)	-7.66, -9.34	Day 7, Night 5	CAM
Contig3509	β-amylase 5 (<i>AMYB5</i>)	2.98	Day 7	CAM
Contig20069	Carbonic anhydrase 2 (<i>CAH2</i>)	9.94	Day 5	CAM
Contig20312	Phosphoenolpyruvate carboxylase (<i>PEPC</i>) family protein	3.23	Night 5	CAM
Contig5856	Phosphoenolpyruvate carboxylase kinase 1 (<i>PPCK1</i>)	3.64	Night 7	CAM
comp24500_c0_seq1	Phosphoenolpyruvate carboxylase 1 (<i>PEPCI</i>)	2.63	Night 7	CAM; GC
comp14048_c0_seq1	Basic helix-loop-helix (bHLH) DNA-binding superfamily	-7.96	Day 6	GC; TF
Mcr002150.001	NAC domain containing protein 83	-9.76	Night 5	GC; TF
comp13165_c0_seq1	AGAMOUS-like 8	3.34	Night 7	TF
comp21521_c0_seq1	NAC domain containing protein 17	4.83	Night 5	TF
Contig16446	GRAS family transcription factor	-2.91	Night 5	TF
Contig18172	Duplicated homeodomain-like superfamily protein	3.56	Night 5	TF
Contig20720	WRKY DNA-binding protein 26	4.17, 2.83	Day 7 & Night 7	TF
Contig2750	DNA-binding bromodomain-containing protein	-8.22	Day 5	TF
Contig5045	FAR1-related sequence 5	4.16	Night 5	TF
Contig7210	BEL1-like homeodomain 7	2.49	Night 7	TF
Contig9771	Homeobox 7	2.63	Night 7	TF
Mcr003484.001	Zinc finger (CCCH-type) family protein	4.74, 5.33	Day 7 & Night 7	TF
Mcr008625.003	Telomere repeat binding factor 1	5.08	Day 6	TF
Mcr009671.000	BES1-interacting MYC-like protein 2	-9.14	Day 7	TF
Mcr010456.002	Basic helix-loop-helix (bHLH) DNA-binding superfamily protein	-3.92	Night 5	TF
Mcr010936.001	Integrase-type DNA-binding superfamily protein	4.63	Night 5	TF
Mcr014957.003	Basic helix-loop-helix (bHLH) DNA-binding superfamily protein	10.40	Night 5	TF
Mcr016258.030	Squamosa promoter binding protein-like 1	9.49	Night 5	TF

^a shows transcripts involved in either CAM mode, expressed in guard cell (GC), or transcription factor (TF)

Figure Legends

Figure 1. Leaf titratable acidity and morphological phenotype of *Mesembryanthemum crystallinum* under control and 500 mM NaCl treatment conditions. Four-week old plants were subjected to control (water) and 500 mM NaCl treatment for 4 to 28 days. **(A)** Measurement of leaf acidity was made at the start (8 am) of the photoperiod. Each bar represents mean of four replicates $\pm$ standard error (SE). Asterisks indicate statistical difference at p -value < 0.05 between the control and salt-treated plants as determined by t-tests. At the 7th day, the p -value is 0.0062. **(B)** Images of control and 7-day salt-treated plants.

Figure 2. Day/night profiles of leaf net CO₂ exchange and stomatal aperture in leaves of *M. crystallinum* grown under control (4-week old + 5 to 7 days with water) and salt treatment conditions (4-week old + 5 to 7 days with 500 mM NaCl). **(A)** Net CO₂ exchange. The dark bars on the x-axis represent dark periods and each gas exchange profile is the average of three biological replicates. Day 28 CAM plants were included as positive controls. **(B)** Stomatal aperture of control and salt-treated plants. Data are mean $\pm$ SE of three independent experiments, each with 60 – 80 stomata for a total of at least 180 stomata. Asterisks indicate statistical difference at p -value < 0.05 .

Figure 3. Expression profiles of CAM marker genes and PEPC activities during the C₃ to CAM transition at 2 am and 4 am. **(A)** Transcript levels of *PEPC*, *PPCK*, *CCAI*, *GTF1*, *ABII* and *PGK3* were determined by qRT-PCR. Error bars show the SE ($n \geq 3$). Asterisks indicate a significant difference between control and salt-treated plants (Student's t -test; p -value < 0.05). **(B)**

PEPC enzyme activities were measured in three biological replicates. Bars represent standard deviation (SD) of the means. Two-way ANOVA and Tukey's test were used for the PEPC activity analysis between different time points and different treated plants. Day 28 CAM plants were used as positive controls.

Figure 4. “Biological Process” functional Gene Ontology (GO) term classifications of DEGs. Please refer to Table S4 for detailed information of the DEGs.

Figure 5. Overlap of DEGs identified at different time points. (A) Overlap of DEGs from different day time points. (B) Overlap of DEGs from different night time points. (C) Overlap of DEGs from day and night.

Figure 6. Clustering analysis of transcripts differentially expressed in guard cells of the C₃ to CAM transition plants. *k*-means clustering algorithm (*k* = 8) were used. *n* indicates the number of transcripts in each cluster.

Supporting Information

Table S1. List of primers used in this study.

Table S2. Expression analysis of CAM-related, circadian-related and guard cell signaling related genes in leaf samples using real-time RT-PCR.

Table S3. List of genes showing reversed expression patterns under control versus salt stress conditions in the guard cell RNA-seq dataset. The expression difference is expressed as log 2 value of fold change between day and night. Cells with "light orange" and "light blue" colors represent up-regulated in day and night samples at adjusted p-value < 0.05, respectively

Table S4. Functional categorization of 495 transcripts significantly differentially expressed in guard cells in response to salt stress. Significantly change is expressed as a Log 2 value of fold change of salt/control samples.

Figure S1. Stomatal aperture in leaves of C₃ (4-week + 21-day water control) and CAM mode ice plants (4-week + 21-day salt treatment) under light and dark conditions. **(A)** Representative images showing stomatal aperture; **(B)** Stomatal aperture in leaves of C₃ and CAM plants under light and dark. Data are mean ± SE of three independent experiments ($n = 3$) with 60 – 80 stomata for each replicate (i.e., a total of at least 180 stomata for each experiment). Two-way ANOVA and Tukey's test were used for stomatal aperture analysis between the C₃ and CAM plants. For the images and stomatal aperture measurement, the epidermal peels were directly obtained from the plants, followed measuring the the width and length of the stomata under microscope.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6