Title:

Short-term evolution under copper stress increases probability of plasmid uptake

Author list:

Uli Klümper^{1,2,3,*}, Arnaud Maillard^{1,4,5}, Elze Hesse^{1,3}, Florian Bayer^{1,3}, Stineke van Houte^{1,3}, Ben Longdon¹, Will Gaze^{2,3}, Angus Buckling^{1,3}

¹ College of Life and Environmental Sciences, University of Exeter, Penryn, Cornwall, United Kingdom

² European Centre for Environment and Human Health, University of Exeter Medical School, Truro, Cornwall, United Kingdom

³ Environment & Sustainability Institute, University of Exeter, Penryn, Cornwall, United Kingdom

⁴ Ecole Normale Supérieure, ENS Ulm, Paris, France

⁵ UPMC, Sorbonne Universities, Paris, France

* corresponding author:

Uli Klümper

College of Life and Environmental Sciences

ESI Building

University of Exeter

TR109FE Penryn, Cornwall

United Kingdom

Email: u.klumper@exeter.ac.uk

Phone: (+44)7497497338

ORCID: 0000-0002-4169-6548

25 **Abstract:**

26 Understanding plasmid transfer dynamics remains a key knowledge gap in the mitigation of antibiotic
 27 resistance gene spread. Direct effects of exposure to stressors on plasmid uptake are well monitored.
 28 However, it remains untested whether evolution of strains under stress conditions modulates subsequent
 29 plasmid uptake. Here, we evolved a compost derived microbial community for six weeks under copper
 30 stress and non-exposed control conditions. We then tested the ability of isolated clones from both
 31 treatments to take up the broad host range plasmid pJKK5 from an *E.coli* donor strain. Clones pre-adapted
 32 to copper displayed a significantly increased probability to be permissive towards the plasmid compared to
 33 those isolated from the control treatment. Further, increased phylogenetic distance to the donor strain was
 34 significantly and negatively correlated with plasmid uptake probabilities across both treatments.

35

36 Text

37 Conjugal plasmids serve as main means of bacterial evolutionary adaptation to environmental stressors
 38 (Norman et al., 2009). The spread of plasmids encoding antibiotic resistance is considered a major threat to
 39 human health (WHO, 2014). Crucially, understanding plasmid spread dynamics remains a key knowledge
 40 gap (Smalla et al., 2018). Direct exposure to environmental stressors such as antibiotics (Slager et al., 2014),
 41 non-antibiotic pharmaceuticals (Wang et al., 2018) or metals (Klümper et al., 2017) can modulate
 42 immediate plasmid uptake in single strains and across microbial communities. This effect can either
 43 originate from direct selection or as a by-product of a general stress response. While immediate stress
 44 effects on plasmid uptake are well monitored, it remains untested whether ecological or evolutionary
 45 selection under stress conditions results in phenotypes with intrinsically higher plasmid permissiveness.
 46 There is evidence that stress or other environmental change can select for increased mutation (Pal et al.,
 47 2007) and recombination (Cooper, 2007) rate in bacteria. We therefore hypothesize that more permissive
 48 bacteria might also be favoured, as a result of horizontally acquired adaptive stress resistance.

49 Here, we tested if evolution in a microbial community exposed to metal stress has an effect on the plasmid
 50 uptake ability of individual clones. To infer a causal relationship between exposure to copper and
 51 subsequent plasmid uptake, we set up experimental compost communities in sterile compost following the
 52 protocol of Hesse et al. (2018). Hence, all treatments started off with the same community and level of
 53 permissiveness. Microcosms were incubated (75% humidity, 26°C), and twice (after 1 and 21 days)
 54 supplemented with either 2 ml filter-sterilized 0.25M CuSO₄ or ddH₂O. We here tested a total of 66 clones
 55 (27 copper, 39 control) that were isolated following 6 weeks of evolution. Copper tolerance increased
 56 significantly in clones pre-adapted to copper (Hesse et al., 2018). Clones were 16S sequenced using the 27F
 57 primer and a phylogenetic tree was constructed using mothur v1.41.1 (Schloss et al., 2009). Isolates
 58 belonged to 4 different phyla: Actinobacteria, Bacteroidetes, Firmicutes and three classes of Proteobacteria

68

To test permissiveness towards broad host range plasmid pKJK5 (Klümper et al., 2015; Li et al., 2018) each strain was mixed at 1:1 ratio with donor strain *Escherichia coli* MG1655::Km^R-Lpp-mCherry hosting the IncP-1 ϵ conjugative plasmid pKJK5::gfpmut3b (Bahl et al., 2007; Klümper et al., 2014). Experiments were carried out in LB medium in absence of any selective pressure, centrifuged for 2 minutes at 10000xg to ensure cell-to-cell contact and incubated (24 h; 28°C). Cells were harvested, resuspended and inoculated on M9 minimal media plates supplemented with 10 mM citrate and 10 μ g/mL tetracycline. Citrate as the single carbon source counter-selects against the *E.coli* donor strain, while tetracycline selects for acquisition of the tetracycline resistance encoding plasmid. Upon successful growth, green fluorescence, repressed in the donor strain but expressed upon transfer in transconjugants, was confirmed using fluorescence microscopy. Out of the 66 strains 42 were able to grow on citrate medium and displayed susceptibility to tetracycline. These were included in subsequent analysis with 71.4% permissive to plasmid pKJK5 (Figure 1). However, permissiveness differed strongly across phyla. Out of 25 proteobacterial strains, belonging to the same phylum as the *E.coli* donor, 22 (88%) took up the plasmid, while only 47% of gram positive strains (8/17) were permissive.

We subsequently fitted a logistic regression model (Figure 2) with the isolates evolutionary background and phylogenetic distance to the donor strain as explanatory variables to predict plasmid uptake probability. Both copper background (ANOVA χ^2 -test, p=0.0416, dF=39) and phylogenetic distance (ANOVA χ^2 -test, p=0.0033, dF=40) proved statistically significant in predicting plasmid uptake. However, while increasing distance to the donor strain had a negative effect on plasmid uptake probabilities, strains pre-adapted to copper were more likely to take up pKJK5 compared to non-adapted strains (Figure 2).

Subsequent work on single tractable species will explore the exact mechanisms underlying increased permissiveness. If, as a direct consequence of stress, more highly permissive bacteria were to survive better, they should also intrinsically host more plasmids, consequentially making them less permissive due to plasmid incompatibility or entry exclusion (Garcillán-Barcia and de la Cruz, 2008). This suggests that higher permissiveness might likely be an indirect by-product due to genetic changes in for example stress response, membrane permeability or pilus expression. Further, immunity towards plasmids could be evolutionary lost; CRISPR-Cas or less specific abortive infection systems can be lost under conditions when they bear immunity to horizontally transferred, potentially beneficial genes (Jiang et al., 2013).

Therefore, the conjugative host range can be increased under metal stress. The conjugative host range is generally assumed as much broader than the persistence host range (De Gelder et al., 2007). Consequently, the genomic signature of IncP-type plasmids suggests Proteobacteria as their main evolutionary hosts (Suzuki et al., 2010), which we also found to display far higher probabilities of plasmid uptake.

In strains with a higher degree of phylogenetic distance to the donor stability of pKJK5 might be very low and thus lost within few generations. However, long-term metal stress has been proven to elevate the retention of plasmids, even for those not coding for metal resistance (Smets et al., 2003). Even though plasmid-host co-evolution might reach an epidemiological dead end in some, likely more distant strains, such spill over remains crucial for propagation of antibiotic resistance. Unique events of recombination with the chromosome could happen before plasmid loss, especially since pKJK5 like many plasmids hosts a highly recombinative integrative element (Bahl et al., 2007). An increase of plasmid uptake ability and potentially retention time under metal stress conditions would consequently increase the likelihood and extent of such recombination events and foster the spread of antibiotic resistance.

124 **Acknowledgments**

125 UK received funding from the European Union's Horizon 2020 research and innovation program under
 126 Marie Skłodowska-Curie grant agreement no. 751699. UK, AB and WG were supported through an
 127 MRC/BBSRC grant (MR/N007174/1). AM received internship funding from the "Conseil Régional d'Ile de
 128 France".

129

130 **Competing interests**

131 The authors declare no competing interests.

132

133 **Author contributions**

134 UK, AM, SvH, WG and AB conceived the study and designed experiments; AM performed permissiveness
 135 assays with support from UK; EH performed evolution experiment and strain isolation; FB performed
 136 molecular work and sequencing; UK, AM, EH, BL analysed data; UK and AB wrote the manuscript.

137

138 **Competing interests**

139 The authors declare no competing interests.

140

141 **Materials & Correspondence**

142 All correspondence and material requests should be addressed to UK.

References

- Bahl, M.I., Hansen, L.H., Goesmann, A., and Sørensen, S.J. (2007) The multiple antibiotic resistance IncP-1 plasmid pKJK5 isolated from a soil environment is phylogenetically divergent from members of the previously established α , β and δ sub-groups. *Plasmid* **58**: 31–43.
- Cooper, T.F. (2007) Recombination speeds adaptation by reducing competition between beneficial mutations in populations of *Escherichia coli*. *PLoS Biol.* **5**: e225.
- Garcillán-Barcia, M.P. and de la Cruz, F. (2008) Why is entry exclusion an essential feature of conjugative plasmids? *Plasmid* **60**: 1–18.
- De Gelder, L., Ponciano, J.M., Joyce, P., and Top, E.M. (2007) Stability of a promiscuous plasmid in different hosts: No guarantee for a long-term relationship. *Microbiology* **153**: 452–463.
- Giller, K.E., Witter, E., and McGrath, S.P. (1998) Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: A review. *Soil Biol. Biochem.* **30**: 1389–1414.
- Hesse, E., O’Brien, S., Tromas, N., Bayer, F., Luján, A.M., van Veen, E.M., et al. (2018) Ecological selection of siderophore-producing microbial taxa in response to heavy metal contamination. *Ecol. Lett.* **21**: 117–127.
- Jiang, W., Maniv, I., Arain, F., Wang, Y., Levin, B.R., and Marraffini, L.A. (2013) Dealing with the Evolutionary Downside of CRISPR Immunity: Bacteria and Beneficial Plasmids. *PLoS Genet.* **9**: e1003844.
- Klümper, U., Riber, L., Dechesne, A., Sannazzarro, A., Hansen, L.H., Sørensen, S.J., and Smets, B.F. (2015) Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community. *ISME J.* **9**: 934–945.
- Klümper, U., Dechesne, A., Riber, L., Brandt, K.K., Gülay, A., Sørensen, S.J., and Smets, B.F. (2017) Metal stressors consistently modulate bacterial conjugal plasmid uptake potential in a phylogenetically conserved manner. *ISME J.* **11**: 152–165.
- Klümper, U., Dechesne, A., and Smets, B.F. (2014) Protocol for evaluating the permissiveness of bacterial communities toward conjugal plasmids by quantification and isolation of transconjugants. In, *Hydrocarbon and Lipid Microbiology Protocols, Springer Protocols Handbook*. Humana Press, pp. 275–288.
- Li, L., Dechesne, A., He, Z., Madsen, J.S., Nesme, J., Sørensen, S.J., and Smets, B.F. (2018) Estimating the transfer range of plasmids encoding antimicrobial resistance in a wastewater treatment plant microbial community. *Environ. Sci. Technol. Lett.* **5**: 260–265.
- Norman, A., Hansen, L.H., and Sørensen, S.J. (2009) Conjugative plasmids: vessels of the communal gene pool. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **364**: 2275–2289.
- Pal, C., Maciá, M.D., Oliver, A., Schachar, I., and Buckling, A. (2007) Coevolution with viruses drives the evolution of bacterial mutation rates. *Nature* **450**: 1079–1081.
- Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., et al. (2009) Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* **75**: 7537–7541.
- Slager, J., Kjos, M., Attaiech, L., and Veening, J.W. (2014) Antibiotic-induced replication stress triggers bacterial competence by increasing gene dosage near the origin. *Cell* **157**: 395–406.
- Smalla, K., Cook, K., Djordjevic, S.P., Klümper, U., and Gillings, M. (2018) Environmental dimensions of antibiotic resistance: Assessment of basic science gaps. *FEMS Microbiol. Ecol.*
- Smets, B.F., Morrow, J.B., and Pinedo, C.A. (2003) Plasmid introduction in metal-stressed, subsurface-

185 derived microcosms: Plasmid fate and community response. *Appl. Environ. Microbiol.* **69**: 4087–4097.

186 Suzuki, H., Yano, H., Brown, C.J., and Top, E.M. (2010) Predicting plasmid promiscuity based on genomic

187 signature. *J. Bacteriol.* **192**: 6045–6055.

188 Wang, Y., Lu, J., Mao, L., Li, J., Yuan, Z., Bond, P.L., and Guo, J. (2018) Antiepileptic drug carbamazepine

189 promotes horizontal transfer of plasmid-borne multi-antibiotic resistance genes within and across

190 bacterial genera. *ISME J.* **1**.

191 WHO (2014) Antimicrobial Resistance Global Report on Surveillance.

192