Peer Community In Ecology

RESEARCH ARTICLE

Field assessment of precocious maturation in salmon parr using ultrasound imaging

Cite as: Nevoux M, Marchand F, Forget G, Huteau D, Tremblay J, and Destouches J-P. Field assessment of precocious maturation in salmon parr using ultrasound imaging. bioRxiv 425561, ver. 3 peer-reviewed and recommended by PCI Ecology (2019). DOI: 10.1101/425561

Published: 25th April 2019

Recommender:
Jean-Olivier Irsson

Reviewers:
Hervé Capra and one anonymous reviewer

Correspondence:
marie.nevoux@inra.fr

Marie Nevoux^{1,2}, Frédéric Marchand^{2,3}, Guillaume Forget^{1,2}, Dominique Huteau^{1,2}, Julien Tremblay^{2,3}, Jean-Pierre Destouches^{2,3}

¹ INRA, Agrocampus-Ouest, Ecology and Ecosystem Health, 65 rue de Saint-Brieuc, Rennes, France

² Management of Aquatic Ecosystems in Agricultural Watersheds, INRA, Agence Française pour la Biodiversité, Rennes, France

³ INRA, Ecologie et Ecotoxicologie Aquatique, 65 rue de Saint-Brieuc, Rennes, France

This article has been peer-reviewed and recommended by:
Peer Community in Ecology (DOI: 10.24072/pci.ecology.100021)

ABSTRACT

Salmonids are characterized by a large diversity of life histories, but their study is often limited by the imperfect observation of the true state of an individual in the wild. Challenged by the need to reduce uncertainty of empirical data, recent development in medical imaging techniques offered new opportunities to assess precocious maturation in Atlantic salmon parr. Traditional phenotypic (external) examination and ultrasound (internal) examination were compared and recommendations on fish handling and ultrasound image interpretation are provided. By allowing to see the unseen, portable ultrasound imaging offers great opportunities for ecological studies in the wild, such as the assessment of individual sexual maturation.

Keywords: Atlantic salmon *Salmo salar*, ecology, empirical study, fisheries, individual heterogeneity, life history trait, maturation, ultrasound scan

Introduction

Long term monitoring of animal populations in the wild is of great interest for ecology and management (Clutton-Brock and Sheldon 2010). In a context of global change, such long term monitoring programs contribute to highlight major changes in population composition, dynamics and abundance over the last decades (Parmesan 2006). In the North Pacific, analyses confirmed climate-related shifts in the abundance of most salmonid species since the 1940's, associated with reported ecosystem regime shifts (Irvine and Fukuwaka 2011). In the North Atlantic, the widespread decline in salmon abundance was associated with a decrease in both survival and age at maturity (Chaput 2012). This global change makes it particularly important to be able to predict population structure and abundance in the future under different climatic scenarios. However, predictive population dynamics models are difficult to parameterize and often return high level of uncertainty, which hinder their use by managers and policy makers. Stock assessments form the basis for catch advice for salmon fisheries, but they mask regional differences and annual river-specific stock assessments are only available for some 25% of the rivers (Chaput 2012). This is partly because field records are imperfect observation of the true state of an individual (Genovart et al. 2012). For instance, maturation status and gender is rarely known with certainty in fish monitoring programs without invasive or lethal methods. Thus, field studies are challenged by the need to reduce uncertainty in individual-level data.

Atlantic salmon (*Salmo salar*) is an anadromous species that reproduces in freshwater and migrates at sea to grow and undergo sexual maturation. However, in this and other salmonid species, some young males (parrs) can mature precociously in freshwater (Fleming 1996). It is proposed that the intensity of intra-sexual competition for mates and environmental variability in growth opportunities, determines a polygenic threshold that triggers precocious maturation (Hutchings and Myers 1994). Mature male parrs attempt to 'sneak' access to large anadromous females for mating (Fleming 1996). They can contribute substantially to the reproduction (Moran et al. 1996) but, as maturation seems to be traded against survival, mature parr have a low probability to migrate at sea (Buoro et al. 2010). The occurrence of this alternative reproduction strategy is highly variable over time, space, and age classes (Baglinière and Maisse 1985; Myers et al. 1986) and may be driven by frequency dependent selection, i.e. the reproductive success of precocious parrs increases with decreasing frequency of precocious maturation in the population (Berejikian et al. 2010). Yet, precise quantification of this phenotype remains rare in empirical studies and in salmon population dynamic models. Furthermore, precocious maturation is virtually ignored by stock assessment models and in the management of Atlantic salmon

fisheries (e.g. ICES 2013). This means that mature parrs are included neither in the stock nor in the recruitment figures. This might bias stock-recruitment relationships and underestimate freshwater survival as well as salmon production. For instance, using more than 280,000 records from Little Codroy River parrs, Myers (1984) estimated that 60% of the stock of adult male salmon in Newfoundland was lost because of increased mortality due to precocious maturation. Caswell et al. (1984) demonstrated that the effect of reproduction by parrs on population growth rate was always greater than that of reproduction by adults. It is suspected that this lack of consideration for mature parrs is due to the complexity of the life cycle of Atlantic salmon. But it could also be due to the difficulty of quantifying the occurrence of precocious maturation in wild salmon populations.

In the field, precocious maturity in parrs is traditionally assessed by pressing the abdomen of the fish to extract milt (sperm). However, not all mature males may be producing milt at the time of capture. Moreover, based on the general aspect of the fish, experienced fieldworkers sometimes suspect maturity in some individuals that do not excrete milt upon abdominal pressure. However, researchers are often reluctant to use such expert opinions in their analysis due to the potential for incorrect diagnosis and strong operator bias.

Ultrasound imaging, which gave rise to significant advances in human medicine, offers reliable visualization of internal organs. Ultrasound imaging is routinely used in aquaculture as it provides an accurate, non-invasive, non-lethal method that is more accurate than visual methods, to determine gender or gonadal growth in fish for example (Novelo and Tiersch 2012). In adult salmon, although gonads of immature males may be difficult to discern, gonads of females are always visible and hence sexing is possible by deduction (Martin et al. 1983; Mattson 1991). However the ovaries and testes of the juvenile (immature) salmon seem difficult to identify (Martin et al. 1983).

The development of portable devices offers an opportunity to bring ultrasound machines in the field. This paper illustrates how ultrasound scanners can increase the range of biological traits that can be monitored in long-term salmonid populations surveys. In particular, the maturation status of salmon parr (i.e. freshwater resident salmon) was assessed in the field using traditional phenotypic examination and ultrasound examination. The two approaches are compared and it is tested whether biological or operator-related factors affect the uncertainty in external phenotypic assessment of parr maturity. Finally, recommendation on fish handling and on the interpretation of ultrasound images is provided.

Methods

Salmon monitoring

Atlantic salmon monitoring takes place every year in September or October in the river Oir, Normandy, France (Marchand et al. 2018). Young individuals (parrs) are captured using a standard electrofishing protocol. They are then placed in a light anesthetic solution, measured, weighed, scanned, and returned to the river within 30 min of their capture. Age is assessed through scale reading. In 2015, 2016, and 2017, a total of 850 salmon was examined (366 parrs of age 0 and 484 parrs of age 1) for maturity using a traditional, phenotypic (i.e. external), approach and ultrasound imaging. All parrs were caught within a period of 8 days in early October. All the animal experimentation in this study was performed according to French legislation and under licence APAFIS-201602051204637. The data is available in Appendix 1.

Traditional phenotypic examination

First, the field operator assesses the maturation state of parrs by gently pressing the side of the fish. The excretion of milt is indicative of a mature male, while the absence of milt can be encountered in a maturing male whose gonads do not yet produce milt, an immature male, or a female (females are all immature at this stage). The field operator may also report that some fish *look like* mature parrs: they display phenotypic traits similar to those of mature fish (e.g. colour pattern, body shape) but do not excrete milt. This phenotypic assessment relies less on a fixed set of quantifiable parameters than on a general, and subjective, appreciation of the fish, which is specific to each operator.

Ultrasound examination

Then, the ultrasound operator screens all salmon parrs for precocious maturation using a portable ultrasound scanner M-Turbo (Sonosite) with a 5-10 MHz linear transducer. The default setting, for muscular examination, is selected. As water perfectly transmits ultrasounds, the use of ultrasound transmission gel is not needed. Fish are placed on their side in a tank of freshwater and the transducer is operated in the water, keeping it 1-2cm above the belly of the fish. Sagittal images of the abdominal cavity are produced, i.e. the transducer is aligned with the lateral line. The maturation state of each individual is directly assessed from live images on the ultrasound monitor. Parrs are classified as mature when gonads are detected, or immature when the gonads are too small to be detected. Ultrasound snapshots can be stored on the ultrasound machine and then exported to a USB memory stick for post-processing.

Comparison of phenotypic and ultrasound methods

The comparison of the proportion of parrs that are recorded each year as mature or immature by each approach is done with a χ^2 test. The number of mature parrs is defined as the sum of individuals whose gonads are detected with the ultrasound and individuals whose gonads are not detected with the ultrasound but that produce milt. Because the probability to detect precocious maturation with the phenotypic method depends on the probability for a mature male to produce milt at the time of capture, an analysis is conducted to investigate which biological factor drive this probability. Empirical records show a large interannual variability in the proportion of maturing parrs (Baglinière and Maisse 1985; Myers et al. 1986). Similarly, the year of capture is expected to affect milting within mature parrs. Thus, a generalized linear model (GLM) with a binomial distribution of errors is used to test for a year effect. Then, a mixed model with year as a random intercept and a binomial distribution of errors is used to test for the effect of age, body length, and their interaction. A GLM is also run to test whether the probability to miss a mature parr that does not produce milt is affected by age, body length, operator identity, and time of day (testing for operator tiredness). The date of capture is not considered because of the low variance in this variable. The significance of the above-mentioned effects is assessed using the z-value. All models are run in R (R Development Core Team 2018).

Results and discussion

Ultrasound assessment of precocious maturity

A trained ultrasound operator conducts ultrasound examination for maturation assessment in ca. 5 seconds. In immature individuals, the liver, stomach, caeca, and intestine can be identified (Figure 1a). Note that all internal organs appear more clearly on live images, during ultrasound examination, than on the snapshots displayed here. Nevertheless, the heterogeneous filling and uneven granularity of the abdominal cavity are diagnostic of immature individuals, as well as the concave shape of the cavity, which appears depressed on both sides. In mature parrs (precocious males), gonads virtually fill the whole abdominal cavity and ultrasound snapshots depict a homogeneous and finely granulated pattern (Figure 1b, c); digestive organs are hardly visible and the cavity is convex. In mature males that do not produce milt (Figure 1b), gonads appear less developed and fill the cavity to a lesser extent than in males producing milt (Figure 1c), which is consistent with a lower degree of maturity in the former. Interestingly, ultrasound imaging provides the same objective diagnosis of maturation in males that do not produce milt (Figure 1b) and ones that do (Figure 1c). It also discriminates truly mature parrs from well-fed individuals, which the operator can mistake for mature males upon visual inspection because of their round belly.

Figure 1. Assessing maturity in salmon parr using phenotypic view (left panel) and ultrasound imaging (right panel): a) immature one-year-old parr, b) mature young-of-the-year parr that does not produce milt (precocious male), and c) mature one-year-old parr that produces milt (precocious male) in river Oir. On ultrasound snapshots, the head of the fish is on the left but not shown, the left flank is towards the bottom of the snapshot and the right towards the top. Gonads are identified by the letter “G”.

Comparison of phenotypic and ultrasound methods

The proportion of mature parrs detected by the phenotypic method is much lower and more variable (63.8 %, SD = 40.7%) than with the ultrasound method (95.4%, SD = 7.1%, $p = 0.199$). On average 33.4% of mature males do not produce milt. A strong year effect is detected ($p < 0.001$) in the probability to produce milt at the time of capture in mature males. When accounting for this year effect as a random factor, a strong increase with age (estimate = 5.582, SE = 1.031, $p < 0.001$) and a negative interaction with body length within each age class (estimate = -0.031, SE = 0.008, $p < 0.001$) are

detected. This means that a higher proportion of mature males produce milt in autumn in old parrs than in young ones and, within a given age class in small parrs than in large ones.

Because the probability to detect precocious maturation with the phenotypic method depends on the probability for a mature male to produce milt at the time of capture, our result highlights that both intrinsic and environmental factors can affect the level of detection of the precocious state. The ultrasound method shows that only 16.8% and 13.2% of the mature males are identified as precocious parrs at the time of capture in 2015 and 2017, while this proportion rises to 87.6% in 2016 (Table 1). Empirical evidence have shown an increase in the proportion of mature parrs as the parrs are getting older (Baglinière and Maisse 1985; Myers et al. 1986). But this study suggests that mature parrs develop and produce milt earlier in the season than young ones, which may accentuate even further the age-specific pattern detected above, with the phenotypic method.

In our study, the maturation state of 106 individuals is uncertain according to the phenotypic method, i.e. individuals that look like mature parrs but do not produce milt. However, following ultrasound examination, 89.6 % of those parrs are indeed found to be mature. This result acknowledges the high level of expertise of our trained operators to assess the maturation state of parrs from external phenotypic examination. Conversely, gonads are not detected in 10.4% of those parrs (false positive) and the field operators do not detect milt nor suspect maturity in 5.6 % of the actually mature males (false negative). This later proportion differs between the two operators ($p = 0.008$). None of the other variables has a significant effect. This highlights how difficult it is to assess parr maturation from external examination in the field in a fully reliable and replicable manner.

Contrary to the phenotypic method, it is assumed that the ultrasound method cannot produce false positives, because gonads large enough to be detected by the ultrasound equipment are only found in mature parrs. The ultrasound operator missed 2.7% of the males producing milt (false negative). The sample size is too small to assess the potential effect of external factors on this proportion through multivariate analysis. Still, this shows that ultrasound imaging is still subject to a low degree of uncertainty in maturity assessment. The correct interpretation of ultrasound images depends on the skill of the operator. Indeed, some training is required to operate the transducer in a fluent way and navigate between different anatomic layers. In this regard, a good knowledge of the internal structure for the species of interest is a prerequisite. This improves the quality of the ultrasound snapshots and minimizes misinterpretation issues. In hindsight, it is also recommended that all ultrasounds

should be examined and interpreted by multiple operators. This ensures a quick examination and thus minimizes the impact on fish wellbeing.

Table 1. Comparison of the number of records using external phenotypic observation (rows) and ultrasound imaging (columns) to assess maturation in parrs each year.

Year	Phenotype method \ ultrasound method	Gonads detected with ultrasound	Gonads not detected with ultrasound
2015	Parr producing milt	17	0
	Parr not producing milt that looks like a mature parr	74	8
	Parr not producing milt that looks like an immature parr	10	73
2016	Parr producing milt	181	2
	Parr not producing milt that looks like a mature parr	20	3
	Parr not producing milt that looks like an immature parr	6	251
2017	Parr producing milt	23	4
	Parr not producing milt that looks like a mature parr	1	0
	Parr not producing milt that looks like an immature parr	3	174

Conclusion

Assessing precocious maturation in parrs from phenotypic observation can reach a high level of accuracy in trained operators. However, this method relies on some subjective criteria and gut feeling, which render the level of accuracy in the data difficult to quantify. By allowing to see the unseen, portable ultrasound imaging offers great opportunities for ecological studies in the wild. As illustrated in this paper, key phenotypic traits become accessible and help better characterize the true biological state of individuals. Ultrasound imaging is an objective, easily transferable and replicable approach to investigate precocious maturity in parrs, as it relies on key diagnostic features. A naïve operator can gain a good expertise and assess precocious maturation with the ultrasound method within a day, whereas more training would be required to achieve a similar level of expertise with the phenotypic method. In the end, this study calls for the assessment of precocious maturation in Atlantic salmon parrs using the two methods together: 1) testing whether parrs produce milt is a quick examination and provides an objective and undeniable evidence of maturity, 2) ultrasound imaging should remove any ambiguity about the state of maturation in parrs that do not produce milt. Reducing uncertainty in empirical data this way should offer new opportunities for further research on this alternative breeding strategy in salmon and improve our understanding of this key biological process.

Acknowledgements

This preprint has been peer-reviewed and recommended by Peer Community In Ecology (<https://dx.doi.org/10.24072/pci.ecology.100021>).

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article.

References

- Baglinière, J.-L., and Maisse, G. 1985. Precocious maturation and smoltification in wild Atlantic salmon in the Armorican massif, France. *Aquaculture* 45(1): 249–263. doi:10.1016/0044-8486(85)90274-1.
- Berejikian, B.A., Van Doornik, D.M., Endicott, R.C., Hoffnagle, T.L., Tezak, E.P., Moore, M.E., and Atkins, J. 2010. Mating success of alternative male phenotypes and evidence for frequency-dependent selection in Chinook salmon, *Oncorhynchus tshawytscha*. *Can. J. Fish. Aquat. Sci.* 67(12): 1933–1941. doi:10.1139/F10-112.
- Buoro, M., Prévost, E., and Gimenez, O. 2010. Investigating Evolutionary Trade-Offs in Wild Populations of Atlantic Salmon (*Salmo salar*): Incorporating Detection Probabilities and Individual Heterogeneity. *Evolution* 64(9): 2629–2642. doi:10.1111/j.1558-5646.2010.01029.x.
- Caswell, H., Naiman, R.J., and Morin, R. 1984. Evaluating the consequence of reproduction in complex salmonid life cycles. *Aquaculture* 43: 123–134.
- Chaput, G. 2012. Overview of the status of Atlantic salmon (*Salmo salar*) in the North Atlantic and trends in marine mortality. *ICES J. Mar. Sci.* 69(9): 1538–1548. doi:10.1093/icesjms/fss013.
- Clutton-Brock, T., and Sheldon, B.C. 2010. Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends Ecol. Evol.* 25(10): 562–573.
- Fleming, I.A. 1996. Reproductive strategies of Atlantic salmon: ecology and evolution. *Rev. Fish Biol. Fish.* 6(4): 379–416. doi:10.1007/BF00164323.
- Genovart, M., Pradel, R., and Oro, D. 2012. Exploiting uncertain ecological fieldwork data with multi-event capture–recapture modelling: an example with bird sex assignment. *J. Anim. Ecol.*: no-no. doi:10.1111/j.1365-2656.2012.01991.x.

- Hutchings, J.A., and Myers, R.A. 1994. The evolution of alternative mating strategies in variable environments. *Evol. Ecol.* 8(3): 256–268.
- ICES. 2013. Report of the Working Group on North Atlantic Salmon (WGNAS). Copenhagen, Denmark.
- Irvine, J.R., and Fukuwaka, M. 2011. Pacific salmon abundance trends and climate change. *ICES J. Mar. Sci.* 68(6): 1122–1130. doi:10.1093/icesjms/fsq199.
- Marchand, F., Azam, D., Delanoë, R., Destouches, J.-P., Tremblay, J., Baglinière, J.-L., Nevoux, M., Prévost, E., and Azam, D. 2018. Abundances and biological traits of fish sampled by electrofishing (except specific abundance indices) on the River Oir since 1988 (France). doi:10.15468/yaikgn.
- Martin, R.W., Myers, J., Sower, S.A., Phillips, D.J., and Mcauley, C. 1983. Ultrasonic Imaging, a Potential Tool for Sex Determination of Live Fish. *North Am. J. Fish. Manag.* 3(3): 258–264. doi:10.1577/1548-8659(1983)3<258:UIAPTF>2.0.CO;2.
- Mattson, N.S. 1991. A new method to determine sex and gonad size in live fishes by using ultrasonography. *J. Fish Biol.* 39(5): 673–677. doi:10.1111/j.1095-8649.1991.tb04397.x.
- Moran, P., Pendás, A.M., Beall, E., and García-Vázquez, E. 1996. Genetic assessment of the reproductive success of Atlantic salmon precocious parr by means of VNTR loci. *Heredity* 77(6): 655–660. doi:10.1038/hdy.1996.193.
- Myers, R.A. 1984. Demographic Consequences of Precocious Maturation of Atlantic Salmon (*Salmo salar*). *Can. J. Fish. Aquat. Sci.* 41(9): 1349–1353. doi:10.1139/f84-165.
- Myers, R.A., Hutchings, J.A., and Gibson, R.J. 1986. Variation in Male Parr Maturation Within and Among Populations of Atlantic Salmon, *Salmo salar*. *Can. J. Fish. Aquat. Sci.* 43(6): 1242–1248. doi:10.1139/f86-154.
- Novelo, N.D., and Tiersch, T.R. 2012. A Review of the Use of Ultrasonography in Fish Reproduction. *North Am. J. Aquac.* 74(2): 169–181. doi:10.1080/15222055.2012.672370.
- Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.* 37(1): 637–669.
- R Development Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available from <http://www.R-project.org>.

Appendix

Appendix 1: Individual records of date and hour of capture, body size, age, phenotypic observation and ultrasound examination for the 850 salmon parrs captured in the river Oir for this study.

Date	Hour	Size (mm)	Age	Phenotype maturation	Phenotype operator	Ultrasound maturation
05/10/2015	10	95	1	milt	FM	mature
05/10/2015	10	115	1	milt	FM	mature
05/10/2015	10	102	1	milt	FM	mature
05/10/2015	10	113	1	milt	FM	mature
05/10/2015	10	125	1	milt	FM	mature
05/10/2015	10	90	1	milt	FM	mature
05/10/2015	10	111	1	milt	FM	mature
05/10/2015	10	104	1	milt	FM	mature
05/10/2015	10	105	1	milt	FM	mature
05/10/2015	10	117	1	milt	FM	mature
05/10/2015	10	105	1	milt	FM	mature
05/10/2015	11	120	1	milt	FM	mature
05/10/2015	11	109	1	milt	FM	mature
05/10/2015	11	111	1	immature	FM	immature
05/10/2015	11	110	1	immature	FM	immature
05/10/2015	11	116	1	immature	FM	immature
05/10/2015	11	124	1	immature	FM	mature
05/10/2015	11	117	1	immature	FM	immature
05/10/2015	11	112	1	immature	FM	immature
05/10/2015	12	79	0	immature	FM	mature
05/10/2015	12	71	0	mature	FM	mature
05/10/2015	12	123	1	immature	FM	immature
05/10/2015	12	121	1	immature	FM	immature
05/10/2015	12	107	1	immature	FM	immature
05/10/2015	12	120	1	mature	FM	mature
05/10/2015	12	130	1	immature	FM	immature
05/10/2015	12	77	0	mature	FM	mature
05/10/2015	12	87	0	immature	FM	immature
05/10/2015	12	75	0	mature	FM	mature
05/10/2015	14	112	1	immature	FM	immature
05/10/2015	13	102	1	immature	FM	immature
05/10/2015	13	112	1	immature	FM	immature

05/10/2015	13	107	1	mature	FM	mature
05/10/2015	14	61	0	mature	FM	mature
05/10/2015	13	100	1	immature	FM	mature
05/10/2015	13	102	1	immature	FM	immature
05/10/2015	14	115	1	mature	FM	mature
05/10/2015	13	93	1	mature	FM	mature
05/10/2015	13	104	1	immature	FM	mature
05/10/2015	13	125	1	immature	FM	immature
05/10/2015	14	110	1	immature	FM	immature
05/10/2015	13	93	1	mature	FM	mature
05/10/2015	13	121	1	immature	FM	immature
05/10/2015	13	108	1	mature	FM	mature
05/10/2015	13	92	1	mature	FM	mature
05/10/2015	13	112	1	immature	FM	immature
05/10/2015	13	100	1	mature	FM	mature
05/10/2015	13	75	0	mature	FM	mature
05/10/2015	15	105	1	mature	FM	mature
05/10/2015	15	121	1	mature	FM	mature
05/10/2015	15	112	1	immature	FM	immature
05/10/2015	15	76	0	mature	FM	mature
05/10/2015	15	100	1	mature	FM	mature
05/10/2015	15	109	1	mature	FM	mature
05/10/2015	15	72	0	mature	FM	mature
05/10/2015	15	108	1	immature	FM	immature
05/10/2015	15	94	1	mature	FM	mature
05/10/2015	15	68	0	immature	FM	immature
05/10/2015	15	103	1	mature	FM	mature
05/10/2015	15	119	1	mature	FM	mature
05/10/2015	15	116	1	mature	FM	mature
05/10/2015	15	71	0	mature	FM	mature
05/10/2015	15	122	1	immature	FM	immature
05/10/2015	15	110	1	immature	FM	immature
05/10/2015	15	110	1	mature	FM	mature
05/10/2015	15	120	1	immature	FM	immature
05/10/2015	16	103	1	mature	FM	mature
05/10/2015	17	115	1	immature	FM	immature
05/10/2015	16	91	1	milt	FM	mature
05/10/2015	16	100	1	immature	FM	mature
05/10/2015	17	66	0	mature	FM	mature
05/10/2015	16	72	0	immature	FM	mature

05/10/2015	16	122	1	mature	FM	mature
05/10/2015	16	127	1	immature	FM	immature
05/10/2015	17	79	0	mature	FM	immature
05/10/2015	16	115	1	immature	FM	immature
05/10/2015	16	101	1	milt	FM	mature
05/10/2015	17	102	1	immature	FM	immature
05/10/2015	17	65	0	milt	FM	mature
05/10/2015	17	122	1	immature	FM	immature
05/10/2015	17	110	1	immature	FM	immature
05/10/2015	17	102	1	immature	FM	immature
05/10/2015	17	103	1	immature	FM	immature
05/10/2015	16	110	1	immature	FM	immature
05/10/2015	16	106	1	mature	FM	mature
05/10/2015	16	71	0	mature	FM	immature
05/10/2015	17	70	0	immature	FM	immature
06/10/2015	10	99	1	immature	FM	mature
06/10/2015	10	117	1	immature	FM	immature
06/10/2015	11	111	1	immature	FM	immature
06/10/2015	11	119	1	immature	FM	immature
06/10/2015	10	102	1	immature	FM	immature
06/10/2015	10	115	1	mature	FM	mature
06/10/2015	10	125	1	mature	FM	mature
06/10/2015	10	106	1	mature	FM	mature
06/10/2015	10	100	1	mature	FM	mature
06/10/2015	09	110	1	mature	FM	mature
06/10/2015	10	70	0	mature	FM	mature
06/10/2015	10	116	1	immature	FM	immature
06/10/2015	10	123	1	immature	FM	immature
06/10/2015	10	71	0	immature	FM	immature
06/10/2015	10	110	1	mature	FM	mature
06/10/2015	10	97	1	mature	FM	mature
06/10/2015	10	109	1	mature	FM	mature
06/10/2015	11	84	0	immature	FM	mature
06/10/2015	12	104	1	immature	FM	immature
06/10/2015	12	79	0	mature	FM	mature
06/10/2015	11	125	1	immature	FM	immature
06/10/2015	11	109	1	immature	FM	immature
06/10/2015	12	105	1	immature	FM	mature
06/10/2015	12	72	0	mature	FM	mature
06/10/2015	12	112	1	immature	FM	immature

06/10/2015	11	113	1	immature	FM	immature
06/10/2015	14	110	1	immature	FM	immature
06/10/2015	15	138	1	immature	FM	immature
06/10/2015	15	119	1	immature	FM	immature
06/10/2015	15	112	1	mature	FM	mature
06/10/2015	14	110	1	immature	FM	immature
06/10/2015	13	118	1	immature	FM	immature
06/10/2015	14	120	1	immature	FM	immature
06/10/2015	13	115	1	immature	FM	immature
06/10/2015	15	115	1	immature	FM	immature
06/10/2015	16	65	0	mature	FM	mature
06/10/2015	15	112	1	immature	FM	immature
06/10/2015	15	110	1	immature	FM	immature
06/10/2015	16	80	0	immature	FM	immature
06/10/2015	15	74	0	mature	FM	mature
06/10/2015	15	69	0	mature	FM	immature
06/10/2015	15	97	1	immature	FM	immature
06/10/2015	15	114	1	mature	FM	mature
06/10/2015	15	127	1	immature	FM	immature
06/10/2015	16	106	1	mature	FM	mature
06/10/2015	17	106	1	mature	FM	mature
06/10/2015	18	102	1	mature	FM	mature
06/10/2015	16	87	0	immature	FM	immature
06/10/2015	17	121	1	mature	FM	mature
06/10/2015	16	105	1	immature	FM	immature
06/10/2015	17	108	1	immature	FM	immature
06/10/2015	17	103	1	immature	FM	immature
06/10/2015	17	122	1	immature	FM	immature
06/10/2015	16	107	1	immature	FM	immature
06/10/2015	17	108	1	immature	FM	immature
06/10/2015	17	105	1	mature	FM	mature
06/10/2015	17	105	1	immature	FM	immature
06/10/2015	17	121	1	immature	FM	immature
06/10/2015	17	113	1	immature	FM	immature
06/10/2015	17	104	1	immature	FM	immature
06/10/2015	17	72	0	mature	FM	mature
06/10/2015	10	94	1	immature	FM	immature
06/10/2015	18	102	1	immature	FM	immature
06/10/2015	10	97	1	mature	FM	mature
07/10/2015	10	122	1	mature	FM	immature

07/10/2015	11	70	0	mature	FM	mature
07/10/2015	11	72	0	mature	FM	mature
07/10/2015	11	108	1	mature	FM	immature
07/10/2015	11	77	0	mature	FM	mature
07/10/2015	12	113	1	mature	FM	mature
07/10/2015	12	121	1	mature	FM	immature
07/10/2015	12	81	0	mature	FM	mature
07/10/2015	14	111	1	mature	FM	mature
07/10/2015	14	101	1	mature	FM	mature
07/10/2015	13	100	1	mature	FM	mature
07/10/2015	14	110	1	mature	FM	mature
07/10/2015	16	71	0	mature	FM	mature
08/10/2015	10	115	1	mature	FM	mature
08/10/2015	09	78	0	mature	FM	mature
08/10/2015	11	75	0	mature	FM	mature
08/10/2015	11	133	1	mature	FM	mature
08/10/2015	09	89	1	mature	FM	mature
08/10/2015	09	100	1	immature	FM	mature
08/10/2015	09	80	0	mature	FM	mature
08/10/2015	10	124	1	mature	FM	immature
08/10/2015	10	118	1	milt	FM	mature
08/10/2015	10	73	0	mature	FM	mature
08/10/2015	11	67	0	immature	FM	immature
08/10/2015	12	81	0	mature	FM	mature
08/10/2015	11	125	1	mature	FM	mature
08/10/2015	12	114	1	mature	FM	mature
08/10/2015	13	122	1	mature	FM	mature
08/10/2015	12	82	0	mature	FM	immature
08/10/2015	12	100	1	mature	FM	mature
08/10/2015	13	72	0	mature	FM	mature
03/10/2016	08	116	1	milt	FM	mature
03/10/2016	08	113	1	milt	FM	mature
03/10/2016	08	106	1	immature	FM	immature
03/10/2016	08	72	0	immature	FM	immature
03/10/2016	08	77	0	immature	FM	immature
03/10/2016	08	74	0	immature	FM	immature
03/10/2016	09	125	1	immature	FM	immature
03/10/2016	09	70	0	immature	FM	immature
03/10/2016	09	119	1	milt	FM	mature
03/10/2016	09	113	1	immature	FM	immature

03/10/2016	09	68	0	immature	FM	immature
03/10/2016	09	100	1	milt	FM	mature
03/10/2016	09	120	1	immature	FM	immature
03/10/2016	09	113	1	immature	FM	immature
03/10/2016	09	69	0	immature	FM	immature
03/10/2016	09	105	1	immature	FM	immature
03/10/2016	09	74	0	immature	FM	immature
03/10/2016	09	128	1	immature	FM	immature
03/10/2016	09	117	1	milt	FM	mature
03/10/2016	09	83	0	immature	FM	immature
03/10/2016	09	108	1	milt	FM	mature
03/10/2016	09	133	1	milt	FM	mature
03/10/2016	09	126	1	milt	FM	mature
03/10/2016	09	118	1	mature	FM	mature
03/10/2016	10	116	1	milt	FM	mature
03/10/2016	10	99	1	milt	FM	mature
03/10/2016	10	72	0	immature	FM	immature
03/10/2016	10	76	0	milt	FM	mature
03/10/2016	10	113	1	milt	FM	mature
03/10/2016	10	97	0	immature	FM	immature
03/10/2016	10	88	0	immature	FM	immature
03/10/2016	10	85	0	milt	FM	mature
03/10/2016	10	69	0	immature	FM	immature
03/10/2016	10	84	0	immature	FM	immature
03/10/2016	10	105	1	mature	FM	mature
03/10/2016	10	115	1	milt	FM	mature
03/10/2016	10	111	1	milt	FM	mature
03/10/2016	10	113	1	milt	FM	mature
03/10/2016	10	119	1	milt	FM	mature
03/10/2016	10	79	0	immature	FM	immature
03/10/2016	10	105	1	immature	FM	immature
03/10/2016	10	111	1	milt	FM	mature
03/10/2016	10	107	1	immature	FM	immature
03/10/2016	10	107	1	milt	FM	mature
03/10/2016	10	115	1	milt	FM	mature
03/10/2016	10	119	1	milt	FM	mature
03/10/2016	10	65	0	immature	FM	immature
03/10/2016	10	128	1	immature	FM	immature
03/10/2016	10	117	1	milt	FM	mature
03/10/2016	11	112	1	milt	FM	mature

03/10/2016	11	123	1	milt	FM	mature
03/10/2016	11	122	1	mature	FM	immature
03/10/2016	11	115	1	immature	FM	immature
03/10/2016	11	109	1	milt	FM	mature
03/10/2016	11	74	0	immature	FM	immature
03/10/2016	11	68	0	immature	FM	immature
03/10/2016	11	108	1	milt	FM	mature
03/10/2016	11	67	0	immature	FM	immature
03/10/2016	11	80	0	milt	FM	mature
03/10/2016	11	115	1	immature	FM	immature
03/10/2016	11	104	1	milt	FM	mature
03/10/2016	11	118	1	immature	FM	immature
03/10/2016	11	103	1	immature	FM	immature
03/10/2016	11	94	1	milt	FM	mature
03/10/2016	11	103	1	immature	FM	immature
03/10/2016	11	69	0	immature	FM	immature
03/10/2016	11	68	0	immature	FM	immature
03/10/2016	11	68	0	immature	FM	immature
03/10/2016	11	124	1	immature	FM	immature
03/10/2016	11	97	1	milt	FM	mature
03/10/2016	11	106	1	milt	FM	mature
03/10/2016	11	72	0	immature	FM	immature
03/10/2016	11	109	1	milt	FM	mature
03/10/2016	11	113	1	immature	FM	immature
03/10/2016	11	66	0	immature	FM	immature
03/10/2016	11	101	1	milt	FM	mature
03/10/2016	11	111	1	immature	FM	immature
03/10/2016	11	106	1	mature	FM	mature
03/10/2016	11	106	1	immature	FM	immature
03/10/2016	12	102	1	immature	FM	immature
03/10/2016	12	115	1	immature	FM	immature
03/10/2016	12	66	0	immature	FM	immature
03/10/2016	12	119	1	immature	FM	immature
03/10/2016	12	66	0	immature	FM	immature
03/10/2016	12	69	0	immature	FM	immature
03/10/2016	12	67	0	immature	FM	immature
03/10/2016	12	99	1	mature	FM	mature
03/10/2016	12	104	1	milt	FM	mature
03/10/2016	12	111	1	milt	FM	mature
03/10/2016	12	105	1	immature	FM	immature

03/10/2016	12	70	0	immature	FM	immature
03/10/2016	12	64	0	immature	FM	immature
03/10/2016	12	67	0	immature	FM	immature
03/10/2016	12	111	1	milt	FM	mature
03/10/2016	12	96	1	milt	FM	mature
03/10/2016	12	119	1	milt	FM	mature
03/10/2016	12	106	1	milt	FM	mature
03/10/2016	12	128	1	milt	FM	mature
03/10/2016	12	67	0	immature	FM	mature
03/10/2016	12	116	1	milt	FM	mature
03/10/2016	12	112	1	milt	FM	mature
03/10/2016	12	71	0	immature	FM	immature
03/10/2016	12	100	1	milt	FM	mature
03/10/2016	12	68	0	immature	FM	immature
03/10/2016	12	107	1	immature	FM	immature
03/10/2016	12	69	0	immature	FM	immature
03/10/2016	12	67	0	immature	FM	immature
03/10/2016	12	64	0	immature	FM	immature
03/10/2016	12	75	0	immature	FM	immature
03/10/2016	12	75	0	milt	FM	mature
03/10/2016	12	106	1	milt	FM	mature
03/10/2016	12	72	0	immature	FM	immature
03/10/2016	12	106	1	immature	FM	immature
03/10/2016	12	102	1	immature	FM	immature
03/10/2016	12	74	0	immature	FM	immature
03/10/2016	12	81	0	milt	FM	mature
03/10/2016	12	71	0	immature	FM	immature
03/10/2016	12	110	1	immature	FM	immature
03/10/2016	12	124	1	milt	FM	mature
03/10/2016	12	107	1	milt	FM	mature
03/10/2016	12	95	1	milt	FM	mature
03/10/2016	12	100	1	milt	FM	mature
03/10/2016	12	71	0	immature	FM	immature
03/10/2016	12	69	0	immature	FM	immature
03/10/2016	12	68	0	immature	FM	immature
03/10/2016	12	64	0	immature	FM	immature
03/10/2016	13	68	0	immature	FM	immature
03/10/2016	13	80	0	immature	FM	immature
03/10/2016	13	118	1	milt	FM	mature
03/10/2016	13	117	1	immature	FM	immature

03/10/2016	13	118	1	milt	FM	mature
03/10/2016	13	118	1	immature	FM	immature
03/10/2016	13	64	0	immature	FM	immature
03/10/2016	13	95	1	milt	FM	mature
03/10/2016	13	108	1	immature	FM	immature
03/10/2016	13	109	1	immature	FM	immature
03/10/2016	13	70	0	immature	FM	immature
03/10/2016	13	73	0	immature	FM	immature
03/10/2016	13	71	0	immature	FM	immature
03/10/2016	13	69	0	immature	FM	immature
03/10/2016	13	71	0	immature	FM	immature
03/10/2016	13	67	0	immature	FM	immature
03/10/2016	13	70	0	immature	FM	immature
03/10/2016	13	62	0	immature	FM	immature
03/10/2016	14	70	0	immature	FM	immature
03/10/2016	14	110	1	immature	FM	immature
03/10/2016	14	105	1	milt	FM	mature
03/10/2016	14	116	1	immature	FM	immature
03/10/2016	14	103	1	milt	FM	mature
03/10/2016	14	82	0	immature	FM	immature
03/10/2016	14	83	0	immature	FM	immature
03/10/2016	14	68	0	immature	FM	immature
03/10/2016	14	74	0	immature	FM	immature
03/10/2016	14	62	0	immature	FM	immature
03/10/2016	14	67	0	immature	FM	immature
03/10/2016	14	80	0	immature	FM	immature
03/10/2016	14	92	1	milt	FM	mature
03/10/2016	14	113	1	milt	FM	mature
03/10/2016	14	98	1	milt	FM	mature
03/10/2016	14	102	1	mature	FM	mature
03/10/2016	14	119	1	immature	FM	immature
03/10/2016	14	55	0	immature	FM	immature
03/10/2016	14	63	0	immature	FM	immature
03/10/2016	14	67	0	immature	FM	immature
03/10/2016	14	71	0	immature	FM	immature
03/10/2016	14	77	0	immature	FM	immature
03/10/2016	14	73	0	immature	FM	immature
03/10/2016	14	105	1	immature	FM	immature
03/10/2016	14	125	1	milt	FM	mature
03/10/2016	14	70	0	immature	FM	immature

03/10/2016	14	66	0	immature	FM	immature
03/10/2016	14	66	0	immature	FM	immature
03/10/2016	14	68	0	immature	FM	immature
03/10/2016	14	97	1	milt	FM	mature
03/10/2016	14	59	0	immature	FM	immature
03/10/2016	14	100	1	mature	FM	mature
03/10/2016	14	82	0	mature	FM	mature
03/10/2016	14	69	0	immature	FM	immature
03/10/2016	14	62	0	immature	FM	immature
03/10/2016	14	114	1	milt	FM	mature
03/10/2016	14	72	0	mature	FM	mature
03/10/2016	14	71	0	milt	FM	mature
03/10/2016	14	129	1	milt	FM	mature
03/10/2016	14	78	0	immature	FM	immature
03/10/2016	14	69	0	mature	FM	mature
03/10/2016	14	72	0	immature	FM	immature
03/10/2016	14	79	0	immature	FM	immature
03/10/2016	14	72	0	immature	FM	immature
03/10/2016	14	83	0	immature	FM	immature
03/10/2016	14	71	0	immature	FM	mature
03/10/2016	14	68	0	immature	FM	immature
03/10/2016	14	72	0	immature	FM	immature
03/10/2016	15	67	0	immature	FM	immature
03/10/2016	15	113	1	mature	FM	mature
03/10/2016	15	76	0	immature	FM	immature
03/10/2016	15	69	0	immature	FM	mature
03/10/2016	15	69	0	mature	FM	mature
03/10/2016	15	63	0	immature	FM	immature
03/10/2016	15	106	1	milt	FM	mature
03/10/2016	15	102	1	immature	FM	immature
03/10/2016	15	75	0	immature	FM	immature
03/10/2016	15	66	0	immature	FM	immature
03/10/2016	15	72	0	immature	FM	immature
03/10/2016	15	122	1	milt	FM	mature
03/10/2016	15	116	1	milt	FM	mature
03/10/2016	15	104	1	milt	FM	mature
03/10/2016	15	104	1	milt	FM	mature
03/10/2016	15	107	1	milt	FM	mature
03/10/2016	15	79	0	immature	FM	immature
03/10/2016	15	94	0	immature	FM	immature

03/10/2016	15	112	1	immature	FM	immature
03/10/2016	15	117	1	immature	FM	immature
03/10/2016	15	67	0	immature	FM	immature
03/10/2016	15	73	0	immature	FM	immature
03/10/2016	15	68	0	immature	FM	immature
03/10/2016	15	74	0	immature	FM	immature
03/10/2016	15	69	0	immature	FM	immature
03/10/2016	15	131	1	milt	FM	mature
03/10/2016	15	112	1	milt	FM	mature
03/10/2016	15	119	1	milt	FM	mature
03/10/2016	15	81	0	immature	FM	immature
03/10/2016	15	118	1	immature	FM	immature
03/10/2016	15	108	1	milt	FM	mature
03/10/2016	15	109	1	milt	FM	mature
03/10/2016	15	113	1	immature	FM	immature
03/10/2016	15	109	1	immature	FM	immature
03/10/2016	16	91	0	immature	FM	immature
03/10/2016	16	82	0	immature	FM	immature
03/10/2016	16	79	0	milt	FM	mature
04/10/2016	08	82	0	immature	FM	immature
04/10/2016	08	126	1	milt	FM	mature
04/10/2016	08	113	1	immature	FM	immature
04/10/2016	08	116	1	immature	FM	mature
04/10/2016	08	83	0	immature	FM	immature
04/10/2016	09	84	0	immature	FM	immature
04/10/2016	09	111	1	milt	FM	mature
04/10/2016	09	81	0	milt	FM	mature
04/10/2016	09	83	0	mature	FM	mature
04/10/2016	09	128	1	immature	FM	immature
04/10/2016	09	133	1	immature	FM	immature
04/10/2016	09	88	0	immature	FM	immature
04/10/2016	09	115	1	milt	FM	mature
04/10/2016	09	124	1	milt	FM	mature
04/10/2016	09	79	0	milt	FM	mature
04/10/2016	09	94	0	immature	FM	immature
04/10/2016	09	106	1	milt	FM	mature
04/10/2016	09	120	1	milt	FM	mature
04/10/2016	09	118	1	immature	FM	immature
04/10/2016	09	113	1	milt	FM	mature
04/10/2016	09	104	1	milt	FM	mature

04/10/2016	09	106	1	milt	FM	mature
04/10/2016	09	95	1	milt	FM	mature
04/10/2016	10	74	0	mature	FM	immature
04/10/2016	10	112	1	milt	FM	mature
04/10/2016	10	80	0	immature	FM	immature
04/10/2016	10	82	0	immature	FM	immature
04/10/2016	10	73	0	immature	FM	immature
04/10/2016	10	74	0	immature	FM	immature
04/10/2016	10	115	1	mature	FM	mature
04/10/2016	10	73	0	mature	FM	immature
04/10/2016	10	74	0	immature	FM	immature
04/10/2016	10	65	0	immature	FM	immature
04/10/2016	10	122	1	milt	FM	mature
04/10/2016	10	105	1	milt	FM	mature
04/10/2016	10	99	1	milt	FM	mature
04/10/2016	10	113	1	milt	FM	mature
04/10/2016	10	111	1	milt	FM	mature
04/10/2016	10	114	1	milt	FM	mature
04/10/2016	10	75	0	mature	FM	mature
04/10/2016	10	79	0	immature	FM	immature
04/10/2016	10	115	1	immature	FM	immature
04/10/2016	10	89	0	immature	FM	immature
04/10/2016	10	81	0	immature	FM	immature
04/10/2016	10	69	0	immature	FM	immature
04/10/2016	10	110	1	immature	FM	immature
04/10/2016	10	128	1	immature	FM	immature
04/10/2016	10	97	1	milt	FM	mature
04/10/2016	10	109	1	immature	FM	immature
04/10/2016	10	104	1	immature	FM	immature
04/10/2016	10	73	0	immature	FM	immature
04/10/2016	10	109	1	immature	FM	immature
04/10/2016	10	112	1	milt	FM	mature
04/10/2016	10	112	1	milt	FM	mature
04/10/2016	10	112	1	milt	FM	mature
04/10/2016	10	100	1	milt	FM	mature
04/10/2016	10	112	1	milt	FM	mature
04/10/2016	10	108	1	mature	FM	mature
04/10/2016	10	110	1	milt	FM	mature
04/10/2016	10	110	1	immature	FM	immature
04/10/2016	10	97	1	milt	FM	mature

04/10/2016	10	71	0	immature	FM	immature
04/10/2016	10	72	0	immature	FM	immature
04/10/2016	10	108	1	milt	FM	mature
04/10/2016	10	114	1	milt	FM	mature
04/10/2016	11	119	1	milt	FM	mature
04/10/2016	11	77	0	immature	FM	immature
04/10/2016	11	129	1	immature	FM	immature
04/10/2016	11	108	1	milt	FM	mature
04/10/2016	12	106	1	immature	FM	immature
04/10/2016	12	118	1	immature	FM	immature
04/10/2016	12	110	1	immature	FM	immature
04/10/2016	12	70	0	immature	FM	immature
04/10/2016	12	114	1	immature	FM	immature
04/10/2016	12	112	1	milt	FM	mature
04/10/2016	12	104	1	immature	FM	immature
04/10/2016	12	106	1	milt	FM	mature
04/10/2016	12	113	1	milt	FM	mature
04/10/2016	12	102	1	milt	FM	mature
04/10/2016	12	63	0	immature	FM	immature
04/10/2016	12	74	0	mature	FM	mature
04/10/2016	12	101	1	milt	FM	mature
04/10/2016	12	75	0	mature	FM	mature
04/10/2016	12	69	0	immature	FM	immature
04/10/2016	12	115	1	immature	FM	immature
04/10/2016	13	107	1	milt	FM	mature
04/10/2016	13	99	1	immature	FM	immature
04/10/2016	13	73	0	immature	FM	immature
04/10/2016	13	96	1	milt	FM	mature
04/10/2016	13	114	1	milt	FM	mature
04/10/2016	13	95	1	milt	FM	mature
04/10/2016	13	73	0	immature	FM	immature
04/10/2016	13	107	1	milt	FM	mature
04/10/2016	13	71	0	milt	FM	immature
04/10/2016	13	111	1	immature	FM	immature
04/10/2016	13	111	1	milt	FM	mature
04/10/2016	13	113	1	immature	FM	immature
04/10/2016	13	106	1	milt	FM	immature
04/10/2016	13	105	1	immature	FM	immature
04/10/2016	13	100	1	milt	FM	mature
04/10/2016	13	69	0	immature	FM	immature

04/10/2016	13	110	1	milt	FM	mature
04/10/2016	13	111	1	immature	FM	immature
04/10/2016	13	119	1	milt	FM	mature
04/10/2016	13	115	1	immature	FM	immature
04/10/2016	13	98	1	milt	FM	mature
04/10/2016	14	120	1	immature	FM	immature
04/10/2016	14	93	1	milt	FM	mature
04/10/2016	14	118	1	immature	FM	immature
04/10/2016	14	101	1	milt	FM	mature
04/10/2016	14	114	1	mature	FM	mature
04/10/2016	14	122	1	milt	FM	mature
04/10/2016	14	105	1	milt	FM	mature
04/10/2016	14	99	1	milt	FM	mature
04/10/2016	14	117	1	milt	FM	mature
04/10/2016	14	112	1	milt	FM	mature
04/10/2016	14	110	1	immature	FM	immature
04/10/2016	14	110	1	mature	FM	mature
04/10/2016	14	106	1	milt	FM	mature
04/10/2016	14	102	1	milt	FM	mature
04/10/2016	14	107	1	immature	FM	immature
04/10/2016	14	124	1	immature	FM	immature
04/10/2016	14	75	0	immature	FM	immature
04/10/2016	14	121	1	milt	FM	mature
04/10/2016	14	114	1	immature	FM	immature
04/10/2016	14	117	1	immature	FM	immature
04/10/2016	14	109	1	milt	FM	mature
04/10/2016	14	114	1	immature	FM	immature
04/10/2016	14	105	1	immature	FM	immature
04/10/2016	14	110	1	immature	FM	immature
04/10/2016	14	114	1	immature	FM	immature
04/10/2016	14	102	1	milt	FM	mature
04/10/2016	14	111	1	milt	FM	mature
04/10/2016	14	109	1	immature	FM	immature
04/10/2016	14	121	1	milt	FM	mature
04/10/2016	14	122	1	immature	FM	immature
04/10/2016	15	114	1	immature	FM	immature
04/10/2016	15	102	1	milt	FM	mature
04/10/2016	15	127	1	milt	FM	mature
04/10/2016	15	114	1	immature	FM	immature
04/10/2016	15	100	1	immature	FM	immature

04/10/2016	15	104	1	milt	FM	mature
04/10/2016	15	95	1	immature	FM	immature
04/10/2016	15	113	1	immature	FM	immature
04/10/2016	15	111	1	immature	FM	immature
04/10/2016	15	113	1	immature	FM	immature
04/10/2016	15	105	1	immature	FM	immature
04/10/2016	15	119	1	immature	FM	immature
04/10/2016	15	104	1	milt	FM	mature
04/10/2016	15	106	1	immature	FM	immature
04/10/2016	15	104	1	immature	FM	immature
04/10/2016	15	106	1	milt	FM	mature
04/10/2016	15	113	1	milt	FM	mature
05/10/2016	08	112	1	milt	FM	mature
05/10/2016	08	102	1	milt	FM	mature
05/10/2016	08	102	1	milt	FM	mature
05/10/2016	08	110	1	milt	FM	mature
05/10/2016	08	114	1	milt	FM	mature
05/10/2016	08	99	1	milt	FM	mature
05/10/2016	08	104	1	immature	FM	immature
05/10/2016	08	110	1	milt	FM	mature
05/10/2016	08	110	1	immature	FM	immature
05/10/2016	08	105	1	milt	FM	mature
05/10/2016	08	120	1	immature	FM	immature
05/10/2016	08	124	1	immature	FM	immature
05/10/2016	09	120	1	immature	FM	immature
05/10/2016	09	113	1	milt	FM	mature
05/10/2016	09	110	1	milt	FM	mature
05/10/2016	09	92	1	milt	FM	mature
05/10/2016	09	112	1	milt	FM	mature
05/10/2016	09	107	1	immature	FM	immature
05/10/2016	09	119	1	immature	FM	immature
05/10/2016	09	108	1	immature	FM	immature
05/10/2016	09	125	1	immature	FM	immature
05/10/2016	10	123	1	milt	FM	mature
05/10/2016	10	127	1	immature	FM	immature
05/10/2016	10	127	1	immature	FM	immature
05/10/2016	10	120	1	milt	FM	mature
05/10/2016	10	115	1	milt	FM	mature
05/10/2016	10	125	1	immature	FM	immature
05/10/2016	10	123	1	milt	FM	mature

05/10/2016	10	121	1	milt	FM	mature
05/10/2016	10	115	1	immature	FM	immature
05/10/2016	11	115	1	milt	FM	mature
05/10/2016	11	119	1	immature	FM	immature
05/10/2016	11	101	1	milt	FM	mature
05/10/2016	11	121	1	milt	FM	mature
05/10/2016	11	112	1	milt	FM	mature
05/10/2016	11	129	1	milt	FM	mature
05/10/2016	11	113	1	milt	FM	mature
05/10/2016	11	121	1	milt	FM	mature
05/10/2016	11	125	1	immature	FM	immature
05/10/2016	11	115	1	immature	FM	immature
05/10/2016	11	125	1	milt	FM	mature
05/10/2016	11	107	1	milt	FM	mature
05/10/2016	11	114	1	immature	FM	immature
05/10/2016	11	117	1	milt	FM	mature
05/10/2016	11	128	1	immature	FM	immature
05/10/2016	11	117	1	milt	FM	mature
05/10/2016	11	114	1	milt	FM	mature
05/10/2016	11	124	1	immature	FM	immature
05/10/2016	11	123	1	immature	FM	immature
05/10/2016	11	121	1	milt	FM	mature
05/10/2016	11	108	1	milt	FM	mature
05/10/2016	11	113	1	immature	FM	immature
05/10/2016	11	113	1	milt	FM	mature
05/10/2016	11	109	1	milt	FM	mature
05/10/2016	11	109	1	milt	FM	mature
05/10/2016	11	129	1	immature	FM	immature
05/10/2016	11	112	1	immature	FM	immature
05/10/2016	11	111	1	immature	FM	immature
05/10/2016	11	107	1	immature	FM	immature
05/10/2016	11	118	1	immature	FM	immature
05/10/2016	11	113	1	immature	FM	immature
05/10/2016	11	101	1	milt	FM	mature
05/10/2016	11	123	1	immature	FM	immature
05/10/2016	11	114	1	immature	FM	immature
05/10/2016	11	106	1	milt	FM	mature
05/10/2016	11	120	1	milt	FM	mature
05/10/2016	11	111	1	immature	FM	immature
05/10/2016	11	111	1	immature	FM	mature

05/10/2016	11	117	1	milt	FM	mature
05/10/2016	11	109	1	immature	FM	immature
05/10/2016	11	109	1	milt	FM	mature
05/10/2016	11	122	1	immature	FM	immature
05/10/2016	11	114	1	milt	FM	mature
05/10/2016	11	114	1	immature	FM	immature
05/10/2016	11	113	1	immature	FM	mature
05/10/2016	12	118	1	mature	FM	mature
05/10/2016	12	111	1	immature	FM	immature
05/10/2016	12	117	1	immature	FM	immature
05/10/2016	12	107	1	immature	FM	immature
05/10/2016	12	113	1	immature	FM	immature
05/10/2016	12	107	1	immature	FM	immature
09/10/2017	09	99	0	immature	JT	immature
09/10/2017	09	111	1	milt	JT	mature
09/10/2017	09	92	0	immature	JT	immature
09/10/2017	09	74	0	immature	JT	immature
09/10/2017	09	97	0	immature	JT	immature
09/10/2017	09	92	0	immature	JT	immature
09/10/2017	09	66	0	immature	JT	immature
09/10/2017	09	109	0	immature	JT	immature
09/10/2017	09	75	0	immature	JT	immature
09/10/2017	10	75	0	immature	JT	immature
09/10/2017	10	89	0	immature	JT	immature
09/10/2017	10	78	0	immature	JT	immature
09/10/2017	10	61	0	immature	JT	immature
09/10/2017	10	85	0	immature	JT	immature
09/10/2017	10	73	0	immature	JT	immature
09/10/2017	10	91	0	milt	JT	mature
09/10/2017	10	73	0	immature	JT	immature
09/10/2017	10	101	0	immature	JT	immature
09/10/2017	10	87	0	immature	JT	immature
09/10/2017	10	78	0	immature	JT	immature
09/10/2017	10	101	0	immature	JT	immature
09/10/2017	10	76	0	immature	JT	immature
09/10/2017	10	65	0	immature	JT	immature
09/10/2017	10	70	0	immature	JT	immature
09/10/2017	10	95	0	immature	JT	immature
09/10/2017	10	78	0	immature	JT	immature
09/10/2017	10	67	0	immature	JT	immature

09/10/2017	10	93	0	immature	JT	immature
09/10/2017	10	71	0	mature	JT	mature
09/10/2017	10	76	0	immature	JT	immature
09/10/2017	10	76	0	immature	JT	immature
09/10/2017	10	90	0	immature	JT	immature
09/10/2017	10	100	0	immature	JT	immature
09/10/2017	10	98	0	immature	JT	immature
09/10/2017	10	73	0	immature	JT	immature
09/10/2017	10	69	0	immature	JT	immature
09/10/2017	10	87	0	immature	JT	immature
09/10/2017	11	95	0	milt	JT	mature
09/10/2017	11	74	0	milt	JT	mature
09/10/2017	11	93	0	immature	JT	immature
09/10/2017	11	82	0	milt	JT	mature
09/10/2017	11	107	0	immature	JT	immature
09/10/2017	11	129	1	immature	JT	immature
09/10/2017	11	129	1	immature	JT	immature
09/10/2017	11	98	0	immature	JT	immature
09/10/2017	11	66	0	immature	JT	immature
09/10/2017	11	82	0	milt	JT	mature
09/10/2017	11	71	0	immature	JT	immature
09/10/2017	11	83	0	immature	JT	immature
09/10/2017	11	97	0	immature	JT	immature
09/10/2017	11	69	0	immature	JT	immature
09/10/2017	11	61	0	immature	JT	immature
09/10/2017	11	81	0	milt	JT	immature
09/10/2017	11	74	0	immature	JT	immature
09/10/2017	11	61	0	immature	JT	immature
09/10/2017	11	71	0	immature	JT	immature
09/10/2017	11	75	0	immature	JT	immature
09/10/2017	11	70	0	immature	JT	immature
09/10/2017	11	70	0	immature	JT	immature
09/10/2017	11	72	0	milt	JT	mature
09/10/2017	11	64	0	immature	JT	immature
09/10/2017	11	85	0	immature	JT	immature
09/10/2017	11	81	0	immature	JT	immature
09/10/2017	11	75	0	immature	JT	immature
09/10/2017	11	87	0	immature	JT	immature
09/10/2017	11	88	0	immature	JT	immature
09/10/2017	11	125	1	immature	JT	immature

09/10/2017	11	88	0	immature	JT	immature
09/10/2017	11	76	0	immature	JT	immature
09/10/2017	12	73	0	immature	JT	immature
09/10/2017	12	75	0	immature	JT	immature
09/10/2017	12	73	0	immature	JT	immature
09/10/2017	12	124	1	immature	JT	immature
09/10/2017	12	131	1	immature	JT	immature
09/10/2017	12	85	0	immature	JT	immature
09/10/2017	12	86	0	immature	JT	immature
09/10/2017	12	122	1	immature	JT	immature
09/10/2017	12	79	0	immature	JT	immature
09/10/2017	12	67	0	immature	JT	immature
09/10/2017	12	86	0	immature	JT	immature
09/10/2017	12	69	0	immature	JT	immature
09/10/2017	12	86	0	immature	JT	immature
09/10/2017	12	71	0	immature	JT	immature
09/10/2017	12	80	0	immature	JT	immature
09/10/2017	12	87	0	immature	JT	immature
09/10/2017	12	105	1	immature	JT	immature
09/10/2017	12	115	1	milt	JT	mature
09/10/2017	12	79	0	immature	JT	immature
09/10/2017	12	91	0	immature	JT	immature
09/10/2017	12	73	0	immature	JT	immature
09/10/2017	12	91	0	immature	JT	immature
09/10/2017	12	84	0	immature	JT	immature
09/10/2017	12	81	0	immature	JT	immature
09/10/2017	12	71	0	immature	JT	immature
09/10/2017	12	91	0	immature	JT	immature
09/10/2017	12	88	0	immature	JT	immature
09/10/2017	12	77	0	immature	JT	immature
09/10/2017	12	71	0	immature	JT	immature
09/10/2017	12	88	0	immature	JT	immature
09/10/2017	12	71	0	immature	JT	immature
09/10/2017	12	75	0	immature	JT	mature
09/10/2017	12	73	0	immature	JT	immature
09/10/2017	12	81	0	milt	JT	mature
09/10/2017	13	100	0	immature	JT	immature
09/10/2017	13	62	0	immature	JT	immature
09/10/2017	13	100	0	immature	JT	immature
09/10/2017	13	86	0	immature	JT	immature

09/10/2017	13	89	0	immature	JT	immature
09/10/2017	13	89	0	immature	JT	immature
09/10/2017	13	87	0	immature	JT	immature
09/10/2017	13	67	0	immature	JT	immature
09/10/2017	13	72	0	immature	JT	immature
09/10/2017	13	76	0	immature	JT	immature
09/10/2017	13	76	0	immature	JT	immature
09/10/2017	13	66	0	immature	JT	immature
09/10/2017	13	122	1	milt	JT	mature
09/10/2017	13	119	1	immature	JT	immature
09/10/2017	13	108	1	milt	JT	mature
09/10/2017	13	112	1	immature	JT	immature
09/10/2017	14	122	1	immature	JT	immature
09/10/2017	14	90	0	immature	JT	immature
09/10/2017	14	123	1	milt	JT	mature
09/10/2017	14	71	0	immature	JT	immature
09/10/2017	14	74	0	immature	JT	immature
09/10/2017	14	99	0	immature	JT	immature
09/10/2017	14	69	0	immature	JT	immature
09/10/2017	14	84	0	immature	JT	immature
09/10/2017	14	120	1	immature	JT	immature
09/10/2017	14	83	0	immature	JT	immature
09/10/2017	14	96	0	immature	JT	immature
09/10/2017	14	56	0	immature	JT	immature
09/10/2017	14	73	0	milt	JT	immature
09/10/2017	14	80	0	immature	JT	immature
09/10/2017	15	104	0	immature	JT	immature
09/10/2017	15	122	1	milt	JT	mature
09/10/2017	15	77	0	immature	JT	immature
09/10/2017	15	118	1	immature	JT	immature
09/10/2017	15	115	1	immature	JT	immature
09/10/2017	15	96	0	immature	JT	immature
09/10/2017	15	107	1	milt	JT	mature
09/10/2017	15	82	0	immature	JT	immature
09/10/2017	15	97	0	immature	JT	immature
09/10/2017	15	91	0	immature	JT	immature
09/10/2017	15	85	0	immature	JT	immature
09/10/2017	16	89	0	milt	JT	mature
09/10/2017	16	93	0	immature	JT	immature
09/10/2017	16	74	0	immature	JT	immature

09/10/2017	16	99	0	immature	JT	immature
09/10/2017	16	88	0	immature	JT	immature
09/10/2017	16	87	0	milt	JT	mature
09/10/2017	16	101	0	immature	JT	immature
09/10/2017	16	95	0	immature	JT	immature
09/10/2017	16	93	0	immature	JT	immature
09/10/2017	16	93	0	immature	JT	immature
09/10/2017	16	75	0	immature	JT	mature
10/10/2017	09	96	0	immature	JT	immature
10/10/2017	09	128	1	milt	JT	mature
10/10/2017	09	96	0	immature	JT	immature
10/10/2017	09	100	1	immature	JT	immature
10/10/2017	10	120	1	milt	JT	immature
10/10/2017	10	85	0	immature	JT	immature
10/10/2017	10	79	0	immature	JT	mature
10/10/2017	10	81	0	immature	JT	immature
10/10/2017	10	106	0	immature	JT	immature
10/10/2017	10	86	0	milt	JT	mature
10/10/2017	10	91	0	immature	JT	immature
10/10/2017	10	100	0	immature	JT	immature
10/10/2017	10	95	0	immature	JT	immature
10/10/2017	10	90	0	immature	JT	immature
10/10/2017	10	85	0	immature	JT	immature
10/10/2017	10	113	1	immature	JT	immature
10/10/2017	10	70	0	immature	JT	immature
10/10/2017	10	84	0	milt	JT	mature
10/10/2017	11	90	0	immature	JT	immature
10/10/2017	11	104	0	immature	JT	immature
10/10/2017	11	90	0	immature	JT	immature
10/10/2017	11	104	0	immature	JT	immature
10/10/2017	11	86	0	immature	JT	immature
10/10/2017	11	82	0	immature	JT	immature
10/10/2017	11	87	0	immature	JT	immature
10/10/2017	11	126	1	immature	JT	immature
10/10/2017	11	95	0	immature	JT	immature
10/10/2017	11	114	1	milt	JT	mature
10/10/2017	11	92	0	immature	JT	immature
10/10/2017	11	74	0	milt	JT	mature
10/10/2017	11	89	0	immature	JT	immature
10/10/2017	11	89	0	immature	JT	immature

10/10/2017	12	86	0	immature	JT	immature
10/10/2017	12	92	0	immature	JT	immature
10/10/2017	12	98	0	milt	JT	mature
10/10/2017	12	88	0	immature	JT	immature
10/10/2017	12	95	0	immature	JT	immature
10/10/2017	12	74	0	immature	JT	immature
10/10/2017	12	93	0	immature	JT	immature
10/10/2017	12	81	0	milt	JT	mature
10/10/2017	12	92	0	immature	JT	immature
10/10/2017	12	82	0	immature	JT	immature
10/10/2017	12	79	0	immature	JT	immature
10/10/2017	12	88	0	immature	JT	immature
10/10/2017	12	82	0	immature	JT	immature
10/10/2017	12	81	0	immature	JT	immature
10/10/2017	12	81	0	immature	JT	immature
10/10/2017	12	75	0	immature	JT	immature
10/10/2017	14	126	1	milt	JT	immature
11/10/2017	11	116	0	immature	JT	immature